

$a_1(1640)$ $I^G(J^{PC}) = 1^+(1^{++})$

OMMITTED FROM SUMMARY TABLE

Seen in the amplitude analysis of the $3\pi^0$ system produced in $\bar{p}p \rightarrow 4\pi^0$. Possibly seen in the study of the hadronic structure in decay $\tau \rightarrow 3\pi\nu_\tau$ (ABREU 98G). Needs confirmation.

 $a_1(1640)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$1640 \pm 12 \pm 30$	1 BAKER 99	SPEC	$1.94 \bar{p}p \rightarrow 4\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1670 ± 90	BELLINI 85	SPEC	$40 \pi^- A \rightarrow \pi^-\pi^+\pi^- A$

¹ Using preliminary CBAR data. **$a_1(1640)$ WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$300 \pm 22 \pm 40$	2 BAKER 99	SPEC	$1.94 \bar{p}p \rightarrow 4\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
300 ± 100	BELLINI 85	SPEC	$40 \pi^- A \rightarrow \pi^-\pi^+\pi^- A$

² Using preliminary CBAR data. **$a_1(1640)$ DECAY MODES**

Mode
$\Gamma_1 f_2(1270)\pi$
$\Gamma_2 \sigma\pi$

 $a_1(1640)$ BRANCHING RATIOS

$\Gamma(f_2(1270)\pi)/\Gamma(\sigma\pi)$	Γ_1/Γ_2
0.24 ± 0.07	3 BAKER 99 SPEC $1.94 \bar{p}p \rightarrow 4\pi^0$

³ Using preliminary CBAR data. **$a_1(1640)$ REFERENCES**

BAKER 99	PL B449 114	C.A. Baker <i>et al.</i>	
ABREU 98G	PL B426 411	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BELLINI 85	Translated from YAF 41 1223.	G. Bellini <i>et al.</i>	