



$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$ Status: ***

According to the quark model, the Ξ_c^+ (quark content $us\bar{c}$) and Ξ_c^0 form an isospin doublet, and the spin-parity ought to be $J^P = 1/2^+$. None of I , J , or P has actually been measured.

Ξ_c^+ MASS

The fit uses the Ξ_c^+ and Ξ_c^0 mass and mass-difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2466.3 ± 1.4 OUR FIT				
2466.4 ± 1.5 OUR AVERAGE				
2465.8 ± 1.9 ± 2.5	90	FRABETTI	98 E687	γ Be, $\bar{E}_\gamma = 220$ GeV
2467.0 ± 1.6 ± 2.0	147	EDWARDS	96 CLE2	$e^+e^- \approx \gamma(4S)$
2465.1 ± 3.6 ± 1.9	30	ALBRECHT	90F ARG	e^+e^- at $\gamma(4S)$
2467 ± 3 ± 4	23	ALAM	89 CLEO	e^+e^- 10.6 GeV
2466.5 ± 2.7 ± 1.2	5	BARLAG	89C ACCM	π^- Cu 230 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2464.4 ± 2.0 ± 1.4	30	FRABETTI	93B E687	See FRABETTI 98
2459 ± 5 ± 30	56	¹ COTEUS	87 SPEC	$nA \simeq 600$ GeV
2460 ± 25	82	BIAGI	83 SPEC	Σ^- Be 135 GeV

¹ Although COTEUS 87 claims to agree well with BIAGI 83 on the mass and width, there appears to be a discrepancy between the two experiments. BIAGI 83 sees a single peak (stated significance about 6 standard deviations) in the $\Lambda K^- \pi^+ \pi^+$ mass spectrum. COTEUS 87 sees two peaks in the same spectrum, one at the Ξ_c^+ mass, the other 75 MeV lower. The latter is attributed to $\Xi_c^+ \rightarrow \Sigma^0 K^- \pi^+ \pi^+ \rightarrow (\Lambda \gamma) K^- \pi^+ \pi^+$, with the γ unseen. The combined significance of the double peak is stated to be 5.5 standard deviations. But the absence of any trace of a lower peak in BIAGI 83 seems to us to throw into question the interpretation of the lower peak of COTEUS 87.

Ξ_c^+ MEAN LIFE

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
0.33^{+0.06}_{-0.04} OUR AVERAGE				
0.34 ^{+0.07} _{-0.05} ± 0.02	56	FRABETTI	98 E687	γ Be, $\bar{E}_\gamma = 220$ GeV
0.20 ^{+0.11} _{-0.06}	6	BARLAG	89C ACCM	π^- (K^-) Cu 230 GeV
0.40 ^{+0.18} _{-0.12} ± 0.10	102	COTEUS	87 SPEC	$nA \simeq 600$ GeV
0.48 ^{+0.21} _{-0.15} ± 0.20	53	BIAGI	85C SPEC	Σ^- Be 135 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.41 ^{+0.11} _{-0.08} ± 0.02	30	FRABETTI	93B E687	See FRABETTI 98

Ξ_c^+ DECAY MODES

No absolute branching fractions have been measured. The following are branching *ratios* relative to $\Xi^- \pi^+ \pi^+$.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \Lambda K^- \pi^+ \pi^+$	[a] 0.58 ± 0.18	
$\Gamma_2 \Lambda \bar{K}^*(892)^0 \pi^+$	[a,b] < 0.29	90%
$\Gamma_3 \Sigma(1385)^+ K^- \pi^+$	[a,b] < 0.41	90%
$\Gamma_4 \Sigma^+ K^- \pi^+$	[a] 1.18 ± 0.31	
$\Gamma_5 \Sigma^+ \bar{K}^*(892)^0$	[a,b] 0.92 ± 0.30	
$\Gamma_6 \Sigma^0 K^- \pi^+ \pi^+$	[a] 0.49 ± 0.26	
$\Gamma_7 \Xi^0 \pi^+$	[a] 0.55 ± 0.16	
$\Gamma_8 \Xi^- \pi^+ \pi^+$	[a] $\equiv 1.0$	
$\Gamma_9 \Xi(1530)^0 \pi^+$	[a,b] < 0.2	90%
$\Gamma_{10} \Xi^0 \pi^+ \pi^0$	[a] 2.34 ± 0.68	
$\Gamma_{11} \Xi^0 \pi^+ \pi^+ \pi^-$	[a] 1.74 ± 0.50	
$\Gamma_{12} \Xi^0 e^+ \nu_e$	[a] $2.3 \begin{array}{l} +0.7 \\ -0.9 \end{array}$	
$\Gamma_{13} p K^- \pi^+$	[a] 0.20 ± 0.05	

[a] No absolute branching fractions have been measured. The following are branching *ratios* relative to $\Xi^- \pi^+ \pi^+$.

[b] This branching fraction includes all the decay modes of the final-state resonance.

Ξ_c^+ BRANCHING RATIOS

$\Gamma(\Lambda K^- \pi^+ \pi^+)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u>
seen	56
seen	82
² BIAGI 85B looks for but does not see the Ξ_c^+ in $p K^- \bar{K}^0 \pi^+$ ($\Gamma(p K^- \bar{K}^0 \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+) < 0.08$ with 90% CL), $p 2K^- 2\pi^+$ ($\Gamma(p 2K^- 2\pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+) < 0.03$, 90% CL), $\Omega^- K^+ \pi^+$, $\Lambda K^{*0} \pi^+$, and $\Sigma(1385)^+ K^- \pi^+$.	

$\Gamma(\Lambda K^- \pi^+ \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$	Γ_1/Γ_8
<u>VALUE</u>	<u>EVTS</u>
0.58 ± 0.16 ± 0.07	61
BERGFELD	96
CLE2	96
$e^+ e^- \approx \Upsilon(4S)$	

$\Gamma(\Lambda \bar{K}^*(892)^0 \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+)$	Γ_2/Γ_1
Unseen decay modes of the $\bar{K}^*(892)^0$ are included.	
<u>VALUE</u>	<u>CL%</u>
<0.5	90
BERGFELD	96
CLE2	96
$e^+ e^- \approx \Upsilon(4S)$	

$\Gamma(\Sigma(1385)^+ K^- \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+)$

Γ_3/Γ_1

Unseen decay modes of the $\Sigma(1385)^+$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	BERGFELD 96	CLE2	$e^+ e^- \approx \gamma(4S)$

$\Gamma(\Sigma^+ K^- \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$

Γ_4/Γ_8

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.18±0.26±0.17	119	BERGFELD 96	CLE2	$e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.92±0.20±0.07	3 JUN	00 SELX	Σ^- nucleus, 600 GeV	■
0.09 ^{+0.13} _{-0.06} ^{+0.03} _{-0.02}	5 BARLAG	89C ACCM	2 $\Sigma^+ K^- \pi^+$, 3 $\Xi^- \pi^+ \pi^+$	■

³ This JUN 00 result is redundant with other results given below.

$\Gamma(\Sigma^+ \bar{K}^*(892)^0)/\Gamma(\Xi^- \pi^+ \pi^+)$

Γ_5/Γ_8

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.92±0.27±0.14	61 BERGFELD 96	CLE2	$e^+ e^- \approx \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	59 Avery	95 CLE2	$e^+ e^- \approx \gamma(4S)$
------	----------	---------	------------------------------

$\Gamma(\Sigma^0 K^- \pi^+ \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+)$

Γ_6/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.84±0.36	47 COTEUS 87	SPEC	$nA \simeq 600$ GeV	

⁴ See, however, the note on the COTEUS 87 Ξ_c^+ mass measurement.

$\Gamma(\Xi^0 \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$

Γ_7/Γ_8

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.55±0.13±0.09	39 EDWARDS 96	CLE2	$e^+ e^- \approx \gamma(4S)$	

$\Gamma(\Xi^- \pi^+ \pi^+)/\Gamma_{\text{total}}$

Γ_8/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	131 BERGFELD 96	CLE2	$e^+ e^- \approx \gamma(4S)$	
seen	160 Avery	95 CLE2	$e^+ e^- \approx \gamma(4S)$	
seen	30 FRABETTI 93B E687	E687	γBe , $\bar{E}_\gamma = 220$ GeV	
seen	30 ALBRECHT 90F ARG	ARG	$e^+ e^-$ at $\gamma(4S)$	
seen	23 ALAM 89 CLEO	CLEO	$e^+ e^-$ 10.6 GeV	

$\Gamma(\Xi(1530)^0 \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$

Γ_9/Γ_8

Unseen decay modes of the $\Xi(1530)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.2	90 BERGFELD 96	CLE2	$e^+ e^- \approx \gamma(4S)$	

$\Gamma(\Xi^0 \pi^+ \pi^0)/\Gamma(\Xi^- \pi^+ \pi^+)$

Γ_{10}/Γ_8

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
2.34±0.57±0.37	81 EDWARDS 96	CLE2	$e^+ e^- \approx \gamma(4S)$	

$\Gamma(\Xi(1530)^0 \pi^+)/\Gamma(\Xi^0 \pi^+ \pi^0)$ Γ_9/Γ_{10}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.3	90	EDWARDS 96	CLE2	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Xi^0 \pi^+ \pi^+ \pi^-)/\Gamma(\Xi^- \pi^+ \pi^+)$ Γ_{11}/Γ_8

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.74±0.42±0.27	57	EDWARDS 96	CLE2	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Xi^0 e^+ \nu_e)/\Gamma(\Xi^- \pi^+ \pi^+)$ Γ_{12}/Γ_8

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.3±0.6±0.3	41	ALEXANDER 95B	CLE2	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(p K^- \pi^+)/\Gamma(\Sigma^+ K^- \pi^+)$ Γ_{13}/Γ_4

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.22±0.06±0.03	76	JUN 00	SELX	Σ^- nucleus, 600 GeV

 $\Gamma(p K^- \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$ Γ_{13}/Γ_8

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.20±0.04±0.02	76	JUN 00	SELX	Σ^- nucleus, 600 GeV

 Ξ_c^+ REFERENCES

JUN	00	PRL 84 1857	S.Y. Jun <i>et al.</i>	(FNAL SELEX Collab.)
FRAEBETTI	98	PL B427 211	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
BERGFELD	96	PL B365 431	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
EDWARDS	96	PL B373 261	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER	95B	PRL 74 3113	J. Alexander <i>et al.</i>	(CLEO Collab.)
Also	95E	PRL 75 4155 (erratum)	J. Alexander <i>et al.</i>	(CLEO Collab.)
AVERY	95	PRL 75 4364	P. Avery <i>et al.</i>	(CLEO Collab.)
FRAEBETTI	93B	PRL 70 1381	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALBRECHT	90F	PL B247 121	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALAM	89	PL B226 401	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BARLAG	89C	PL B233 522	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
COTÉUS	87	PRL 59 1530	P. Côtéus <i>et al.</i>	(FNAL E400 Collab.)
BIAGI	85B	ZPHY C28 175	S.F. Biagi <i>et al.</i>	(CERN WA62 Collab.)
BIAGI	85C	PL 150B 230	S.F. Biagi <i>et al.</i>	(CERN WA62 Collab.)
BIAGI	83	PL 122B 455	S.F. Biagi <i>et al.</i>	(CERN WA62 Collab.)