



$I(J^P) = 0(?)$

J^P is natural, width and decay modes consistent with 1^- .

$D_s^{*\pm}$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2112.4 ± 0.7 OUR FIT	Error includes scale factor of 1.1.		
2106.6 ± 2.1 ± 2.7	¹ BLAYLOCK 87 MRK3 $e^+ e^- \rightarrow D_s^\pm \gamma X$		

¹ Assuming D_s^\pm mass = 1968.7 ± 0.9 MeV.

$m_{D_s^{*\pm}} - m_{D_s^\pm}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
143.8 ± 0.4 OUR FIT				
143.9 ± 0.4 OUR AVERAGE				
143.76 ± 0.39 ± 0.40		GRONBERG 95	CLE2	$e^+ e^-$
144.22 ± 0.47 ± 0.37		BROWN 94	CLE2	$e^+ e^-$
142.5 ± 0.8 ± 1.5		² ALBRECHT 88	ARG	$e^+ e^- \rightarrow D_s^\pm \gamma X$
139.5 ± 8.3 ± 9.7	60	AIHARA 84D	TPC	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
143.0 ± 18.0	8	ASRATYAN 85	HLBC	FNAL 15-ft, $\nu^2 H$
110 ± 46		BRANDELIK 79	DASP	$e^+ e^- \rightarrow D_s^\pm \gamma X$

² Result includes data of ALBRECHT 84B.

$D_s^{*\pm}$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.9	90	GRONBERG 95	CLE2	$e^+ e^-$
< 4.5	90	ALBRECHT 88	ARG	$E_{cm}^{ee} = 10.2$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.9	90	BROWN 94	CLE2	$e^+ e^-$
< 22	90	BLAYLOCK 87	MRK3	$e^+ e^- \rightarrow D_s^\pm \gamma X$

D_s^{*+} DECAY MODES

D_s^{*-} modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad D_s^+ \gamma$	(94.2±2.5) %
$\Gamma_2 \quad D_s^+ \pi^0$	(5.8±2.5) %

CONSTRAINED FIT INFORMATION

An overall fit to a branching ratio uses 1 measurements and one constraint to determine 2 parameters. The overall fit has a $\chi^2 = 0.0$ for 0 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c} x_2 \\ \boxed{-100} \\ x_1 \end{array}$$

D_s^{*+} BRANCHING RATIOS

$\Gamma(D_s^+ \gamma) / \Gamma_{\text{total}}$	Γ_1 / Γ
VALUE 0.942±0.026 OUR FIT	

• • • We do not use the following data for averages, fits, limits, etc. • • •

	DOCUMENT ID	TECN	COMMENT
seen	ASRATYAN 91	HLBC	$\bar{\nu}_\mu$ Ne
seen	ALBRECHT 88	ARG	$e^+ e^- \rightarrow D_s^\pm \gamma X$
seen	AIHARA 84D		
seen	ALBRECHT 84B		
seen	BRANDELIK 79		

$\Gamma(D_s^+ \pi^0) / \Gamma(D_s^+ \gamma)$	Γ_2 / Γ_1
VALUE 0.062±0.029 OUR FIT	
0.062 ^{+0.020} _{-0.018} ±0.022	GRONBERG 95 CLE2 $e^+ e^-$

$D_s^{*\pm}$ REFERENCES

GRONBERG 95	PRL 75 3232	J. Gronberg <i>et al.</i>	(CLEO Collab.)
BROWN 94	PR D50 1884	D. Brown <i>et al.</i>	(CLEO Collab.)
ASRATYAN 91	PL B257 525	A.E. Asratyan <i>et al.</i>	(ITEP, BELG, SACL+)
ALBRECHT 88	PL B207 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BLAYLOCK 87	PRL 58 2171	G.T. Blaylock <i>et al.</i>	(Mark III Collab.)
ASRATYAN 85	PL 156B 441	A.E. Asratyan <i>et al.</i>	(ITEP, SERP)
AIHARA 84D	PRL 53 2465	H. Aihara <i>et al.</i>	(TPC Collab.)
ALBRECHT 84B	PL 146B 111	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BRANDELIK 79	PL 80B 412	R. Brandelik <i>et al.</i>	(DASP Collab.)

———— OTHER RELATED PAPERS ——

KAMAL 92 PL B284 421
BRANDELIK 78C PL 76B 361
BRANDELIK 77B PL 70B 132

A.N. Kamal, Q.P. Xu
R. Brandelik *et al.*
R. Brandelik *et al.*

(ALBE)
(DASP Collab.)
(DASP Collab.)
