

$I(J^P) = 0(\frac{1}{2}^+)$ Status: ***

In the quark model, a Λ_b^0 is an isospin-0 $ud\bar{b}$ state. The lowest Λ_b^0 ought to have $J^P = 1/2^+$. None of I , J , or P have actually been measured.

Λ_b^0 MASS


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5624 ± 9 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			
5621 ± 4 ± 3	1	ABE	97B CDF	$p\bar{p}$ at 1.8 TeV
5668 ± 16 ± 8	4	2 ABREU	96N DLPH	$e^+e^- \rightarrow Z$
5614 ± 21 ± 4	4	2 BUSKULIC	96L ALEP	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	3	ABE	93B CDF	Sup. by ABE 97B
5640 ± 50 ± 30	16	4 ALBAJAR	91E UA1	$p\bar{p}$ 630 GeV
5640 $^{+100}_{-210}$	52	BARI	91 SFM	$\Lambda_b^0 \rightarrow p D^0 \pi^-$
5650 $^{+150}_{-200}$	90	BARI	91 SFM	$\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \pi^-$

¹ ABE 97B observed 38 events above a background 18 ± 1.6 events in the mass range $5.60\text{--}5.65 \text{ GeV}/c^2$, a significance of > 3.4 standard deviations.

² Uses 4 fully reconstructed Λ_b events.

³ ABE 93B states that, based on the signal claimed by ALBAJAR 91E, CDF should have found $30 \pm 23 \Lambda_b^0 \rightarrow J/\psi(1S)\Lambda$ events. Instead, CDF found not more than 2 events.

⁴ ALBAJAR 91E claims 16 ± 5 events above a background of 9 ± 1 events, a significance of about 5 standard deviations.



Λ_b^0 MEAN LIFE

These are actually measurements of the average lifetime of weakly decaying b baryons weighted by generally unknown production rates, branching fractions, and detection efficiencies. Presumably, the mix is mainly Λ_b^0 , with some Ξ_b^0 and Ξ_b^- .

See b -baryon Admixture section for data on b -baryon mean life average over species of b -baryon particles.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account corrections between the measurements and asymmetric lifetime errors.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.230±0.074 OUR EVALUATION				
1.22 $+0.22$ -0.18	± 0.04	5 ABAZOV	05C D0	$p\bar{p}$ at 1.96 TeV
1.11 $+0.19$ -0.18	± 0.05	6 ABREU	99W DLPH	$e^+ e^- \rightarrow Z$
1.29 $+0.24$ -0.22	± 0.06	6 ACKERSTAFF	98G OPAL	$e^+ e^- \rightarrow Z$
1.21 ± 0.11		6 BARATE	98D ALEP	$e^+ e^- \rightarrow Z$
1.32 ± 0.15	± 0.07	7 ABE	96M CDF	$p\bar{p}$ at 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.19	$\begin{array}{c} +0.21 \\ -0.18 \end{array}$	$\begin{array}{c} +0.07 \\ -0.08 \end{array}$	ABREU	96D DLPH	Repl. by ABREU 99W
1.14	$\begin{array}{c} +0.22 \\ -0.19 \end{array}$	± 0.07	69	AKERS	95K OPAL Repl. by ACKER-STAFF 98G
1.02	$\begin{array}{c} +0.23 \\ -0.18 \end{array}$	± 0.06	44	BUSKULIC	95L ALEP Repl. by BARATE 98D

⁵ Measured mean life using fully reconstructed $\Lambda_b^0 \rightarrow J/\psi \Lambda$ decays.

⁶ Measured using $\Lambda_c \ell^-$ and $\Lambda \ell^+ \ell^-$.

⁷ Excess $\Lambda_c \ell^-$, decay lengths.

$\tau_{\Lambda_b^0}/\tau_{B^0}$ MEAN LIFE RATIO

$\tau_{\Lambda_b^0}/\tau_{B^0}$ (direct measurements)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.87^{+0.17}_{-0.14} \pm 0.03$	8 ABAZOV	05C D0	$p\bar{p}$ at 1.96 TeV

⁸ Measured mean life ratio using fully reconstructed decays.

Λ_b^0 DECAY MODES

These branching fractions are actually an average over weakly decaying b -baryons weighted by their production rates in Z decay (or high-energy $p\bar{p}$), branching ratios, and detection efficiencies. They scale with the LEP b -baryon production fraction $B(b \rightarrow b\text{-baryon})$ and are evaluated for our value $B(b \rightarrow b\text{-baryon}) = (10.0 \pm 2.0)\%$.

The branching fractions $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{anything})$ and $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})$ are not pure measurements because the underlying measured products of these with $B(b \rightarrow b\text{-baryon})$ were used to determine $B(b \rightarrow b\text{-baryon})$, as described in the note “Production and Decay of b -Flavored Hadrons.”

For inclusive branching fractions, e.g., $B \rightarrow D^\pm \text{anything}$, the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $J/\psi(1S)\Lambda$	$(4.7 \pm 2.8) \times 10^{-4}$	
Γ_2 $p D^0 \pi^-$		
Γ_3 $\Lambda_c^+ \pi^-$	seen	
Γ_4 $\Lambda_c^+ a_1(1260)^-$	seen	
Γ_5 $\Lambda_c^+ \pi^+ \pi^- \pi^-$		
Γ_6 $\Lambda K^0 2\pi^+ 2\pi^-$		

Γ_7	$\Lambda_c^+ \ell^- \bar{\nu}_\ell$ anything	[a]	(9.1±2.3) %	
Γ_8	$\Lambda_c^+ \ell^- \bar{\nu}_\ell$		(5.0 $^{+1.9}_{-1.4}$) %	
Γ_9	$\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell$		(5.6±3.1) %	
Γ_{10}	$p h^-$	[b]	< 2.3 $\times 10^{-5}$	90%
Γ_{11}	$p \pi^-$		< 5.0 $\times 10^{-5}$	90%
Γ_{12}	$p K^-$		< 5.0 $\times 10^{-5}$	90%
Γ_{13}	$\Lambda \gamma$		< 1.3 $\times 10^{-3}$	90%

[a] Not a pure measurement. See note at head of Λ_b^0 Decay Modes.

[b] Here h^- means π^- or K^- .

Λ_b^0 BRANCHING RATIOS

$\Gamma(J/\psi(1S)\Lambda)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.7± 2.1± 1.9		⁹ ABE	97B CDF	$p\bar{p}$ at 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

180. ±110. ±36. 16 ¹⁰ ALBAJAR 91E UA1 $J/\psi(1S) \rightarrow \mu^+ \mu^-$

⁹ ABE 97B reports $(0.037 \pm 0.017(\text{stat}) \pm 0.007(\text{sys}))\%$ for $B(b \rightarrow b\text{-baryon}) = 0.1$ and for $B(B^0 \rightarrow J/\psi(1S) K_S^0) = 0.037\%$. We rescale to our PDG 98 best value $B(b \rightarrow b\text{-baryon}) = (10.1^{+3.9}_{-3.1})\%$ and $B(B^0 \rightarrow J/\psi(1S) K_S^0) = (0.044 \pm 0.006)\%$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰ ALBAJAR 91E reports $(180 \pm 110) \times 10^{-4}$ for $B(\bar{b} \rightarrow b\text{-baryon}) = 0.10$. We rescale to our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (10.0 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p D^0 \pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	52	BARI	91 SFM	$D^0 \rightarrow K^- \pi^+$
seen		BASILE	81 SFM	$D^0 \rightarrow K^- \pi^+$

$\Gamma(\Lambda_c^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	3	ABREU	96N DLPH	$\Lambda_c^+ \rightarrow p K^- \pi^+$
seen	4	BUSKULIC	96L ALEP	$\Lambda_c^+ \rightarrow p K^- \pi^+, p \bar{K}^0, \Lambda \pi^+ \pi^+ \pi^-$

$\Gamma(\Lambda_c^+ a_1(1260)^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	1	ABREU	96N DLPH	$\Lambda_c^+ \rightarrow p K^- \pi^+, a_1^- \rightarrow \rho^0 \pi^- \rightarrow \pi^+ \pi^- \pi^-$

$\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	90	BARI	91 SFM	$\Lambda_c^+ \rightarrow p K^- \pi^+$

$\Gamma(\Lambda K^0 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	4	¹¹ ARENTON	86 FMPS	$\Lambda K_S^0 2\pi^+ 2\pi^-$

¹¹ See the footnote to the ARENTON 86 mass value.

$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}$ Γ_7/Γ

The values and averages in this section serve only to show what values result if one assumes our $B(b \rightarrow b\text{-baryon})$. They cannot be thought of as measurements since the underlying product branching fractions were also used to determine $B(b \rightarrow b\text{-baryon})$ as described in the note on "Production and Decay of *b*-Flavored Hadrons."

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.091±0.023 OUR AVERAGE

0.086±0.016±0.017 12 BARATE 98D ALEP $e^+ e^- \rightarrow Z$

0.118^{+0.040}_{-0.033}±0.024 29 ABREU 95S DLPH $e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.075±0.018±0.015 55 BUSKULIC 95L ALEP Repl. by BARATE 98D

0.15 ± 0.06 ± 0.03 21 BUSKULIC 92E ALEP $\Lambda_c^+ \rightarrow p K^- \pi^+$

¹² BARATE 98D reports $[B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = 0.0086 \pm 0.0007 \pm 0.0014$. We divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (10.0 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Measured using $\Lambda_c^+ \ell^-$ and $\Lambda \ell^+ \ell^-$.

¹³ ABREU 95S reports $[B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = 0.0118 \pm 0.0026 \pm 0.0031$. We divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (10.0 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁴ BUSKULIC 95L reports $[B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = 0.00755 \pm 0.0014 \pm 0.0012$. We divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (10.0 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁵ BUSKULIC 92E reports $[B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = 0.015 \pm 0.0035 \pm 0.0045$. We divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (10.0 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by BUSKULIC 95L.

$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.050^{+0.011}_{-0.008}^{+0.016}_{-0.012}	16 ABDALLAH	04A DLPH	$e^+ e^- \rightarrow Z^0$

¹⁶ Derived from a combined likelihood and event rate fit to the distribution of the Isgur-Wise variable and using HQET. The slope of the form factor is measured to be $\rho^2 = 2.03 \pm 0.46^{+0.72}_{-1.00}$.

$\Gamma(\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)/\Gamma_{\text{total}}$ Γ_9/Γ VALUE**0.056^{+0.031}_{-0.030}**DOCUMENT ID

17 ABDALLAH 04A DLPH

TECNCOMMENT $e^+ e^- \rightarrow Z^0$

17 Derived from the fraction of $\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell) / (\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell) + \Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)) = 0.47^{+0.10+0.07}_{-0.08-0.06}$.

 $\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)/[\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell) + \Gamma(\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)]$ $\Gamma_8/(\Gamma_8+\Gamma_9)$ VALUE**0.47^{+0.10+0.07}_{-0.08-0.06}**DOCUMENT ID

ABDALLAH 04A DLPH

TECNCOMMENT $e^+ e^- \rightarrow Z^0$ $\Gamma(p h^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ VALUE**<2.3 × 10⁻⁵**CL%

90

DOCUMENT ID

18 ACOSTA 050 CDF

TECNCOMMENT $p\bar{p}$ at 1.96 TeV

18 Assumes $f_{\Lambda} / f_d = 0.25$, and equal momentum distribution for Λ_b and B mesons.

 $\Gamma(p \pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ VALUE**<5.0 × 10⁻⁵**CL%

90

DOCUMENT ID

19 BUSKULIC 96V ALEP

TECNCOMMENT $e^+ e^- \rightarrow Z$

19 BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

 $\Gamma(p K^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ VALUE**<5.0 × 10⁻⁵**CL%

90

DOCUMENT ID

20 BUSKULIC 96V ALEP

TECNCOMMENT $e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $<3.6 \times 10^{-4}$ CL%

90

DOCUMENT ID

21 ADAM 96D DLPH

TECNCOMMENT $e^+ e^- \rightarrow Z$

20 BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

21 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

 $\Gamma(\Lambda \gamma)/\Gamma_{\text{total}}$ Γ_{13}/Γ VALUE**<1.3 × 10⁻³**CL%

90

DOCUMENT ID

ACOSTA 02G CDF

TECNCOMMENT $p\bar{p}$ at 1.8 TeV Λ_b^0 REFERENCES

ABAZOV	05C	PRL 94 102001
ACOSTA	05O	PR D72 051104R
ABDALLAH	04A	PL B585 63
ACOSTA	02G	PR D66 112002
ABREU	99W	EPJ C10 185
ACKERSTAFF	98G	PL B426 161
BARATE	98D	EPJ C2 197
PDG	98	EPJ C3 1
ABE	97B	PR D55 1142
ABE	96M	PRL 77 1439
ABREU	96D	ZPHY C71 199
ABREU	96N	PL B374 351
ADAM	96D	ZPHY C72 207

V.M. Abazov <i>et al.</i>	(D0 Collab.)
D. Acosta <i>et al.</i>	(CDF Collab.)
J. Abdallah <i>et al.</i>	(DELPHI Collab.)
D. Acosta <i>et al.</i>	(CDF Collab.)
P. Abreu <i>et al.</i>	(DELPHI Collab.)
K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
R. Barate <i>et al.</i>	(ALEPH Collab.)
C. Caso <i>et al.</i>	
F. Abe <i>et al.</i>	(CDF Collab.)
F. Abe <i>et al.</i>	(CDF Collab.)
P. Abreu <i>et al.</i>	(DELPHI Collab.)
P. Abreu <i>et al.</i>	(DELPHI Collab.)
W. Adam <i>et al.</i>	(DELPHI Collab.)

BUSKULIC	96L	PL B380 442	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96V	PL B384 471	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	
ABREU	95S	ZPHY C68 375	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	95K	PL B353 402	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	95L	PL B357 685	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABE	93B	PR D47 R2639	F. Abe <i>et al.</i>	(CDF Collab.)
BUSKULIC	92E	PL B294 145	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ALBAJAR	91E	PL B273 540	C. Albajar <i>et al.</i>	(UA1 Collab.)
BARI	91	NC 104A 1787	G. Bari <i>et al.</i>	(CERN R422 Collab.)
ARENTON	86	NP B274 707	M.W. Arenton <i>et al.</i>	(ARIZ, NDAM, VAND)
BASILE	81	LNC 31 97	M. Basile <i>et al.</i>	(CERN R415 Collab.)