

$N(1440) P_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ****$$

Most of the results published before 1975 were last included in our 1982 edition, Physics Letters **111B** 1 (1982). Some further obsolete results published before 1980 were last included in our 2006 edition, Journal of Physics, G **33** 1 (2006).

$N(1440)$ BREIT-WIGNER MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1420 to 1470 (≈ 1440) OUR ESTIMATE			
1485.0 \pm 1.2	ARNDT	06	DPWA $\pi N \rightarrow \pi N, \eta N$
1462 \pm 10	MANLEY	92	IPWA $\pi N \rightarrow \pi N \& N\pi\pi$
1440 \pm 30	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1410 \pm 12	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1468.0 \pm 4.5	ARNDT	04	DPWA $\pi N \rightarrow \pi N, \eta N$
1518 \pm 5	PENNER	02C	DPWA Multichannel
1479 \pm 80	VRANA	00	DPWA Multichannel
1463 \pm 7	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
1467	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1421 \pm 18	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
1465	LI	93	IPWA $\gamma N \rightarrow \pi N$
1471	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$
1411	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
1380	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
1390	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

$N(1440)$ BREIT-WIGNER WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
200 to 450 (≈ 300) OUR ESTIMATE			
284 \pm 18	ARNDT	06	DPWA $\pi N \rightarrow \pi N, \eta N$
391 \pm 34	MANLEY	92	IPWA $\pi N \rightarrow \pi N \& N\pi\pi$
340 \pm 70	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
135 \pm 10	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
360 \pm 26	ARNDT	04	DPWA $\pi N \rightarrow \pi N, \eta N$
668 \pm 41	PENNER	02C	DPWA Multichannel
490 \pm 120	VRANA	00	DPWA Multichannel
360 \pm 20	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
440	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
250 \pm 63	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
315	LI	93	IPWA $\gamma N \rightarrow \pi N$
545 \pm 170	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$
334	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
200	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
200	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

N(1440) POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1350 to 1380 (\approx 1365) OUR ESTIMATE			
1359	³ ARNDT	06	DPWA $\pi N \rightarrow \pi N, \eta N$
1385	⁴ HOEHLER	93	SPED $\pi N \rightarrow \pi N$
1375 \pm 30	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1357	⁵ ARNDT	04	DPWA $\pi N \rightarrow \pi N, \eta N$
1383	VRANA	00	DPWA Multichannel
1346	⁶ ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1360	⁷ ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
1370	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$
1381 or 1379	⁸ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
1360 or 1333	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

–2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
160 to 220 (\approx 190) OUR ESTIMATE			
162	³ ARNDT	06	DPWA $\pi N \rightarrow \pi N, \eta N$
164	⁴ HOEHLER	93	SPED $\pi N \rightarrow \pi N$
180 \pm 40	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
160	⁵ ARNDT	04	DPWA $\pi N \rightarrow \pi N, \eta N$
316	VRANA	00	DPWA Multichannel
176	⁶ ARNDT	95	DPWA $\pi N \rightarrow N\pi$
252	⁷ ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
228	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$
209 or 210	⁸ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
167 or 234	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

N(1440) ELASTIC POLE RESIDUE

MODULUS $|r|$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
38	³ ARNDT	06	DPWA $\pi N \rightarrow \pi N, \eta N$
40	HOEHLER	93	SPED $\pi N \rightarrow \pi N$
52 \pm 5	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
36	⁵ ARNDT	04	DPWA $\pi N \rightarrow \pi N, \eta N$
42	⁶ ARNDT	95	DPWA $\pi N \rightarrow N\pi$
109	⁷ ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
74	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$

PHASE θ

VALUE ($^\circ$)	DOCUMENT ID	TECN	COMMENT
- 98	³ ARNDT	06	DPWA $\pi N \rightarrow \pi N, \eta N$
-100 \pm 35	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-102	⁵ ARNDT	04	DPWA $\pi N \rightarrow \pi N, \eta N$
-101	⁶ ARNDT	95	DPWA $\pi N \rightarrow N\pi$
- 93	⁷ ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
- 84	CUTKOSKY	90	IPWA $\pi N \rightarrow \pi N$

N(1440) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	0.55 to 0.75
Γ_2 $N\eta$	
Γ_3 $N\pi\pi$	30-40 %
Γ_4 $\Delta\pi$	20-30 %
Γ_5 $\Delta(1232)\pi, P$ -wave	
Γ_6 $N\rho$	<8 %
Γ_7 $N\rho, S=1/2, P$ -wave	
Γ_8 $N\rho, S=3/2, P$ -wave	
Γ_9 $N(\pi\pi)_{S\text{-wave}}^{I=0}$	5-10 %
Γ_{10} $p\gamma$	0.035-0.048 %
Γ_{11} $p\gamma, \text{helicity}=1/2$	0.035-0.048 %
Γ_{12} $n\gamma$	0.009-0.032 %
Γ_{13} $n\gamma, \text{helicity}=1/2$	0.009-0.032 %

N(1440) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.55 to 0.75 OUR ESTIMATE				
0.787 \pm 0.016	ARNDT	06	DPWA $\pi N \rightarrow \pi N, \eta N$	
0.69 \pm 0.03	MANLEY	92	IPWA $\pi N \rightarrow \pi N \& N\pi\pi$	
0.68 \pm 0.04	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$	
0.51 \pm 0.05	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.750 \pm 0.024	ARNDT	04	DPWA $\pi N \rightarrow \pi N, \eta N$	
0.57 \pm 0.01	PENNER	02C	DPWA Multichannel	
0.72 \pm 0.05	VRANA	00	DPWA Multichannel	
0.68	ARNDT	95	DPWA $\pi N \rightarrow N\pi$	
0.56 \pm 0.08	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$	

$\Gamma(N\eta)/\Gamma_{\text{total}}$					Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.00±0.01	VRANA	00	DPWA	Multichannel	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1440) \rightarrow \Delta(1232)\pi$, <i>P-wave</i>					$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
+0.37 to +0.41 OUR ESTIMATE					
+0.39±0.02	MANLEY	92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
+0.41	^{1,9} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$	
+0.37	² LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$	

$\Gamma(\Delta(1232)\pi)$, <i>P-wave</i> / Γ_{total}					Γ_5/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.16±0.01	VRANA	00	DPWA	Multichannel	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1440) \rightarrow N\rho$, <i>S=1/2, P-wave</i>					$(\Gamma_1\Gamma_7)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
±0.07 to ±0.25 OUR ESTIMATE					
-0.11	^{1,9} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$	
+0.23	² LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$	

$\Gamma(N\rho, S=1/2)$, <i>P-wave</i> / Γ_{total}					Γ_7/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.00±0.01	VRANA	00	DPWA	Multichannel	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1440) \rightarrow N\rho$, <i>S=3/2, P-wave</i>					$(\Gamma_1\Gamma_8)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
+0.18	^{1,9} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1440) \rightarrow N(\pi\pi)_{S=0}^{I=0}$, <i>S-wave</i>					$(\Gamma_1\Gamma_9)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
±0.17 to ±0.25 OUR ESTIMATE					
+0.24±0.03	MANLEY	92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
-0.18	^{1,9} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$	
-0.23	² LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$	

$\Gamma(N(\pi\pi)_{S=0}^{I=0})/\Gamma_{\text{total}}$					Γ_9/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.12±0.01	VRANA	00	DPWA	Multichannel	

N(1440) PHOTON DECAY AMPLITUDES

N(1440) → pγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
−0.065 ±0.004 OUR ESTIMATE			
−0.063 ±0.005	ARNDT	96	IPWA γ N → π N
−0.069 ±0.018	CRAWFORD	83	IPWA γ N → π N
−0.063 ±0.008	AWAJI	81	DPWA γ N → π N
−0.069 ±0.004	ARAI	80	DPWA γ N → π N (fit 1)
−0.066 ±0.004	ARAI	80	DPWA γ N → π N (fit 2)
−0.079 ±0.009	BRATASHEV...	80	DPWA γ N → π N
−0.068 ±0.015	CRAWFORD	80	DPWA γ N → π N
−0.0584 ±0.0148	ISHII	80	DPWA Compton scattering
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
−0.087	PENNER	02D	DPWA Multichannel
−0.085 ±0.003	LI	93	IPWA γ N → π N
−0.129	¹⁰ WADA	84	DPWA Compton scattering

N(1440) → nγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.040 ±0.010 OUR ESTIMATE			
0.045 ±0.015	ARNDT	96	IPWA γ N → π N
0.037 ±0.010	AWAJI	81	DPWA γ N → π N
0.030 ±0.003	FUJII	81	DPWA γ N → π N
0.023 ±0.009	ARAI	80	DPWA γ N → π N (fit 1)
0.019 ±0.012	ARAI	80	DPWA γ N → π N (fit 2)
0.056 ±0.015	CRAWFORD	80	DPWA γ N → π N
−0.029 ±0.035	TAKEDA	80	DPWA γ N → π N
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.121	PENNER	02D	DPWA Multichannel
0.085 ±0.006	LI	93	IPWA γ N → π N

N(1440) FOOTNOTES

- ¹ LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to π N → Nππ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ² From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ³ ARNDT 06 also finds a second-sheet pole with real part = 1388 MeV, −2 × imaginary part = 165 MeV, and residue with modulus 86 MeV and phase = −46 degrees.
- ⁴ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of π N elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- ⁵ ARNDT 04 also finds a second-sheet pole with real part = 1385 MeV, −2 × imaginary part = 166 MeV, and residue with modulus 82 MeV and phase = −51°.
- ⁶ ARNDT 95 also finds a second-sheet pole with real part = 1383 MeV, −2 × imaginary part = 210 MeV, and residue with modulus 92 MeV and phase = −54°.
- ⁷ ARNDT 91 (Soln SM90) also finds a second-sheet pole with real part = 1413 MeV, −2 × imaginary part = 256 MeV, and residue = (78−153i) MeV.

⁸ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

⁹ LONGACRE 77 considers this coupling to be well determined.

¹⁰ WADA 84 is inconsistent with other analyses; see the Note on N and Δ Resonances.

N(1440) REFERENCES

For early references, see Physics Letters **111B** 1 (1982).

ARNDT	06	PR C74 045205	R.A. Arndt <i>et al.</i>	(GWU)
PDG	06	JPG 33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
ARNDT	04	PR C69 035213	R.A. Arndt <i>et al.</i>	(GWU, TRIU)
PENNER	02C	PR C66 055211	G. Penner, U. Mosel	(GIES)
PENNER	02D	PR C66 055212	G. Penner, U. Mosel	(GIES)
VRANA	00	PRPL 328 181	T.P. Vrana, S.A. Dytman,, T.-S.H. Lee	(PITT+)
ARNDT	96	PR C53 430	R.A. Arndt, I.I. Strakovsky, R.L. Workman	(VPI)
ARNDT	95	PR C52 2120	R.A. Arndt <i>et al.</i>	(VPI, BRCO)
BATINIC	95	PR C51 2310	M. Batinic <i>et al.</i>	(BOSK, UCLA)
Also		PR C57 1004 (erratum)	M. Batinic <i>et al.</i>	
HOEHLER	93	πN Newsletter 9 1	G. Hohler	(KARL)
LI	93	PR C47 2759	Z.J. Li <i>et al.</i>	(VPI)
MANLEY	92	PR D45 4002	D.M. Manley, E.M. Saleski	(KENT) IJP
Also		PR D30 904	D.M. Manley <i>et al.</i>	(VPI)
ARNDT	91	PR D43 2131	R.A. Arndt <i>et al.</i>	(VPI, TELE) IJP
CUTKOSKY	90	PR D42 235	R.E. Cutkosky, S. Wang	(CMU)
WADA	84	NP B247 313	Y. Wada <i>et al.</i>	(INUS)
CRAWFORD	83	NP B211 1	R.L. Crawford, W.T. Morton	(GLAS)
PDG	82	PL 111B 1	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	N. Awaji, R. Kajikawa	(NAGO)
Also		NP B197 365	K. Fujii <i>et al.</i>	(NAGO)
FUJII	81	NP B187 53	K. Fujii <i>et al.</i>	(NAGO, OSAK)
ARAI	80	Toronto Conf. 93	I. Arai	(INUS)
Also		NP B194 251	I. Arai, H. Fujii	(INUS)
BRATASHEV...	80	NP B166 525	A.S. Bratashvsky <i>et al.</i>	(KFTI)
CRAWFORD	80	Toronto Conf. 107	R.L. Crawford	(GLAS)
CUTKOSKY	80	Toronto Conf. 19	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
Also		PR D20 2839	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
ISHII	80	NP B165 189	T. Ishii <i>et al.</i>	(KYOT, INUS)
TAKEDA	80	NP B168 17	H. Takeda <i>et al.</i>	(TOKY, INUS)
HOEHLER	79	PDAT 12-1	G. Hohler <i>et al.</i>	(KARLT) IJP
Also		Toronto Conf. 3	R. Koch	(KARLT) IJP
LONGACRE	78	PR D17 1795	R.S. Longacre <i>et al.</i>	(LBL, SLAC)
LONGACRE	77	NP B122 493	R.S. Longacre, J. Dolbeau	(SACL) IJP
Also		NP B108 365	J. Dolbeau <i>et al.</i>	(SACL) IJP
LONGACRE	75	PL 55B 415	R.S. Longacre <i>et al.</i>	(LBL, SLAC) IJP
