

# K<sub>1</sub>(1650)

$$I(J^P) = \frac{1}{2}(1^+)$$

## OMITTED FROM SUMMARY TABLE

This entry contains various peaks in strange meson systems ( $K^+ \phi$ ,  $K \pi \pi$ ) reported in partial-wave analysis in the 1600–1900 mass region.

### K<sub>1</sub>(1650) MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>1650±50</b>	FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 1840	ARMSTRONG	83	OMEG -	18.5 $K^- p \rightarrow 3K p$
~ 1800	DAUM	81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$

### K<sub>1</sub>(1650) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>150±50</b>	FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 250	DAUM	81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$

### K<sub>1</sub>(1650) DECAY MODES

Mode	
$\Gamma_1$	$K \pi \pi$
$\Gamma_2$	$K \phi$

### K<sub>1</sub>(1650) REFERENCES

FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)
ARMSTRONG	83	NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)