K^0_S

$I(J^P) = \frac{1}{2}(0^-)$

K^0_S MEAN LIFE

For earlier measurements, beginning with BOLDT 58B, see our 1986 edition, Physics Letters **170B** 130 (1986).

OUR FIT is described in the note on "*CP* violation in K_L decays" in the K^0_L Particle Listings. The result labeled "OUR FIT Assuming *CPT*" ["OUR FIT Not assuming *CPT*"] includes all measurements except those with the comment "Not assuming *CPT*" ["Assuming *CPT*"]. Measurements with neither comment do not assume *CPT* and enter both fits.

VALUE (10^{-10} s)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
0.8953 ±0.0005 OUR FIT	Error includes scale factor of 1.1. Assuming *CPT*				
0.8958 ±0.0005 OUR FIT	Not assuming *CPT*				
0.8965 ±0.0007		[1,2] ALAVI-HARATI	03	KTEV	Assuming *CPT*
0.8958 ±0.0013		[2,3] ALAVI-HARATI	03	KTEV	Not assuming *CPT*
0.89598±0.00048±0.00051	16M	LAI	02C	NA48	
0.8971 ±0.0021		BERTANZA	97	NA31	
0.8941 ±0.0014 ±0.0009		SCHWINGEN...	95	E773	Assuming *CPT*
0.8929 ±0.0016		GIBBONS	93	E731	Assuming *CPT*
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.8920 ±0.0044	214k	GROSSMAN	87	SPEC	
0.905 ±0.007		[4] ARONSON	82B	SPEC	
0.881 ±0.009	26k	ARONSON	76	SPEC	
0.8926 ±0.0032 ±0.0002		[5] CARITHERS	75	SPEC	
0.8937 ±0.0048	6M	GEWENIGER	74B	ASPK	
0.8958 ±0.0045	50k	[6] SKJEGGEST...	72	HBC	
0.856 ±0.008	19994	[7] DONALD	68B	HBC	
0.872 ±0.009	20000	[7,8] HILL	68	DBC	

[1] This ALAVI-HARATI 03 fit has Δm and τ_S free but constrains ϕ_{+-} to the Superweak value, i.e. assumes *CPT*. This τ_S value is correlated with their $\Delta m = m_{K^0_L} - m_{K^0_S}$ measurement in the K^0_L listings. The correlation coefficient $\rho(\tau_S,\Delta m) = -0.396$.

[2] The two ALAVI-HARATI 03 values use the same data. The first enters the "assuming *CPT*" fit and the second enters the "not assuming *CPT*" fit.

[3] This ALAVI-HARATI 03 fit has Δm, ϕ_{+-}, and τ_{K_S} free. See ϕ_{+-} in the "K_L *CP* violation" section for correlation information.

[4] ARONSON 82 find that K^0_S mean life may depend on the kaon energy.

[5] CARITHERS 75 measures the Δm dependence of the total decay rate (inverse mean life) to be $\Gamma(K^0_S) = \left[(1.122 \pm 0.004) + 0.16(\Delta m - 0.5348)/\Delta m\right] 10^{10}$/s, or, in terms of mean life, CARITHERS 75 measures $\tau_S = (0.8913 \pm 0.0032) - 0.238\ [\Delta m - 0.5348]$ (10^{-10} s). We have adjusted the measurement to use our best values of ($\Delta m = 0.5292 \pm 0.0009$) ($10^{10}\ \hbar\ \mathrm{s}^{-1}$). Our first error is their experiment's error and our second error is the systematic error from using our best values.

[6] HILL 68 has been changed by the authors from the published value (0.865 ± 0.009) because of a correction in the shift due to η_{+-}. SKJEGGESTAD 72 and HILL 68 give detailed discussions of systematics encountered in this type of experiment.

[7] Pre-1971 experiments are excluded from the average because of disagreement with later more precise experiments.

[8] HILL 68 has been changed by the authors from the published value (0.865 ± 0.009) because of a correction in the shift due to η_{+-}. SKJEGGESTAD 72 and HILL 68 give detailed discussions of systematics encountered in this type of experiment.

K^0_S DECAY MODES

	Mode			Fraction (Γ_i/Γ)	Confidence level
	Hadronic modes				
Γ_1	$\pi^0\pi^0$			(30.69 ± 0.05) %	
Γ_2	$\pi^+\pi^-$			(69.20 ± 0.05) %	
Γ_3	$\pi^+\pi^-\pi^0$			$(3.5\ ^{+1.1}_{-0.9})\times10^{-7}$	
	Modes with photons or $\ell\overline{\ell}$ pairs				
Γ_4	$\pi^+\pi^-\gamma$		[*a*,*b*]	$(1.79\pm0.05)\times10^{-3}$	
Γ_5	$\pi^+\pi^-e^+e^-$			$(4.69\pm0.30)\times10^{-5}$	
Γ_6	$\pi^0\gamma\gamma$		[*b*]	$(4.9\pm1.8)\times10^{-8}$	
Γ_7	$\gamma\gamma$			$(2.71\pm0.06)\times10^{-6}$	
	Semileptonic modes				
Γ_8	$\pi^\pm e^\mp\nu_e$		[*c*]	$(7.04\pm0.09)\times10^{-4}$	
Γ_9	$\pi^\pm\mu^\mp\nu_\mu$		[*c*,*d*]	$(4.69\pm0.06)\times10^{-4}$	
	***CP* violating (*CP*) and $\Delta S=1$ weak neutral current (*S1*) modes**				
Γ_{10}	$3\pi^0$	*CP*		$<1.2\times10^{-7}$	90%
Γ_{11}	$\mu^+\mu^-$	*S1*		$<3.2\times10^{-7}$	90%
Γ_{12}	e^+e^-	*S1*		$<1.4\times10^{-7}$	90%
Γ_{13}	$\pi^0e^+e^-$	*S1*	[*b*]	$(3.0\ ^{+1.5}_{-1.2})\times10^{-9}$	
Γ_{14}	$\pi^0\mu^+\mu^-$	*S1*		$(2.9\ ^{+1.5}_{-1.2})\times10^{-9}$	

[*a*] Most of this radiative mode, the low-momentum γ part, is also included in the parent mode listed without γ's.

[*b*] See the Particle Listings below for the energy limits used in this measurement.

[*c*] The value is for the sum of the charge states or particle/antiparticle states indicated.

[*d*] Not a measurement. Calculated as $0.666\cdot B(\pi^\pm e^\mp\nu_e)$.

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 4 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 =$ 0.1 for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\left\langle \delta x_i \delta x_j \right\rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	−100		
x_8	−6	4	
x_9	−6	4	100
	x_1	x_2	x_8

K^0_S DECAY RATES

$\Gamma(\pi^\pm e^\mp \nu_e)$ — Γ_8

VALUE (10^6 s^{-1})	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •					
8.1 ±1.6	75	[9] AKHMETSHIN	99	CMD2	Tagged K^0_S using $\phi \to K^0_L K^0_S$
7.50±0.08		[10] PDG	98		
seen		BURGUN	72	HBC	$K^+ p \to K^0 p \pi^+$
9.3 ±2.5		AUBERT	65	HLBC	ΔS=ΔQ, *CP* cons. not assumed

[9] AKHMETSHIN 99 is from a measured branching ratio B($K^0_S \to \pi e \nu_e$)= $(7.2 \pm 1.4) \times 10^{-4}$ and $\tau_{K^0_S} = (0.8934 \pm 0.0008) \times 10^{-10}$ s. Not independent of measured branching ratio.

[10] PDG 98 from K^0_L measurements, assuming that ΔS=ΔQ in K^0 decay so that $\Gamma(K^0_S \to \pi^\pm e^\mp \nu_e) = \Gamma(K^0_L \to \pi^\pm e^\mp \nu_e)$.

$\Gamma(\pi^\pm \mu^\mp \nu_\mu)$ — Γ_9

VALUE (10^6 s^{-1})	*DOCUMENT ID*	
• • • We do not use the following data for averages, fits, limits, etc. • • •		
5.25±0.07	[11] PDG	98

[11] PDG 98 from K^0_L measurements, assuming that ΔS=ΔQ in K^0 decay so that $\Gamma(K^0_S \to \pi^\pm \mu^\mp \nu_\mu) = \Gamma(K^0_L \to \pi^\pm \mu^\mp \nu_\mu)$.

K^0_S BRANCHING RATIOS

Hadronic modes

$\Gamma(\pi^0\pi^0)/\Gamma_{total}$ — Γ_1/Γ

VALUE	EVTS	DOCUMENT ID		TECN
0.3069±0.0005 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.335 ±0.014	1066	BROWN	63	HLBC
0.288 ±0.021	198	CHRETIEN	63	HLBC
0.30 ±0.035		BROWN	61	HLBC

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$ — Γ_2/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.6920±0.0005 OUR FIT					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.670 ±0.010	3447	DOYLE	69	HBC	$\pi^- p \to \Lambda K^0$

$\Gamma(\pi^+\pi^-)/\Gamma(\pi^0\pi^0)$ — Γ_2/Γ_1

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
2.255 ±0.005 OUR FIT					
2.2549±0.0054		[12] AMBROSINO	06C	KLOE	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.2555±0.0012±0.0054		[13] AMBROSINO	06C	KLOE	
2.236 ±0.003 ±0.015	766k	[13] ALOISIO	02B	KLOE	
2.11 ±0.09	1315	EVERHART	76	WIRE	$\pi^- p \to \Lambda K^0$
2.169 ±0.094	16k	COWELL	74	OSPK	$\pi^- p \to \Lambda K^0$
2.16 ±0.08	4799	HILL	73	DBC	$K^+ d \to K^0 p p$
2.22 ±0.10	3068	[14] ALITTI	72	HBC	$K^+ p \to \pi^+ p K^0$
2.22 ±0.08	6380	MORSE	72B	DBC	$K^+ n \to K^0 p$
2.10 ±0.11	701	[15] NAGY	72	HLBC	$K^+ n \to K^0 p$
2.22 ±0.095	6150	[16] BALTAY	71	HBC	$K p \to K^0$ neutrals
2.282 ±0.043	7944	[17] MOFFETT	70	OSPK	$K^+ n \to K^0 p$
2.12 ±0.17	267	[15] BOZOKI	69	HLBC	
2.285 ±0.055	3016	[17] GOBBI	69	OSPK	$K^+ n \to K^0 p$
2.10 ±0.06	3700	MORFIN	69	HLBC	$K^+ n \to K^0 p$

[12] This result combines AMBROSINO 06C KLOE 2001-02 data with ALOISIO 02B KLOE 2000 data. $K^0_S \to \pi^+\pi^-$ fully inclusive.

[13] Includes radiative decays $\pi^+\pi^-\gamma$.

[14] The directly measured quantity is $K^0_S \to \pi^+\pi^-/\text{all } K^0 = 0.345 \pm 0.005$.

[15] NAGY 72 is a final result which includes BOZOKI 69.

[16] The directly measured quantity is $K^0_S \to \pi^+\pi^-/\text{all } \overline{K}^0 = 0.345 \pm 0.005$.

[17] MOFFETT 70 is a final result which includes GOBBI 69.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\rm total}$ Γ_3/Γ

VALUE (units 10^{-7})	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
$\mathbf{3.5^{+1.1}_{-0.9}}$ **OUR AVERAGE**					
$4.7^{+2.2}_{-1.7}{}^{+1.7}_{-1.5}$		[18] BATLEY	05	NA48	
$2.5^{+1.3}_{-1.0}{}^{+0.5}_{-0.6}$	500k	[19] ADLER	97B	CPLR	
$4.8^{+2.2}_{-1.6}\pm1.1$		[20] ZOU	96	E621	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$4.1^{+2.5}_{-1.9}{}^{+0.5}_{-0.6}$		[21] ADLER	96E	CPLR	Sup. by ADLER 97B
$3.9^{+5.4}_{-1.8}{}^{+0.9}_{-0.7}$		[22] THOMSON	94	E621	Sup. by ZOU 96

[18] BATLEY 05 is obtained by measuring the interference parameters in K_S, $K_L \rightarrow \pi^+\pi^-\pi^0$: Re($\lambda$) = $0.038 \pm 0.008 \pm 0.006$ and Im(λ) = $-0.013 \pm 0.005 \pm 0.004$; the correlation coeff. between Re(λ) and Im(λ) is 0.66 (statistical only).

[19] ADLER 97B find the *CP*-conserving parameters Re(λ) = $(28 \pm 7 \pm 3) \times 10^{-3}$, Im($\lambda$) = $(-10 \pm 8 \pm 2) \times 10^{-3}$. They estimate B($K^0_S \rightarrow \pi^+\pi^-\pi^0$) from Re($\lambda$) and the K^0_L decay parameters. See also ANGELOPOULOS 98C.

[20] ZOU 96 is from the the measured quantities $|\rho_{+-0}| = 0.039^{+0.009}_{-0.006} \pm 0.005$ and $\phi_\rho = (-9 \pm 18)^\circ$.

[21] ADLER 96E is from the measured quantities Re(λ) = $0.036 \pm 0.010^{+0.002}_{-0.003}$ and Im(λ) consistent with zero. Note that the quantity λ is the same as ρ_{+-0} used in other footnotes.

[22] THOMSON 94 calculates this branching ratio from their measurements $|\rho_{+-0}| = 0.035^{+0.019}_{-0.011} \pm 0.004$ and $\phi_\rho = (-59 \pm 48)^\circ$ where $|\rho_{+-0}|e^{i\phi_\rho}$ = A($K^0_S \rightarrow \pi^+\pi^-\pi^0$, $I = 2$)/A($K^0_L \rightarrow \pi^+\pi^-\pi^0$).

Modes with photons or $\ell\overline{\ell}$ pairs

$\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-)$ Γ_4/Γ_2

VALUE (units 10^{-3})	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
2.59±0.08 OUR AVERAGE					
2.56±0.09	1286	RAMBERG	93	E731	p_γ >50 MeV/c
2.68±0.15		[23] TAUREG	76	SPEC	p_γ >50 MeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.10±0.22	3723	RAMBERG	93	E731	p_γ >20 MeV/c
3.0 ±0.6	29	[24] BOBISUT	74	HLBC	p_γ >40 MeV/c
2.8 ±0.6		[25] BURGUN	73	HBC	p_γ >50 MeV/c

[23] TAUREG 76 find direct emission contribution <0.06, CL = 90%.

[24] BOBISUT 74 not included in average because p_γ cut differs. Estimates direct emission contribution to be 0.5 or less, CL = 95%.

[25] BURGUN 73 estimates that direct emission contribution is 0.3 ± 0.6.

$\Gamma(\pi^+\pi^- e^+ e^-)/\Gamma_{total}$ — Γ_5/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID		TECN	COMMENT
4.69±0.30	676	[26] LAI	03C	NA48	1998+1999 data
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.71±0.23±0.22	620	[26,27] LAI	03C	NA48	1999 data
4.5 ±0.7 ±0.4	56	LAI	00B	NA48	1998 data

[26] Uses normalization BR($K_L \to \pi^+\pi^-\pi^0$)*BR($\pi^0 \to e^+e^-$) = $(1.505 \pm 0.047) \times 10^{-3}$ from our 2000 Edition.
[27] Second error is 0.16(syst)±0.15(norm) combined in quadrature.

$\Gamma(\pi^0\gamma\gamma)/\Gamma_{total}$ — Γ_6/Γ

VALUE (units 10^{-8})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
4.9±1.6±0.9		17	[28] LAI	04	NA48	$m^2_{\gamma\gamma}/m^2_K > 0.2$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<33	90		LAI	03B	NA48	$m^2_{\gamma\gamma}/m^2_K > 0.2$

[28] Spectrum also measured and found consistent with the one generated by a constant matrix element.

$\Gamma(\gamma\gamma)/\Gamma_{total}$ — Γ_7/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID		TECN
2.713±0.063±0.005		7.5k	[29] LAI	03	NA48
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.58 ±0.36 ±0.22		149	LAI	00	NA48
2.2 ±1.1		16	[30] BARR	95B	NA31
2.4 ±0.9		35	[31] BARR	95B	NA31
< 13	90		BALATS	89	SPEC
2.4 ±1.2		19	BURKHARDT	87	NA31
<133	90		BARMIN	86B	XEBC

[29] LAI 03 reports [B($K^0_S \to \gamma\gamma$) / B($K^0_S \to \pi^0\pi^0$)] = $(8.84 \pm 0.18 \pm 0.10) \times 10^{-6}$. We multiply by our best value B($K^0_S \to \pi^0\pi^0$) = $(30.69 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
[30] BARR 95B result is calculated using B($K_L \to \gamma\gamma$) = $(5.86 \pm 0.17) \times 10^{-4}$.
[31] BARR 95B quotes this as the combined BARR 95B + BURKHARDT 87 result after rescaling BURKHARDT 87 to use same branching ratios and lifetimes as BARR 95B.

Semileptonic modes

$\Gamma(\pi^\pm e^\mp \nu_e)/\Gamma_{total}$ — Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
7.04±0.09 OUR FIT					
7.04±0.09 OUR AVERAGE					
7.05±0.09	13k	[32] AMBROSINO	06E	KLOE	Not fitted
6.91±0.34±0.15	624	[33] ALOISIO	02	KLOE	Tagged K^0_S using $\phi \to K^0_L K^0_S$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.2 ±1.4	75	AKHMETSHIN	99	CMD2	Tagged K^0_S using $\phi \to K^0_L K^0_S$

[32] Obtained by imposing Σ_i B($K^0_S \rightarrow i$) = 1, where i runs over all the four branching ratios $\pi^+\pi^-$, $\pi^0\pi^0$, $\pi e \nu$, and $\pi \mu \nu$. Input value of B($K^0_S \rightarrow \pi^+\pi^-$) / B($K^0_S \rightarrow \pi^0\pi^0$) from AMBROSINO 06C is used. To derive $\Gamma(K^0_S \rightarrow \pi^+ \mu \nu)$ / $\Gamma(K^0_S \rightarrow \pi^+ e \nu)$, lepton universality is assumed, radiative corrections from ANDRE 04 are used, and phase space integrals are taken from KTeV, ALEXOPOULOS 04A. This branching fraction enters our fit via their $\Gamma(\pi^\pm e^\mp \nu_e)$ / $\Gamma(\pi^+\pi^-)$ branching ratio measurement.

[33] Uses the PDG 00 value for B($K^0_S \rightarrow \pi^+\pi^-$).

$\Gamma(\pi^\pm \mu^\mp \nu_\mu)/\Gamma_{total}$ — Γ_9/Γ

The PDG 06 value below has not been measured but is computed to be 0.666 times the $K_S \rightarrow \pi^\pm e^\mp \nu_e$ branching fraction. It is included in the fit that constrains the four branching ratios $\pi^+\pi^-$, $\pi^0\pi^0$, $\pi e \nu$, and $\pi \mu \nu$ to sum to 1. This treatment, used by AMBROSINO 06E, is preferable to our previous practice of constraining the $\pi^+\pi^-$ and $\pi^0\pi^0$ modes to sum to 1. The 0.666 factor is obtained from AMBROSINO 06E and assumes lepton universality, radiative corrections from ANDRE 04, and phase space integrals from KTeV, ALEXOPOULOS 04A.

VALUE (units 10^{-4})	DOCUMENT ID		COMMENT
4.69 ±0.06 OUR FIT			
4.691±0.001±0.060	[34] PDG	06	calculated from $\pi^\pm e^\mp \nu_e$

[34] The PDG 06 value is computed to be $B_{PDG06}(\pi \mu \nu) = 0.666\ B_{FIT}(\pi e \nu)$. The first error specifies the arbitrarily small error, 0.001×10^{-4}, on $B_{PDG06}(\pi \mu \nu)$ for fixed $B_{FIT}(\pi e \nu)$. The second error is that due to the uncertainty in $B_{FIT}(\pi e \nu)$.

$\Gamma(\pi^\pm e^\mp \nu_e)/\Gamma(\pi^+\pi^-)$ — Γ_8/Γ_2

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN
10.18±0.13 OUR FIT				
10.19±0.11±0.07	13k	AMBROSINO	06E	KLOE

CP violating (*CP*) and $\Delta S = 1$ weak neutral current (*S1*) modes

$\Gamma(3\pi^0)/\Gamma_{total}$ — Γ_{10}/Γ

Violates *CP* conservation.

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID		TECN
< 1.2	90	37.8M	AMBROSINO	05B	KLOE
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 7.4	90	4.9M	[35] LAI	05A	NA48
<140	90	7M	ACHASOV	99D	SND
<190	90	17300	[36] ANGELOPO...	98B	CPLR
<370	90		BARMIN	83	HLBC

[35] LAI 05A value is obtained from their bound on $|\eta_{000}|$ (not assuming *CPT*) and B($K^0_L \rightarrow 3\pi^0$) = 0.211 ± 0.003, and PDG 04 values for K^0_L and K^0_S lifetimes. If *CPT* is assumed then B$(K^0_S \rightarrow 3\pi^0)_{CPT} < 2.3 \times 10^{-7}$ at 90% CL

[36] ANGELOPOULOS 98B is from Im(η_{000}) = −0.05 ± 0.12 ± 0.05, assuming Re(η_{000}) = Re(ϵ) = 1.635×10^{-3} and using the value B($K^0_L \rightarrow \pi^0\pi^0\pi^0$) = 0.2112 ± 0.0027.

$\Gamma(\mu^+\mu^-)/\Gamma_{total}$ — Γ_{11}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN
<0.032	90	GJESDAL	73	ASPK
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.7	90	HYAMS	69B	OSPK

$\Gamma(e^+e^-)/\Gamma_{total}$ — Γ_{12}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
< 1.4	90		ANGELOPO...	97	CPLR	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 28	90	0	BLICK	94	CNTR	Hyperon facility
<100	90		BARMIN	86	XEBC	

$\Gamma(\pi^0 e^+e^-)/\Gamma_{total}$ — Γ_{13}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-9})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
$\mathbf{3.0^{+1.5}_{-1.2}\pm0.2}$		7	[37] BATLEY	03	NA48	m_{ee} >0.165 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 140	90		LAI	01	NA48	
< 1100	90	0	BARR	93B	NA31	
<45000	90		GIBBONS	88	E731	

[37] BATLEY 03 extrapolate also to the full kinematical region using a constant form factor and a vector matrix element. The resulting branching ratio is $(5.8^{+2.9}_{-2.4}) \times 10^{-9}$.

$\Gamma(\pi^0\mu^+\mu^-)/\Gamma_{total}$ — Γ_{14}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-9})	EVTS	DOCUMENT ID		TECN	COMMENT
$\mathbf{2.9^{+1.5}_{-1.2}\pm0.2}$	6	[38] BATLEY	04A	NA48	NA48/1 K^0_S beam

[38] Background estimate is $0.22^{+0.18}_{-0.11}$ events. Branching ratio assumes a vector matrix element and unit form factor.

K^0_S FORM FACTORS

For discussion, see note on $K_{\ell3}$ form factors in the $K^\pm$ section of the Particle Listings above. Because the semileptonic branching fraction is smaller in K^0_S than K^0_L by the ratio of the mean lives, the K^0_S semileptonic form factor has so far been measured only in the K_{e3} mode using the linear expansion $f_+(t) = f_+(0)\ (1 + \lambda_+ t\ /m^2_{\pi^+})$, which gives the vector form factor $f_+(t)$ relative to its value at $t = 0$.

λ_+ (LINEAR ENERGY DEPENDENCE OF f_+ IN K^0_{e3} DECAY)

VALUE (units 10^{-2})	*EVTS*	*DOCUMENT ID*		*TECN*
3.39±0.41	15k	AMBROSINO	06E	KLOE

A REVIEW GOES HERE – Check our WWW List of Reviews

CP-VIOLATION PARAMETERS IN K^0_S DECAY

$A_S = [\ \Gamma(K^0_S \to \pi^- e^+ \nu_e)$ - $\Gamma(K^0_S \to \pi^+ e^- \overline{\nu}_e)\ ]$ / SUM

Such asymmetry violates *CP*. If *CPT* is assumed then $A_S = 2\ \mathrm{Re}(\epsilon)$.

VALUE (units 10^{-3})	*EVTS*	*DOCUMENT ID*		*TECN*
1.5± 9.6±2.9	13k	AMBROSINO	06E	KLOE

——— PARAMETERS FOR $K^0_S \to 3\pi$ DECAY ———

$\mathrm{Im}(\eta_{+-0})^2 = \Gamma(K^0_S \to \pi^+\pi^-\pi^0$, *CP*-violating) / $\Gamma(K^0_L \to \pi^+\pi^-\pi^0)$

CPT assumed valid (i.e. $\mathrm{Re}(\eta_{+-0}) \simeq 0$).

VALUE	*CL%*	*EVTS*	*DOCUMENT ID*		*TECN*
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.23	90	601	[39] BARMIN	85	HLBC
<0.12	90	384	METCALF	72	ASPK

[39] BARMIN 85 find $\mathrm{Re}(\eta_{+-0}) = (0.05 \pm 0.17)$ and $\mathrm{Im}(\eta_{+-0}) = (0.15 \pm 0.33)$. Includes events of BALDO-CEOLIN 75.

$\mathrm{Im}(\eta_{+-0}) = \mathrm{Im}(\mathrm{A}(K^0_S \to \pi^+\pi^-\pi^0$, *CP*-violating) / $\mathrm{A}(K^0_L \to \pi^+\pi^-\pi^0))$

VALUE	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
$\mathbf{-0.002 \pm 0.009^{+0.002}_{-0.001}}$	500k	[40] ADLER	97B	CPLR	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
−0.002±0.018±0.003	137k	[41] ADLER	96D	CPLR	Sup. by ADLER 97B
−0.015±0.017±0.025	272k	[42] ZOU	94	SPEC	

[40] ADLER 97B also find $\mathrm{Re}(\eta_{+-0}) = -0.002 \pm 0.007^{+0.004}_{-0.001}$. See also ANGELOPOULOS 98C.

[41] The ADLER 96D fit also yields $\mathrm{Re}(\eta_{+-0}) = 0.006 \pm 0.013 \pm 0.001$ with a correlation $+0.66$ between real and imaginary parts. Their results correspond to $|\eta_{+-0}| < 0.037$ with 90% CL.

[42] ZOU 94 use theoretical constraint $\mathrm{Re}(\eta_{+-0}) = \mathrm{Re}(\epsilon) = 0.0016$. Without this constraint they find $\mathrm{Im}(\eta_{+-0}) = 0.019 \pm 0.061$ and $\mathrm{Re}(\eta_{+-0}) = 0.019 \pm 0.027$.

$\mathrm{Im}(\eta_{000})^2 = \Gamma(K^0_S \to 3\pi^0)$ / $\Gamma(K^0_L \to 3\pi^0)$

CPT assumed valid (i.e. $\mathrm{Re}(\eta_{000}) \simeq 0$). This limit determines branching ratio $\Gamma(3\pi^0)/\Gamma_{total}$ above.

VALUE	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.1	90	632	[43] BARMIN	83	HLBC	
<0.28	90		[44] GJESDAL	74B	SPEC	Indirect meas.

[43] BARMIN 83 find $\mathrm{Re}(\eta_{000}) = (-0.08 \pm 0.18)$ and $\mathrm{Im}(\eta_{000}) = (-0.05 \pm 0.27)$. Assuming *CPT* invariance they obtain the limit quoted above.

[44] GJESDAL 74B uses $K2\pi$, $K_{\mu 3}$, and K_{e3} decay results, unitarity, and *CPT*. Calculates $|(\eta_{000})| = 0.26 \pm 0.20$. We convert to upper limit.

$\mathrm{Im}(\eta_{000}) = \mathrm{Im}(A(K^0_S \to \pi^0\pi^0\pi^0)/A(K^0_L \to \pi^0\pi^0\pi^0))$

$K^0_S \to \pi^0\pi^0\pi^0$ violates *CP* conservation, in contrast to $K^0_S \to \pi^+\pi^-\pi^0$ which has a *CP*-conserving part.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
$(-0.1 \pm 1.6) \times 10^{-2}$ OUR AVERAGE					
$0.000 \pm 0.009 \pm 0.013$	4.9M	[45] LAI	05A	NA48	Assumes *CPT*
$-0.05 \pm 0.12 \pm 0.05$	17300	[46] ANGELOPO...	98B	CPLR	Assumes *CPT*

[45] LAI 05A assumes $\mathrm{Re}(\eta_{000})=\mathrm{Re}(\epsilon)=1.66 \times 10^{-3}$. The equivalent limit is $|\eta_{000}|_{CPT} < 0.025$ at 90% CL Without assuming *CPT* invariance, they obtain $\mathrm{Re}(\eta_{000})=-0.002 \pm 0.011 \pm 0.015$ and $\mathrm{Im}(\eta_{000})=-0.003 \pm 0.013 \pm 0.017$ with a statistical correlation coefficient of 0.77 and an overall correlation coefficient of 0.57 between imaginary and real part. The equivalent limit is $|\eta_{000}| < 0.045$ at 90% CL

[46] ANGELOPOULOS 98B assumes $\mathrm{Re}(\eta_{000}) = \mathrm{Re}(\epsilon) = 1.635 \times 10^{-3}$. Without assuming *CPT* invariance, they obtain $\mathrm{Re}(\eta_{000}) = 0.18 \pm 0.14 \pm 0.06$ and $\mathrm{Im}(\eta_{000}) = 0.15 \pm 0.20 \pm 0.03$.

$|\eta_{000}| = |A(K^0_S \to 3\pi^0)/A(K^0_L \to 3\pi^0)|$

A non-zero value violates CP invariance.

VALUE	CL%	EVTS	DOCUMENT ID		TECN
<0.018	90	37.8M	AMBROSINO	05B	KLOE
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.045	90	4.9M	LAI	05A	NA48

DECAY-PLANE ASYMMETRY IN $\pi^+\pi^-e^+e^-$ DECAYS

This is the *CP*-violating asymmetry

$$A = \frac{N_{\sin\phi\cos\phi>0.0} - N_{\sin\phi\cos\phi<0.0}}{N_{\sin\phi\cos\phi>0.0} + N_{\sin\phi\cos\phi<0.0}}$$

where ϕ is the angle between the e^+e^- and $\pi^+\pi^-$ planes in the K^0_S rest frame.

CP asymmetry *A* in $K^0_S \to \pi^+\pi^-e^+e^-$

VALUE (%)	DOCUMENT ID		TECN	COMMENT
-1.1 ± 4.1	LAI	03C	NA48	1998+1999 data
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.5 \pm 4.0 \pm 1.6$	LAI	03C	NA48	1999 data

K^0_S REFERENCES

AMBROSINO	06C	EPJ C48 767	F. Ambrosino et al.	(KLOE Collab.)
AMBROSINO	06E	PL B636 173	F. Ambrosino et al.	(KLOE Collab.)
PDG	06	JPG 33 1	W.-M. Yao et al.	(PDG Collab.)
AMBROSINO	05B	PL B619 61	F. Ambrosino et al.	(KLOE Collab.)
BATLEY	05	PL B630 31	J.R. Batley et al.	(NA48 Collab.)
LAI	05A	PL B610 165	A. Lai et al.	(CERN NA48 Collab.)
ALEXOPOU...	04A	PR D70 092007	T. Alexopoulos et al.	(FNAL KTeV Collab.)
ANDRE	04	hep-ph/0406006	T. Andre	(EFI)
BATLEY	04A	PL B599 197	J.R. Batley et al.	(NA48 Collab.)
LAI	04	PL B578 276	A. Lai et al.	(CERN NA48 Collab.)
PDG	04	PL B592 1	S. Eidelman et al.	
ALAVI-HARATI	03	PR D67 012005	A. Alavi-Harati et al.	(FNAL KTeV Collab.)
Also		PR D70 079904 (Errat.)	A. Alavi-Harati et al.	(FNAL KTeV Collab.)
BATLEY	03	PL B576 43	J.R. Batley et al.	(CERN NA48 Collab.)
LAI	03	PL B551 7	A. Lai et al.	(CERN NA48 Collab.)
LAI	03B	PL B556 105	A. Lai et al.	(CERN NA48 Collab.)
LAI	03C	EPJ C30 33	A. Lai et al.	(CERN NA48 Collab.)
ALOISIO	02	PL B535 37	A. Aloisio et al.	(KLOE Collab.)
ALOISIO	02B	PL B538 21	A. Aloisio et al.	(KLOE Collab.)
LAI	02C	PL B537 28	A. Lai et al.	(CERN NA48 Collab.)
LAI	01	PL B514 253	A. Lai et al.	(CERN NA48 Collab.)
LAI	00	PL B493 29	A. Lai et al.	(CERN NA48 Collab.)
LAI	00B	PL B496 137	A. Lai et al.	(CERN NA48 Collab.)
PDG	00	EPJ C15 1	D.E. Groom et al.	
ACHASOV	99D	PL B459 674	M.N. Achasov et al.	
AKHMETSHIN	99	PL B456 90	R.R. Akhmetshin et al.	(Novosibirsk CMD-2 Collab.)
ANGELOPO...	98B	PL B425 391	A. Angelopoulos et al.	(CPLEAR Collab.)
ANGELOPO...	98C	EPJ C5 389	A. Angelopoulos et al.	(CPLEAR Collab.)
PDG	98	EPJ C3 1	C. Caso et al.	
ADLER	97B	PL B407 193	R. Adler et al.	(CPLEAR Collab.)
ANGELOPO...	97	PL B413 232	A. Angelopoulos et al.	(CPLEAR Collab.)
BERTANZA	97	ZPHY C73 629	L. Bertanza	(PISA, CERN, EDIN, MANZ, ORSAY+)
ADLER	96D	PL B370 167	R. Adler et al.	(CPLEAR Collab.)
ADLER	96E	PL B374 313	R. Adler et al.	(CPLEAR Collab.)
ZOU	96	PL B369 362	Y. Zou et al.	(RUTG, MINN, MICH)
BARR	95B	PL B351 579	G.D. Barr et al.	(CERN, EDIN, MANZ, LALO+)
SCHWINGEN...	95	PRL 74 4376	B. Schwingenheuer et al.	(EFI, CHIC+)
BLICK	94	PL B334 234	A.M. Blick et al.	(SERP, JINR)
THOMSON	94	PL B337 411	G.B. Thomson et al.	(RUTG, MINN, MICH)
ZOU	94	PL B329 519	Y. Zou et al.	(RUTG, MINN, MICH)
BARR	93B	PL B304 381	G.D. Barr et al.	(CERN, EDIN, MANZ, LALO+)
GIBBONS	93	PRL 70 1199	L.K. Gibbons et al.	(FNAL E731 Collab.)
Also		PR D55 6625	L.K. Gibbons et al.	(FNAL E731 Collab.)
RAMBERG	93	PRL 70 2525	E. Ramberg et al.	(FNAL E731 Collab.)
BALATS	89	SJNP 49 828	M.Y. Balats et al.	(ITEP)
		Translated from YAF 49 1332.		
GIBBONS	88	PRL 61 2661	L.K. Gibbons et al.	(FNAL E731 Collab.)
BURKHARDT	87	PL B199 139	H. Burkhardt et al.	(CERN, EDIN, MANZ+)
GROSSMAN	87	PRL 59 18	N. Grossman et al.	(MINN, MICH, RUTG)
BARMIN	86	SJNP 44 622	V.V. Barmin et al.	(ITEP)
		Translated from YAF 44 965.		
BARMIN	86B	NC 96A 159	V.V. Barmin et al.	(ITEP, PADO)
PDG	86B	PL 170B 130	M. Aguilar-Benitez et al.	(CERN, CIT+)
BARMIN	85	NC 85A 67	V.V. Barmin et al.	(ITEP, PADO)
Also		SJNP 41 759	V.V. Barmin et al.	(ITEP)
		Translated from YAF 41 1187.		
BARMIN	83	PL 128B 129	V.V. Barmin et al.	(ITEP, PADO)
Also		SJNP 39 269	V.V. Barmin et al.	(ITEP, PADO)
		Translated from YAF 39 428.		
ARONSON	82	PRL 48 1078	S.H. Aronson et al.	(BNL, CHIC, STAN+)
ARONSON	82B	PRL 48 1306	S.H. Aronson et al.	(BNL, CHIC, PURD)
Also		PL 116B 73	E. Fischbach et al.	(PURD, BNL, CHIC)
Also		PR D28 476	S.H. Aronson et al.	(BNL, CHIC, PURD)
Also		PR D28 495	S.H. Aronson et al.	(BNL, CHIC, PURD)
ARONSON	76	NC 32A 236	S.H. Aronson et al.	(WISC, EFI, UCSD+)
EVERHART	76	PR D14 661	G.C. Everhart et al.	(PENN)
TAUREG	76	PL 65B 92	H. Taureg et al.	(HEIDH, CERN, DORT)
BALDO-...	75	NC 25A 688	M. Baldo-Ceolin et al.	(PADO, WISC)
CARITHERS	75	PRL 34 1244	W.C.J. Carithers et al.	(COLU, NYU)

BOBISUT	74	LNC 11 646	F. Bobisut *et al.*	(PADO)
COWELL	74	PR D10 2083	P.L. Cowell *et al.*	(STON, COLU)
GEWENIGER	74B	PL 48B 487	C. Geweniger *et al.*	(CERN, HEIDH)
GJESDAL	74B	PL 52B 119	S. Gjesdal *et al.*	(CERN, HEIDH)
BURGUN	73	PL 46B 481	G. Burgun *et al.*	(SACL, CERN)
GJESDAL	73	PL 44B 217	S. Gjesdal *et al.*	(CERN, HEIDH)
HILL	73	PR D8 1290	D.G. Hill *et al.*	(BNL, CMU)
ALITTI	72	PL 39B 568	J. Alitti, E. Lesquoy, A. Muller	(SACL)
BURGUN	72	NP B50 194	G. Burgun *et al.*	(SACL, CERN, OSLO)
METCALF	72	PL 40B 703	M. Metcalf *et al.*	(CERN, IPN, WIEN)
MORSE	72B	PRL 28 388	R. Morse *et al.*	(COLO, PRIN, UMD)
NAGY	72	NP B47 94	E. Nagy, F. Telbisz, G. Vesztergombi	(BUDA)
Also		PL 30B 498	G. Bozoki *et al.*	(BUDA)
SKJEGGEST...	72	NP B48 343	O. Skjeggestad *et al.*	(OSLO, CERN, SACL)
BALTAY	71	PRL 27 1678	C. Baltay *et al.*	(COLU)
Also		Thesis Nevis 187	W.A. Cooper	(COLU)
MOFFETT	70	BAPS 15 512	R. Moffett *et al.*	(ROCH)
BOZOKI	69	PL 30B 498	G. Bozoki *et al.*	(BUDA)
DOYLE	69	Thesis UCRL 18139	J.C. Doyle	(LRL)
GOBBI	69	PRL 22 682	B. Gobbi *et al.*	(ROCH)
HYAMS	69B	PL 29B 521	B.D. Hyams *et al.*	(CERN, MPIM)
MORFIN	69	PRL 23 660	J.G. Morfin, D. Sinclair	(MICH)
DONALD	68B	PL 27B 58	R.A. Donald *et al.*	(LIVP, CERN, IPNP+)
HILL	68	PR 171 1418	D.G. Hill *et al.*	(BNL, CMU)
AUBERT	65	PL 17 59	B. Aubert *et al.*	(EPOL, ORSAY)
BROWN	63	PR 130 769	J.L. Brown *et al.*	(LRL, MICH)
CHRETIEN	63	PR 131 2208	M. Chretien *et al.*	(BRAN, BROW, HARV+)
BROWN	61	NC 19 1155	J.L. Brown *et al.*	(MICH)
BOLDT	58B	PRL 1 150	E. Boldt, D.O. Caldwell, Y. Pal	(MIT)

OTHER RELATED PAPERS

LITTENBERG	93	ARNPS 43 729	L.S. Littenberg, G. Valencia	(BNL, FNAL)
Rare and Radiative Kaon Decays				
BATTISTON	92	PRPL 214 293	R. Battiston *et al.*	(PGIA, CERN, TRSTT)
Status and Perspectives of K Decay Physics				
TRILLING	65B	UCRL 16473	G.N. Trilling	(LRL)
Updated from 1965 Argonne Conference, page 115.				
CRAWFORD	62	CERN Conf. 827	F.S. Crawford	(LRL)
FITCH	61	NC 22 1160	V.L. Fitch, P.A. Piroue, R.B. Perkins	(PRIN+)
GOOD	61	PR 124 1223	R.H. Good *et al.*	(LRL)
BIRGE	60	Rochester Conf. 601	R.W. Birge *et al.*	(LRL, WISC)
MULLER	60	PRL 4 418	F. Muller *et al.*	(LRL, BNL)