$B^{\pm}$

$I(J^P) = \frac{1}{2}(0^-)$

Quantum numbers not measured. Values shown are quark-model predictions.

See also the $B^{\pm}/B^0$ ADMIXTURE and $B^{\pm}/B^0/B^0_s/b$-baryon ADMIXTURE sections.

$B^{\pm}$ MASS

The fit uses m_{B^+}, $(m_{B^0} - m_{B^+})$, and m_{B^0} to determine m_{B^+}, m_{B^0}, and the mass difference.

VALUE (MeV)	EVTS	DOCUMENT ID		TECN	COMMENT
5279.13±0.31 OUR FIT					
5279.1 ±0.4 OUR AVERAGE					
5279.10±0.41±0.36		[1] ACOSTA	06	CDF	$p\overline{p}$ at 1.96 TeV
5279.1 ±0.4 ±0.4	526	[2] CSORNA	00	CLE2	$e^+e^- \to \Upsilon(4S)$
5279.1 ±1.7 ±1.4	147	ABE	96B	CDF	$p\overline{p}$ at 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
5278.8 ±0.54±2.0	362	ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$
5278.3 ±0.4 ±2.0		BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
5280.5 ±1.0 ±2.0		[3] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$
5275.8 ±1.3 ±3.0	32	ALBRECHT	87C	ARG	$e^+e^- \to \Upsilon(4S)$
5278.2 ±1.8 ±3.0	12	[4] ALBRECHT	87D	ARG	$e^+e^- \to \Upsilon(4S)$
5278.6 ±0.8 ±2.0		BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[1] Uses exclusively reconstructed final states containing a $J/\psi \to \mu^+\mu^-$ decays.

[2] CSORNA 00 uses fully reconstructed 526 $B^+ \to J/\psi^{(\prime)} K^+$ events and invariant masses without beam constraint.

[3] ALBRECHT 90J assumes 10580 for $\Upsilon(4S)$ mass. Supersedes ALBRECHT 87C and ALBRECHT 87D.

[4] Found using fully reconstructed decays with $J/\psi(1S)$. ALBRECHT 87D assume $m_{\Upsilon(4S)}$ = 10577 MeV.

$B^{\pm}$ MEAN LIFE

See $B^{\pm}/B^0/B^0_s/b$-baryon ADMIXTURE section for data on B-hadron mean life averaged over species of bottom particles.

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at http://www.slac.stanford.edu/xorg/hfag/. The averaging/rescaling procedure takes into account correlations between the measurements and asymmetric lifetime errors.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID		TECN	COMMENT
1.638±0.011 OUR EVALUATION					
1.635±0.011±0.011		[5] ABE	05B	BELL	$e^+e^- \to \Upsilon(4S)$
1.624±0.014±0.018		[6] ABDALLAH	04E	DLPH	$e^+e^- \to Z$
1.636±0.058±0.025		[7] ACOSTA	02C	CDF	$p\overline{p}$ at 1.8 TeV
1.673±0.032±0.023		[8] AUBERT	01F	BABR	$e^+e^- \to \Upsilon(4S)$
1.648±0.049±0.035		[9] BARATE	00R	ALEP	$e^+e^- \to Z$

$1.643\pm0.037\pm0.025$		[10] ABBIENDI	99J	OPAL	$e^+e^- \to Z$
$1.637\pm0.058^{+0.045}_{-0.043}$		[9] ABE	98Q	CDF	$p\overline{p}$ at 1.8 TeV
$1.66\pm0.06\pm0.03$		[10] ACCIARRI	98S	L3	$e^+e^- \to Z$
$1.66\pm0.06\pm0.05$		[10] ABE	97J	SLD	$e^+e^- \to Z$
$1.58^{+0.21}_{-0.18}{}^{+0.04}_{-0.03}$	94	[7] BUSKULIC	96J	ALEP	$e^+e^- \to Z$
$1.61\pm0.16\pm0.12$		[9,11] ABREU	95Q	DLPH	$e^+e^- \to Z$
$1.72\pm0.08\pm0.06$		[12] ADAM	95	DLPH	$e^+e^- \to Z$
$1.52\pm0.14\pm0.09$		[9] AKERS	95T	OPAL	$e^+e^- \to Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.695\pm0.026\pm0.015$		[8] ABE	02H	BELL	Repl. by ABE 05B
$1.68\pm0.07\pm0.02$		[7] ABE	98B	CDF	Repl. by ACOSTA 02C
$1.56\pm0.13\pm0.06$		[9] ABE	96C	CDF	Repl. by ABE 98Q
$1.58\pm0.09\pm0.03$		[13] BUSKULIC	96J	ALEP	$e^+e^- \to Z$
$1.58\pm0.09\pm0.04$		[9] BUSKULIC	96J	ALEP	Repl. by BARATE 00R
1.70 ± 0.09		[14] ADAM	95	DLPH	$e^+e^- \to Z$
$1.61\pm0.16\pm0.05$	148	[7] ABE	94D	CDF	Repl. by ABE 98B
$1.30^{+0.33}_{-0.29}\pm0.16$	92	[9] ABREU	93D	DLPH	Sup. by ABREU 95Q
$1.56\pm0.19\pm0.13$	134	[12] ABREU	93G	DLPH	Sup. by ADAM 95
$1.51^{+0.30}_{-0.28}{}^{+0.12}_{-0.14}$	59	[9] ACTON	93C	OPAL	Sup. by AKERS 95T
$1.47^{+0.22}_{-0.19}{}^{+0.15}_{-0.14}$	77	[9] BUSKULIC	93D	ALEP	Sup. by BUSKULIC 96J

[5] Measurement performed using a combined fit of *CP*-violation, mixing and lifetimes.
[6] Measurement performed using an inclusive reconstruction and B flavor identification technique.
[7] Measured mean life using fully reconstructed decays.
[8] Events are selected in which one B meson is fully reconstructed while the second B meson is reconstructed inclusively.
[9] Data analyzed using $D/D^*\ell X$ event vertices.
[10] Data analyzed using charge of secondary vertex.
[11] ABREU 95Q assumes $B(B^0 \to D^{**-}\ell^+\nu_\ell) = 3.2 \pm 1.7\%$.
[12] Data analyzed using vertex-charge technique to tag B charge.
[13] Combined result of $D/D^*\ell X$ analysis and fully reconstructed B analysis.
[14] Combined ABREU 95Q and ADAM 95 result.

B^+ DECAY MODES

B^- modes are charge conjugates of the modes below. Modes which do not identify the charge state of the B are listed in the $B^\pm/B^0$ ADMIXTURE section.

The branching fractions listed below assume 50% $B^0\overline{B}^0$ and 50% B^+B^- production at the $\Upsilon(4S)$. We have attempted to bring older measurements up to date by rescaling their assumed $\Upsilon(4S)$ production ratio to 50:50 and their assumed D, D_s, D^*, and ψ branching ratios to current values whenever this would affect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

For inclusive branching fractions, *e.g.*, $B \to D^{\pm}$ anything, the values usually are multiplicities, not branching fractions. They can be greater than one.

	Mode		Fraction (Γ_i/Γ)	Scale factor/ Confidence level
	Semileptonic and leptonic modes			
Γ_1	$\ell^+ \nu_\ell$ anything	[a]	(10.99 ± 0.28) %	
Γ_2	$e^+ \nu_e X_c$		(10.8 ± 0.4) %	
Γ_3	$\overline{D}^0 \ell^+ \nu_\ell$	[a]	(2.15 ± 0.22) %	
Γ_4	$\overline{D}^*(2007)^0 \ell^+ \nu_\ell$	[a]	(6.5 ± 0.5) %	
Γ_5	$\overline{D}_1(2420)^0 \ell^+ \nu_\ell$		$(5.6 \pm 1.6) \times 10^{-3}$	
Γ_6	$\overline{D}_2^*(2460)^0 \ell^+ \nu_\ell$		$< 8 \times 10^{-3}$	CL=90%
Γ_7	$D^- \pi^+ \ell^+ \nu_\ell$		$(5.2 \pm 1.0) \times 10^{-3}$	
Γ_8	$D^{*-} \pi^+ \ell^+ \nu_\ell$		$(6.3 \pm 1.5) \times 10^{-3}$	
Γ_9	$\pi^0 \ell^+ \nu_\ell$		$(7.7 \pm 1.2) \times 10^{-5}$	
Γ_{10}	$\pi^0 e^+ \nu_e$			
Γ_{11}	$\eta \ell^+ \nu_\ell$		$(8 \pm 4) \times 10^{-5}$	
Γ_{12}	$\omega \ell^+ \nu_\ell$	[a]	$(1.3 \pm 0.6) \times 10^{-4}$	
Γ_{13}	$\omega \mu^+ \nu_\mu$			
Γ_{14}	$\rho^0 \ell^+ \nu_\ell$	[a]	$(1.28 \pm 0.18) \times 10^{-4}$	
Γ_{15}	$p \overline{p} e^+ \nu_e$		$< 5.2 \times 10^{-3}$	CL=90%
Γ_{16}	$e^+ \nu_e$		$< 9.8 \times 10^{-6}$	CL=90%
Γ_{17}	$\mu^+ \nu_\mu$		$< 1.7 \times 10^{-6}$	CL=90%
Γ_{18}	$\tau^+ \nu_\tau$		$(1.8 \pm 0.7) \times 10^{-4}$	
Γ_{19}	$e^+ \nu_e \gamma$		$< 2.0 \times 10^{-4}$	CL=90%
Γ_{20}	$\mu^+ \nu_\mu \gamma$		$< 5.2 \times 10^{-5}$	CL=90%
	Inclusive modes			
Γ_{21}	$D^0 X$		(8.6 ± 0.7) %	
Γ_{22}	$\overline{D}^0 X$		(79 ± 4) %	
Γ_{23}	$D^+ X$		(2.5 ± 0.5) %	
Γ_{24}	$D^- X$		(9.9 ± 1.2) %	
Γ_{25}	$D_s^+ X$		$(7.9 ^{+1.4}_{-1.3})$ %	
Γ_{26}	$D_s^- X$		$(1.10 ^{+0.45}_{-0.32})$ %	
Γ_{27}	$\Lambda_c^+ X$		$(2.1 ^{+0.9}_{-0.6})$ %	
Γ_{28}	$\overline{\Lambda}_c^- X$		$(2.8 ^{+1.1}_{-0.9})$ %	

Γ_{29}	$\overline{c}X$		(97	± 4	) %
Γ_{30}	cX		(23.4	$^{+2.2}_{-1.8}$	) %
Γ_{31}	$\overline{c}cX$		(120	± 6	) %

D, D^*, or D_s modes

Γ_{32}	$\overline{D}^0\pi^+$		(4.84	± 0.15	) $\times 10^{-3}$	
Γ_{33}	$D_{CP(+1)}\pi^+$	[b]	(4.0	± 0.8	) $\times 10^{-3}$	
Γ_{34}	$D_{CP(-1)}\pi^+$	[b]	(3.6	± 0.8	) $\times 10^{-3}$	
Γ_{35}	$\overline{D}^0\rho^+$		(1.34	± 0.18	) %	
Γ_{36}	$\overline{D}^0K^+$		(4.02	± 0.21	) $\times 10^{-4}$	
Γ_{37}	$D_{CP(+1)}K^+$	[b]	(3.6	± 0.5	) $\times 10^{-4}$	
Γ_{38}	$D_{CP(-1)}K^+$	[b]	(3.5	± 0.5	) $\times 10^{-4}$	
Γ_{39}	$[K^-\pi^+]_DK^+$	[c]				
Γ_{40}	$[K^+\pi^-]_DK^+$	[c]				
Γ_{41}	$[K^-\pi^+]_DK^*(892)^+$	[c]				
Γ_{42}	$[K^+\pi^-]_DK^*(892)^+$	[c]				
Γ_{43}	$[K^-\pi^+]_D\pi^+$	[c]	(1.7	± 0.5	) $\times 10^{-5}$	
Γ_{44}	$[\pi^+\pi^-\pi^0]_DK^-$		(5.5	± 1.2	) $\times 10^{-6}$	
Γ_{45}	$\overline{D}^0K^*(892)^+$		(5.3	± 0.4	) $\times 10^{-4}$	
Γ_{46}	$D_{CP(-1)}K^*(892)^+$	[b]	(1.7	± 0.7	) $\times 10^{-4}$	
Γ_{47}	$D_{CP(+1)}K^*(892)^+$	[b]	(5.2	± 1.2	) $\times 10^{-4}$	
Γ_{48}	$\overline{D}^0K^+\overline{K}^0$		(5.5	± 1.6	) $\times 10^{-4}$	
Γ_{49}	$\overline{D}^0K^+\overline{K}^*(892)^0$		(7.5	± 1.7	) $\times 10^{-4}$	
Γ_{50}	$\overline{D}^0\pi^+\pi^+\pi^-$		(1.1	± 0.4	) %	
Γ_{51}	$\overline{D}^0\pi^+\pi^+\pi^-$ nonresonant		(5	± 4	) $\times 10^{-3}$	
Γ_{52}	$\overline{D}^0\pi^+\rho^0$		(4.2	± 3.0	) $\times 10^{-3}$	
Γ_{53}	$\overline{D}^0a_1(1260)^+$		(4	± 4	) $\times 10^{-3}$	
Γ_{54}	$\overline{D}^0\omega\pi^+$		(4.1	± 0.9	) $\times 10^{-3}$	
Γ_{55}	$D^*(2010)^-\pi^+\pi^+$		(1.35	± 0.22	) $\times 10^{-3}$	
Γ_{56}	$D^-\pi^+\pi^+$		(1.02	± 0.16	) $\times 10^{-3}$	
Γ_{57}	D^+K^0		< 5.0		$\times 10^{-6}$	CL=90%
Γ_{58}	$\overline{D}^*(2007)^0\pi^+$		(5.19	± 0.26	) $\times 10^{-3}$	
Γ_{59}	$\overline{D}^{*0}_{CP(+1)}\pi^+$	[d]				
Γ_{60}	$D^{*0}_{CP(-1)}\pi^+$	[d]				
Γ_{61}	$\overline{D}^*(2007)^0\omega\pi^+$		(4.5	± 1.2	) $\times 10^{-3}$	
Γ_{62}	$\overline{D}^*(2007)^0\rho^+$		(9.8	± 1.7	) $\times 10^{-3}$	
Γ_{63}	$\overline{D}^*(2007)^0K^+$		(4.16	± 0.33	) $\times 10^{-4}$	
Γ_{64}	$\overline{D}^{*0}_{CP(+1)}K^+$	[d]				
Γ_{65}	$\overline{D}^{*0}_{CP(-1)}K^+$	[d]				
Γ_{66}	$\overline{D}^*(2007)^0K^*(892)^+$		(8.1	± 1.4	) $\times 10^{-4}$	
Γ_{67}	$\overline{D}^*(2007)^0K^+\overline{K}^0$		< 1.06		$\times 10^{-3}$	CL=90%
Γ_{68}	$\overline{D}^*(2007)^0K^+K^*(892)^0$		(1.5	± 0.4	) $\times 10^{-3}$	

Γ_{69}	$\overline{D}^*(2007)^0\pi^+\pi^+\pi^-$		$(1.03\pm0.12)\ \%$	
Γ_{70}	$\quad\overline{D}^*(2007)^0 a_1(1260)^+$		$(1.9\pm0.5)\ \%$	
Γ_{71}	$\overline{D}^*(2007)^0\pi^-\pi^+\pi^+\pi^0$		$(1.8\pm0.4)\ \%$	
Γ_{72}	$\overline{D}^{*0}3\pi^+2\pi^-$		$(5.7\pm1.2)\times10^{-3}$	
Γ_{73}	$D^*(2010)^+\pi^0$		$<1.7\times10^{-4}$	CL=90%
Γ_{74}	$D^*(2010)^+K^0$		$<9.0\times10^{-6}$	CL=90%
Γ_{75}	$D^*(2010)^-\pi^+\pi^+\pi^0$		$(1.5\pm0.7)\ \%$	
Γ_{76}	$D^*(2010)^-\pi^+\pi^+\pi^+\pi^-$		$(2.6\pm0.4)\times10^{-3}$	
Γ_{77}	$\overline{D}^{**0}\pi^+$	[e]	$(5.9\pm1.3)\times10^{-3}$	
Γ_{78}	$\overline{D}^*_1(2420)^0\pi^+$		$(1.5\pm0.6)\times10^{-3}$	S=1.3
Γ_{79}	$\overline{D}_1(2420)^0\pi^+\times\mathrm{B}(\overline{D}^0_1\to\overline{D}^0\pi^+\pi^-)$		$(1.9^{+0.5}_{-0.6})\times10^{-4}$	
Γ_{80}	$\overline{D}^*_2(2462)^0\pi^+\times\mathrm{B}(\overline{D}^*_2(2462)^0\to D^-\pi^+)$		$(3.4\pm0.8)\times10^{-4}$	
Γ_{81}	$\overline{D}^*_0(2400)^0\pi^+\times\mathrm{B}(\overline{D}^*_0(2400)^0\to D^-\pi^+)$		$(6.1\pm1.9)\times10^{-4}$	
Γ_{82}	$\overline{D}_1(2421)^0\pi^+\times\mathrm{B}(\overline{D}_1(2421)^0\to D^{*-}\pi^+)$		$(6.8\pm1.5)\times10^{-4}$	
Γ_{83}	$\overline{D}^*_2(2462)^0\pi^+\times\mathrm{B}(\overline{D}^*_2(2462)^0\to D^{*-}\pi^+)$		$(1.8\pm0.5)\times10^{-4}$	
Γ_{84}	$\overline{D}'_1(2427)^0\pi^+\times\mathrm{B}(\overline{D}'_1(2427)^0\to D^{*-}\pi^+)$		$(5.0\pm1.2)\times10^{-4}$	
Γ_{85}	$\overline{D}_1(2420)^0\pi^+\times\mathrm{B}(\overline{D}^0_1\to\overline{D}^{*0}\pi^+\pi^-)$		$<6\times10^{-6}$	CL=90%
Γ_{86}	$\overline{D}^*_1(2420)^0\rho^+$		$<1.4\times10^{-3}$	CL=90%
Γ_{87}	$\overline{D}^*_2(2460)^0\pi^+$		$<1.3\times10^{-3}$	CL=90%
Γ_{88}	$\overline{D}^*_2(2460)^0\pi^+\times\mathrm{B}(\overline{D}^{*0}_2\to\overline{D}^{*0}\pi^+\pi^-)$		$<2.2\times10^{-5}$	CL=90%
Γ_{89}	$\overline{D}^*_2(2460)^0\rho^+$		$<4.7\times10^{-3}$	CL=90%
Γ_{90}	$\overline{D}^0 D^+_s$		$(10.0\pm1.7)\times10^{-3}$	
Γ_{91}	$D_{s0}(2317)^+\overline{D}^0\times\mathrm{B}(D_{s0}(2317)^+\to D^+_s\pi^0)$		$(7.3^{+2.2}_{-1.7})\times10^{-4}$	
Γ_{92}	$D_{s0}(2317)^+\overline{D}^0\times\mathrm{B}(D_{s0}(2317)^+\to D^{*+}_s\gamma)$		$<7.6\times10^{-4}$	CL=90%
Γ_{93}	$D_{s0}(2317)^+\overline{D}^*(2010)^0\times\mathrm{B}(D_{s0}(2317)^+\to D^+_s\pi^0)$		$(9\pm7)\times10^{-4}$	
Γ_{94}	$D_{sJ}(2457)^+\overline{D}^0$		$(3.1^{+1.0}_{-0.9})\times10^{-3}$	
Γ_{95}	$D_{sJ}(2457)^+\overline{D}^0\times\mathrm{B}(D_{sJ}(2457)^+\to D^+_s\gamma)$		$(4.6^{+1.3}_{-1.1})\times10^{-4}$	

Γ_{96}	$D_{sJ}(2457)^+ \overline{D}^0 \times$ B$(D_{sJ}(2457)^+ \to D_s^+ \pi^+ \pi^-)$	<	2.2		$\times 10^{-4}$	CL=90%
Γ_{97}	$D_{sJ}(2457)^+ \overline{D}^0 \times$ B$(D_{sJ}(2457)^+ \to D_s^+ \pi^0)$	<	2.7		$\times 10^{-4}$	CL=90%
Γ_{98}	$D_{sJ}(2457)^+ \overline{D}^0 \times$ B$(D_{sJ}(2457)^+ \to D_s^{*+} \gamma)$	<	9.8		$\times 10^{-4}$	CL=90%
Γ_{99}	$D_{sJ}(2457)^+ \overline{D}^*(2007)^0$	(	1.20	± 0.30	) %	
Γ_{100}	$D_{sJ}(2457)^+ \overline{D}^*(2007)^0 \times$ B$(D_{sJ}(2457)^+ \to D_s^+ \gamma)$	(	1.4	${}^{+0.7}_{-0.6}$	) $\times 10^{-3}$	
Γ_{101}	$\overline{D}^0 D_{sJ}(2536)^+ \times$ B$(D_{sJ}(2536)^+ \to D^*(2007)^0 K^+)$	<	2		$\times 10^{-4}$	CL=90%
Γ_{102}	$\overline{D}^*(2007)^0 D_{sJ}(2536)^+ \times$ B$(D_{sJ}(2536)^+ \to D^*(2007)^0 K^+)$	<	7		$\times 10^{-4}$	CL=90%
Γ_{103}	$\overline{D}^0 D_{sJ}(2573)^+ \times$ B$(D_{sJ}(2573)^+ \to D^0 K^+)$	<	2		$\times 10^{-4}$	CL=90%
Γ_{104}	$\overline{D}^*(2007)^0 D_{sJ}(2573)^+ \times$ B$(D_{sJ}(2573)^+ \to D^0 K^+)$	<	5		$\times 10^{-4}$	CL=90%
Γ_{105}	$\overline{D}^0 D_s^{*+}$	(	7.6	± 1.6	) $\times 10^{-3}$	
Γ_{106}	$\overline{D}^*(2007)^0 D_s^+$	(	8.2	± 1.7	) $\times 10^{-3}$	
Γ_{107}	$\overline{D}^*(2007)^0 D_s^{*+}$	(	1.71	± 0.24	) %	
Γ_{108}	$D_s^{(*)+} \overline{D}^{**0}$	(	2.7	± 1.2	) %	
Γ_{109}	$\overline{D}^*(2007)^0 D^*(2010)^+$	(	8.1	± 1.7	) $\times 10^{-4}$	
Γ_{110}	$\overline{D}^0 D^*(2010)^+ + \overline{D}^*(2007)^0 D^+$	<	1.30		%	CL=90%
Γ_{111}	$\overline{D}^0 D^*(2010)^+$	(	3.9	± 0.5	) $\times 10^{-4}$	
Γ_{112}	$\overline{D}^0 D^+$	(	4.2	± 0.6	) $\times 10^{-4}$	
Γ_{113}	$\overline{D}^0 D^+ K^0$	<	2.8		$\times 10^{-3}$	CL=90%
Γ_{114}	$D^+ \overline{D}^*(2007)^0$	(	6.3	± 1.7	) $\times 10^{-4}$	
Γ_{115}	$\overline{D}^*(2007)^0 D^+ K^0$	<	6.1		$\times 10^{-3}$	CL=90%
Γ_{116}	$\overline{D}^0 \overline{D}^*(2010)^+ K^0$	(	5.2	± 1.2	) $\times 10^{-3}$	
Γ_{117}	$\overline{D}^*(2007)^0 D^*(2010)^+ K^0$	(	7.8	± 2.6	) $\times 10^{-3}$	
Γ_{118}	$\overline{D}^0 D^0 K^+$	(	1.37	± 0.32	) $\times 10^{-3}$	S=1.5
Γ_{119}	$\overline{D}^*(2007)^0 D^0 K^+$	<	3.8		$\times 10^{-3}$	CL=90%
Γ_{120}	$\overline{D}^0 D^*(2007)^0 K^+$	(	4.7	± 1.0	) $\times 10^{-3}$	
Γ_{121}	$\overline{D}^*(2007)^0 D^*(2007)^0 K^+$	(	5.3	± 1.6	) $\times 10^{-3}$	
Γ_{122}	$D^- D^+ K^+$	<	4		$\times 10^{-4}$	CL=90%
Γ_{123}	$D^- D^*(2010)^+ K^+$	<	7		$\times 10^{-4}$	CL=90%
Γ_{124}	$D^*(2010)^- D^+ K^+$	(	1.5	± 0.4	) $\times 10^{-3}$	
Γ_{125}	$D^*(2010)^- D^*(2010)^+ K^+$	<	1.8		$\times 10^{-3}$	CL=90%

Γ_{126}	$(\overline{D}+\overline{D}^*)(D+D^*)K$	(3.5 ± 0.6) %			
Γ_{127}	$D_s^+\pi^0$	(1.6 ± 0.5) $\times 10^{-5}$			
Γ_{128}	$D_s^{*+}\pi^0$	$<$ 2.6	$\times 10^{-4}$	CL=90%	
Γ_{129}	$D_s^+\eta$	$<$ 4	$\times 10^{-4}$	CL=90%	
Γ_{130}	$D_s^{*+}\eta$	$<$ 6	$\times 10^{-4}$	CL=90%	
Γ_{131}	$D_s^+\rho^0$	$<$ 3.0	$\times 10^{-4}$	CL=90%	
Γ_{132}	$D_s^{*+}\rho^0$	$<$ 4	$\times 10^{-4}$	CL=90%	
Γ_{133}	$D_s^+\omega$	$<$ 4	$\times 10^{-4}$	CL=90%	
Γ_{134}	$D_s^{*+}\omega$	$<$ 6	$\times 10^{-4}$	CL=90%	
Γ_{135}	$D_s^+ a_1(1260)^0$	$<$ 1.8	$\times 10^{-3}$	CL=90%	
Γ_{136}	$D_s^{*+} a_1(1260)^0$	$<$ 1.3	$\times 10^{-3}$	CL=90%	
Γ_{137}	$D_s^+\phi$	$<$ 1.9	$\times 10^{-6}$	CL=90%	
Γ_{138}	$D_s^{*+}\phi$	$<$ 1.2	$\times 10^{-5}$	CL=90%	
Γ_{139}	$D_s^+\overline{K}^0$	$<$ 8	$\times 10^{-4}$	CL=90%	
Γ_{140}	$D_s^{*+}\overline{K}^0$	$<$ 9	$\times 10^{-4}$	CL=90%	
Γ_{141}	$D_s^+\overline{K}^*(892)^0$	$<$ 4	$\times 10^{-4}$	CL=90%	
Γ_{142}	$D_s^{*+}\overline{K}^*(892)^0$	$<$ 3.5	$\times 10^{-4}$	CL=90%	
Γ_{143}	$D_s^-\pi^+K^+$	$<$ 7	$\times 10^{-4}$	CL=90%	
Γ_{144}	$D_s^{*-}\pi^+K^+$	$<$ 9.6	$\times 10^{-4}$	CL=90%	
Γ_{145}	$D_s^-\pi^+K^*(892)^+$	$<$ 5	$\times 10^{-3}$	CL=90%	
Γ_{146}	$D_s^{*-}\pi^+K^*(892)^+$	$<$ 7	$\times 10^{-3}$	CL=90%	

Charmonium modes

Γ_{147}	$\eta_c K^+$	(9.1 ± 1.3) $\times 10^{-4}$			
Γ_{148}	$\eta_c' K^+$	(3.4 ± 1.8) $\times 10^{-4}$			
Γ_{149}	$J/\psi(1S)K^+$	(1.007 ± 0.035) $\times 10^{-3}$			
Γ_{150}	$J/\psi(1S)K^+\pi^+\pi^-$	(1.07 ± 0.19) $\times 10^{-3}$			S=1.9
Γ_{151}	$h_c(1P)K^+ \times$ B($h_c(1P) \to J/\psi\pi^+\pi^-$)	$<$ 3.4	$\times 10^{-6}$	CL=90%	
Γ_{152}	$X(3872)K^+$	$<$ 3.2	$\times 10^{-4}$	CL=90%	
Γ_{153}	$X(3872)K^+ \times$ B($X \to J/\psi\pi^+\pi^-$)	(1.14 ± 0.20) $\times 10^{-5}$			
Γ_{154}	$X(3872)K^+ \times$ B($X \to J/\psi\gamma$)	(3.3 ± 1.0) $\times 10^{-6}$			
Γ_{155}	$X(3872)K^+ \times$ B($X(3872) \to D^0\overline{D}^0$)	$<$ 6.0	$\times 10^{-5}$	CL=90%	
Γ_{156}	$X(3872)K^+ \times$ B($X(3872) \to D^+D^-$)	$<$ 4.0	$\times 10^{-5}$	CL=90%	
Γ_{157}	$X(3872)K^+ \times$ B($X(3872) \to D^0\overline{D}^0\pi^0$)	(1.0 ± 0.4) $\times 10^{-4}$			
Γ_{158}	$X(3872)K^+ \times$ B($X(3872) \to J/\psi(1S)\eta$)	$<$ 7.7	$\times 10^{-6}$	CL=90%	

Γ_{159}	$X(3872)^+ K^0 \times$ B$(X(3872)^+ \to J/\psi(1S)\pi^+\pi^0)$	[f] $<$	2.2		$\times 10^{-5}$	CL=90%
Γ_{160}	$X(4260)^0 K^+ \times$ B$(X^0 \to J/\psi\pi^+\pi^-)$	$<$	2.9		$\times 10^{-5}$	CL=95%
Γ_{161}	$X(3945)^0 K^+ \times$ B$(X^0 \to J/\psi\gamma)$	$<$	1.4		$\times 10^{-5}$	CL=90%
Γ_{162}	$Z(3930)^0 K^+ \times$ B$(Z^0 \to J/\psi\gamma)$	$<$	2.5		$\times 10^{-6}$	CL=90%
Γ_{163}	$J/\psi(1S) K^*(892)^+$	(	1.41	± 0.08	$) \times 10^{-3}$	
Γ_{164}	$J/\psi(1S) K(1270)^+$	(	1.8	± 0.5	$) \times 10^{-3}$	
Γ_{165}	$J/\psi(1S) K(1400)^+$	$<$	5		$\times 10^{-4}$	CL=90%
Γ_{166}	$J/\psi(1S)\eta K^+$	(	1.08	± 0.33	$) \times 10^{-4}$	
Γ_{167}	$J/\psi(1S)\phi K^+$	(	5.2	± 1.7	$) \times 10^{-5}$	S=1.2
Γ_{168}	$J/\psi(1S)\pi^+$	(	4.9	± 0.6	$) \times 10^{-5}$	S=1.5
Γ_{169}	$J/\psi(1S)\rho^+$	$<$	7.7		$\times 10^{-4}$	CL=90%
Γ_{170}	$J/\psi(1S) a_1(1260)^+$	$<$	1.2		$\times 10^{-3}$	CL=90%
Γ_{171}	$J/\psi(1S) p\overline{\Lambda}$	(	1.18	± 0.31	$) \times 10^{-5}$	
Γ_{172}	$J/\psi(1S)\overline{\Sigma}^0 p$	$<$	1.1		$\times 10^{-5}$	CL=90%
Γ_{173}	$J/\psi(1S) D^+$	$<$	1.2		$\times 10^{-4}$	CL=90%
Γ_{174}	$J/\psi(1S)\overline{D}^0\pi^+$	$<$	2.5		$\times 10^{-5}$	CL=90%
Γ_{175}	$\psi(2S) K^+$	(	6.48	± 0.35	$) \times 10^{-4}$	
Γ_{176}	$\psi(2S) K^*(892)^+$	(	6.7	± 1.4	$) \times 10^{-4}$	S=1.3
Γ_{177}	$\psi(2S) K^+\pi^+\pi^-$	(	1.9	± 1.2	$) \times 10^{-3}$	
Γ_{178}	$\psi(3770) K^+$	(	4.9	± 1.3	$) \times 10^{-4}$	
Γ_{179}	$\psi(3770) K^+ \times$ B$(\psi(3770) \to D^0\overline{D}^0)$	(	3.4	± 0.9	$) \times 10^{-4}$	
Γ_{180}	$\psi(3770) K^+ \times$ B$(\psi(3770) \to D^+ D^- K^+)$	(	1.4	± 0.8	$) \times 10^{-4}$	
Γ_{181}	$\chi_{c0}\pi^+ \times$B$(\chi_{c0} \to \pi^+\pi^-)$	$<$	3		$\times 10^{-7}$	CL=90%
Γ_{182}	$\chi_{c0}(1P) K^+$	(	1.40	$^{+0.23}_{-0.19}$	$) \times 10^{-4}$	
Γ_{183}	$\chi_{c0} K^*(892)^+$	$<$	2.86		$\times 10^{-3}$	CL=90%
Γ_{184}	$\chi_{c2} K^+$	$<$	2.9		$\times 10^{-5}$	CL=90%
Γ_{185}	$\chi_{c2} K^*(892)^+$	$<$	1.2		$\times 10^{-5}$	CL=90%
Γ_{186}	$\chi_{c1}(1P)\pi^+$	(	2.2	± 0.5	$) \times 10^{-5}$	
Γ_{187}	$\chi_{c1}(1P) K^+$	(	4.9	± 0.5	$) \times 10^{-4}$	S=1.5
Γ_{188}	$\chi_{c1}(1P) K^*(892)^+$	(	3.6	± 0.9	$) \times 10^{-4}$	
Γ_{189}	$h_c K^+$	$<$	3.8		$\times 10^{-5}$	

K or *K** modes

Γ_{190}	$K^0\pi^+$	(	2.30	± 0.12	$) \times 10^{-5}$	
Γ_{191}	$K^+\pi^0$	(	1.21	± 0.08	$) \times 10^{-5}$	
Γ_{192}	$\eta' K^+$	(	6.97	± 0.28	$) \times 10^{-5}$	
Γ_{193}	$\eta' K^*(892)^+$	(	4.9	± 2.0	$) \times 10^{-6}$	
Γ_{194}	ηK^+	(	2.6	± 0.6	$) \times 10^{-6}$	S=1.3
Γ_{195}	$\eta K^*(892)^+$	(	1.93	± 0.22	$) \times 10^{-5}$	

Γ_{196}	$\eta K_0^*(1430)^+$		$(1.8 \pm 0.4) \times 10^{-5}$		
Γ_{197}	$\eta K_2^*(1430)^+$		$(9.1 \pm 3.0) \times 10^{-6}$		
Γ_{198}	ωK^+		$(6.8 \pm 0.9) \times 10^{-6}$	S=1.6	
Γ_{199}	$\omega K^*(892)^+$	$<$	3.4×10^{-6}	CL=90%	
Γ_{200}	$a_0^+ K^0$	$<$	3.9×10^{-6}	CL=90%	
Γ_{201}	$a_0^0 K^+$	$<$	2.5×10^{-6}	CL=90%	
Γ_{202}	$K^*(892)^0 \pi^+$		$(1.09 \pm 0.18) \times 10^{-5}$	S=2.1	
Γ_{203}	$K^*(892)^+ \pi^0$		$(6.9 \pm 2.4) \times 10^{-6}$		
Γ_{204}	$K^+ \pi^- \pi^+$		$(5.5 \pm 0.7) \times 10^{-5}$	S=2.6	
Γ_{205}	$K^+ \pi^- \pi^+$ nonresonant		$(6 {}^{+6}_{-4}) \times 10^{-6}$	S=6.1	
Γ_{206}	$K^+ f_0(980) \times$ B$(f_0 \to \pi^+ \pi^-)$		$(9.2 {}^{+0.8}_{-1.1}) \times 10^{-6}$		
Γ_{207}	$f_2(1270)^0 K^+$		$(1.3 {}^{+0.4}_{-0.5}) \times 10^{-6}$		
Γ_{208}	$f_0(1370)^0 K^+ \times$ B$(f_0(1370)^0 \to \pi^+ \pi^-)$	$<$	1.07×10^{-5}	CL=90%	
Γ_{209}	$\rho^0(1450) K^+ \times$ B$(\rho^0(1450) \to \pi^+ \pi^-)$	$<$	1.17×10^{-5}	CL=90%	
Γ_{210}	$f_0(1500) K^+ \times$ B$(f_0(1500) \to \pi^+ \pi^-)$	$<$	4.4×10^{-6}	CL=90%	
Γ_{211}	$f_2'(1525) K^+ \times$ B$(f_2'(1525) \to \pi^+ \pi^-)$	$<$	3.4×10^{-6}	CL=90%	
Γ_{212}	$K^+ \rho^0$		$(4.2 \pm 0.5) \times 10^{-6}$		
Γ_{213}	$K_0^*(1430)^0 \pi^+$		$(4.7 \pm 0.5) \times 10^{-5}$		
Γ_{214}	$K_2^*(1430)^0 \pi^+$	$<$	6.9×10^{-6}	CL=90%	
Γ_{215}	$K^*(1410)^0 \pi^+$	$<$	4.5×10^{-5}	CL=90%	
Γ_{216}	$K^*(1680)^0 \pi^+$	$<$	1.2×10^{-5}	CL=90%	
Γ_{217}	$K^- \pi^+ \pi^+$	$<$	1.8×10^{-6}	CL=90%	
Γ_{218}	$K^- \pi^+ \pi^+$ nonresonant	$<$	5.6×10^{-5}	CL=90%	
Γ_{219}	$K_1(1400)^0 \pi^+$	$<$	2.6×10^{-3}	CL=90%	
Γ_{220}	$K^0 \pi^+ \pi^0$	$<$	6.6×10^{-5}	CL=90%	
Γ_{221}	$K^0 \rho^+$	$<$	4.8×10^{-5}	CL=90%	
Γ_{222}	$K^*(892)^+ \pi^+ \pi^-$		$(7.5 \pm 1.0) \times 10^{-5}$		
Γ_{223}	$K^*(892)^+ \rho^0$	$<$	6.1×10^{-6}	CL=90%	
Γ_{224}	$K^*(892)^+ f_0(980)$		$(5.2 \pm 1.3) \times 10^{-6}$		
Γ_{225}	$K^*(892)^0 \rho^+$		$(9.2 \pm 1.5) \times 10^{-6}$		
Γ_{226}	$K^*(892)^+ K^*(892)^0$	$<$	7.1×10^{-5}	CL=90%	
Γ_{227}	$K_1(1400)^+ \rho^0$	$<$	7.8×10^{-4}	CL=90%	
Γ_{228}	$K_2^*(1430)^+ \rho^0$	$<$	1.5×10^{-3}	CL=90%	
Γ_{229}	$K^+ \overline{K}^0$		$(1.28 \pm 0.30) \times 10^{-6}$		
Γ_{230}	$\overline{K}^0 K^+ \pi^0$	$<$	2.4×10^{-5}	CL=90%	
Γ_{231}	$K^+ K_S^0 K_S^0$		$(1.15 \pm 0.13) \times 10^{-5}$		
Γ_{232}	$K_S^0 K_S^0 \pi^+$	$<$	3.2×10^{-6}	CL=90%	

Γ_{233}	$K^+ K^- \pi^+$	<	6.3		$\times 10^{-6}$	CL=90%
Γ_{234}	$K^+ K^- \pi^+$ nonresonant	<	7.5		$\times 10^{-5}$	CL=90%
Γ_{235}	$K^+ K^+ \pi^-$	<	1.3		$\times 10^{-6}$	CL=90%
Γ_{236}	$K^+ K^+ \pi^-$ nonresonant	<	8.79		$\times 10^{-5}$	CL=90%
Γ_{237}	$K^+ K^*(892)^0$	<	5.3		$\times 10^{-6}$	CL=90%
Γ_{238}	$K^+ f_J(2220)$					
Γ_{239}	$K^{*+} \pi^+ K^-$	<	1.18		$\times 10^{-5}$	CL=90%
Γ_{240}	$K^{*+} K^+ \pi^-$	<	6.1		$\times 10^{-6}$	CL=90%
Γ_{241}	$K^+ K^- K^+$	(	3.37	± 0.22	) $\times 10^{-5}$	S=1.4
Γ_{242}	$K^+ \phi$	(	8.3	± 0.7	) $\times 10^{-6}$	
Γ_{243}	$f_0(980) K^+ \times$ B($f_0(980) \rightarrow K^+ K^-$)	<	2.9		$\times 10^{-6}$	CL=90%
Γ_{244}	$a_2(1320) K^+ \times$ B($a_2(1320) \rightarrow K^+ K^-$)	<	1.1		$\times 10^{-6}$	CL=90%
Γ_{245}	$f'_2(1525) K^+ \times$ B($f'_2(1525) \rightarrow K^+ K^-$)	<	4.9		$\times 10^{-6}$	CL=90%
Γ_{246}	$X_0(1550) K^+ \times$ B($X_0(1550) \rightarrow K^+ K^-$)	(	4.3	± 0.7	) $\times 10^{-6}$	
Γ_{247}	$\phi(1680) K^+ \times$ B($\phi(1680) \rightarrow K^+ K^-$)	<	8		$\times 10^{-7}$	CL=90%
Γ_{248}	$f_0(1710) K^+ \times$ B($f_0(1710) \rightarrow K^+ K^-$)	(	1.7	± 1.0	) $\times 10^{-6}$	
Γ_{249}	$K^+ K^- K^+$ nonresonant	(	2.8	$^{+0.9}_{-1.6}$	) $\times 10^{-5}$	S=3.3
Γ_{250}	$K^*(892)^+ K^+ K^-$	(	3.6	± 0.5	) $\times 10^{-5}$	
Γ_{251}	$K^*(892)^+ \phi$	(	9.6	± 3.0	) $\times 10^{-6}$	S=1.9
Γ_{252}	$K_1(1400)^+ \phi$	<	1.1		$\times 10^{-3}$	CL=90%
Γ_{253}	$K_2^*(1430)^+ \phi$	<	3.4		$\times 10^{-3}$	CL=90%
Γ_{254}	$K^+ \phi \phi$	(	4.9	$^{+2.4}_{-2.2}$	) $\times 10^{-6}$	S=2.9
Γ_{255}	$\eta' \eta' K^+$	<	2.5		$\times 10^{-5}$	CL=90%
Γ_{256}	$K^*(892)^+ \gamma$	(	4.03	± 0.26	) $\times 10^{-5}$	
Γ_{257}	$K_1(1270)^+ \gamma$	(	4.3	± 1.3	) $\times 10^{-5}$	
Γ_{258}	$\eta K^+ \gamma$	(	9.4	± 1.1	) $\times 10^{-6}$	
Γ_{259}	$\eta' K^+ \gamma$	<	4.2		$\times 10^{-6}$	CL=90%
Γ_{260}	$\phi K^+ \gamma$	(	3.5	± 0.6	) $\times 10^{-6}$	
Γ_{261}	$K^+ \pi^- \pi^+ \gamma$	(	2.50	± 0.28	) $\times 10^{-5}$	
Γ_{262}	$K^*(892)^0 \pi^+ \gamma$	(	2.0	$^{+0.7}_{-0.6}$	) $\times 10^{-5}$	
Γ_{263}	$K^+ \rho^0 \gamma$	<	2.0		$\times 10^{-5}$	CL=90%
Γ_{264}	$K^+ \pi^- \pi^+ \gamma$ nonresonant	<	9.2		$\times 10^{-6}$	CL=90%
Γ_{265}	$K_1(1400)^+ \gamma$	<	1.5		$\times 10^{-5}$	CL=90%
Γ_{266}	$K_2^*(1430)^+ \gamma$	(	1.4	± 0.4	) $\times 10^{-5}$	

Γ_{267}	$K^*(1680)^+\gamma$	<	1.9	$\times 10^{-3}$	CL=90%
Γ_{268}	$K_3^*(1780)^+\gamma$	<	3.9	$\times 10^{-5}$	CL=90%
Γ_{269}	$K_4^*(2045)^+\gamma$	<	9.9	$\times 10^{-3}$	CL=90%

Light unflavored meson modes

Γ_{270}	$\rho^+\gamma$		$(8.8\ {}^{+2.9}_{-2.5})$	$\times 10^{-7}$	
Γ_{271}	$\pi^+\pi^0$		$(5.5\ \pm 0.6)$	$\times 10^{-6}$	
Γ_{272}	$\pi^+\pi^+\pi^-$		$(1.62\ \pm 0.15)$	$\times 10^{-5}$	
Γ_{273}	$\rho^0\pi^+$		$(8.7\ \pm 1.1)$	$\times 10^{-6}$	
Γ_{274}	$\pi^+ f_0(980)\times$ B($f_0(980) \to \pi^+\pi^-$)	<	3.0	$\times 10^{-6}$	CL=90%
Γ_{275}	$\pi^+ f_2(1270)$		$(8.2\ \pm 2.5)$	$\times 10^{-6}$	
Γ_{276}	$\rho(1450)^0\pi^+$	<	2.3	$\times 10^{-6}$	CL=90%
Γ_{277}	$f_0(1370)\pi^+\times$B($f_0(1370) \to \pi^+\pi^-$)	<	3.0	$\times 10^{-6}$	CL=90%
Γ_{278}	$f_0(600)\pi^+\times$B($f_0(600) \to \pi^+\pi^-$)	<	4.1	$\times 10^{-6}$	CL=90%
Γ_{279}	$\pi^+\pi^-\pi^+$ nonresonant	<	4.6	$\times 10^{-6}$	CL=90%
Γ_{280}	$\pi^+\pi^0\pi^0$	<	8.9	$\times 10^{-4}$	CL=90%
Γ_{281}	$\rho^+\pi^0$		$(1.20\ \pm 0.19)$	$\times 10^{-5}$	
Γ_{282}	$\pi^+\pi^-\pi^+\pi^0$	<	4.0	$\times 10^{-3}$	CL=90%
Γ_{283}	$\rho^+\rho^0$		$(1.8\ \pm 0.4)$	$\times 10^{-5}$	S=1.5
Γ_{284}	$\rho^+ f_0(980)\times$B($f_0(980) \to \pi^+\pi^-$)	<	1.9	$\times 10^{-6}$	CL=90%
Γ_{285}	$a_1(1260)^+\pi^0$	<	1.7	$\times 10^{-3}$	CL=90%
Γ_{286}	$a_1(1260)^0\pi^+$	<	9.0	$\times 10^{-4}$	CL=90%
Γ_{287}	$\omega\pi^+$		$(6.7\ \pm 0.6)$	$\times 10^{-6}$	
Γ_{288}	$\omega\rho^+$		$(1.06\ {}^{+0.26}_{-0.23})$	$\times 10^{-5}$	
Γ_{289}	$\eta\pi^+$		$(4.9\ \pm 0.5)$	$\times 10^{-6}$	
Γ_{290}	$\eta'\pi^+$		$(2.6\ \pm 1.1)$	$\times 10^{-6}$	S=2.0
Γ_{291}	$\eta'\rho^+$		$(8.7\ {}^{+3.9}_{-3.1})$	$\times 10^{-6}$	
Γ_{292}	$\eta\rho^+$		$(8.4\ \pm 2.2)$	$\times 10^{-6}$	
Γ_{293}	$\phi\pi^+$	<	2.4	$\times 10^{-7}$	CL=90%
Γ_{294}	$\phi\rho^+$	<	1.6	$\times 10^{-5}$	
Γ_{295}	$a_0^0\pi^+$	<	5.8	$\times 10^{-6}$	CL=90%
Γ_{296}	$\pi^+\pi^+\pi^+\pi^-\pi^-$	<	8.6	$\times 10^{-4}$	CL=90%
Γ_{297}	$\rho^0 a_1(1260)^+$	<	6.2	$\times 10^{-4}$	CL=90%
Γ_{298}	$\rho^0 a_2(1320)^+$	<	7.2	$\times 10^{-4}$	CL=90%
Γ_{299}	$\pi^+\pi^+\pi^+\pi^-\pi^-\pi^0$	<	6.3	$\times 10^{-3}$	CL=90%
Γ_{300}	$a_1(1260)^+ a_1(1260)^0$	<	1.3	%	CL=90%

Charged particle ($h^{\pm}$) modes

$h^{\pm} = K^{\pm}$ or $\pi^{\pm}$

Γ_{301}	$h^+\pi^0$		(	1.6	$^{+0.7}_{-0.6}$	) $\times 10^{-5}$	
Γ_{302}	ωh^+		(	1.38	$^{+0.27}_{-0.24}$	) $\times 10^{-5}$	
Γ_{303}	$h^+ X^0$ (Familon)		$<$	4.9		$\times 10^{-5}$	CL=90%

Baryon modes

Γ_{304}	$p\overline{p}\pi^+$		(	3.1	$^{+0.8}_{-0.7}$	) $\times 10^{-6}$	
Γ_{305}	$p\overline{p}\pi^+$ nonresonant		$<$	5.3		$\times 10^{-5}$	CL=90%
Γ_{306}	$p\overline{p}\pi^+\pi^+\pi^-$		$<$	5.2		$\times 10^{-4}$	CL=90%
Γ_{307}	$p\overline{p}K^+$		(	5.6	± 1.0	) $\times 10^{-6}$	S=2.4
Γ_{308}	$\Theta(1710)^{++}\overline{p} \times$ B$(\Theta(1710)^{++} \to pK^+)$	[g]	$<$	9.1		$\times 10^{-8}$	CL=90%
Γ_{309}	$f_J(2220)K^+ \times$ B$(f_J(2220) \to p\overline{p})$	[g]	$<$	4.1		$\times 10^{-7}$	CL=90%
Γ_{310}	$p\overline{\Lambda}(1520)$		$<$	1.5		$\times 10^{-6}$	CL=90%
Γ_{311}	$p\overline{p}K^+$ nonresonant		$<$	8.9		$\times 10^{-5}$	CL=90%
Γ_{312}	$p\overline{p}K^*(892)^+$		(	1.03	$^{+0.38}_{-0.33}$	) $\times 10^{-5}$	
Γ_{313}	$p\overline{\Lambda}$		$<$	4.9		$\times 10^{-7}$	CL=90%
Γ_{314}	$p\overline{\Lambda}\gamma$		(	2.2	± 0.6	) $\times 10^{-6}$	
Γ_{315}	$p\overline{\Sigma}\gamma$		$<$	4.6		$\times 10^{-6}$	CL=90%
Γ_{316}	$p\overline{\Lambda}\pi^+\pi^-$		$<$	2.0		$\times 10^{-4}$	CL=90%
Γ_{317}	$\Lambda\overline{\Lambda}\pi^+$		$<$	2.8		$\times 10^{-6}$	CL=90%
Γ_{318}	$\Lambda\overline{\Lambda}K^+$		(	2.9	$^{+1.0}_{-0.8}$	) $\times 10^{-6}$	
Γ_{319}	$\overline{\Delta}^0 p$		$<$	3.8		$\times 10^{-4}$	CL=90%
Γ_{320}	$\Delta^{++}\overline{p}$		$<$	1.5		$\times 10^{-4}$	CL=90%
Γ_{321}	$D^+ p\overline{p}$		$<$	1.5		$\times 10^{-5}$	CL=90%
Γ_{322}	$D^*(2010)^+ p\overline{p}$		$<$	1.5		$\times 10^{-5}$	CL=90%
Γ_{323}	$\overline{\Lambda}_c^- p\pi^+$		(	2.1	± 0.6	) $\times 10^{-4}$	
Γ_{324}	$\overline{\Lambda}_c^- \Delta(1232)^{++}$		$<$	1.9		$\times 10^{-5}$	CL=90%
Γ_{325}	$\overline{\Lambda}_c^- \Delta_X(1600)^{++}$		(	5.9	± 1.9	) $\times 10^{-5}$	
Γ_{326}	$\overline{\Lambda}_c^- \Delta_X(2420)^{++}$		(	4.7	± 1.6	) $\times 10^{-5}$	
Γ_{327}	$(\overline{\Lambda}_c^- p)_s \pi^+$	[h]	(	3.9	± 1.3	) $\times 10^{-5}$	
Γ_{328}	$\overline{\Lambda}_c^- p\pi^+\pi^0$		(	1.8	± 0.6	) $\times 10^{-3}$	
Γ_{329}	$\overline{\Lambda}_c^- p\pi^+\pi^+\pi^-$		(	2.3	± 0.7	) $\times 10^{-3}$	
Γ_{330}	$\overline{\Lambda}_c^- p\pi^+\pi^+\pi^-\pi^0$		$<$	1.34		%	CL=90%
Γ_{331}	$\Lambda_c^+\Lambda_c^- K^+$		(	7	± 4	) $\times 10^{-4}$	
Γ_{332}	$\overline{\Sigma}_c(2455)^0 p$		(	3.7	± 1.3	) $\times 10^{-5}$	
Γ_{333}	$\overline{\Sigma}_c(2520)^0 p$		$<$	2.7		$\times 10^{-5}$	CL=90%

Γ_{334}	$\overline{\Sigma}_c(2455)^0 p\pi^0$			(	4.4	± 1.8	$) \times 10^{-4}$	
Γ_{335}	$\overline{\Sigma}_c(2455)^0 p\pi^-\pi^+$			(	4.4	± 1.7	$) \times 10^{-4}$	
Γ_{336}	$\overline{\Sigma}_c(2455)^{--} p\pi^+\pi^+$			(	2.8	± 1.2	$) \times 10^{-4}$	
Γ_{337}	$\overline{\Lambda}_c(2593)^- / \overline{\Lambda}_c(2625)^- p\pi^+$			$<$	1.9		$\times 10^{-4}$	CL=90%
Γ_{338}	$\overline{\Xi}_c^0 \Lambda_c^+ \times$ B($\overline{\Xi}_c^0 \to \ \overline{\Xi}^+\pi^-$)			(	5.6	$^{+2.7}_{-2.4}$	$) \times 10^{-5}$	
Γ_{339}	$\overline{\Xi}_c^0 \Lambda_c^+ \times$ B($\overline{\Xi}_c^0 \to \ \Lambda K^+\pi^-$)			(	4.0	± 1.6	$) \times 10^{-5}$	

Lepton Family number (*LF*) or Lepton number (*L*) violating modes, or $\Delta B = 1$ weak neutral current (*B1*) modes

Γ_{340}	$\pi^+ e^+ e^-$	*B1*		$<$	3.9		$\times 10^{-3}$	CL=90%
Γ_{341}	$\pi^+ \mu^+ \mu^-$	*B1*		$<$	9.1		$\times 10^{-3}$	CL=90%
Γ_{342}	$\pi^+ \nu\overline{\nu}$	*B1*		$<$	1.0		$\times 10^{-4}$	CL=90%
Γ_{343}	$K^+ e^+ e^-$	*B1*		(	4.9	± 1.0	$) \times 10^{-7}$	
Γ_{344}	$K^+ \mu^+ \mu^-$	*B1*		(	3.9	$^{+1.0}_{-0.9}$	$) \times 10^{-7}$	
Γ_{345}	$K^+ \ell^+ \ell^-$	*B1*	[a]	(	4.4	$^{+0.8}_{-0.7}$	$) \times 10^{-7}$	S=1.1
Γ_{346}	$K^+ \overline{\nu}\nu$	*B1*		$<$	5.2		$\times 10^{-5}$	CL=90%
Γ_{347}	$K^*(892)^+ e^+ e^-$	*B1*		(	8	± 8	$) \times 10^{-7}$	
Γ_{348}	$K^*(892)^+ \mu^+ \mu^-$	*B1*		(	8	$^{+6}_{-4}$	$) \times 10^{-7}$	
Γ_{349}	$K^*(892)^+ \ell^+ \ell^-$	*B1*	[a]	(	7	± 5	$) \times 10^{-7}$	
Γ_{350}	$\pi^+ e^+ \mu^-$	*LF*		$<$	6.4		$\times 10^{-3}$	CL=90%
Γ_{351}	$\pi^+ e^- \mu^+$	*LF*		$<$	6.4		$\times 10^{-3}$	CL=90%
Γ_{352}	$K^+ e^+ \mu^-$	*LF*		$<$	9.1		$\times 10^{-8}$	CL=90%
Γ_{353}	$K^+ e^- \mu^+$	*LF*		$<$	1.3		$\times 10^{-7}$	CL=90%
Γ_{354}	$K^+ e^\pm \mu^\mp$	*LF*		$<$	9.1		$\times 10^{-8}$	CL=90%
Γ_{355}	$K^*(892)^+ e^+ \mu^-$	*LF*		$<$	1.3		$\times 10^{-6}$	CL=90%
Γ_{356}	$K^*(892)^+ e^- \mu^+$	*LF*		$<$	9.9		$\times 10^{-7}$	CL=90%
Γ_{357}	$K^*(892)^+ e^\pm \mu^\mp$	*LF*		$<$	1.4		$\times 10^{-7}$	CL=90%
Γ_{358}	$\pi^- e^+ e^+$	*L*		$<$	1.6		$\times 10^{-6}$	CL=90%
Γ_{359}	$\pi^- \mu^+ \mu^+$	*L*		$<$	1.4		$\times 10^{-6}$	CL=90%
Γ_{360}	$\pi^- e^+ \mu^+$	*L*		$<$	1.3		$\times 10^{-6}$	CL=90%
Γ_{361}	$\rho^- e^+ e^+$	*L*		$<$	2.6		$\times 10^{-6}$	CL=90%
Γ_{362}	$\rho^- \mu^+ \mu^+$	*L*		$<$	5.0		$\times 10^{-6}$	CL=90%
Γ_{363}	$\rho^- e^+ \mu^+$	*L*		$<$	3.3		$\times 10^{-6}$	CL=90%
Γ_{364}	$K^- e^+ e^+$	*L*		$<$	1.0		$\times 10^{-6}$	CL=90%
Γ_{365}	$K^- \mu^+ \mu^+$	*L*		$<$	1.8		$\times 10^{-6}$	CL=90%
Γ_{366}	$K^- e^+ \mu^+$	*L*		$<$	2.0		$\times 10^{-6}$	CL=90%
Γ_{367}	$K^*(892)^- e^+ e^+$	*L*		$<$	2.8		$\times 10^{-6}$	CL=90%
Γ_{368}	$K^*(892)^- \mu^+ \mu^+$	*L*		$<$	8.3		$\times 10^{-6}$	CL=90%
Γ_{369}	$K^*(892)^- e^+ \mu^+$	*L*		$<$	4.4		$\times 10^{-6}$	CL=90%

[*a*] An ℓ indicates an *e* or a μ mode, not a sum over these modes.

[*b*] An *CP*($\pm$1) indicates the *CP*=+1 and *CP*=−1 eigenstates of the D^0-$\overline{D}^0$ system.

[*c*] *D* denotes D^0 or $\overline{D}^0$.

[*d*] D^{*0}_{CP+} decays into $D^0\pi^0$ with the D^0 reconstructed in *CP*-even eigenstates K^+K^- and $\pi^+\pi^-$.

[*e*] $\overline{D}^{**}$ represents an excited state with mass 2.2 < M < 2.8 GeV/c^2.

[*f*] $X(3872)^+$ is a hypothetical charged partner of the $X(3872)$.

[*g*] $\Theta(1710)^{++}$ is a possible narrow pentaquark state and $G(2220)$ is a possible glueball resonance.

[*h*] $(\overline{\Lambda}_c^- p)_s$ denotes a low-mass enhancement near 3.35 GeV/c^2.

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 11 measurements and one constraint to determine 3 parameters. The overall fit has a χ^2 = 7.2 for 9 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\left\langle\delta x_i \delta x_j\right\rangle/(\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_{168}	24
	x_{149}

B^+ BRANCHING RATIOS

$\Gamma(\ell^+\nu_\ell\,\text{anything})/\Gamma_{\text{total}}$ Γ_1/Γ

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at http://www.slac.stanford.edu/xorg/hfag/. The averaging/rescaling procedure takes into account correlations between the measurements.

VALUE (units 10^{-2})	*DOCUMENT ID*		*TECN*	*COMMENT*
10.99±0.28 OUR EVALUATION				
10.76±0.32 OUR AVERAGE Error includes scale factor of 1.1.				
11.17±0.25±0.28	[15] URQUIJO	07	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
10.28±0.26±0.39	[16] AUBERT,B	06Y	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
10.25±0.57±0.65	[17] ARTUSO	97	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
11.15±0.26±0.41	[18] OKABE	05	BELL	Repl. by URQUIJO 07
10.1 ±1.8 ±1.5	ATHANAS	94	CLE2	Sup. by ARTUSO 97

[15] URQUIJO 07 report a measurement of (10.34 ± 023 ± 0.25)% for the partial branching fraction of $B^+ \to e^+ \nu_e X_c$ decay with electron energy above 0.6 GeV. We converted the result to $B^+ \to e^+ \nu_e X$ branching fraction.

[16] The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame. The best precision on the ratio is achieved for a momentum threshold of 1.0 GeV: B($B^+ \to e^+ \nu_e X$) / B($B^0 \to e^+ \nu_e X$) = 1.074 ± 0.041 ± 0.026.

[17] ARTUSO 97 uses partial reconstruction of $B \to D^* \ell \nu_\ell$ and inclusive semileptonic branching ratio from BARISH 96B (0.1049 ± 0.0017 ± 0.0043).

[18] The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame, and their ratio of B($B^+ \to e^+ \nu_e X$)/B($B^0 \to e^+ \nu_e X$) = 1.08 ± 0.05 ± 0.02.

$\Gamma(e^+ \nu_e X_c)/\Gamma_{total}$ Γ_2/Γ

VALUE (units 10^{-2})	DOCUMENT ID		TECN	COMMENT
10.79±0.25±0.27	[19] URQUIJO	07	BELL	$e^+ e^- \to \Upsilon(4S)$

[19] Measure the independent B^+ and B^0 partial branching fractions with electron threshold energies of 0.4 GeV.

$\Gamma(\overline{D}^0 \ell^+ \nu_\ell)/\Gamma_{total}$ Γ_3/Γ

$\ell = e$ or μ, not sum over e and μ modes.

VALUE	DOCUMENT ID		TECN	COMMENT
0.0215±0.0022 OUR AVERAGE				
0.0221±0.0013±0.0019	[20] BARTELT	99	CLE2	$e^+ e^- \to \Upsilon(4S)$
0.016 ±0.006 ±0.003	[21] FULTON	91	CLEO	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0194±0.0015±0.0034	[22] ATHANAS	97	CLE2	Repl. by BARTELT 99

[20] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[21] FULTON 91 assumes equal production of $B^0 \overline{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$.

[22] ATHANAS 97 uses missing energy and missing momentum to reconstruct neutrino.

$\Gamma(\overline{D}^*(2007)^0 \ell^+ \nu_\ell)/\Gamma_{total}$ Γ_4/Γ

$\ell = e$ or μ, not sum over e and μ modes.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.065 ±0.005 OUR AVERAGE					
0.0650±0.0020±0.0043		[23] ADAM	03	CLE2	$e^+ e^- \to \Upsilon(4S)$
0.066 ±0.016 ±0.015		[24] ALBRECHT	92C	ARG	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0650±0.0020±0.0043		[25] BRIERE	02	CLE2	$e^+ e^- \to \Upsilon(4S)$
0.0513±0.0054±0.0064	302	[26] BARISH	95	CLE2	Repl. by ADAM 03
seen	398	[27] SANGHERA	93	CLE2	$e^+ e^- \to \Upsilon(4S)$
0.041 ±0.008 $^{+0.008}_{-0.009}$		[28] FULTON	91	CLEO	$e^+ e^- \to \Upsilon(4S)$
0.070 ±0.018 ±0.014		[29] ANTREASYAN	90B	CBAL	$e^+ e^- \to \Upsilon(4S)$

[23] Simultaneous measurements of both $B^0 \rightarrow D^*(2010)^- \ell\nu$ and $B^+ \rightarrow \overline{D}(2007)^0 \ell\nu$.

[24] ALBRECHT 92C reports $0.058 \pm 0.014 \pm 0.013$. We rescale using the method described in STONE 94 but with the updated PDG 94 B($D^0 \rightarrow K^- \pi^+$). Assumes equal production of $B^0 \overline{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$.

[25] The results are based on the same analysis and data sample reported in ADAM 03.

[26] BARISH 95 use B($D^0 \rightarrow K^- \pi^+$) = (3.91 ± 0.08 ± 0.17)% and B($D^{*0} \rightarrow D^0 \pi^0$) = (63.6 ± 2.3 ± 3.3)%.

[27] Combining $\overline{D}^{*0} \ell^+ \nu_\ell$ and $\overline{D}^{*-} \ell^+ \nu_\ell$ SANGHERA 93 test $V-A$ structure and fit the decay angular distributions to obtain $A_{FB} = 3/4*(\Gamma^- - \Gamma^+)/\Gamma = 0.14 \pm 0.06 \pm 0.03$. Assuming a value of V_{cb}, they measure V, A_1, and A_2, the three form factors for the $D^* \ell \nu_\ell$ decay, where results are slightly dependent on model assumptions.

[28] Assumes equal production of $B^0 \overline{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$. Uncorrected for D and D^* branching ratio assumptions.

[29] ANTREASYAN 90B is average over B and $\overline{D}^*(2010)$ charge states.

$\Gamma(\overline{D}_1(2420)^0 \ell^+ \nu_\ell)/\Gamma_{total}$ — Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0056±0.0013±0.0009	[30] ANASTASSOV 98	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[30] ANASTASSOV 98 result is derived from the measurement of B($B^+ \rightarrow \overline{D}_1^0 \ell^+ \nu_\ell$)×B($\overline{D}_1^0 \rightarrow D^{*+} \pi^-$) = (0.373 ± 0.085 ± 0.052 ± 0.024)% by assuming B($\overline{D}_1^0 \rightarrow D^{*+} \pi^-$) = 67%, where the third error includes theoretical uncertainties.

$\Gamma(\overline{D}_2^*(2460)^0 \ell^+ \nu_\ell)/\Gamma_{total}$ — Γ_6/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-3}$	90	[31] ANASTASSOV 98	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[31] ANASTASSOV 98 result is derived from the measurement of B($B^+ \rightarrow \overline{D}_2^{*0} \ell^+ \nu_\ell$)×B($\overline{D}_2^{*0} \rightarrow D^{*+} \pi^-$) < 0.16% at 90% CL by assuming B($\overline{D}_2^{*0} \rightarrow D^{*+} \pi^-$) = 20%.

$\Gamma(D^- \pi^+ \ell^+ \nu_\ell)/\Gamma_{total}$ — Γ_7/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
5.2±0.9±0.5	[32] LIVENTSEV 05	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[32] LIVENTSEV 05 reports [B($B^+ \rightarrow D^- \pi^+ \ell^+ \nu_\ell$) / B($B^0 \rightarrow D^- \ell^+ \nu_\ell$)] = 0.25 ± 0.03 ± 0.03. We multiply by our best value B($B^0 \rightarrow D^- \ell^+ \nu_\ell$) = $(2.08 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^{*-} \pi^+ \ell^+ \nu_\ell)/\Gamma_{total}$ — Γ_8/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.3±1.5±0.2	[33,34] LIVENTSEV 05	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[33] Excludes D^{*+} contribution to $D\pi$ modes.

[34] LIVENTSEV 05 reports [B($B^+ \rightarrow D^{*-} \pi^+ \ell^+ \nu_\ell$) / B($B^0 \rightarrow D^*(2010)^- \ell^+ \nu_\ell$)] = 0.12 ± 0.02 ± 0.02. We multiply by our best value B($B^0 \rightarrow D^*(2010)^- \ell^+ \nu_\ell$) = $(5.29 \pm 0.19) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0 \ell^+ \nu_\ell)/\Gamma_{total}$ — Γ_9/Γ

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at http://www.slac.stanford.edu/xorg/hfag/. The averaging/rescaling procedure takes into account correlations between the measurements.

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
0.77±0.10±0.07 OUR EVALUATION				
0.75±0.09 OUR AVERAGE				
0.77±0.14±0.08	[35] HOKUUE	07	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
0.74±0.05±0.10	[36] AUBERT,B	05O	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

[35] The signal events are tagged by a second B meson reconstructed in the semileptonic mode $B \rightarrow D^{(*)} \ell \nu_\ell$.

[36] B^+ and B^0 decays combined assuming isospin symmetry. Systematic errors include both experimental and form-factor uncertainties.

$\Gamma(\pi^0 e^+ \nu_e)/\Gamma_{total}$ — Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.9±0.2±0.2		[37] ALEXANDER	96T	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<22	90	ANTREASYAN	90B	CBAL	$e^+ e^- \rightarrow \Upsilon(4S)$

[37] Derived based in the reported B^0 result by assuming isospin symmetry: $\Gamma(B^0 \rightarrow \pi^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \pi^0 \ell^+ \nu)$.

$\Gamma(\eta \ell^+ \nu_\ell)/\Gamma_{total}$ — Γ_{11}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
0.84±0.31±0.18	[38] ATHAR	03	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[38] ATHAR 03 reports systematic errors 0.16 ± 0.09, which are experimental systematic and systematic due to model dependence. We combine these in quadrature.

$\Gamma(\omega \ell^+ \nu_\ell)/\Gamma_{total}$ — Γ_{12}/Γ

$\ell = e$ or μ, not sum over e and μ modes.

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
1.3±0.4±0.4		[39] SCHWANDA	04	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.1	90	[40] BEAN	93B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[39] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[40] BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\rho^0 \ell^+ \nu_\ell)$ and $\Gamma(\rho^- \ell^+ \nu_\ell)$ with this result, they obtain a limit $<(1.6–2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \omega \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8–0.13$ at 90% CL is derived as well.

$\Gamma(\omega \mu^+ \nu_\mu)/\Gamma_{total}$ — Γ_{13}/Γ

VALUE	DOCUMENT ID		TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	[41] ALBRECHT	91C	ARG

[41] In ALBRECHT 91C, one event is fully reconstructed providing evidence for the $b \rightarrow u$ transition.

$\Gamma(\rho^0 \ell^+ \nu_\ell)/\Gamma_{total}$ — Γ_{14}/Γ

$\ell = e$ or μ, not sum over e and μ modes.

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
1.28±0.18 OUR AVERAGE					
$1.33\pm0.23\pm0.18$		[42] HOKUUE	07	BELL	$e^+e^- \to \Upsilon(4S)$
$1.16\pm0.11\pm0.30$		[43] AUBERT,B	05O	BABR	$e^+e^- \to \Upsilon(4S)$
$1.34\pm0.15^{+0.28}_{-0.32}$		[44] BEHRENS	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1.40\pm0.21^{+0.32}_{-0.33}$		[44] BEHRENS	00	CLE2	$e^+e^- \to \Upsilon(4S)$
$1.2\ \pm0.2\ ^{+0.3}_{-0.4}$		[44] ALEXANDER	96T	CLE2	$e^+e^- \to \Upsilon(4S)$
<2.1	90	[45] BEAN	93B	CLE2	$e^+e^- \to \Upsilon(4S)$

[42] The signal events are tagged by a second B meson reconstructed in the semileptonic mode $B \to D^{(*)}\ell\nu_\ell$.

[43] B^+ and B^0 decays combined assuming isospin symmetry. Systematic errors include both experimental and form-factor uncertainties.

[44] Derived based in the reported B^0 result by assuming isospin symmetry: $\Gamma(B^0 \to \rho^-\ell^+\nu) = 2\Gamma(B^+ \to \rho^0\ell^+\nu) \approx 2\Gamma(B^+ \to \omega\ell^+\nu)$.

[45] BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\omega^0\ell^+\nu_\ell)$ and $\Gamma(\rho^-\ell^+\nu_\ell)$ with this result, they obtain a limit $<(1.6–2.7)\times10^{-4}$ at 90% CL for $B^+ \to \rho^0\ell^+\nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8–0.13$ at 90% CL is derived as well.

$\Gamma(p\overline{p}e^+\nu_e)/\Gamma_{total}$ — Γ_{15}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<5.2\times10^{-3}$	90	[46] ADAM	03B	CLE2	$e^+e^- \to \Upsilon(4S)$

[46] Based on phase-space model; if $V{-}A$ model is used, the 90% CL upper limit becomes $<1.2\times10^{-3}$.

$\Gamma(e^+\nu_e)/\Gamma_{total}$ — Γ_{16}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$<$ 9.8	90	[47] SATOYAMA	07	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<15	90	ARTUSO	95	CLE2	$e^+e^- \to \Upsilon(4S)$

[47] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\mu^+\nu_\mu)/\Gamma_{total}$ — Γ_{17}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$<$ 1.7	90	[48] SATOYAMA	07	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<$ 6.6	90	AUBERT	04O	BABR	$e^+e^- \to \Upsilon(4S)$
<21	90	ARTUSO	95	CLE2	$e^+e^- \to \Upsilon(4S)$

[48] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\tau^+\nu_\tau)/\Gamma_{total}$ — Γ_{18}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.79^{+0.56}_{-0.49}{}^{+0.46}_{-0.51}}$		[49] IKADO	06	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<$ 2.6	90	[49] AUBERT	06K	BABR	$e^+e^- \to \Upsilon(4S)$
$<$ 4.2	90	[49] AUBERT,B	05B	BABR	Repl. by AUBERT 06K
$<$ 8.3	90	[50] BARATE	01E	ALEP	$e^+e^- \to Z$
$<$ 8.4	90	[49] BROWDER	01	CLE2	$e^+e^- \to \Upsilon(4S)$
$<$ 5.7	90	[51] ACCIARRI	97F	L3	$e^+e^- \to Z$
$<$104	90	[52] ALBRECHT	95D	ARG	$e^+e^- \to \Upsilon(4S)$
$<$ 22	90	ARTUSO	95	CLE2	$e^+e^- \to \Upsilon(4S)$
$<$ 18	90	[53] BUSKULIC	95	ALEP	$e^+e^- \to Z$

[49] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[50] The energy-flow and b-tagging algorithms were used.
[51] ACCIARRI 97F uses missing-energy technique and $f(b \to B^-) = (38.2 \pm 2.5)\%$.
[52] ALBRECHT 95D use full reconstruction of one B decay as tag.
[53] BUSKULIC 95 uses same missing-energy technique as in $\overline{b} \to \tau^+\nu_\tau X$, but analysis is restricted to endpoint region of missing-energy distribution.

$\Gamma(e^+\nu_e\gamma)/\Gamma_{total}$ — Γ_{19}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<2.0 \times 10^{-4}}$	90	[54] BROWDER	97	CLE2	$e^+e^- \to \Upsilon(4S)$

[54] BROWDER 97 uses the hermiticity of the CLEO II detector to reconstruct the neutrino energy and momentum.

$\Gamma(\mu^+\nu_\mu\gamma)/\Gamma_{total}$ — Γ_{20}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<5.2 \times 10^{-5}}$	90	[55] BROWDER	97	CLE2	$e^+e^- \to \Upsilon(4S)$

[55] BROWDER 97 uses the hermiticity of the CLEO II detector to reconstruct the neutrino energy and momentum.

$\Gamma(D^0 X)/\Gamma_{total}$ — Γ_{21}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.086±0.006±0.004	[56] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.098±0.009±0.006	[56] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[56] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\overline{D}^0 X)/\Gamma_{total}$ — Γ_{22}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.786 \pm 0.016^{+0.034}_{-0.033}}$	[57] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.793 \pm 0.025^{+0.045}_{-0.044}$	[57] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[57] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D^0 X)/[\Gamma(D^0 X)+\Gamma(\overline{D}^0 X)]$ — $\Gamma_{21}/(\Gamma_{21}+\Gamma_{22})$

VALUE	DOCUMENT ID		TECN	COMMENT
0.098±0.007±0.001	AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.110±0.010±0.003	AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

$\Gamma(D^+ X)/\Gamma_{\text{total}}$ — Γ_{23}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.025±0.005±0.002	[58] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.038±0.009±0.005	[58] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[58] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D^- X)/\Gamma_{\text{total}}$ — Γ_{24}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.099±0.008±0.009	[59] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.098±0.012±0.014	[59] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[59] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D^+ X)/[\Gamma(D^+ X)+\Gamma(D^- X)]$ — $\Gamma_{23}/(\Gamma_{23}+\Gamma_{24})$

VALUE	DOCUMENT ID		TECN	COMMENT
0.204±0.035±0.001	AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.278±0.052±0.009	AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

$\Gamma(D_s^+ X)/\Gamma_{\text{total}}$ — Γ_{25}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.079 \pm 0.006^{+0.013}_{-0.011}}$	[60] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.143 \pm 0.016^{+0.051}_{-0.034}$	[60] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[60] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D_s^- X)/\Gamma_{\text{total}}$ — Γ_{26}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.011^{+0.004}_{-0.003}{}^{+0.002}_{-0.001}}$		[61] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.022	90	[61] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[61] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D_s^+ X)/[\Gamma(D_s^+ X)+\Gamma(D_s^- X)]$ — $\Gamma_{25}/(\Gamma_{25}+\Gamma_{26})$

VALUE	DOCUMENT ID		TECN	COMMENT
0.884±0.038±0.002	AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.966±0.039±0.012	AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

$\Gamma(D_s^- X)/[\Gamma(D_s^+ X)+\Gamma(D_s^- X)]$ — $\Gamma_{26}/(\Gamma_{25}+\Gamma_{26})$

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.126	90	AUBERT,BE	04B	BABR	$e^+e^- \to \Upsilon(4S)$

$\Gamma(\Lambda_c^+ X)/\Gamma_{total}$ — Γ_{27}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.021\pm0.005^{+0.008}_{-0.004}}$	[62] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.029\pm0.008^{+0.011}_{-0.007}$	[62] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[62] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\overline{\Lambda}_c^- X)/\Gamma_{total}$ — Γ_{28}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.028\pm0.005^{+0.010}_{-0.007}}$	[63] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.035\pm0.008^{+0.013}_{-0.009}$	[63] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[63] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\Lambda_c^+ X)/[\Gamma(\Lambda_c^+ X)+\Gamma(\overline{\Lambda}_c^- X)]$ — $\Gamma_{27}/(\Gamma_{27}+\Gamma_{28})$

VALUE	DOCUMENT ID		TECN	COMMENT
0.427±0.071±0.001	AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.452±0.090±0.003	AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

$\Gamma(\overline{c} X)/\Gamma_{total}$ — Γ_{29}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.968\pm0.019^{+0.041}_{-0.039}}$	[64] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.983\pm0.030^{+0.054}_{-0.051}$	[64] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[64] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(cX)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.234 \pm 0.012 ^{+0.018}_{-0.014}}$	[65] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.330 \pm 0.022 ^{+0.055}_{-0.037}$	[65] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[65] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\overline{c}cX)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.202 \pm 0.023 ^{+0.053}_{-0.049}}$	[66] AUBERT	07N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.313 \pm 0.037 ^{+0.088}_{-0.075}$	[66] AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

[66] Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\overline{D}^0\pi^+)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
4.84±0.15 OUR AVERAGE					
$4.90 \pm 0.07 \pm 0.22$		[67] AUBERT	07H	BABR	$e^+e^- \to \Upsilon(4S)$
$5.3 \pm 0.6 \pm 0.3$		[68] ABULENCIA	06J	CDF	$p\overline{p}$ at 1.96 TeV
$4.49 \pm 0.21 \pm 0.23$		[69] AUBERT,BE	06J	BABR	$e^+e^- \to \Upsilon(4S)$
$4.97 \pm 0.12 \pm 0.29$		[67,70] AHMED	02B	CLE2	$e^+e^- \to \Upsilon(4S)$
$5.0 \pm 0.7 \pm 0.6$	54	[71] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
$5.4 \ ^{+1.8}_{-1.5} \ ^{+1.2}_{-0.9}$	14	[72] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$4.84 \pm 0.27 \pm 0.09$		[73] AUBERT,B	04P	BABR	Repl. by AUBERT 07H
$5.5 \pm 0.4 \pm 0.5$	304	[74] ALAM	94	CLE2	Repl. by AHMED 02B
$2.0 \pm 0.8 \pm 0.6$	12	[71] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$
$1.9 \pm 1.0 \pm 0.6$	7	[75] ALBRECHT	88K	ARG	$e^+e^- \to \Upsilon(4S)$

[67] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[68] ABULENCIA 06J reports $[B(B^+ \to \overline{D}^0\pi^+) / B(B^0 \to D^-\pi^+)] = 1.97 \pm 0.10 \pm 0.21$. We multiply by our best value $B(B^0 \to D^-\pi^+) = (2.68 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[69] Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.

[70] AHMED 02B reports an additional uncertainty on the branching ratios to account for 4.5% uncertainty on relative production of B^0 and B^+, which is not included here.

[71] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D.

[72] BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

[73] AUBERT,B 04P reports $[B(B^+ \to \overline{D}^0\pi^+) \times B(D^0 \to K^-\pi^+)] = (1.846 \pm 0.032 \pm 0.097) \times 10^{-4}$. We divide by our best value $B(D^0 \to K^-\pi^+) = (3.82 \pm 0.07) \times 10^{-2}$.

Our first error is their experiment's error and our second error is the systematic error from using our best value.

[74] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute B($D^0 \to K^-\pi^+$) and the PDG 1992 B($D^0 \to K^-\pi^+\pi^0$)/B($D^0 \to K^-\pi^+$) and B($D^0 \to K^-\pi^+\pi^+\pi^-$)/B($D^0 \to K^-\pi^+$).

[75] ALBRECHT 88K assumes $B^0\overline{B}^0$:B^+B^- ratio is 45:55. Superseded by ALBRECHT 90J.

$\Gamma(\overline{D}^0\rho^+)/\Gamma_{total}$ — Γ_{35}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0134±0.0018 OUR AVERAGE					
0.0135±0.0012±0.0015	212	[76] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$
0.013 ±0.004 ±0.004	19	[77] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.021 ±0.008 ±0.009	10	[78] ALBRECHT	88K	ARG	$e^+e^- \to \Upsilon(4S)$

[76] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute B($D^0 \to K^-\pi^+$) and the PDG 1992 B($D^0 \to K^-\pi^+\pi^0$)/B($D^0 \to K^-\pi^+$) and B($D^0 \to K^-\pi^+\pi^+\pi^-$)/B($D^0 \to K^-\pi^+$).

[77] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D.

[78] ALBRECHT 88K assumes $B^0\overline{B}^0$:B^+B^- ratio is 45:55.

$\Gamma(\overline{D}^0K^+)/\Gamma_{total}$ — Γ_{36}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
4.02±0.21 OUR AVERAGE				
4.02±0.20±0.13	[79] AUBERT	04N	BABR	$e^+e^- \to \Upsilon(4S)$
$4.8\ ^{+0.8}_{-0.6}\ \pm0.2$	[80] BORNHEIM	03	CLE2	$e^+e^- \to \Upsilon(4S)$
3.7 ±0.4 ±0.1	[81,82] SWAIN	03	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.6 ±0.6 ±0.1	[81,83] ABE	03D	BELL	Repl. by SWAIN 03
4.19±0.57±0.40	[84] ABE	01I	BELL	Repl. by ABE 03D
2.92±0.80±0.28	[85] ATHANAS	98	CLE2	Repl. by BORNHEIM 03

[79] AUBERT 04N reports [B($B^+ \to \overline{D}^0K^+$) / B($B^+ \to \overline{D}^0\pi^+$)] = $(831 \pm 35 \pm 20) \times 10^{-4}$. We multiply by our best value B($B^+ \to \overline{D}^0\pi^+$) = $(4.84 \pm 0.15) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[80] BORNHEIM 03 reports [B($B^+ \to \overline{D}^0K^+$) / B($B^+ \to \overline{D}^0\pi^+$)] = $(990^{+140}_{-120}{}^{+70}_{-60}) \times 10^{-4}$. We multiply by our best value B($B^+ \to \overline{D}^0\pi^+$) = $(4.84 \pm 0.15) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[81] Flavor specific D^0 meson is reconstructed via $D^0 \to K^-\pi^+$.

[82] SWAIN 03 reports [B($B^+ \to \overline{D}^0K^+$) / B($B^+ \to \overline{D}^0\pi^+$)] = $(770 \pm 50 \pm 60) \times 10^{-4}$. We multiply by our best value B($B^+ \to \overline{D}^0\pi^+$) = $(4.84 \pm 0.15) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[83] ABE 03D reports [B($B^+ \to \overline{D}^0K^+$) / B($B^+ \to \overline{D}^0\pi^+$)] = $(940 \pm 90 \pm 70) \times 10^{-4}$. We multiply by our best value B($B^+ \to \overline{D}^0\pi^+$) = $(4.84 \pm 0.15) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[84] ABE 01I reports B($B^+ \to \overline{D}^0 K^+$)/B($B^+ \to \overline{D}^0 \pi^+$)= 0.079 ± 0.009 ± 0.006. We multiply by our best value B($B^+ \to \overline{D}^0 \pi^+$)= (5.3 ± 0.5) × 10^{-3}. Our first error is their experiment's error and the second error is systematic error from using our best value.

[85] ATHANAS 98 reports [B($B^+ \to \overline{D}^0 K^+$)]/[B($B^+ \to \overline{D}^0 \pi^+$)] =0.055 ± 0.014 ± 0.005. We multiply by our best value B($B^+ \to \overline{D}^0 \pi^+$)= (5.3 ± 0.5) × 10^{-3}. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{CP(+1)} K^+)/\Gamma_{\text{total}}$ — Γ_{37}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
3.6±0.5±0.2	[86] AUBERT	06J	BABR	$e^+ e^- \to \Upsilon(4S)$

[86] AUBERT 06J reports [B($B^+ \to D_{CP(+1)} K^+$) / B($B^+ \to \overline{D}^0 K^+$)] = 0.90 ± 0.12 ± 0.04. We multiply by our best value B($B^+ \to \overline{D}^0 K^+$) = (4.02 ± 0.21) × 10^{-4}. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{CP(-1)} K^+)/\Gamma_{\text{total}}$ — Γ_{38}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
3.5±0.4±0.2	[87] AUBERT	06J	BABR	$e^+ e^- \to \Upsilon(4S)$

[87] AUBERT 06J reports [B($B^+ \to D_{CP(-1)} K^+$) / B($B^+ \to \overline{D}^0 K^+$)] = 0.86 ± 0.10 ± 0.05. We multiply by our best value B($B^+ \to \overline{D}^0 K^+$) = (4.02 ± 0.21) × 10^{-4}. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{CP(+1)} K^+)/\Gamma(D_{CP(+1)} \pi^+)$ — Γ_{37}/Γ_{33}

VALUE	DOCUMENT ID		TECN	COMMENT
0.091±0.012 OUR AVERAGE				
0.094±0.015±0.007	[88] ABE	06	BELL	$e^+ e^- \to \Upsilon(4S)$
0.088±0.016±0.005	[89] AUBERT	04N	BABR	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.125±0.036±0.010	[89] ABE	03D	BELL	Repl. by SWAIN 03
0.093±0.018±0.008	[89] SWAIN	03	BELL	Repl. by ABE 06

[88] Reports a double ratio of B($B^+ \to D_{CP(+1)} K^+$)/B($B^+ \to D_{CP(+1)} \pi^+$) and B($B^+ \to \overline{D}^0 K^+$)/B($B^+ \to \overline{D}^0 \pi^+$), 1.13 ± 0.16 ± 0.08. We multiply by our best value of B($B^+ \to \overline{D}^0 K^+$)/B($B^+ \to \overline{D}^0 \pi^+$) = 0.083 ± 0.006. Our first error is their experiment's error and the second error is systematic error from using our best value.

[89] CP=+1 eigenstate of $D^0 \overline{D}^0$ system is reconstructed via $K^+ K^-$ and $\pi^+ \pi^-$.

$\Gamma(D_{CP(-1)} K^+)/\Gamma(D_{CP(-1)} \pi^+)$ — Γ_{38}/Γ_{34}

VALUE	DOCUMENT ID		TECN	COMMENT
0.097±0.016±0.007	[90] ABE	06	BELL	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.119±0.028±0.006	[91] ABE	03D	BELL	Repl. by SWAIN 03
0.108±0.019±0.007	[91] SWAIN	03	BELL	Repl. by ABE 06

[90] Reports a double ratio of B($B^+ \to D_{CP(-1)} K^+$)/B($B^+ \to D_{CP(-1)} \pi^+$) and B($B^+ \to \overline{D}^0 K^+$)/B($B^+ \to \overline{D}^0 \pi^+$), $1.17 \pm 0.14 \pm 0.14$. We multiply by our best value of B($B^+ \to \overline{D}^0 K^+$)/B($B^+ \to \overline{D}^0 \pi^+$) $= 0.083 \pm 0.006$. Our first error is their experiment's error and the second error is systematic error from using our best value.

[91] $CP{=}-1$ eigenstate of $D^0\overline{D}^0$ system is reconstructed via $K^0_S\pi^0$, $K^0_S\omega$, $K^0_S\phi$, $K^0_S\eta$, and $K^0_S\eta'$.

$\Gamma([K^-\pi^+]_D K^+)/\Gamma([K^+\pi^-]_D K^+)$ — Γ_{39}/Γ_{40}

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.029	90	[92] AUBERT	05G	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.044	90	[93] SAIGO	05	BELL	$e^+e^- \to \Upsilon(4S)$
<0.026	90	[94] AUBERT,B	04L	BABR	Repl. by AUBERT 05G

[92] AUBERT 05G extract a constraint on the magnitude of the ratio of amplitudes $|$A($B^+ \to D^0 K^+$) / A($B^+ \to \overline{D}^0 K^+$)$|$ < 0.23 at 90% CL (Bayesian). Similar measurements from $B^+ \to D^{*0} K^+$ are also reported.

[93] SAIGO 05 extract a constraint on the magnitude of the ratio of amplitudes $|$A($B^+ \to D^0 K^+$) / A($B^+ \to \overline{D}^0 K^+$)$|$ < 0.27 at 90% CL.

[94] AUBERT,B 04L extract a constraint on the magnitude of the ratio of amplitudes $|$A($B^+ \to D^0 K^+$)/A($B^+ \to \overline{D}^0 K^+$)$|$ < 0.22 at 90% CL.

$\Gamma([K^-\pi^+]_D K^*(892)^+)/\Gamma([K^+\pi^-]_D K^*(892)^+)$ — Γ_{41}/Γ_{42}

VALUE	DOCUMENT ID		TECN	COMMENT
0.046±0.031±0.008	AUBERT,B	05V	BABR	$e^+e^- \to \Upsilon(4S)$

$\Gamma([K^-\pi^+]_D \pi^+)/\Gamma_{\text{total}}$ — Γ_{43}/Γ

VALUE (units 10^{-5})	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.73^{+0.51}_{-0.46}\pm0.03}$	[95] SAIGO	05	BELL	$e^+e^- \to \Upsilon(4S)$

[95] SAIGO 05 reports [B($B^+ \to [K^-\pi^+]_D \pi^+$) × B($D^0 \to K^-\pi^+$)] $= (6.6^{+1.9}_{-1.7} \pm 0.5) \times 10^{-7}$. We divide by our best value B($D^0 \to K^-\pi^+$) $= (3.82 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma([\pi^+\pi^-\pi^0]_D K^-)/\Gamma_{\text{total}}$ — Γ_{44}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
5.5±1.0±0.7	[96] AUBERT,B	05T	BABR	$e^+e^- \to \Upsilon(4S)$

[96] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma([K^-\pi^+]_D \pi^+)/\Gamma(\overline{D}^0\pi^+)$ — Γ_{43}/Γ_{32}

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
$\mathbf{3.5^{+1.0}_{-0.9}\pm0.2}$	SAIGO	05	BELL	$e^+e^- \to \Upsilon(4S)$

$\Gamma(\overline{D}^0 K^*(892)^+)/\Gamma_{total}$ Γ_{45}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
5.3 ±0.4 OUR AVERAGE				
5.29±0.30±0.34	[97] AUBERT	06Z	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
6.1 ±1.6 ±1.7	[97] MAHAPATRA	02	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
6.3 ±0.7 ±0.5	[97] AUBERT	04Q	BABR	Repl. by AUBERT 06Z

[97] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{CP(-1)} K^*(892)^+)/\Gamma_{total}$ Γ_{46}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
1.7±0.7±0.1	[98] AUBERT,B	05U	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

[98] AUBERT,B 05U reports $[B(B^+ \rightarrow D_{CP(-1)} K^*(892)^+) / B(B^+ \rightarrow \overline{D}^0 K^*(892)^+)] = 0.325 \pm 0.13 \pm 0.04$. We multiply by our best value $B(B^+ \rightarrow \overline{D}^0 K^*(892)^+) = (5.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{CP(+1)} K^*(892)^+)/\Gamma_{total}$ Γ_{47}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
5.2±1.1±0.4	[99] AUBERT,B	05U	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

[99] AUBERT,B 05U reports $[B(B^+ \rightarrow D_{CP(+1)} K^*(892)^+) / B(B^+ \rightarrow \overline{D}^0 K^*(892)^+)] = 0.98 \pm 0.20 \pm 0.055$. We multiply by our best value $B(B^+ \rightarrow \overline{D}^0 K^*(892)^+) = (5.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\overline{D}^0 K^+ \overline{K}^0)/\Gamma_{total}$ Γ_{48}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
5.5±1.4±0.8	[100] DRUTSKOY	02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[100] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^0 K^+ \overline{K}^*(892)^0)/\Gamma_{total}$ Γ_{49}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
7.5±1.3±1.1	[101] DRUTSKOY	02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[101] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^0 \pi^+ \pi^+ \pi^-)/\Gamma_{total}$ Γ_{50}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.0115±0.0029±0.0021	[102] BORTOLETTO	92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

[102] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma(\overline{D}^0 \pi^+ \pi^+ \pi^- \text{ nonresonant})/\Gamma_{total}$ Γ_{51}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.0051±0.0034±0.0023	[103] BORTOLETTO	92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

[103] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma(\overline{D}^0 \pi^+ \rho^0)/\Gamma_{\text{total}}$ — Γ_{52}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0042±0.0023±0.0020	[104] BORTOLETTO 92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

[104] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma(\overline{D}^0 a_1(1260)^+)/\Gamma_{\text{total}}$ — Γ_{53}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0045±0.0019±0.0031	[105] BORTOLETTO 92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

[105] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

$\Gamma(\overline{D}^0 \omega \pi^+)/\Gamma_{\text{total}}$ — Γ_{54}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0041±0.0007±0.0006	[106] ALEXANDER 01B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[106] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega\pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-\ 5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

$\Gamma(D^*(2010)^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ — Γ_{55}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.35±0.22 OUR AVERAGE					
1.25±0.08±0.22			[107] ABE 04D	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
1.9 ±0.7 ±0.3		14	[108] ALAM 94	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
2.6 ±1.4 ±0.7		11	[109] ALBRECHT 90J	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
$2.4\ ^{+1.7}_{-1.6}\ ^{+1.0}_{-0.6}$		3	[110] BEBEK 87	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<4.	90		[111] BORTOLETTO 92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
5. ±2. ±3.		7	[112] ALBRECHT 87C	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

[107] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[108] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0\pi^+)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

[109] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D.

[110] BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

[111] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$. The authors also find the product branching fraction into $D^{**}\pi$ followed by $D^{**} \rightarrow D^*(2010)\pi$ to be $0.0014^{+0.0008}_{-0.0006} \pm 0.0003$ where D^{**} represents all orbitally excited D mesons.

[112] ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0\overline{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^-\pi^+\pi^+)/\Gamma_{total}$ Γ_{56}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
1.02±0.04±0.15			[113] ABE	04D	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<1.4	90		[114] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$
<7	90		[115] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
$2.5\ ^{+4.1}_{-2.3}\ ^{+2.4}_{-0.8}$		1	[116] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[113] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[114] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \to K^-\pi^+\pi^+)$.

[115] BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D. The product branching fraction into $D_0^*(2340)\pi$ followed by $D_0^*(2340) \to D\pi$ is < 0.005 at 90%CL and into $D_2^*(2460)$ followed by $D_2^*(2460) \to D\pi$ is < 0.004 at 90%CL.

[116] BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. $B(D^- \to K^+\pi^-\pi^-) = (9.1 \pm 1.3 \pm 0.4)\%$ is assumed.

$\Gamma(D^+K^0)/\Gamma_{total}$ Γ_{57}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<5.0	90	[117] AUBERT,B	05E	BABR	$e^+e^- \to \Upsilon(4S)$

[117] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^*(2007)^0\pi^+)/\Gamma_{total}$ Γ_{58}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
5.19±0.26 OUR AVERAGE					
5.52±0.17±0.42		[118] AUBERT	07H	BABR	$e^+e^- \to \Upsilon(4S)$
5.5 ±0.4 ±0.2		[119,120] AUBERT,BE	06J	BABR	$e^+e^- \to \Upsilon(4S)$
4.34±0.47±0.18		[121] BRANDENB...	98	CLE2	$e^+e^- \to \Upsilon(4S)$
5.2 ±0.7 ±0.7	71	[122] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$
7.2 ±1.8 ±1.6		[123] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
4.0 ±1.4 ±1.2	9	[123] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.7 ±4.4		[124] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[118] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[119] AUBERT,BE 06J reports $[B(B^+ \to \overline{D}^*(2007)^0\pi^+) / B(B^+ \to \overline{D}^0\pi^+)] = 1.14 \pm 0.07 \pm 0.04$. We multiply by our best value $B(B^+ \to \overline{D}^0\pi^+) = (4.84 \pm 0.15) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[120] Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.

[121] BRANDENBURG 98 assume equal production of B^+ and B^0 at $\Upsilon(4S)$ and use the D^* reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of $B(D^* \to D\pi)$.

[122] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \to D^0\pi^0)$ and absolute $B(D^0 \to K^-\pi^+)$ and the PDG 1992 $B(D^0 \to K^-\pi^+\pi^0)/B(D^0 \to K^-\pi^+)$ and $B(D^0 \to K^-\pi^+\pi^+\pi^-)/B(D^0 \to K^-\pi^+)$.

[123] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

[124] This is a derived branching ratio, using the inclusive pion spectrum and other two-body B decays. BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0 \overline{B}^0$.

$\Gamma(\overline{D}^*(2007)^0 \omega \pi^+)/\Gamma_{total}$ Γ_{61}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.0045±0.0010±0.0007	[125] ALEXANDER	01B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[125] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-\ 5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

$\Gamma(\overline{D}^*(2007)^0 \rho^+)/\Gamma_{total}$ Γ_{62}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0098±0.0017 OUR AVERAGE					
0.0098±0.0006±0.0017		[126] CSORNA	03	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.010 ±0.006 ±0.004	7	[127] ALBRECHT	90J	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0168±0.0021±0.0028	86	[128] ALAM	94	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[126] Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$ resonance. The second error combines the systematic and theoretical uncertainties in quadrature. CSORNA 03 includes data used in ALAM 94. A full angular fit to three complex helicity amplitudes is performed.

[127] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

[128] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$. The nonresonant $\pi^+ \pi^0$ contribution under the ρ^+ is negligible.

$\Gamma(\overline{D}^*(2007)^0 K^+)/\Gamma_{total}$ Γ_{63}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
4.16±0.33 OUR AVERAGE				
$4.22^{+0.30}_{-0.26} \pm 0.21$	[129] AUBERT	05N	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
3.59±0.97±0.31	[130] ABE	01I	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[129] AUBERT 05N reports $[B(B^+ \rightarrow \overline{D}^*(2007)^0 K^+) \ / \ B(B^+ \rightarrow \overline{D}^*(2007)^0 \pi^+)] = 0.0813 \pm 0.0040^{+0.0042}_{-0.0031}$. We multiply by our best value $B(B^+ \rightarrow \overline{D}^*(2007)^0 \pi^+) = (5.19 \pm 0.26) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[130] ABE 01I reports $B(B^+ \rightarrow \overline{D}^*(2007)^0 K^+)/B(B^+ \rightarrow \overline{D}^*(2007)^0 \pi^+) = 0.078 \pm 0.019 \pm 0.009$. We multiply by our best value $B(B^+ \rightarrow \overline{D}^*(2007)^0 \pi^+) = (4.6 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and the second error is systematic error from using our best value.

$\Gamma(\overline{D}^{*0}_{CP(+1)} K^+)/\Gamma(\overline{D}^{*0}_{CP(+1)} \pi^+)$ Γ_{64}/Γ_{59}

VALUE	DOCUMENT ID		TECN	COMMENT
0.095±0.017 OUR AVERAGE				
0.11 ±0.02 ±0.02	[131] ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$
0.086±0.021±0.007	[132] AUBERT	05N	BABR	$e^+e^- \to \Upsilon(4S)$

[131] Reports a double ratio of B($B^+ \to (D^*_{CP(+1)})^0 K^+$)/B($B^+ \to (D^*_{CP(+1)})^0 \pi^+$) and B($B^+ \to \overline{D}^{*0} K^+$)/B($B^+ \to \overline{D}^{*0} \pi^+$), $1.41 \pm 0.25 \pm 0.06$. We multiply by our best value of B($B^+ \to \overline{D}^{*0} K^+$)/B($B^+ \to \overline{D}^{*0} \pi^+$) $= 0.080 \pm 0.011$. Our first error is their experiment's error and the second error is systematic error from using our best value.

[132] Uses $D^{*0} \to D^0 \pi^0$ with D^0 reconstructed in the CP-even eigenstates $K^+ K^-$ and $\pi^+ \pi^-$.

$\Gamma(\overline{D}^{*0}_{CP(-1)} K^+)/\Gamma(D^{*0}_{CP(-1)} \pi^+)$ Γ_{65}/Γ_{60}

VALUE	DOCUMENT ID		TECN	COMMENT
0.09±0.03 ±0.01	[133] ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$

[133] Reports a double ratio of B($B^+ \to (D^*_{CP(-1)})^0 K^+$)/B($B^+ \to (D^*_{CP(-1)})^0 \pi^+$) and B($B^+ \to \overline{D}^{*0} K^+$)/B($B^+ \to \overline{D}^{*0} \pi^+$), $1.15 \pm 0.31 \pm 0.12$. We multiply by our best value of B($B^+ \to \overline{D}^{*0} K^+$)/B($B^+ \to \overline{D}^{*0} \pi^+$) $= 0.080 \pm 0.011$. Our first error is their experiment's error and the second error is systematic error from using our best value.

$\Gamma(\overline{D}^*(2007)^0 K^*(892)^+)/\Gamma_{total}$ Γ_{66}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
8.1±1.4 OUR AVERAGE				
8.3±1.1±1.0	[134] AUBERT	04K	BABR	$e^+e^- \to \Upsilon(4S)$
7.2±2.2±2.6	[135] MAHAPATRA	02	CLE2	$e^+e^- \to \Upsilon(4S)$

[134] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[135] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and an unpolarized final state.

$\Gamma(\overline{D}^*(2007)^0 K^+ \overline{K}^0)/\Gamma_{total}$ Γ_{67}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<10.6	90	[136] DRUTSKOY	02	BELL	$e^+e^- \to \Upsilon(4S)$

[136] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^*(2007)^0 K^+ K^*(892)^0)/\Gamma_{total}$ Γ_{68}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
15.3±3.1±2.9	[137] DRUTSKOY	02	BELL	$e^+e^- \to \Upsilon(4S)$

[137] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^*(2007)^0 \pi^+ \pi^+ \pi^-)/\Gamma_{total}$ Γ_{69}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID		TECN	COMMENT
1.03 ±0.12 OUR AVERAGE					
1.055±0.047±0.129		[138] MAJUMDER	04	BELL	$e^+e^- \to \Upsilon(4S)$
0.94 ±0.20 ±0.17	48	[139,140] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$

[138] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[139] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II B($D^*(2007)^0 \to D^0\pi^0$) and absolute B($D^0 \to K^-\pi^+$) and the PDG 1992 B($D^0 \to K^-\pi^+\pi^0$)/B($D^0 \to K^-\pi^+$) and B($D^0 \to K^-\pi^+\pi^+\pi^-$)/B($D^0 \to K^-\pi^+$).

[140] The three pion mass is required to be between 1.0 and 1.6 GeV consistent with an a_1 meson. (If this channel is dominated by a_1^+, the branching ratio for $\overline{D}^{*0} a_1^+$ is twice that for $\overline{D}^{*0}\pi^+\pi^+\pi^-$.)

$\Gamma(\overline{D}^*(2007)^0 a_1(1260)^+)/\Gamma_{total}$ — Γ_{70}/Γ

VALUE		DOCUMENT ID		TECN	COMMENT
0.0188±0.0040±0.0034	[141,142]	ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$

[141] ALAM 94 value is twice their $\Gamma(\overline{D}^*(2007)^0\pi^+\pi^+\pi^-)/\Gamma_{total}$ value based on their observation that the three pions are dominantly in the $a_1(1260)$ mass range 1.0 to 1.6 GeV.

[142] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II B($D^*(2007)^0 \to D^0\pi^0$) and absolute B($D^0 \to K^-\pi^+$) and the PDG 1992 B($D^0 \to K^-\pi^+\pi^0$)/B($D^0 \to K^-\pi^+$) and B($D^0 \to K^-\pi^+\pi^+\pi^-$)/B($D^0 \to K^-\pi^+$).

$\Gamma(\overline{D}^*(2007)^0\pi^-\pi^+\pi^+\pi^0)/\Gamma_{total}$ — Γ_{71}/Γ

VALUE		DOCUMENT ID		TECN	COMMENT
0.0180±0.0024±0.0027	[143]	ALEXANDER	01B	CLE2	$e^+e^- \to \Upsilon(4S)$

[143] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega\pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

$\Gamma(\overline{D}^{*0}3\pi^+2\pi^-)/\Gamma_{total}$ — Γ_{72}/Γ

VALUE (units 10^{-3})		DOCUMENT ID		TECN	COMMENT
5.67±0.91±0.85	[144]	MAJUMDER	04	BELL	$e^+e^- \to \Upsilon(4S)$

[144] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^+\pi^0)/\Gamma_{total}$ — Γ_{73}/Γ

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
<0.00017	90	[145]	BRANDENB...	98	CLE2	$e^+e^- \to \Upsilon(4S)$

[145] BRANDENBURG 98 assume equal production of B^+ and B^0 at $\Upsilon(4S)$ and use the D^* partial reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of B($D^* \to D\pi$).

$\Gamma(D^*(2010)^+K^0)/\Gamma_{total}$ — Γ_{74}/Γ

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
$<9.0 \times 10^{-6}$	90	[146]	AUBERT,B	05E	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<9.5 \times 10^{-5}$	90	[146]	GRITSAN	01	CLE2	$e^+e^- \to \Upsilon(4S)$

[146] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^0)/\Gamma_{total}$ Γ_{75}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0152±0.0071±0.0001	26	[147] ALBRECHT	90J	ARG	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.043 ±0.013 ±0.026	24	[148] ALBRECHT	87C	ARG	$e^+ e^- \to \Upsilon(4S)$

[147] ALBRECHT 90J reports $0.018 \pm 0.007 \pm 0.005$ for $B(D^*(2010)^+ \to D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \to D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D.

[148] ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \to B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \to B^0 \overline{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^+ \pi^-)/\Gamma_{total}$ Γ_{76}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
2.56±0.26±0.33		[149] MAJUMDER	04	BELL	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<10	90	[150] ALBRECHT	90J	ARG	$e^+ e^- \to \Upsilon(4S)$

[149] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[150] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

$\Gamma(\overline{D}^{**0} \pi^+)/\Gamma_{total}$ Γ_{77}/Γ

D^{**0} represents an excited state with mass $2.2 < M < 2.8$ GeV/c^2.

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
5.9±1.3±0.2	[151,152] AUBERT,BE	06J	BABR	$e^+ e^- \to \Upsilon(4S)$

[151] AUBERT,BE 06J reports $[B(B^+ \to \overline{D}^{**0} \pi^+) / B(B^+ \to \overline{D}^0 \pi^+)] = 1.22 \pm 0.13 \pm 0.23$. We multiply by our best value $B(B^+ \to \overline{D}^0 \pi^+) = (4.84 \pm 0.15) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[152] Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.

$\Gamma(\overline{D}_1^*(2420)^0 \pi^+)/\Gamma_{total}$ Γ_{78}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0015±0.0006 OUR AVERAGE Error includes scale factor of 1.3.					
0.0011±0.0005±0.0002	8	[153] ALAM	94	CLE2	$e^+ e^- \to \Upsilon(4S)$
0.0025±0.0007±0.0006		[154] ALBRECHT	94D	ARG	$e^+ e^- \to \Upsilon(4S)$

[153] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \to D^0 \pi^+)$ and absolute $B(D^0 \to K^- \pi^+)$ and the PDG 1992 $B(D^0 \to K^- \pi^+ \pi^0)/B(D^0 \to K^- \pi^+)$ and assuming $B(D_1(2420)^0 \to D^*(2010)^+ \pi^-) = 67\%$.

[154] ALBRECHT 94D assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \to D^0 \pi^+)$ assuming $B(D_1(2420)^0 \to D^*(2010)^+ \pi^-) = 67\%$.

$\Gamma\left(\overline{D}_1(2420)^0\pi^+ \times B(\overline{D}{}_1^0 \to \overline{D}{}^0\pi^+\pi^-)\right)/\Gamma_{total}$ — Γ_{79}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
$1.85\pm0.29^{+0.35}_{-0.55}$	[155] ABE	05A	BELL	$e^+e^- \to \Upsilon(4S)$

[155] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(\overline{D}{}_2^*(2462)^0\pi^+ \times B(\overline{D}{}_2^*(2462)^0 \to D^-\pi^+)\right)/\Gamma_{total}$ — Γ_{80}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
$3.4\pm0.3\pm0.72$	[156] ABE	04D	BELL	$e^+e^- \to \Upsilon(4S)$

[156] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(\overline{D}{}_0^*(2400)^0\pi^+ \times B(\overline{D}{}_0^*(2400)^0 \to D^-\pi^+)\right)/\Gamma_{total}$ — Γ_{81}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
$6.1\pm0.6\pm1.8$	[157] ABE	04D	BELL	$e^+e^- \to \Upsilon(4S)$

[157] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(\overline{D}_1(2421)^0\pi^+ \times B(\overline{D}_1(2421)^0 \to D^{*-}\pi^+)\right)/\Gamma_{total}$ — Γ_{82}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
$6.8\pm0.7\pm1.3$	[158] ABE	04D	BELL	$e^+e^- \to \Upsilon(4S)$

[158] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(\overline{D}{}_2^*(2462)^0\pi^+ \times B(\overline{D}{}_2^*(2462)^0 \to D^{*-}\pi^+)\right)/\Gamma_{total}$ — Γ_{83}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
$1.8\pm0.3\pm0.4$	[159] ABE	04D	BELL	$e^+e^- \to \Upsilon(4S)$

[159] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(\overline{D}{}_1'(2427)^0\pi^+ \times B(\overline{D}{}_1'(2427)^0 \to D^{*-}\pi^+)\right)/\Gamma_{total}$ — Γ_{84}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
$5.0\pm0.4\pm1.1$	[160] ABE	04D	BELL	$e^+e^- \to \Upsilon(4S)$

[160] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(\overline{D}_1(2420)^0\pi^+ \times B(\overline{D}{}_1^0 \to \overline{D}{}^{*0}\pi^+\pi^-)\right)/\Gamma_{total}$ — Γ_{85}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<0.06	90	[161] ABE	05A	BELL	$e^+e^- \to \Upsilon(4S)$

[161] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(\overline{D}{}_1^*(2420)^0\rho^+\right)/\Gamma_{total}$ — Γ_{86}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0014	90	[162] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$

[162] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \to D^0\pi^+)$ assuming $B(D_1(2420)^0 \to D^*(2010)^+\pi^-) = 67\%$.

$\Gamma(\overline{D}{}_2^*(2460)^0 \pi^+)/\Gamma_{\text{total}}$ — Γ_{87}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0013	90	[163] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0028	90	[164] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$
<0.0023	90	[165] ALBRECHT	94D	ARG	$e^+e^- \to \Upsilon(4S)$

[163] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III B($D^+ \to K^-\pi^+\pi^+$) and B($D_2^*(2460)^0 \to D^+\pi^-$) = 30%.

[164] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III B($D^+ \to K^-\pi^+\pi^+$), the CLEO II B($D^*(2010)^+ \to D^0\pi^+$) and B($D_2^*(2460)^0 \to D^*(2010)^+\pi^-$) = 20%.

[165] ALBRECHT 94D assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II B($D^*(2010)^+ \to D^0\pi^+$) and B($D_2^*(2460)^0 \to D^*(2010)^+\pi^-$) = 30%.

$\Gamma(\overline{D}{}_2^*(2460)^0 \pi^+ \times \text{B}(\overline{D}{}_2^{*0} \to \overline{D}{}^{*0}\pi^+\pi^-))/\Gamma_{\text{total}}$ — Γ_{88}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<0.22	90	[166] ABE	05A	BELL	$e^+e^- \to \Upsilon(4S)$

[166] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}{}_2^*(2460)^0 \rho^+)/\Gamma_{\text{total}}$ — Γ_{89}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0047	90	[167] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$
<0.005	90	[168] ALAM	94	CLE2	$e^+e^- \to \Upsilon(4S)$

[167] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III B($D^+ \to K^-\pi^+\pi^+$) and B($D_2^*(2460)^0 \to D^+\pi^-$) = 30%.

[168] ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III B($D^+ \to K^-\pi^+\pi^+$), the CLEO II B($D^*(2010)^+ \to D^0\pi^+$) and B($D_2^*(2460)^0 \to D^*(2010)^+\pi^-$) = 20%.

$\Gamma(\overline{D}{}^0 D_s^+)/\Gamma_{\text{total}}$ — Γ_{90}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0100±0.0017 OUR AVERAGE					
0.0095±0.0020±0.0008		[169] AUBERT	06N	BABR	$e^+e^- \to \Upsilon(4S)$
0.0098±0.0026±0.0009		[170] GIBAUT	96	CLE2	$e^+e^- \to \Upsilon(4S)$
0.014 ±0.008 ±0.001		[171] ALBRECHT	92G	ARG	$e^+e^- \to \Upsilon(4S)$
0.013 ±0.006 ±0.001	5	[172] BORTOLETTO	90	CLEO	$e^+e^- \to \Upsilon(4S)$

[169] AUBERT 06N reports $(0.92 \pm 0.14 \pm 0.18) \times 10^{-2}$ for B($D_s^+ \to \phi\pi^+$) = 0.0462 ± 0.0062. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = $(4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[170] GIBAUT 96 reports 0.0126 ± 0.0022 ± 0.0025 for B($D_s^+ \to \phi\pi^+$) = 0.035. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = $(4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[171] ALBRECHT 92G reports 0.024 ± 0.012 ± 0.004 for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = $(4.5 \pm 0.4) \times 10^{-2}$. Our first error is their

experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 branching ratios, e.g., B($D^0 \rightarrow K^- \pi^+$) = 3.71 ± 0.25%.

[172] BORTOLETTO 90 reports 0.029 ± 0.013 for B($D_s^+ \rightarrow \phi\pi^+$) = 0.02. We rescale to our best value B($D_s^+ \rightarrow \phi\pi^+$) = $(4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma\left(D_{s0}(2317)^+ \overline{D}^0 \times \text{B}(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0)\right)/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$0.73^{+0.22}_{-0.17}$ OUR AVERAGE			
$0.80^{+0.35}_{-0.21} \pm 0.07$	[173,174] AUBERT,B 04S	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.65^{+0.26}_{-0.24} \pm 0.06$	[173,175] KROKOVNY 03B	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[173] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[174] AUBERT,B 04S reports $(1.0 \pm 0.3^{+0.4}_{-0.2}) \times 10^{-3}$ for B($D_s^+ \rightarrow \phi\pi^+$) = 0.036 ± 0.009. We rescale to our best value B($D_s^+ \rightarrow \phi\pi^+$) = $(4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[175] KROKOVNY 03B reports $(0.81^{+0.30}_{-0.27} \pm 0.24) \times 10^{-3}$ for B($D_s^+ \rightarrow \phi\pi^+$) = 0.036 ± 0.009. We rescale to our best value B($D_s^+ \rightarrow \phi\pi^+$) = $(4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma\left(D_{s0}(2317)^+ \overline{D}^0 \times \text{B}(D_{s0}(2317)^+ \rightarrow D_s^{*+} \gamma)\right)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.76	90	[176] KROKOVNY 03B	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[176] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(D_{s0}(2317)^+ \overline{D}^*(2010)^0 \times \text{B}(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0)\right)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$0.9 \pm 0.6^{+0.4}_{-0.3}$	[177] AUBERT,B 04S	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[177] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(D_{sJ}(2457)^+ \overline{D}^0\right)/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$3.1^{+1.0}_{-0.9}$ OUR AVERAGE			
$4.3 \pm 1.6 \pm 1.3$	[178] AUBERT 06N	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$4.6^{+1.8}_{-1.6} \pm 1.0$	[179,180] AUBERT,B 04S	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$2.1^{+1.1}_{-0.9} \pm 0.5$	[179,181] KROKOVNY 03B	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[178] Uses a missing-mass method in the events that one of the B mesons is fully reconstructed.
[179] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[180] AUBERT,B 04S reports $[\mathrm{B}(B^+ \to D_{sJ}(2457)^+ \overline{D}^0) \times \mathrm{B}(D_{s1}(2460)^+ \to D_s^{*+}\pi^0)] = (2.2^{+0.8}_{-0.7} \pm 0.3) \times 10^{-3}$. We divide by our best value $\mathrm{B}(D_{s1}(2460)^+ \to D_s^{*+}\pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
[181] KROKOVNY 03B reports $[\mathrm{B}(B^+ \to D_{sJ}(2457)^+ \overline{D}^0) \times \mathrm{B}(D_{s1}(2460)^+ \to D_s^{*+}\pi^0)] = (1.0^{+0.5}_{-0.4} \pm 0.1) \times 10^{-3}$. We divide by our best value $\mathrm{B}(D_{s1}(2460)^+ \to D_s^{*+}\pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{sJ}(2457)^+ \overline{D}^0 \times \mathrm{B}(D_{sJ}(2457)^+ \to D_s^+\gamma))/\Gamma_{\mathrm{total}}$ — Γ_{95}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
$0.46^{+0.13}_{-0.11}$ OUR AVERAGE				
$0.48^{+0.19}_{-0.13} \pm 0.04$	[182,183] AUBERT,B	04S	BABR	$e^+e^- \to \Upsilon(4S)$
$0.45^{+0.15}_{-0.14} \pm 0.04$	[182,184] KROKOVNY	03B	BELL	$e^+e^- \to \Upsilon(4S)$

[182] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[183] AUBERT,B 04S reports $(0.6 \pm 0.2^{+0.2}_{-0.1}) \times 10^{-3}$ for $\mathrm{B}(D_s^+ \to \phi\pi^+) = 0.036 \pm 0.009$. We rescale to our best value $\mathrm{B}(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
[184] KROKOVNY 03B reports $(0.56^{+0.16}_{-0.15} \pm 0.17) \times 10^{-3}$ for $\mathrm{B}(D_s^+ \to \phi\pi^+) = 0.036 \pm 0.009$. We rescale to our best value $\mathrm{B}(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{sJ}(2457)^+ \overline{D}^0 \times \mathrm{B}(D_{sJ}(2457)^+ \to D_s^+\pi^+\pi^-))/\Gamma_{\mathrm{total}}$ — Γ_{96}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<0.22	90	[185] KROKOVNY	03B	BELL	$e^+e^- \to \Upsilon(4S)$

[185] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{sJ}(2457)^+ \overline{D}^0 \times \mathrm{B}(D_{sJ}(2457)^+ \to D_s^+\pi^0))/\Gamma_{\mathrm{total}}$ — Γ_{97}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<0.27	90	[186] KROKOVNY	03B	BELL	$e^+e^- \to \Upsilon(4S)$

[186] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{sJ}(2457)^+ \overline{D}^0 \times \mathrm{B}(D_{sJ}(2457)^+ \to D_s^{*+}\gamma))/\Gamma_{\mathrm{total}}$ — Γ_{98}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<0.98	90	[187] KROKOVNY	03B	BELL	$e^+e^- \to \Upsilon(4S)$

[187] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{sJ}(2457)^+ \overline{D}^*(2007)^0)/\Gamma_{total}$ — Γ_{99}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
12.0±3.0 OUR AVERAGE				
$11.2 \pm 2.6 \pm 2.0$	[188] AUBERT	06N	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$16.\ ^{+8.}_{-6.} \pm 4.$	[189,190] AUBERT,B	04S	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[188] Uses a missing-mass method in the events that one of the B mesons is fully reconstructed.
[189] AUBERT,B 04S reports $[B(B^+ \rightarrow D_{sJ}(2457)^+ \overline{D}^*(2007)^0) \times B(D_{s1}(2460)^+ \rightarrow D_s^{*+}\pi^0)] = (7.6 \pm 1.7^{+3.2}_{-2.4}) \times 10^{-3}$. We divide by our best value $B(D_{s1}(2460)^+ \rightarrow D_s^{*+}\pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
[190] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{sJ}(2457)^+ \overline{D}^*(2007)^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+\gamma))/\Gamma_{total}$ — Γ_{100}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.4 \pm 0.4^{+0.6}_{-0.4}}$	[191] AUBERT,B	04S	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[191] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^0 D_{sJ}(2536)^+ \times B(D_{sJ}(2536)^+ \rightarrow D^*(2007)^0 K^+))/\Gamma_{total}$ — Γ_{101}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<2	90	AUBERT	03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\overline{D}^*(2007)^0 D_{sJ}(2536)^+ \times B(D_{sJ}(2536)^+ \rightarrow D^*(2007)^0 K^+))/\Gamma_{total}$ — Γ_{102}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<7	90	AUBERT	03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\overline{D}^0 D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+))/\Gamma_{total}$ — Γ_{103}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<2	90	AUBERT	03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\overline{D}^*(2007)^0 D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+))/\Gamma_{total}$ — Γ_{104}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<5	90	AUBERT	03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\overline{D}^0 D_s^{*+})/\Gamma_{total}$ — Γ_{105}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.0076±0.0016 OUR AVERAGE				
$0.0079 \pm 0.0017 \pm 0.0007$	[192] AUBERT	06N	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.0068 \pm 0.0025 \pm 0.0006$	[193] GIBAUT	96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$0.010 \pm 0.007 \pm 0.001$	[194] ALBRECHT	92G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[192] AUBERT 06N reports $(0.77 \pm 0.15 \pm 0.13) \times 10^{-2}$ for $B(D_s^+ \to \phi\pi^+) = 0.0462 \pm 0.0062$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[193] GIBAUT 96 reports $0.0087 \pm 0.0027 \pm 0.0017$ for $B(D_s^+ \to \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[194] ALBRECHT 92G reports $0.016 \pm 0.012 \pm 0.003$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 branching ratios, e.g., $B(D^0 \to K^-\pi^+) = 3.71 \pm 0.25\%$.

$\Gamma(\overline{D}^*(2007)^0 D_s^+)/\Gamma_{total}$ $\qquad\qquad$ Γ_{106}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.0082±0.0017 OUR AVERAGE				
0.0078±0.0018±0.0007	[195] AUBERT	06N	BABR	$e^+e^- \to \Upsilon(4S)$
0.011 ±0.004 ±0.001	[196] GIBAUT	96	CLE2	$e^+e^- \to \Upsilon(4S)$
0.008 ±0.006 ±0.001	[197] ALBRECHT	92G	ARG	$e^+e^- \to \Upsilon(4S)$

[195] AUBERT 06N reports $(0.76 \pm 0.15 \pm 0.13) \times 10^{-2}$ for $B(D_s^+ \to \phi\pi^+) = 0.0462 \pm 0.0062$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[196] GIBAUT 96 reports $0.0140 \pm 0.0043 \pm 0.0035$ for $B(D_s^+ \to \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[197] ALBRECHT 92G reports $0.013 \pm 0.009 \pm 0.002$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 and $D^*(2007)^0$ branching ratios, e.g., $B(D^0 \to K^-\pi^+) = 3.71 \pm 0.25\%$ and $B(D^*(2007)^0 \to D^0\pi^0) = 55 \pm 6\%$.

$\Gamma(\overline{D}^*(2007)^0 D_s^{*+})/\Gamma_{total}$ $\qquad\qquad$ Γ_{107}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.0171±0.0024 OUR AVERAGE				
0.0167±0.0019±0.0015	[198] AUBERT	06N	BABR	$e^+e^- \to \Upsilon(4S)$
0.024 ±0.009 ±0.002	[199] GIBAUT	96	CLE2	$e^+e^- \to \Upsilon(4S)$
0.019 ±0.010 ±0.002	[200] ALBRECHT	92G	ARG	$e^+e^- \to \Upsilon(4S)$

[198] AUBERT 06N reports $(1.62 \pm 0.22 \pm 0.18) \times 10^{-2}$ for $B(D_s^+ \to \phi\pi^+) = 0.0462 \pm 0.0062$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[199] GIBAUT 96 reports $0.0310 \pm 0.0088 \pm 0.0065$ for $B(D_s^+ \to \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[200] ALBRECHT 92G reports $0.031 \pm 0.016 \pm 0.005$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

Assumes PDG 1990 D^0 and $D^*(2007)^0$ branching ratios, e.g., B($D^0 \to K^- \pi^+$) = 3.71 ± 0.25% and B($D^*(2007)^0 \to D^0 \pi^0$) = 55 ± 6%.

$\Gamma(D_s^{(*)+} \overline{D}^{**0})/\Gamma_{total}$ — Γ_{108}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
(2.73 ±0.93 ±0.68) × 10^{-2}	[201] AHMED	00B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[201] AHMED 00B reports their experiment's uncertainties (±0.78 ± 0.48 ± 0.68)%, where the first error is statistical, the second is systematic, and the third is the uncertainty in the $D_s \to \phi\pi$ branching fraction. We combine the first two in quadrature.

$\Gamma(\overline{D}^*(2007)^0 D^*(2010)^+)/\Gamma_{total}$ — Γ_{109}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
8.1±1.2±1.2		[202] AUBERT,B	06A	BABR	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<110	90	BARATE	98Q	ALEP	$e^+ e^- \to Z$

[202] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$[\Gamma(\overline{D}^0 D^*(2010)^+) + \Gamma(\overline{D}^*(2007)^0 D^+)]/\Gamma_{total}$ — Γ_{110}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<130 (CL = 90%) [<13000 × 10^{-6} (CL = 90%) OUR 2007 BEST LIMIT]					
<130	90	BARATE	98Q	ALEP	$e^+ e^- \to Z$

$\Gamma(\overline{D}^0 D^*(2010)^+)/\Gamma_{total}$ — Γ_{111}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
3.9 ±0.5 OUR AVERAGE				
3.6 ±0.5 ±0.4	[203] AUBERT,B	06A	BABR	$e^+ e^- \to \Upsilon(4S)$
4.57±0.71±0.56	[203] MAJUMDER	05	BELL	$e^+ e^- \to \Upsilon(4S)$

[203] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^0 D^+)/\Gamma_{total}$ — Γ_{112}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
4.2 ±0.6 OUR AVERAGE					
3.8 ±0.6 ±0.5		[204] AUBERT,B	06A	BABR	$e^+ e^- \to \Upsilon(4S)$
4.83±0.78±0.58		[204] MAJUMDER	05	BELL	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<67	90	BARATE	98Q	ALEP	$e^+ e^- \to Z$

[204] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^0 D^+ K^0)/\Gamma_{total}$ — Γ_{113}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<2.8	90	[205] AUBERT	03X	BABR	$e^+ e^- \to \Upsilon(4S)$

[205] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^+ \overline{D}^*(2007)^0)/\Gamma_{total}$ — Γ_{114}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
6.3±1.4±1.0	[206] AUBERT,B	06A	BABR	$e^+ e^- \to \Upsilon(4S)$

[206] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\overline{D}^*(2007)^0 D^+ K^0\big)/\Gamma_{total}$ Γ_{115}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<6.1	90	[207] AUBERT 03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[207] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\overline{D}^0 \overline{D}^*(2010)^+ K^0\big)/\Gamma_{total}$ Γ_{116}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$\mathbf{5.2^{+1.0}_{-0.9}\pm0.7}$	[208] AUBERT 03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[208] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\overline{D}^*(2007)^0 D^*(2010)^+ K^0\big)/\Gamma_{total}$ Γ_{117}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$\mathbf{7.8^{+2.3}_{-2.1}\pm1.4}$	[209] AUBERT 03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[209] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\overline{D}^0 D^0 K^+\big)/\Gamma_{total}$ Γ_{118}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.37±0.32 OUR AVERAGE	Error includes scale factor of 1.5.		
1.17±0.21±0.15	[210] CHISTOV 04	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
1.9 ±0.3 ±0.3	[210] AUBERT 03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[210] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\overline{D}^*(2007)^0 D^0 K^+\big)/\Gamma_{total}$ Γ_{119}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3.8	90	[211] AUBERT 03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[211] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\overline{D}^0 D^*(2007)^0 K^+\big)/\Gamma_{total}$ Γ_{120}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
4.7±0.7±0.7	[212] AUBERT 03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[212] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\overline{D}^*(2007)^0 D^*(2007)^0 K^+\big)/\Gamma_{total}$ Γ_{121}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$\mathbf{5.3^{+1.1}_{-1.0}\pm1.2}$	[213] AUBERT 03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[213] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(D^- D^+ K^+\big)/\Gamma_{total}$ Γ_{122}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	[214] AUBERT 03X	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.90	90	[214] CHISTOV 04	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[214] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- D^*(2010)^+ K^+)/\Gamma_{total}$ — Γ_{123}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<0.7	90	[215] AUBERT	03X	BABR	$e^+e^- \to \Upsilon(4S)$

[215] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^- D^+ K^+)/\Gamma_{total}$ — Γ_{124}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
1.5±0.3±0.2	[216] AUBERT	03X	BABR	$e^+e^- \to \Upsilon(4S)$

[216] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^- D^*(2010)^+ K^+)/\Gamma_{total}$ — Γ_{125}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<1.8	90	[217] AUBERT	03X	BABR	$e^+e^- \to \Upsilon(4S)$

[217] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma((\overline{D}+\overline{D}^*)(D+D^*)K)/\Gamma_{total}$ — Γ_{126}/Γ

VALUE (units 10^{-2})	DOCUMENT ID		TECN	COMMENT
3.5±0.3±0.5	[218] AUBERT	03X	BABR	$e^+e^- \to \Upsilon(4S)$

[218] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_s^+\pi^0)/\Gamma_{total}$ — Γ_{127}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.6^{+0.6}_{-0.5}\pm0.1}$		[219] AUBERT	07M	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<16	90	[220] ALEXANDER	93B	CLE2	$e^+e^- \to \Upsilon(4S)$

[219] AUBERT 07M reports $[B(B^+ \to D_s^+\pi^0) \times B(D_s^+ \to \phi\pi^+)] = (7.0^{+2.4}_{-2.1}{}^{+0.6}_{-0.8}) \times 10^{-7}$. We divide by our best value $B(D_s^+ \to \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[220] ALEXANDER 93B reports $< 2.0 \times 10^{-4}$ for $B(D_s^+ \to \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.045$.

$[\Gamma(D_s^+\pi^0) + \Gamma(D_s^{*+}\pi^0)]/\Gamma_{total}$ — $(\Gamma_{127}+\Gamma_{128})/\Gamma$

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0005	90	[221] ALBRECHT	93E	ARG	$e^+e^- \to \Upsilon(4S)$

[221] ALBRECHT 93E reports $< 0.9 \times 10^{-3}$ for $B(D_s^+ \to \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.045$.

$\Gamma(D_s^{*+}\pi^0)/\Gamma_{total}$ — Γ_{128}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.00026	90	[222] ALEXANDER	93B	CLE2	$e^+e^- \to \Upsilon(4S)$

[222] ALEXANDER 93B reports $< 3.2 \times 10^{-4}$ for $B(D_s^+ \to \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \to \phi\pi^+) = 0.045$.

$\Gamma(D_s^+ \eta)/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0004	90	[223] ALEXANDER	93B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[223] ALEXANDER 93B reports $< 4.6 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^{*+} \eta)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0006	90	[224] ALEXANDER	93B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[224] ALEXANDER 93B reports $< 7.5 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.00030	90	[225] ALEXANDER	93B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[225] ALEXANDER 93B reports $< 3.7 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\left[\Gamma(D_s^+ \rho^0) + \Gamma(D_s^+ \overline{K}^*(892)^0)\right]/\Gamma_{\text{total}}$ $(\Gamma_{131}+\Gamma_{141})/\Gamma$

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0020	90	[226] ALBRECHT	93E	ARG	$e^+ e^- \to \Upsilon(4S)$

[226] ALBRECHT 93E reports $< 3.4 \times 10^{-3}$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^{*+} \rho^0)/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0004	90	[227] ALEXANDER	93B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[227] ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\left[\Gamma(D_s^{*+} \rho^0) + \Gamma(D_s^{*+} \overline{K}^*(892)^0)\right]/\Gamma_{\text{total}}$ $(\Gamma_{132}+\Gamma_{142})/\Gamma$

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0012	90	[228] ALBRECHT	93E	ARG	$e^+ e^- \to \Upsilon(4S)$

[228] ALBRECHT 93E reports $< 2.0 \times 10^{-3}$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^+ \omega)/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0004	90	[229] ALEXANDER	93B	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0020	90	[230] ALBRECHT	93E	ARG	$e^+ e^- \to \Upsilon(4S)$

[229] ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

[230] ALBRECHT 93E reports $< 3.4 \times 10^{-3}$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^{*+}\omega)/\Gamma_{total}$ — Γ_{134}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0006	90	[231] ALEXANDER	93B	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0011	90	[232] ALBRECHT	93E	ARG	$e^+e^- \to \Upsilon(4S)$

[231] ALEXANDER 93B reports $< 6.8 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

[232] ALBRECHT 93E reports $< 1.9 \times 10^{-3}$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^+ a_1(1260)^0)/\Gamma_{total}$ — Γ_{135}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0018	90	[233] ALBRECHT	93E	ARG	$e^+e^- \to \Upsilon(4S)$

[233] ALBRECHT 93E reports $< 3.0 \times 10^{-3}$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^{*+} a_1(1260)^0)/\Gamma_{total}$ — Γ_{136}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0013	90	[234] ALBRECHT	93E	ARG	$e^+e^- \to \Upsilon(4S)$

[234] ALBRECHT 93E reports $< 2.2 \times 10^{-3}$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^+\phi)/\Gamma_{total}$ — Γ_{137}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.9 × 10⁻⁶	90	[235] AUBERT	06F	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0010	90	[236] ALBRECHT	93E	ARG	$e^+e^- \to \Upsilon(4S)$
<0.00026	90	[237] ALEXANDER	93B	CLE2	$e^+e^- \to \Upsilon(4S)$

[235] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[236] ALBRECHT 93E reports $< 1.7 \times 10^{-3}$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

[237] ALEXANDER 93B reports $< 3.1 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^{*+}\phi)/\Gamma_{total}$ Γ_{138}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.2 × 10^{-5}	90	[238] AUBERT	06F	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0013	90	[239] ALBRECHT	93E	ARG	$e^+e^- \to \Upsilon(4S)$
<0.00035	90	[240] ALEXANDER	93B	CLE2	$e^+e^- \to \Upsilon(4S)$

[238] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[239] ALBRECHT 93E reports $< 2.1 \times 10^{-3}$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

[240] ALEXANDER 93B reports $< 4.2 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^{+}\overline{K}^0)/\Gamma_{total}$ Γ_{139}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0008	90	[241] ALEXANDER	93B	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0015	90	[242] ALBRECHT	93E	ARG	$e^+e^- \to \Upsilon(4S)$

[241] ALEXANDER 93B reports $< 10.3 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

[242] ALBRECHT 93E reports $< 2.5 \times 10^{-3}$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^{*+}\overline{K}^0)/\Gamma_{total}$ Γ_{140}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0009	90	[243] ALEXANDER	93B	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0019	90	[244] ALBRECHT	93E	ARG	$e^+e^- \to \Upsilon(4S)$

[243] ALEXANDER 93B reports $< 10.9 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

[244] ALBRECHT 93E reports $< 3.1 \times 10^{-3}$ for B($D_s^+ \to \phi\pi^+$) = 0.027. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^{+}\overline{K}^*(892)^0)/\Gamma_{total}$ Γ_{141}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0004	90	[245] ALEXANDER	93B	CLE2	$e^+e^- \to \Upsilon(4S)$

[245] ALEXANDER 93B reports $< 4.4 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^{*+}\overline{K}^*(892)^0)/\Gamma_{total}$ Γ_{142}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.00035	90	[246] ALEXANDER	93B	CLE2	$e^+e^- \to \Upsilon(4S)$

[246] ALEXANDER 93B reports $< 4.3 \times 10^{-4}$ for B($D_s^+ \to \phi\pi^+$) = 0.037. We rescale to our best value B($D_s^+ \to \phi\pi^+$) = 0.045.

$\Gamma(D_s^- \pi^+ K^+)/\Gamma_{total}$ — Γ_{143}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0007	90	[247] ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[247] ALBRECHT 93E reports $< 1.1 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.045$.

$\Gamma(D_s^{*-} \pi^+ K^+)/\Gamma_{total}$ — Γ_{144}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0010	90	[248] ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[248] ALBRECHT 93E reports $< 1.6 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.045$.

$\Gamma(D_s^- \pi^+ K^*(892)^+)/\Gamma_{total}$ — Γ_{145}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.005	90	[249] ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[249] ALBRECHT 93E reports $< 8.6 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.045$.

$\Gamma(D_s^{*-} \pi^+ K^*(892)^+)/\Gamma_{total}$ — Γ_{146}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.007	90	[250] ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[250] ALBRECHT 93E reports $< 1.1 \times 10^{-2}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.045$.

$\Gamma(\eta_c K^+)/\Gamma_{total}$ — Γ_{147}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
0.91±0.13 OUR AVERAGE				
0.87 ± 0.15	[251,252] AUBERT	06E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.4\ ^{+0.3}_{-0.2}\ \pm0.4$	[253] AUBERT,B	05L	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.25\pm0.14\ ^{+0.39}_{-0.40}$	[254] FANG	03	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$0.69\ ^{+0.26}_{-0.21}\pm0.22$	[255] EDWARDS	01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.06\pm0.12\pm0.18$	[252,256] AUBERT,B	04B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[251] Perform measurements of absolute branching fractions using a missing mass technique.

[252] The ratio of $B(B^\pm \rightarrow K^\pm \eta_c)$ $B(\eta_c \rightarrow K\overline{K}\pi) = (7.4 \pm 0.5 \pm 0.7) \times 10^{-5}$ reported in AUBERT,B 04B and $B(B^\pm \rightarrow K^\pm \eta_c) = (8.7 \pm 1.5) \times 10^{-3}$ reported in AUBERT 06E contribute to the determination of $B(\eta_c \rightarrow K\overline{K}\pi)$, which is used by others for normalization.

[253] AUBERT,B 05L reports $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c(1S) \rightarrow p\overline{p})] = (1.8^{+0.3}_{-0.2} \pm 0.2) \times 10^{-6}$. We divide by our best value $B(\eta_c(1S) \rightarrow p\overline{p}) = (1.3 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[254] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[255] EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from B($J/\psi(1S) \to \gamma\eta_c$) in those modes have been accounted for.

[256] AUBERT,B 04B reports [B($B^+ \to \eta_c K^+$) × B($\eta_c(1S) \to K\overline{K}\pi$)] = $(0.074 \pm 0.005 \pm 0.007) \times 10^{-3}$. We divide by our best value B($\eta_c(1S) \to K\overline{K}\pi$) = $(7.0 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta'_c K^+)/\Gamma_{total}$ — Γ_{148}/Γ

VALUE (units 10^{-4})	DOCUMENT ID		TECN	COMMENT
3.4±1.8±0.3	[257] AUBERT	06E	BABR	$e^+e^- \to \Upsilon(4S)$

[257] Perform measurements of absolute branching fractions using a missing mass technique.

$\Gamma(\eta_c K^+)/\Gamma(J/\psi(1S)K^+)$ — $\Gamma_{147}/\Gamma_{149}$

VALUE	DOCUMENT ID		TECN	COMMENT
1.33±0.10±0.43	[258] AUBERT,B	04B	BABR	$e^+e^- \to \Upsilon(4S)$

[258] Uses BABAR measurement of B($B^+ \to J/\psi K^+$) = $(10.1 \pm 0.3 \pm 0.5) \times 10^{-4}$.

$\Gamma(J/\psi(1S)K^+)/\Gamma_{total}$ — Γ_{149}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
10.07± 0.35 OUR FIT					
10.22± 0.35 OUR AVERAGE					
8.1 ± 1.3 ±0.7		[259] AUBERT	06E	BABR	$e^+e^- \to \Upsilon(4S)$
10.61± 0.15±0.48		[260] AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$
10.1 ± 1.0 ±0.3		[261] AUBERT,B	05L	BABR	$e^+e^- \to \Upsilon(4S)$
10.1 ± 0.2 ±0.7		[260] ABE	03B	BELL	$e^+e^- \to \Upsilon(4S)$
10.2 ± 0.8 ±0.7		[260] JESSOP	97	CLE2	$e^+e^- \to \Upsilon(4S)$
9.3 ± 3.1 ±0.1		[262] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
8.1 ± 3.5 ±0.1	6	[263] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
10.1 ± 0.3 ±0.5		[260] AUBERT	02	BABR	Repl. by AUBERT 05J
11.0 ± 1.5 ±0.9	59	[260] ALAM	94	CLE2	Repl. by JESSOP 97
22 ±10 ±2		BUSKULIC	92G	ALEP	$e^+e^- \to Z$
7 ± 4	3	[264] ALBRECHT	87D	ARG	$e^+e^- \to \Upsilon(4S)$
10 ± 7 ±2	3	[265] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$
9 ± 5	3	[266] ALAM	86	CLEO	$e^+e^- \to \Upsilon(4S)$

[259] Perform measurements of absolute branching fractions using a missing mass technique.

[260] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[261] AUBERT,B 05L reports [B($B^+ \to J/\psi(1S)K^+$) × B($J/\psi(1S) \to p\overline{p}$)] = $(2.2 \pm 0.2 \pm 0.1) \times 10^{-6}$. We divide by our best value B($J/\psi(1S) \to p\overline{p}$) = $(2.17 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[262] BORTOLETTO 92 reports $(8 \pm 2 \pm 2) \times 10^{-4}$ for B($J/\psi(1S) \to e^+e^-$) = 0.069 ± 0.009. We rescale to our best value B($J/\psi(1S) \to e^+e^-$) = $(5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[263] ALBRECHT 90J reports $(7 \pm 3 \pm 1) \times 10^{-4}$ for B($J/\psi(1S) \to e^+e^-$) = 0.069 ± 0.009. We rescale to our best value B($J/\psi(1S) \to e^+e^-$) = $(5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[264] ALBRECHT 87D assume $B^+ B^- / B^0 \overline{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.
[265] BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.
[266] ALAM 86 assumes $B^\pm / B^0$ ratio is 60/40.

$\Gamma(J/\psi(1S) K^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{150}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
1.07±0.19 OUR AVERAGE			Error includes scale factor of 1.9. See the ideogram below.			
1.16±0.07±0.09			[267] AUBERT	05R	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.69±0.18±0.12			[268] ACOSTA	02F	CDF	$p\overline{p}$ 1.8 TeV
1.39±0.82±0.01			[269] BORTOLETTO	92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
1.39±0.91±0.01		6	[270] ALBRECHT	87D	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<1.9	90		[271] ALBRECHT	90J	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

[267] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[268] ACOSTA 02F uses as reference of B($B \rightarrow J/\psi(1S) K^+$) = $(10.1 \pm 0.6) \times 10^{-4}$. The second error includes the systematic error and the uncertainties of the branching ratio.
[269] BORTOLETTO 92 reports $(1.2 \pm 0.6 \pm 0.4) \times 10^{-3}$ for B($J/\psi(1S) \rightarrow e^+ e^-$) = 0.069 ± 0.009. We rescale to our best value B($J/\psi(1S) \rightarrow e^+ e^-$) = $(5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[270] ALBRECHT 87D reports $(1.2 \pm 0.8) \times 10^{-3}$ for B($J/\psi(1S) \rightarrow e^+ e^-$) = 0.069 ± 0.009. We rescale to our best value B($J/\psi(1S) \rightarrow e^+ e^-$) = $(5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. They actually report 0.0011 ± 0.0007 assuming $B^+ B^- / B^0 \overline{B}^0$ ratio is 55/45. We rescale to 50/50. Analysis explicitly removes $B^+ \rightarrow \psi(2S) K^+$.
[271] ALBRECHT 90J reports $< 1.6 \times 10^{-3}$ for B($J/\psi(1S) \rightarrow e^+ e^-$) = 0.069. We rescale to our best value B($J/\psi(1S) \rightarrow e^+ e^-$) = 0.0594. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.



$\Gamma\Big(J/\psi(1S)K^+\pi^+\pi^-\Big)/\Gamma_{\text{total}}$ Γ_{150}/Γ

$\Gamma\big(h_c(1P)K^+ \times B(h_c(1P) \to J/\psi\pi^+\pi^-)\big)/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.4 \times 10^{-6}$	90	[272] AUBERT	05R	BABR	$e^+e^- \to \Upsilon(4S)$

[272] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(X(3872)K^+\big)/\Gamma_{\text{total}}$ Γ_{152}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.2 \times 10^{-4}$	90	[273] AUBERT	06E	BABR	$e^+e^- \to \Upsilon(4S)$

[273] Perform measurements of absolute branching fractions using a missing mass technique.

$\Gamma\big(X(3872)K^+ \times B(X \to J/\psi\pi^+\pi^-)\big)/\Gamma_{\text{total}}$ Γ_{153}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
11.4±2.0 OUR AVERAGE				
10.1±2.5±1.0	[274] AUBERT	06	BABR	$e^+e^- \to \Upsilon(4S)$
13.0±2.9±0.7	[275] CHOI	03	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
12.8±4.1	[274] AUBERT	05R	BABR	Repl. by AUBERT 06

[274] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[275] CHOI 03 reports $[B(B^+ \to X(3872)K^+ \times B(X \to J/\psi\pi^+\pi^-))$ / $B(B^+ \to \psi(2S)K^+)] = 0.0200 \pm 0.0038 \pm 0.0023$. We multiply by our best value $B(B^+ \to \psi(2S)K^+) = (6.48 \pm 0.35) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(X(3872)K^+ \times B(X \to J/\psi\gamma))/\Gamma_{total}$ — Γ_{154}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
3.3±1.0±0.3	[276] AUBERT,BE	06M	BABR	$e^+e^- \to \Upsilon(4S)$

[276] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X(3872)K^+ \times B(X(3872) \to D^0\overline{D}^0))/\Gamma_{total}$ — Γ_{155}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<6.0 \times 10^{-5}$	90	[277] CHISTOV	04	BELL	$e^+e^- \to \Upsilon(4S)$

[277] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X(3872)K^+ \times B(X(3872) \to D^+D^-))/\Gamma_{total}$ — Γ_{156}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<4.0 \times 10^{-5}$	90	[278] CHISTOV	04	BELL	$e^+e^- \to \Upsilon(4S)$

[278] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X(3872)K^+ \times B(X(3872) \to D^0\overline{D}^0\pi^0))/\Gamma_{total}$ — Γ_{157}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
$1.02 \pm 0.31^{+0.21}_{-0.29}$		[279] GOKHROO	06	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.6	90	[280] CHISTOV	04	BELL	Repl. by GOKHROO 06

[279] Measure the near-threshold enhancements in the $(D^0\overline{D}^0\pi^0)$ system at a mass $3875.2 \pm 0.7^{+0.3}_{-1.6} \pm 0.8$ MeV/c^2.

[280] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X(3872)K^+ \times B(X(3872) \to J/\psi(1S)\eta))/\Gamma_{total}$ — Γ_{158}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<7.7 \times 10^{-6}$	90	[281] AUBERT	04Y	BABR	$e^+e^- \to \Upsilon(4S)$

[281] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X(3872)^+K^0 \times B(X(3872)^+ \to J/\psi(1S)\pi^+\pi^0))/\Gamma_{total}$ — Γ_{159}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<22	90	[282] AUBERT	05B	BABR	$e^+e^- \to \Upsilon(4S)$

[282] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The isovector-X hypothesis is excluded with a likelihood test at 1×10^{-4} level.

$\Gamma(X(4260)^0K^+ \times B(X^0 \to J/\psi\pi^+\pi^-))/\Gamma_{total}$ — Γ_{160}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<29	95	[283] AUBERT	06	BABR	$e^+e^- \to \Upsilon(4S)$

[283] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X(3945)^0K^+ \times B(X^0 \to J/\psi\gamma))/\Gamma_{total}$ — Γ_{161}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<14	90	[284] AUBERT,BE	06M	BABR	$e^+e^- \to \Upsilon(4S)$

[284] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(Z(3930)^0 K^+ \times B(Z^0 \to J/\psi\gamma))/\Gamma_{total}$ — Γ_{162}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<2.5	90	[285] AUBERT,BE	06M	BABR	$e^+e^- \to \Upsilon(4S)$

[285] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S) K^*(892)^+)/\Gamma_{total}$ — Γ_{163}/Γ

For polarization information see the Listings at the end of the "B^0 Branching Ratios" section.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
1.41 ±0.08 OUR AVERAGE					
1.454±0.047±0.097		[286] AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$
1.28 ±0.07 ±0.14		[286] ABE	02N	BELL	$e^+e^- \to \Upsilon(4S)$
1.41 ±0.23 ±0.24		[286] JESSOP	97	CLE2	$e^+e^- \to \Upsilon(4S)$
1.58 ±0.47 ±0.27		[287] ABE	96H	CDF	$p\overline{p}$ at 1.8 TeV
1.51 ±1.08 ±0.02		[288] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
1.86 ±1.30 ±0.02	2	[289] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.37 ±0.09 ±0.11		[286] AUBERT	02	BABR	Repl. by AUBERT 05J
1.78 ±0.51 ±0.23	13	[286] ALAM	94	CLE2	Sup. by JESSOP 97

[286] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[287] ABE 96H assumes that $B(B^+ \to J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$.
[288] BORTOLETTO 92 reports $(1.3 \pm 0.9 \pm 0.3) \times 10^{-3}$ for $B(J/\psi(1S) \to e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \to e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[289] ALBRECHT 90J reports $(1.6 \pm 1.1 \pm 0.3) \times 10^{-3}$ for $B(J/\psi(1S) \to e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \to e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S) K^*(892)^+)/\Gamma(J/\psi(1S) K^+)$ — $\Gamma_{163}/\Gamma_{149}$

VALUE	DOCUMENT ID		TECN	COMMENT
1.39±0.09 OUR AVERAGE				
1.37±0.05±0.08	AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$
1.45±0.20±0.17	[290] JESSOP	97	CLE2	$e^+e^- \to \Upsilon(4S)$
1.92±0.60±0.17	ABE	96Q	CDF	$p\overline{p}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.37±0.10±0.08	[291] AUBERT	02	BABR	Repl. by AUBERT 05J

[290] JESSOP 97 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The measurement is actually measured as an average over kaon charged and neutral states.
[291] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S) K(1270)^+)/\Gamma_{total}$ — Γ_{164}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
1.80±0.34±0.39	[292] ABE	01L	BELL	$e^+e^- \to \Upsilon(4S)$

[292] Uses the PDG value of $B(B^+ \to J/\psi(1S) K^+) = (1.00 \pm 0.10) \times 10^{-3}$.

$\Gamma(J/\psi(1S)K(1400)^+)/\Gamma(J/\psi(1S)K(1270)^+)$ — $\Gamma_{165}/\Gamma_{164}$

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.30	90	ABE	01L	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(J/\psi(1S)\eta K^+)/\Gamma_{total}$ — Γ_{166}/Γ

VALUE (units 10^{-5})	DOCUMENT ID		TECN	COMMENT
10.8±2.3±2.4	[293] AUBERT	04Y	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[293] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\phi K^+)/\Gamma_{total}$ — Γ_{167}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$(5.2\pm1.7) \times 10^{-5}$ OUR AVERAGE Error includes scale factor of 1.2.				
$(4.4\pm1.4\pm0.5) \times 10^{-5}$	[294] AUBERT	03O	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$(8.8^{+3.5}_{-3.0}\pm1.3) \times 10^{-5}$	[295] ANASTASSOV	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[294] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[295] ANASTASSOV 00 finds 10 events on a background of 0.5 ± 0.2. Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$, a uniform Dalitz plot distribution, isotropic $J/\psi(1S)$ and ϕ decays, and $B(B^+ \rightarrow J/\psi(1S)\phi K^+) = B(B^0 \rightarrow J/\psi(1S)\phi K^0)$.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma_{total}$ — Γ_{168}/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
$(4.9\pm0.6) \times 10^{-5}$ OUR FIT Error includes scale factor of 1.5.				
$(3.8\pm0.6\pm0.3) \times 10^{-5}$	[296] ABE	03B	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[296] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma(J/\psi(1S)K^+)$ — $\Gamma_{168}/\Gamma_{149}$

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.049 ±0.006 OUR FIT Error includes scale factor of 1.5.					
0.053 ±0.004 OUR AVERAGE					
$0.0537\pm0.0045\pm0.0011$		AUBERT	04P	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.05\ ^{+0.019}_{-0.017}\pm0.001$		ABE	96R	CDF	$p\overline{p}$ 1.8 TeV
0.052 ± 0.024		BISHAI	96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.0391\pm0.0078\pm0.0019$		AUBERT	02F	BABR	Repl. by AUBERT 04P
0.043 ± 0.023	5	[297] ALEXANDER	95	CLE2	Sup. by BISHAI 96

[297] Assumes equal production of B^+B^- and $B^0\overline{B}^0$ on $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\rho^+)/\Gamma_{total}$ — Γ_{169}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<7.7 \times 10^{-4}$	90	BISHAI	96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(J/\psi(1S)a_1(1260)^+)/\Gamma_{total}$ — Γ_{170}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.2 \times 10^{-3}$	90	BISHAI	96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(J/\psi(1S)p\overline{\Lambda})/\Gamma_{total}$ Γ_{171}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
11.8±3.1 OUR AVERAGE					
$11.7\pm2.8^{+1.8}_{-2.3}$		[298] XIE	05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$12\ ^{+9}_{-6}$		[298] AUBERT	03K	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<41	90	ZANG	04	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[298] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\overline{\Sigma}^0 p)/\Gamma_{total}$ Γ_{172}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.1 × 10⁻⁵	90	[299] XIE	05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[299] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)D^+)/\Gamma_{total}$ Γ_{173}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<12	90	[300] AUBERT	05U	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[300] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\overline{D}^0\pi^+)/\Gamma_{total}$ Γ_{174}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<2.5	90	[301] ZHANG	05B	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<5.2	90	[301] AUBERT	05R	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[301] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S)K^+)/\Gamma_{total}$ Γ_{175}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
6.48± 0.35 OUR AVERAGE						
4.9 ± 1.6 ±0.4			[302] AUBERT	06E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
6.17± 0.32±0.44			[303] AUBERT	05J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
6.9 ± 0.6			[303] ABE	03B	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
7.8 ± 0.7 ±0.9			[303] RICHICHI	01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5.5 ± 1.0 ±0.6			[304] ABE	98O	CDF	$p\overline{p}$ 1.8 TeV
18 ± 8 ±4		5	[303] ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
6.4 ± 0.5 ±0.8			[303] AUBERT	02	BABR	Repl. by AUBERT 05J
6.1 ± 2.3 ±0.9		7	[303] ALAM	94	CLE2	Repl. by RICHICHI 01
<5	90		[303] BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
22 ±17		3	[305] ALBRECHT	87D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[302] Perform measurements of absolute branching fractions using a missing mass technique.
[303] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[304] ABE 98O reports $[B(B^+ \rightarrow \psi(2S)K^+)]/[B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.558 \pm 0.082 \pm 0.056$. We multiply by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (9.9 \pm 1.0) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
[305] ALBRECHT 87D assume $B^+B^-/B^0\overline{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.

$\Gamma(\psi(2S)K^+)/\Gamma(J/\psi(1S)K^+)$ — $\Gamma_{175}/\Gamma_{149}$

VALUE	DOCUMENT ID		TECN	COMMENT
0.64±0.06±0.07	[306] AUBERT	02	BABR	$e^+e^- \to \Upsilon(4S)$

[306] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S)K^*(892)^+)/\Gamma_{total}$ — Γ_{176}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
6.7 ±1.4 OUR AVERAGE		Error includes scale factor of 1.3.			
5.92±0.85±0.89		[307] AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$
9.2 ±1.9 ±1.2		[307] RICHICHI	01	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<30	90	[307] ALAM	94	CLE2	Repl. by RICHICHI 01
<35	90	[307] BORTOLETTO	92	CLEO	$e^+e^- \to \Upsilon(4S)$
<49	90	[307] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$

[307] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S)K^*(892)^+)/\Gamma(\psi(2S)K^+)$ — $\Gamma_{176}/\Gamma_{175}$

VALUE	DOCUMENT ID		TECN	COMMENT
0.96±0.15±0.09	AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$

$\Gamma(\psi(2S)K^+\pi^+\pi^-)/\Gamma_{total}$ — Γ_{177}/Γ

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0019±0.0011±0.0004	3	[308] ALBRECHT	90J	ARG	$e^+e^- \to \Upsilon(4S)$

[308] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(3770)K^+)/\Gamma_{total}$ — Γ_{178}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
0.49±0.13 OUR AVERAGE				
3.5 ±2.5 ±0.3	[309] AUBERT	06E	BABR	$e^+e^- \to \Upsilon(4S)$
0.48±0.11±0.07	[310] CHISTOV	04	BELL	$e^+e^- \to \Upsilon(4S)$

[309] Perform measurements of absolute branching fractions using a missing mass technique.
[310] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(3770)K^+ \times B(\psi(3770) \to D^0\overline{D}^0))/\Gamma_{total}$ — Γ_{179}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
0.34±0.08±0.05	[311] CHISTOV	04	BELL	$e^+e^- \to \Upsilon(4S)$

[311] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(3770)K^+ \times B(\psi(3770) \to D^+D^-K^+))/\Gamma_{total}$ — Γ_{180}/Γ

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
0.14±0.08±0.02	[312] CHISTOV	04	BELL	$e^+e^- \to \Upsilon(4S)$

[312] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c0}\pi^+ \times B(\chi_{c0} \to \pi^+\pi^-))/\Gamma_{total}$ — Γ_{181}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<0.3	90	[313] AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$

[313] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c0}(1P)K^+)/\Gamma_{\text{total}}$ — Γ_{182}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$1.40^{+0.23}_{-0.19}$ OUR AVERAGE				
$1.84\pm0.32\pm0.31$		[314,315] AUBERT	06O	BABR	$e^+e^- \to \Upsilon(4S)$
$4.6 \pm2.1 \pm0.4$		[316] AUBERT,BE	06M	BABR	$e^+e^- \to \Upsilon(4S)$
$1.12\pm0.12^{+0.30}_{-0.20}$		[315] GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$1.39\pm0.49\pm0.11$		[317] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.8	90	[318] AUBERT	06E	BABR	$e^+e^- \to \Upsilon(4S)$
<8.9	90	[315] AUBERT	05K	BABR	$e^+e^- \to \Upsilon(4S)$
$1.96\pm0.35^{+2.00}_{-0.42}$		[315] GARMASH	05	BELL	Repl. by GARMASH 06
2.7 ± 0.7		[319] AUBERT	04T	BABR	Repl. by AUBERT,B 04P
$3.0 \pm0.8 \pm0.3$		[320] AUBERT,B	04P	BABR	Repl. by AUBERT,B 05N
$6.0\ ^{+2.1}_{-1.8} \pm1.1$		[321] ABE	02B	BELL	Repl. by GARMASH 05
<4.8	90	[322] EDWARDS	01	CLE2	$e^+e^- \to \Upsilon(4S)$

[314] Measured in the $B^+ \to K^+K^-K^+$ decay.
[315] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[316] AUBERT,BE 06M reports $[B(B^+ \to \chi_{c0}(1P)K^+) \times B(\chi_{c0}(1P) \to \gamma J/\psi(1S))] = (6.1 \pm 2.6 \pm 1.1) \times 10^{-6}$. We divide by our best value $B(\chi_{c0}(1P) \to \gamma J/\psi(1S)) = (1.32 \pm 0.11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The significance of the observed signal is 2.4 σ.
[317] AUBERT,B 05N reports $(0.66 \pm 0.22 \pm 0.08) \times 10^{-6}$ for $B(B^+ \to \chi^0_c K^+) \times B(\chi^0_c \to \pi^+\pi^-)$. We compute $B(B^+ \to \chi^0_c K^+)$ using the PDG value $B(\chi^0_c \to \pi^+\pi^-) = (7.1 \pm 0.6) \times 10^{-3}$ and 2/3 for the $\pi^+\pi^-$ fraction.
[318] Perform measurements of absolute branching fractions using a missing mass technique.
[319] The measurement performed using decay channels $\chi^0_c \to \pi^+\pi^-$ and $\chi^0_c \to K^+K^-$. The ratio of the branching ratios for these channels is found to be consistent with world average.
[320] AUBERT 04P reports $B(B^+ \to \chi^0_c K^+) \times B(\chi^0_c \to \pi^+\pi^-) = (1.5 \pm 0.4 \pm 0.1) \times 10^{-6}$ and used PDG value of $B(\chi^0_c \to \pi\pi) = (7.4 \pm 0.8) \times 10^{-3}$ and Clebsh-Gordan coefficient to compute $B(B^\pm - > \chi^0_c K^+)$.
[321] ABE 02B measures the ratio of $B(B^+ \to \chi^0_c K^+)/B(B^+ \to J/\psi(1S)K^+) = 0.60 + 0.21 - 0.18 \pm 0.05 \pm 0.08$, where the third error is due to the uncertainty in the $B(\chi^0_c \to \pi^+\pi^-)$, and uses $B(B^+ \to J/\psi(1S)K^+) = (10.0 \pm 1.0) \times 10^{-4}$ to obtain the result.
[322] EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \to \gamma\eta_c)$ in those modes have been accounted for.

$\Gamma(\chi_{c0}K^*(892)^+)/\Gamma_{\text{total}}$ — Γ_{183}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.86 \times 10^{-3}$	90	[323] AUBERT	05K	BABR	$e^+e^- \to \Upsilon(4S)$

[323] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c2} K^+)/\Gamma_{total}$ Γ_{184}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.9 \times 10^{-5}$	90	[324] SONI	06	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.0 \times 10^{-4}$	90	[325] AUBERT	06E	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<3.0 \times 10^{-5}$	90	[324] AUBERT	05K	BABR	Repl. by AUBERT 06E

[324] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[325] Perform measurements of absolute branching fractions using a missing mass technique.

$\Gamma(\chi_{c2} K^*(892)^+)/\Gamma_{total}$ Γ_{185}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.2 \times 10^{-5}$	90	[326] AUBERT	05K	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.27 \times 10^{-4}$	90	[326] SONI	06	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[326] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P)\pi^+)/\Gamma_{total}$ Γ_{186}/Γ

VALUE (units 10^{-5})	DOCUMENT ID		TECN	COMMENT
2.2±0.4±0.3	[327] KUMAR	06	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[327] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P) K^+)/\Gamma_{total}$ Γ_{187}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
4.9 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.			
8.1 ± 1.4 ±0.7		[328] AUBERT	06E	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
4.9 ± 0.4 ±0.3		[329] AUBERT,BE	06M	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
4.49± 0.19±0.53		[330] SONI	06	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
15.5 ± 5.4 ±2.0		[331] ACOSTA	02F	CDF	$p\overline{p}$ 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
5.79± 0.26±0.65		[330] AUBERT	05J	BABR	Repl. by AUBERT,BE 06M
5.7 ± 0.9 ±0.3		[332] AUBERT	02	BABR	Repl. by AUBERT 05J
9.7 ± 4.0 ±0.9	6	[330] ALAM	94	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
19 ±13 ±6		[333] ALBRECHT	92E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

[328] Perform measurements of absolute branching fractions using a missing mass technique.
[329] AUBERT,BE 06M reports $[B(B^+ \rightarrow \chi_{c1}(1P) K^+) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))] = (1.76 \pm 0.07 \pm 0.12) \times 10^{-4}$. We divide by our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (35.9 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
[330] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[331] ACOSTA 02F uses as reference of $B(B \rightarrow J/\psi(1S) K^+) = (10.1 \pm 0.6) \times 10^{-4}$. The second error includes the systematic error and the uncertainties of the branching ratio.
[332] AUBERT 02 reports $(7.5 \pm 0.9 \pm 0.8) \times 10^{-4}$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (35.9 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[333] ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production and $B(\Upsilon(4S) \rightarrow B^+ B^-) = 50\%$.



$\Gamma\big(\chi_{c1}(1P)K^+\big)/\Gamma_{\text{total}}$ Γ_{187}/Γ

$\Gamma\big(\chi_{c1}(1P)K^+\big)/\Gamma\big(J/\psi(1S)K^+\big)$ $\Gamma_{187}/\Gamma_{149}$

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
0.57±0.06±0.03	[334] AUBERT	02	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[334] AUBERT 02 reports $0.75 \pm 0.08 \pm 0.05$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (35.9 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\chi_{c1}(1P)\pi^+\big)/\Gamma\big(\chi_{c1}(1P)K^+\big)$ $\Gamma_{186}/\Gamma_{187}$

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
0.043±0.008±0.003	[335] KUMAR	06	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[335] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\chi_{c1}(1P)K^*(892)^+\big)/\Gamma_{\text{total}}$ Γ_{188}/Γ

VALUE (units 10^{-4})	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
3.6 ±0.9 OUR AVERAGE					
4.05±0.59±0.95		[336] SONI	06	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
2.94±0.95±0.98		[336] AUBERT	05J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<21	90	[336] ALAM	94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[336] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\chi_{c1}(1P)K^*(892)^+\big)/\Gamma\big(\chi_{c1}(1P)K^+\big)$ $\Gamma_{188}/\Gamma_{187}$

VALUE	*DOCUMENT ID*		*TECN*	*COMMENT*
0.51±0.17±0.16	AUBERT	05J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(h_c K^+)/\Gamma_{total}$ — Γ_{189}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID		TECN	COMMENT
<3.8	90	[337] FANG	06	BELL	$e^+e^- \to \Upsilon(4S)$

[337] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and B($h_c \to \eta_c\gamma$) = 50%.

$\Gamma(K^0\pi^+)/\Gamma_{total}$ — Γ_{190}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
23.0± 1.2 OUR AVERAGE					
$23.9 \pm 1.1 \pm 1.0$		[338] AUBERT,BE	06C	BABR	$e^+e^- \to \Upsilon(4S)$
$22.0 \pm 1.9 \pm 1.1$		[338] CHAO	04	BELL	$e^+e^- \to \Upsilon(4S)$
$18.8^{+3.7+2.1}_{-3.3-1.8}$		[338] BORNHEIM	03	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$26.0 \pm 1.3 \pm 1.0$		[338] AUBERT,BE	05E	BABR	Repl. by AUBERT,BE 06C
$22.3 \pm 1.7 \pm 1.1$		[338] AUBERT	04M	BABR	Repl. by AUBERT,BE 05E
$19.4^{+3.1}_{-3.0} \pm 1.6$		[338] CASEY	02	BELL	Repl. by CHAO 04
$13.7^{+5.7+1.9}_{-4.8-1.8}$		[338] ABE	01H	BELL	Repl. by CASEY 02
$18.2^{+3.3}_{-3.0} \pm 2.0$		[338] AUBERT	01E	BABR	Repl. by AUBERT 04M
$18.2^{+4.6}_{-4.0} \pm 1.6$		[338] CRONIN-HEN...	00	CLE2	Repl. by BORNHEIM 03
$23\ ^{+11}_{-10} \pm 3.6$		GODANG	98	CLE2	Repl. by CRONIN-HENNESSY 00
< 48	90	ASNER	96	CLE2	Repl. by GODANG 98
<190	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$
<100	90	[339] AVERY	89B	CLEO	$e^+e^- \to \Upsilon(4S)$
<680	90	AVERY	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[338] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[339] AVERY 89B reports $< 9 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^+\pi^0)/\Gamma_{total}$ — Γ_{191}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
12.1±0.8 OUR AVERAGE					
$12.0 \pm 0.7 \pm 0.6$		[340] AUBERT	05L	BABR	$e^+e^- \to \Upsilon(4S)$
$12.0 \pm 1.3^{+1.3}_{-0.9}$		[340] CHAO	04	BELL	$e^+e^- \to \Upsilon(4S)$
$12.9^{+2.4+1.2}_{-2.2-1.1}$		[340] BORNHEIM	03	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$12.8^{+1.2}_{-1.1} \pm 1.0$		[340] AUBERT	03L	BABR	Repl. by AUBERT 05L
$13.0^{+2.5}_{-2.4} \pm 1.3$		[340] CASEY	02	BELL	Repl. by CHAO 04
$16.3^{+3.5+1.6}_{-3.3-1.8}$		[340] ABE	01H	BELL	Repl. by CASEY 02
$10.8^{+2.1}_{-1.9} \pm 1.0$		[340] AUBERT	01E	BABR	Repl. by AUBERT 03L

$11.6^{+3.0+1.4}_{-2.7-1.3}$		[340] CRONIN-HEN...00		CLE2	Repl. by BORNHEIM 03
<16	90	GODANG	98	CLE2	Repl. by CRONIN-HENNESSY 00
<14	90	ASNER	96	CLE2	Repl. by GODANG 98

[340] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+\pi^0)/\Gamma(K^0\pi^+)$ $\Gamma_{191}/\Gamma_{190}$

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{2.38^{+0.98+0.39}_{-1.10-0.26}}$	ABE	01H	BELL	$e^+e^- \to \Upsilon(4S)$

$\Gamma(\eta' K^+)/\Gamma_{total}$ Γ_{192}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
69.7± 2.8 OUR AVERAGE				
$69.2\pm 2.2\pm3.7$	[341] SCHUEMANN	06	BELL	$e^+e^- \to \Upsilon(4S)$
$68.9\pm 2.0\pm3.2$	[341] AUBERT	05M	BABR	$e^+e^- \to \Upsilon(4S)$
$80\ ^{+10}_{-\ 9}\ \pm7$	[341] RICHICHI	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$76.9\pm 3.5\pm4.4$	[341] AUBERT	03W	BABR	Repl. by AUBERT 05M
$79\ ^{+12}_{-11}\ \pm9$	[341] ABE	01M	BELL	Repl. by SCHUEMANN 06
$70\ \pm\ 8\ \pm5$	[341] AUBERT	01G	BABR	Repl. by AUBERT 03W
$65\ ^{+15}_{-14}\ \pm9$	BEHRENS	98	CLE2	Repl. by RICHICHI 00

[341] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' K^*(892)^+)/\Gamma_{total}$ Γ_{193}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{4.9^{+1.9}_{-1.7}\pm0.8}$		[342] AUBERT	07E	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<14	90	[342] AUBERT,B	04D	BABR	Repl. by AUBERT 07E
<35	90	[342] RICHICHI	00	CLE2	$e^+e^- \to \Upsilon(4S)$
<13	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

[342] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K^+)/\Gamma_{total}$ Γ_{194}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
2.6±0.6 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			
$3.3\pm0.6\pm0.3$		[343] AUBERT,B	05K	BABR	$e^+e^- \to \Upsilon(4S)$
$2.1\pm0.6\pm0.2$		[343] CHANG	05A	BELL	$e^+e^- \to \Upsilon(4S)$
$2.2^{+2.8}_{-2.2}$		[343] RICHICHI	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$3.4\pm0.8\pm0.2$		[343] AUBERT	04H	BABR	Repl. by AUBERT,B 05K
<14	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

[343] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.



$\Gamma\left(\eta K^+\right)/\Gamma_{\text{total}}$ Γ_{194}/Γ

$\Gamma\left(\eta K^*(892)^+\right)/\Gamma_{\text{total}}$ Γ_{195}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
19.3±2.2 OUR AVERAGE					
$18.9\pm1.8\pm1.3$		[344] AUBERT,B	06H	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$26.4^{+9.6}_{-8.2}\pm3.3$		[344] RICHICHI	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$25.6\pm4.0\pm2.4$		[344] AUBERT,B	04D	BABR	Repl. by AUBERT,B 06H
<30	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

[344] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(\eta K_0^*(1430)^+\right)/\Gamma_{\text{total}}$ Γ_{196}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
18.2±2.6±2.6	[345] AUBERT,B	06H	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[345] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\left(\eta K_2^*(1430)^+\right)/\Gamma_{\text{total}}$ Γ_{197}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
9.1±2.7±1.4	[346] AUBERT,B	06H	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[346] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega K^+)/\Gamma_{total}$ Γ_{198}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
6.8±0.9 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.			
$6.1 \pm 0.6 \pm 0.4$		[347] AUBERT,B 06E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$8.1 \pm 0.6 \pm 0.6$		[347] JEN 06	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$3.2^{+2.4}_{-1.9} \pm 0.8$		[347] JESSOP 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$4.8 \pm 0.8 \pm 0.4$		[347] AUBERT 04H	BABR	Repl. by AUBERT,B 06E
$6.5^{+1.3}_{-1.2} \pm 0.6$		[347] WANG 04A	BELL	Repl. by JEN 06
$9.2^{+2.6}_{-2.3} \pm 1.0$		[347] LU 02	BELL	Repl. by WANG 04A
<4	90	[347] AUBERT 01G	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.5^{+7}_{-6} \pm 2$		[347] BERGFELD 98	CLE2	Repl. by JESSOP 00

[347] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.



$\Gamma(\omega K^*(892)^+)/\Gamma_{total}$ Γ_{199}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.4	90	[348] AUBERT,B 06T	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 7.4	90	[348] AUBERT 05O	BABR	Repl. by AUBERT,B 06T
<87	90	[348] BERGFELD 98	CLE2	

[348] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_0^0 K^+)/\Gamma_{total}$ — Γ_{201}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<2.5	90	[349] AUBERT,BE	04	BABR	$e^+e^- \to \Upsilon(4S)$

[349] Assumes equal production of charged and neutral B mesons from $\Upsilon(4S)$ decays.

$\Gamma(a_0^+ K^0)/\Gamma_{total}$ — Γ_{200}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<3.9	90	[350] AUBERT,BE	04	BABR	$e^+e^- \to \Upsilon(4S)$

[350] Assumes equal production of charged and neutral B mesons from $\Upsilon(4S)$ decays.

$\Gamma(K^*(892)^0\pi^+)/\Gamma_{total}$ — Γ_{202}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
10.9 ±1.8 OUR AVERAGE Error includes scale factor of 2.1.					
$9.67\pm0.64^{+0.81}_{-0.89}$		[351] GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$13.5\pm1.2^{+0.8}_{-0.9}$		[351] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$9.8\pm0.9^{+1.1}_{-1.2}$		[351] GARMASH	05	BELL	Repl. by GARMASH 06
$15.5\pm1.8^{+1.5}_{-4.0}$		[351,352] AUBERT,B	04P	BABR	Repl. by AUBERT,B 05N
$19.4^{+4.2}_{-3.9}{}^{+4.1}_{-7.1}$		[353] GARMASH	02	BELL	Repl. by GARMASH 05
<119	90	[354] ABE	00C	SLD	$e^+e^- \to Z$
< 16	90	[351] JESSOP	00	CLE2	$e^+e^- \to \Upsilon(4S)$
<390	90	[355] ADAM	96D	DLPH	$e^+e^- \to Z$
< 41	90	ASNER	96	CLE2	Repl. by JESSOP 00
<480	90	[355] ABREU	95N	DLPH	Sup. by ADAM 96D
<170	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$
<150	90	[356] AVERY	89B	CLEO	$e^+e^- \to \Upsilon(4S)$
<260	90	AVERY	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[351] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[352] AUBERT 04P also report a branching ratio for $B^+ \to$ "higher $K*$ resonances" π^+, $K* \to K^+\pi^-$, $(25.1 \pm 2.0^{+11.0}_{-\ 5.7}) \times 10^{-6}$.
[353] Uses a reference decay mode $B^+ \to \overline{D}^0\pi^+$ and $\overline{D}^0 \to K^+\pi^-$ with $B(B^+ \to \overline{D}^0\pi^+)\cdot B(\overline{D}^0 \to K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.
[354] ABE 00C assumes $B(Z \to b\overline{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$.
[355] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.
[356] AVERY 89B reports $< 1.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^+\pi^0)/\Gamma_{total}$ — Γ_{203}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
6.9±2.0±1.3		[357] AUBERT	05X	BABR	$e^+e^- -> \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<31	90	[357] JESSOP	00	CLE2	$e^+e^- \to \Upsilon(4S)$
<99	90	ASNER	96	CLE2	Repl. by JESSOP 00

[357] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+\pi^-\pi^+)/\Gamma_{total}$ — Γ_{204}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
55 ±7 OUR AVERAGE	Error includes scale factor of 2.6.			
$48.8\pm1.1\pm3.6$	[358] GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$64.1\pm2.4\pm4.0$	[358] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$46.6\pm2.1\pm4.3$	[358] GARMASH	05	BELL	Repl. by GARMASH 06
$53.6\pm3.1\pm5.1$	[358] GARMASH	04	BELL	Repl. by GARMASH 05
$59.1\pm3.8\pm3.2$	[359] AUBERT	03M	BABR	Repl. by AUBERT,B 05N
$55.6\pm5.8\pm7.7$	[360] GARMASH	02	BELL	Repl. by GARMASH 04

[358] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[359] Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

[360] Uses a reference decay mode $B^+ \to \overline{D}^0\pi^+$ and $\overline{D}^0 \to K^+\pi^-$ with $B(B^+ \to \overline{D}^0\pi^+)\cdot B(\overline{D}^0 \to K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

$\Gamma(K^+\pi^-\pi^+\text{ nonresonant})/\Gamma_{total}$ — Γ_{205}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$6\ ^{+6}_{-4}$ OUR AVERAGE		Error includes scale factor of 6.1.			
$16.9\pm1.3^{+\ 1.7}_{-\ 1.6}$		[361] GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$2.9\pm0.6^{+\ 0.8}_{-\ 0.5}$		[361] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$17.3\pm1.7^{+17.2}_{-\ 8.0}$		[361] GARMASH	05	BELL	Repl. by GARMASH 06
$<$ 17	90	[361] AUBERT,B	04P	BABR	Repl. by AUBERT,B 05N
$<$330	90	[362] ADAM	96D	DLPH	$e^+e^- \to Z$
$<$ 28	90	BERGFELD	96B	CLE2	$e^+e^- \to \Upsilon(4S)$
$<$400	90	[362] ABREU	95N	DLPH	Sup. by ADAM 96D
$<$330	90	ALBRECHT	91E	ARG	$e^+e^- \to \Upsilon(4S)$
$<$190	90	[363] AVERY	89B	CLEO	$e^+e^- \to \Upsilon(4S)$

[361] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[362] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

[363] AVERY 89B reports $< 1.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^+ f_0(980) \times B(f_0 \to \pi^+\pi^-))/\Gamma_{total}$ — Γ_{206}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$9.2\ ^{+0.8}_{-1.1}$ OUR AVERAGE					
$8.78\pm0.82^{+0.85}_{-1.76}$		[364] GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$9.47\pm0.97^{+0.62}_{-0.88}$		[364] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.55\pm1.24^{+1.63}_{-1.18}$		[364] GARMASH	05	BELL	Repl. by GARMASH 06
$9.2\ \pm1.2\ ^{+2.1}_{-2.6}$		[365] AUBERT,B	04P	BABR	Repl. by AUBERT,B 05N
$9.6\ ^{+2.5}_{-2.3}\ ^{+3.7}_{-1.7}$		[366] GARMASH	02	BELL	Repl. by GARMASH 05
<80	90	[367] AVERY	89B	CLEO	$e^+e^- \to \Upsilon(4S)$

[364] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[365] AUBERT,B 04P also reports B($B^+ \to$ "higher f^0 resonances" π^+, $f(980)^0 \to \pi^+\pi^-$) $= (3.2 \pm 1.2^{+6.0}_{-2.9}) \times 10^{-6}$.
[366] Uses a reference decay mode $B^+ \to \overline{D}{}^0\pi^+$ and $\overline{D}{}^0 \to K^+\pi^-$ with B($B^+ \to \overline{D}{}^0\pi^+$)×B($\overline{D}{}^0 \to K^+\pi^-$) $= (20.3 \pm 2.0) \times 10^{-5}$. Only charged pions from the $f_0(980)$ are used.
[367] AVERY 89B reports $< 7\times10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}{}^0$. We rescale to 50%.

$\Gamma\big(f_2(1270)^0 K^+\big)/\Gamma_{\rm total}$ Γ_{207}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.33\pm0.30^{+0.23}_{-0.34}}$		[368] GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<16	90	[369] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$
$<\ 2.3$	90	[370] GARMASH	05	BELL	Repl. by GARMASH 06

[368] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[369] AUBERT,B 05N reports 8.9×10^{-6} at 90% CL for B($B^+ \to f_2(1270)\pi^+$) × B($f_2(1270) \to \pi^+\pi^-$). We rescaled it using the PDG value B($f_2(1270) \to \pi\pi$) = 84.7% and 2/3 for the $K^+\pi^-$ fraction.
[370] GARMASH 05 reports 1.3×10^{-6} at 90% CL for B($B^+ \to f_2(1270)\pi^+$) × B($f_2(1270) \to \pi^+\pi^-$). We rescaled it using the PDG value B($f_2(1270) \to \pi\pi$) = 84.7% and 2/3 for the $\pi^+\pi^-$ fraction.

$\Gamma\big(f_0(1370)^0 K^+ \times {\rm B}(f_0(1370)^0 \to \pi^+\pi^-)\big)/\Gamma_{\rm total}$ Γ_{208}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<10.7 \times 10^{-6}}$	90	[371] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$

[371] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(\rho^0(1450) K^+ \times {\rm B}(\rho^0(1450) \to \pi^+\pi^-)\big)/\Gamma_{\rm total}$ Γ_{209}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<11.7 \times 10^{-6}}$	90	[372] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$

[372] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma\big(f_0(1500) K^+ \times {\rm B}(f_0(1500) \to \pi^+\pi^-)\big)/\Gamma_{\rm total}$ Γ_{210}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<4.4 \times 10^{-6}}$	90	[373] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$

[373] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_2'(1525)K^+ \times B(f_2'(1525) \to \pi^+\pi^-))/\Gamma_{total}$ Γ_{211}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<3.4 × 10^{-6}	90	[374] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$

[374] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+\rho^0)/\Gamma_{total}$ Γ_{212}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
4.2 ±0.5 OUR AVERAGE					
$3.89\pm0.47^{+0.43}_{-0.41}$		[375] GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$5.07\pm0.75^{+0.55}_{-0.88}$		[375] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$4.78\pm0.75^{+1.01}_{-0.97}$		[375] GARMASH	05	BELL	Repl. by GARMASH 06
< 6.2	90	[376] AUBERT,B	04P	BABR	Repl. by AUBERT,B 05N
< 12	90	[377] GARMASH	02	BELL	$e^+e^- \to \Upsilon(4S)$
< 86	90	[378] ABE	00C	SLD	$e^+e^- \to Z$
< 17	90	[375] JESSOP	00	CLE2	$e^+e^- \to \Upsilon(4S)$
<120	90	[379] ADAM	96D	DLPH	$e^+e^- \to Z$
< 19	90	ASNER	96	CLE2	Repl. by JESSOP 00
<190	90	[379] ABREU	95N	DLPH	Sup. by ADAM 96D
<180	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$
< 80	90	[380] AVERY	89B	CLEO	$e^+e^- \to \Upsilon(4S)$
<260	90	AVERY	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[375] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[376] AUBERT 04P reports a central value of $(3.9\pm1.2^{+1.3}_{-3.5})\times10^{-6}$ for this branching ratio.
[377] Uses a reference decay mode $B^+ \to \overline{D}^0\pi^+$ and $\overline{D}^0 \to K^+\pi^-$ with $B(B^+ \to \overline{D}^0\pi^+)\cdot B(\overline{D}^0 \to K^+\pi^-) = (20.3\pm2.0)\times10^{-5}$.
[378] ABE 00C assumes $B(Z \to b\overline{b})$=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.
[379] Assumes production fractions $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.
[380] AVERY 89B reports $< 7\times10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K_0^*(1430)^0\pi^+)/\Gamma_{total}$ Γ_{213}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
47 ±5 OUR AVERAGE				
$51.6\pm1.7^{+\ 7.0}_{-\ 7.5}$	[381] GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$44.4\pm2.2\pm\ 5.3$	[381,382] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$45.0\pm2.9^{+15.0}_{-10.7}$	[381] GARMASH	05	BELL	Repl. by GARMASH 06

[381] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[382] See erratum: AUBERT,BE 06A.

$\Gamma(K_2^*(1430)^0\pi^+)/\Gamma_{total}$ Γ_{214}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
< 6.9	90	[383] GARMASH	05	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 23	90	[384] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$
<680	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$

[383] GARMASH 05 reports 2.3×10^{-6} at 90% CL for B($B^+ \to K_2^*(1430)^0\pi^+$) × B($K_2^*(1430)^0 \to K^+\pi^-$). We rescaled it using the PDG value B($K_2^*(1430)^0 \to K\pi$) = 49.9% and 2/3 for the $K^+\pi^-$ mode.

[384] AUBERT,B 05N reports 7.7×10^{-6} at 90% CL for B($B^+ \to K_2^*(1430)^0\pi^+$) × B($K_2^*(1430)^0 \to K^+\pi^-$). We rescaled it using the PDG value B($K_2^*(1430)^0 \to K\pi$) = 49.9% and 2/3 for the $K^+\pi^-$ fraction.

$\Gamma(K^*(1410)^0\pi^+)/\Gamma_{total}$ Γ_{215}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<45	90	[385] GARMASH	05	BELL	$e^+e^- \to \Upsilon(4S)$

[385] GARMASH 05 reports 2.0×10^{-6} at 90% CL for B($B^+ \to K^*(1410)^0\pi^+$) × B($K^*(1410)^0 \to K^+\pi^-$). We rescaled it using the PDG value B($K^*(1410)^0 \to K\pi$) = 6.6% and 2/3 for the $K^+\pi^-$ mode.

$\Gamma(K^*(1680)^0\pi^+)/\Gamma_{total}$ Γ_{216}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<12	90	[386] GARMASH	05	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<15	90	[387] AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$

[386] GARMASH 05 reports 3.1×10^{-6} at 90% CL for B($B^+ \to K^*(1680)^0\pi^+$) × B($K^*(1680)^0 \to K^+\pi^-$). We rescaled it using the PDG value B($K^*(1680)^0 \to K\pi$) = 38.7% and 2/3 for the $K^+\pi^-$ mode.

[387] AUBERT,B 05N reports 3.8×10^{-6} at 90% CL for B($B^+ \to K^*(1680)^0\pi^+$) × B($K^*(1680)^0 \to K^+\pi^-$). We rescaled it using the PDG value B($K^*(1680)^0 \to K\pi$) = 38.7% and 2/3 for the $K^+\pi^-$ fraction.

$\Gamma(K^-\pi^+\pi^+)/\Gamma_{total}$ Γ_{217}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<1.8	90	[388] AUBERT	03M	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<4.5	90	[389] GARMASH	04	BELL	$e^+e^- \to \Upsilon(4S)$
<7.0	90	[390] GARMASH	02	BELL	$e^+e^- \to \Upsilon(4S)$

[388] Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

[389] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[390] Uses a reference decay mode $B^+ \to \overline{D}^0\pi^+$ and $\overline{D}^0 \to K^+\pi^-$ with B($B^+ \to \overline{D}^0\pi^+$)·B($\overline{D}^0 \to K^+\pi^-$) = $(20.3 \pm 2.0) \times 10^{-5}$.

$\Gamma(K^- \pi^+ \pi^+ \text{nonresonant})/\Gamma_{total}$ — Γ_{218}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<56	90	BERGFELD	96B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_1(1400)^0 \pi^+)/\Gamma_{total}$ — Γ_{219}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<2.6 \times 10^{-3}$	90	ALBRECHT	91B	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^0 \pi^+ \pi^0)/\Gamma_{total}$ — Γ_{220}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<66 \times 10^{-6}$	90	[391] ECKHART	02	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[391] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^0 \rho^+)/\Gamma_{total}$ — Γ_{221}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<48	90	ASNER	96	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)^+ \pi^+ \pi^-)/\Gamma_{total}$ — Γ_{222}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
75.3± 6.0±8.1		[392] AUBERT,B	06U	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1100	90	ALBRECHT	91E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

[392] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+ \rho^0)/\Gamma_{total}$ — Γ_{223}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
< 6.1	90	[393] AUBERT,B	06G	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$10.6^{+3.0}_{-2.6} \pm 2.4$		[393] AUBERT	03V	BABR	Repl. by AUBERT,B 06G
< 74	90	[394] GODANG	02	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<900	90	ALBRECHT	91B	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

[393] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[394] Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 4.9×10^{-5}.

$\Gamma(K^*(892)^+ f_0(980))/\Gamma_{total}$ — Γ_{224}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
5.2±1.2±0.5	[395] AUBERT,B	06G	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

[395] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^0 \rho^+)/\Gamma_{total}$ — Γ_{225}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
9.2±1.5 OUR AVERAGE				
9.6±1.7±1.5	[396] AUBERT,B	06G	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
8.9±1.7±1.2	[396] ZHANG	05D	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[396] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+ K^*(892)^0)/\Gamma_{\text{total}}$ — Γ_{226}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<71	90	[397] GODANG	02	CLE2	$e^+e^- \to \Upsilon(4S)$

[397] Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 4.8×10^{-5}.

$\Gamma(K_1(1400)^+ \rho^0)/\Gamma_{\text{total}}$ — Γ_{227}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<7.8 \times 10^{-4}$	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$

$\Gamma(K_2^*(1430)^+ \rho^0)/\Gamma_{\text{total}}$ — Γ_{228}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.5 \times 10^{-3}$	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$

$\Gamma(K^+ \overline{K}^0)/\Gamma_{\text{total}}$ — Γ_{229}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
1.28±0.30 OUR AVERAGE					
1.61±0.44±0.09		[398] AUBERT,BE	06C	BABR	$e^+e^- \to \Upsilon(4S)$
1.0 ±0.4 ±0.1		[398] ABE	05G	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.5 ±0.5 ±0.1		[398] AUBERT,BE	05E	BABR	Repl. by AUBERT,BE 06C
< 2.5	90	[398] AUBERT	04M	BABR	Repl. by AUBERT,BE 05E
< 3.3	90	[398] CHAO	04	BELL	$e^+e^- \to \Upsilon(4S)$
< 3.3	90	[398] BORNHEIM	03	CLE2	$e^+e^- \to \Upsilon(4S)$
< 2.0	90	[398] CASEY	02	BELL	Repl. by CHAO 04
< 5.0	90	[398] ABE	01H	BELL	$e^+e^- \to \Upsilon(4S)$
< 2.4	90	[398] AUBERT	01E	BABR	$e^+e^- \to \Upsilon(4S)$
< 5.1	90	[398] CRONIN-HEN...	00	CLE2	$e^+e^- \to \Upsilon(4S)$
<21	90	GODANG	98	CLE2	Repl. by CRONIN-HENNESSY 00

[398] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{K}^0 K^+ \pi^0)/\Gamma_{\text{total}}$ — Γ_{230}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<24 \times 10^{-6}$	90	[399] ECKHART	02	CLE2	$e^+e^- \to \Upsilon(4S)$

[399] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ K_S^0 K_S^0)/\Gamma_{\text{total}}$ — Γ_{231}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
11.5±1.3 OUR AVERAGE				
10.7±1.2±1.0	[400] AUBERT,B	04V	BABR	$e^+e^- \to \Upsilon(4S)$
13.4±1.9±1.5	[400] GARMASH	04	BELL	$e^+e^- \to \Upsilon(4S)$

[400] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_S^0 K_S^0 \pi^+)/\Gamma_{\text{total}}$ — Γ_{232}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<3.2	90	[401] GARMASH	04	BELL	$e^+e^- \to \Upsilon(4S)$

[401] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ K^- \pi^+)/\Gamma_{total}$ Γ_{233}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
< 6.3	90	[402] AUBERT	03M	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<13	90	[403] GARMASH	04	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
<12	90	[404] GARMASH	02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[402] Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.
[403] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[404] Uses a reference decay mode $B^+ \rightarrow \overline{D}^0 \pi^+$ and $\overline{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \overline{D}^0 \pi^+) \cdot B(\overline{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

$\Gamma(K^+ K^- \pi^+ \text{nonresonant})/\Gamma_{total}$ Γ_{234}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<75	90	BERGFELD	96B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^+ K^+ \pi^-)/\Gamma_{total}$ Γ_{235}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.3 × 10^{-6}	90	[405] AUBERT	03M	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.4 \times 10^{-6}$	90	[406] GARMASH	04	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$<3.2 \times 10^{-6}$	90	[407] GARMASH	02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[405] Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.
[406] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[407] Uses a reference decay mode $B^+ \rightarrow \overline{D}^0 \pi^+$ and $\overline{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \overline{D}^0 \pi^+) \cdot B(\overline{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

$\Gamma(K^+ K^+ \pi^- \text{nonresonant})/\Gamma_{total}$ Γ_{236}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<87.9	90	ABBIENDI	00B	OPAL	$e^+ e^- \rightarrow Z$

$\Gamma(K^+ K^*(892)^0)/\Gamma_{total}$ Γ_{237}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<5.3 × 10^{-6}	90	[408] JESSOP	00	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.29 \times 10^{-4}$	90	ABBIENDI	00B	OPAL	$e^+ e^- \rightarrow Z$
$<1.38 \times 10^{-4}$	90	[409] ABE	00C	SLD	$e^+ e^- \rightarrow Z$

[408] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[409] ABE 00C assumes $B(Z \rightarrow b\overline{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$.

$\Gamma(K^+ f_J(2220))/\Gamma_{total}$ Γ_{238}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
not seen	[410] HUANG	03	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[410] No evidence is found for such decay and set a limit on $B(B^+ \rightarrow f_J(2220)) \times B(f_J(2220) \rightarrow \phi\phi) < 1.2 \times 10^{-6}$ at 90%CL where the $f_J(2220)$ is a possible glueball state.

$\Gamma(K^{*+}\pi^+K^-)/\Gamma_{\text{total}}$ — Γ_{239}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<11.8	90	[411] AUBERT,B	06U	BABR	$e^+e^- \to \Upsilon(4S)$

[411] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^{*+}K^+\pi^-)/\Gamma_{\text{total}}$ — Γ_{240}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<6.1	90	[412] AUBERT,B	06U	BABR	$e^+e^- \to \Upsilon(4S)$

[412] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+K^-K^+)/\Gamma_{\text{total}}$ — Γ_{241}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
33.7±2.2 OUR AVERAGE		Error includes scale factor of 1.4.			
35.2±0.9±1.6		[413] AUBERT	06O	BABR	$e^+e^- \to \Upsilon(4S)$
30.6±1.2±2.3		[413] GARMASH	05	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
32.8±1.8±2.8		[413] GARMASH	04	BELL	Repl. by GARMASH 05
29.6±2.1±1.6		[414] AUBERT	03M	BABR	Repl. by AUBERT 06O
35.3±3.7±4.5		[415] GARMASH	02	BELL	Repl. by GARMASH 04
<200	90	[416] ADAM	96D	DLPH	$e^+e^- \to Z$
<320	90	[416] ABREU	95N	DLPH	Sup. by ADAM 96D
<350	90	ALBRECHT	91E	ARG	$e^+e^- \to \Upsilon(4S)$

[413] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[414] Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

[415] Uses a reference decay mode $B^+ \to \overline{D}^0\pi^+$ and $\overline{D}^0 \to K^+\pi^-$ with $B(B^+ \to \overline{D}^0\pi^+)\cdot B(\overline{D}^0 \to K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

[416] Assumes B^0 and B^- production fractions of 0.39, and B_s production fraction of 0.12.

$\Gamma(K^+\phi)/\Gamma_{\text{total}}$ — Γ_{242}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
8.3 ±0.7 OUR AVERAGE					
8.4 ±0.7 ±0.7		[417] AUBERT	06O	BABR	$e^+e^- \to \Upsilon(4S)$
7.6 ±1.3 ±0.6		[418] ACOSTA	05J	CDF	$p\overline{p}$ at 1.96 TeV
$9.60 \pm 0.92 ^{+1.05}_{-0.85}$		[417] GARMASH	05	BELL	$e^+e^- \to \Upsilon(4S)$
$5.5 ^{+2.1}_{-1.8} \pm 0.6$		[417] BRIERE	01	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$10.0 ^{+0.9}_{-0.8} \pm 0.5$		[417] AUBERT	04A	BABR	Repl. by AUBERT 06O
9.4 ±1.1 ±0.7		[417] CHEN	03B	BELL	Repl. by GARMASH 05
$14.6 ^{+3.0}_{-2.8} \pm 2.0$		[419] GARMASH	02	BELL	Repl. by CHEN 03B
$7.7 ^{+1.6}_{-1.4} \pm 0.8$		[417] AUBERT	01D	BABR	$e^+e^- \to \Upsilon(4S)$
<144	90	[420] ABE	00C	SLD	$e^+e^- \to Z$
< 5	90	[417] BERGFELD	98	CLE2	

<280	90	[421] ADAM	96D	DLPH	$e^+e^- \to Z$
< 12	90	ASNER	96	CLE2	$e^+e^- \to \Upsilon(4S)$
<440	90	[422] ABREU	95N	DLPH	Sup. by ADAM 96D
<180	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$
< 90	90	[423] AVERY	89B	CLEO	$e^+e^- \to \Upsilon(4S)$
<210	90	AVERY	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[417] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[418] Uses $B(B^+ \to J/\psi K^+) = (1.00 \pm 0.04) \times 10^{-3}$ and $B(J/\psi \to \mu^+\mu^-) = 0.0588 \pm 0.0010$.

[419] Uses a reference decay mode $B^+ \to \overline{D}^0\pi^+$ and $\overline{D}^0 \to K^+\pi^-$ with $B(B^+ \to \overline{D}^0\pi^+) \cdot B(\overline{D}^0 \to K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

[420] ABE 00C assumes $B(Z \to b\overline{b})$=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.

[421] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

[422] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

[423] AVERY 89B reports $< 8 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(f_0(980)K^+ \times B(f_0(980) \to K^+K^-))/\Gamma_{total}$ Γ_{243}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<2.9	90	[424] GARMASH	05	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.5±2.5±1.6		[424] AUBERT	06O	BABR	$e^+e^- \to \Upsilon(4S)$

[424] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_2(1320)K^+ \times B(a_2(1320) \to K^+K^-))/\Gamma_{total}$ Γ_{244}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.1 \times 10^{-6}$	90	[425] GARMASH	05	BELL	$e^+e^- \to \Upsilon(4S)$

[425] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f'_2(1525)K^+ \times B(f'_2(1525) \to K^+K^-))/\Gamma_{total}$ Γ_{245}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<4.9 \times 10^{-6}$	90	[426] GARMASH	05	BELL	$e^+e^- \to \Upsilon(4S)$

[426] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(X_0(1550)K^+ \times B(X_0(1550) \to K^+K^-))/\Gamma_{total}$ Γ_{246}/Γ

$X_0(1550)$ is a possible spin zero state near 1.55 GeV/c^2 invariant mass of $K^+K^+K^-$.

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
4.3±0.6±0.3	[427] AUBERT	06O	BABR	$e^+e^- \to \Upsilon(4S)$

[427] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi(1680)K^+ \times B(\phi(1680) \to K^+K^-))/\Gamma_{total}$ Γ_{247}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<0.8 \times 10^{-6}$	90	[428] GARMASH	05	BELL	$e^+e^- \to \Upsilon(4S)$

[428] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_0(1710)K^+ \times B(f_0(1710) \to K^+K^-))/\Gamma_{total}$ Γ_{248}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
1.7±1.0±0.3	[429] AUBERT	06O	BABR	$e^+e^- \to \Upsilon(4S)$

[429] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+K^-K^+ \text{nonresonant})/\Gamma_{total}$ Γ_{249}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
28^{+9}_{-16} OUR AVERAGE		Error includes scale factor of 3.3.			
$50.0 \pm 6.0 \pm 4.0$		[430] AUBERT	06O	BABR	$e^+e^- \to \Upsilon(4S)$
$24.0 \pm 1.5^{+2.6}_{-6.0}$		[430] GARMASH	05	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<38	90	BERGFELD	96B	CLE2	$e^+e^- \to \Upsilon(4S)$

[430] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+K^+K^-)/\Gamma_{total}$ Γ_{250}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
36.2±3.3±3.6		[431] AUBERT,B	06U	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1600	90	ALBRECHT	91E	ARG	$e^+e^- \to \Upsilon(4S)$

[431] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+\phi)/\Gamma_{total}$ Γ_{251}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
9.6±3.0 OUR AVERAGE		Error includes scale factor of 1.9.			
$12.7^{+2.2}_{-2.0} \pm 1.1$		[432] AUBERT	03V	BABR	$e^+e^- \to \Upsilon(4S)$
$6.7^{+2.1}_{-1.9}{}^{+0.7}_{-1.0}$		[432] CHEN	03B	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$9.7^{+4.2}_{-3.4} \pm 1.7$		[432] AUBERT	01D	BABR	Repl. by AUBERT 03V
< 22.5	90	[432] BRIERE	01	CLE2	$e^+e^- \to \Upsilon(4S)$
< 41	90	[432] BERGFELD	98	CLE2	
< 70	90	ASNER	96	CLE2	$e^+e^- \to \Upsilon(4S)$
<1300	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$

[432] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_1(1400)^+\phi)/\Gamma_{total}$ Γ_{252}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.1 \times 10^{-3}$	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$

$\Gamma(K_2^*(1430)^+\phi)/\Gamma_{total}$ Γ_{253}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<3.4 \times 10^{-3}$	90	ALBRECHT	91B	ARG	$e^+e^- \to \Upsilon(4S)$

$\Gamma(K^+\phi\phi)/\Gamma_{total}$ — Γ_{254}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
$4.9^{+2.4}_{-2.2}$ OUR AVERAGE Error includes scale factor of 2.9.				
$7.5\pm1.0\pm0.7$	[433] AUBERT,BE	06H	BABR	$e^+e^- \to \Upsilon(4S)$
$2.6^{+1.1}_{-0.9}\pm0.3$	[433] HUANG	03	BELL	$e^+e^- \to \Upsilon(4S)$

[433] Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$ and for a $\phi\phi$ invariant mass below 2.85 GeV/c^2.

$\Gamma(\eta'\eta' K^+)/\Gamma_{total}$ — Γ_{255}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<25	90	[434] AUBERT,B	06P	BABR	$e^+e^- \to \Upsilon(4S)$

[434] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+\gamma)/\Gamma_{total}$ — Γ_{256}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
4.03 ± 0.26 OUR AVERAGE					
$3.87\pm0.28\pm0.26$		[435] AUBERT,BE	04A	BABR	$e^+e^- \to \Upsilon(4S)$
$4.25\pm0.31\pm0.24$		[436] NAKAO	04	BELL	$e^+e^- \to \Upsilon(4S)$
$3.76^{+0.89}_{-0.83}\pm0.28$		[436] COAN	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$3.83\pm0.62\pm0.22$		[436] AUBERT	02C	BABR	Repl. by AUBERT,BE 04A
$5.7\pm3.1\pm1.1$		[437] AMMAR	93	CLE2	Repl. by COAN 00
<55	90	[438] ALBRECHT	89G	ARG	$e^+e^- \to \Upsilon(4S)$
<55	90	[438] AVERY	89B	CLEO	$e^+e^- \to \Upsilon(4S)$
<180	90	AVERY	87	CLEO	$e^+e^- \to \Upsilon(4S)$

[435] Uses the production ratio of charged and neutral B from $\Upsilon(4S)$ decays $R^{+/0} = 1.006 \pm 0.048$.
[436] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[437] AMMAR 93 observed 4.1 ± 2.3 events above background.
[438] Assumes the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$.

$\Gamma(K_1(1270)^+\gamma)/\Gamma_{total}$ — Γ_{257}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
$4.3\pm0.9\pm0.9$		[439] YANG	05	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<9.9	90	[439] NISHIDA	02	BELL	Repl. by YANG 05
<730	90	[440] ALBRECHT	89G	ARG	$e^+e^- \to \Upsilon(4S)$

[439] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[440] ALBRECHT 89G reports < 0.0066 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\eta K^+ \gamma)/\Gamma_{total}$ — Γ_{258}/Γ

VALUE (units 10^{-6})		DOCUMENT ID		TECN	COMMENT
9.4±1.1 OUR AVERAGE					
$10.0 \pm 1.3 \pm 0.5$	[441,442]	AUBERT,B	06M	BABR	$e^+e^- \to \Upsilon(4S)$
$8.4 \pm 1.5 ^{+1.2}_{-0.9}$	[442,443]	NISHIDA	05	BELL	$e^+e^- \to \Upsilon(4S)$

[441] $m_{\eta K} < 3.25$ GeV/c^2.
[442] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[443] $m_{\eta K} < 2.4$ GeV/c^2

$\Gamma(\eta' K^+ \gamma)/\Gamma_{total}$ — Γ_{259}/Γ

VALUE (units 10^{-6})	CL%		DOCUMENT ID		TECN	COMMENT
<4.2	90	[444,445]	AUBERT,B	06M	BABR	$e^+e^- \to \Upsilon(4S)$

[444] $m_{\eta' K} < 3.25$ GeV/c^2.
[445] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi K^+ \gamma)/\Gamma_{total}$ — Γ_{260}/Γ

VALUE (units 10^{-6})		DOCUMENT ID		TECN	COMMENT
3.5±0.6 OUR AVERAGE					
$3.5 \pm 0.6 \pm 0.4$	[446]	AUBERT	07Q	BABR	$e^+e^- \to \Upsilon(4S)$
$3.4 \pm 0.9 \pm 0.4$	[446]	DRUTSKOY	04	BELL	$e^+e^- \to \Upsilon(4S)$

[446] Assumes equal production of B^+ and B^0 at $\Upsilon(4S)$.

$\Gamma(K^+ \pi^- \pi^+ \gamma)/\Gamma_{total}$ — Γ_{261}/Γ

VALUE (units 10^{-5})		DOCUMENT ID		TECN	COMMENT
2.50±0.18±0.22	[447]	YANG	05	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$2.4 \pm 0.5 ^{+0.4}_{-0.2}$	[447,448]	NISHIDA	02	BELL	Repl. by YANG 05

[447] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[448] $M_{K\pi\pi} < 2.4$ GeV/c^2.

$\Gamma(K^*(892)^0 \pi^+ \gamma)/\Gamma_{total}$ — Γ_{262}/Γ

VALUE		DOCUMENT ID		TECN	COMMENT
$\mathbf{(2.0^{+0.7}_{-0.6} \pm 0.2) \times 10^{-5}}$	[449,450]	NISHIDA	02	BELL	$e^+e^- \to \Upsilon(4S)$

[449] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[450] $M_{K\pi\pi} < 2.4$ GeV/c^2.

$\Gamma(K^+ \rho^0 \gamma)/\Gamma_{total}$ — Γ_{263}/Γ

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
$\mathbf{<2.0 \times 10^{-5}}$	90	[451,452]	NISHIDA	02	BELL	$e^+e^- \to \Upsilon(4S)$

[451] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[452] $M_{K\pi\pi} < 2.4$ GeV/c^2.

$\Gamma(K^+\pi^-\pi^+\gamma$ nonresonant$)/\Gamma_{total}$ — Γ_{264}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.2\times10^{-6}$	90	[453,454] NISHIDA 02	BELL	$e^+e^-\to\Upsilon(4S)$

[453] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[454] $M_{K\pi\pi} < 2.4$ GeV/c^2.

$\Gamma(K_1(1400)^+\gamma)/\Gamma_{total}$ — Γ_{265}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.5	90	[455] YANG 05	BELL	$e^+e^-\to\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 5.0	90	[455] NISHIDA 02	BELL	Repl. by YANG 05
<220	90	[456] ALBRECHT 89G	ARG	$e^+e^-\to\Upsilon(4S)$

[455] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[456] ALBRECHT 89G reports < 0.0020 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K_2^*(1430)^+\gamma)/\Gamma_{total}$ — Γ_{266}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
1.45±0.40±0.15		[457] AUBERT,B 04U	BABR	$e^+e^-\to\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<140	90	[458] ALBRECHT 89G	ARG	$e^+e^-\to\Upsilon(4S)$

[457] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[458] ALBRECHT 89G reports < 0.0013 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^*(1680)^+\gamma)/\Gamma_{total}$ — Γ_{267}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0019	90	[459] ALBRECHT 89G	ARG	$e^+e^-\to\Upsilon(4S)$

[459] ALBRECHT 89G reports < 0.0017 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K_3^*(1780)^+\gamma)/\Gamma_{total}$ — Γ_{268}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 39	90	[460,461] NISHIDA 05	BELL	$e^+e^-\to\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<5500	90	[462] ALBRECHT 89G	ARG	$e^+e^-\to\Upsilon(4S)$

[460] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[461] Uses B($K_3^*(1780)\to\eta K$) $= 0.11^{+0.05}_{-0.04}$.
[462] ALBRECHT 89G reports < 0.005 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K_4^*(2045)^+\gamma)/\Gamma_{total}$ — Γ_{269}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0099	90	[463] ALBRECHT 89G	ARG	$e^+e^-\to\Upsilon(4S)$

[463] ALBRECHT 89G reports < 0.0090 assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\rho^+\gamma)/\Gamma_{total}$ — Γ_{270}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$0.88^{+0.29}_{-0.25}$ OUR AVERAGE				
$1.10^{+0.37}_{-0.33}\pm0.09$		[464] AUBERT 07L	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.55^{+0.42}_{-0.36}{}^{+0.09}_{-0.08}$		[464] MOHAPATRA 06	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.9\ {}^{+0.6}_{-0.5}\ \pm0.1$	90	[464] AUBERT 05	BABR	Repl. by AUBERT 07L
$<$ 2.2	90	[464] MOHAPATRA 05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 2.1	90	[464] AUBERT 04C	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$<$13	90	[464,465] COAN 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[464] Assumes equal production of B^+ and B^0 at $\Upsilon(4S)$.
[465] No evidence for a nonresonant $K\pi\gamma$ contamination was seen; the central value assumes no contamination.

$\Gamma(\pi^+\pi^0)/\Gamma_{total}$ — Γ_{271}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
5.5 ± 0.6 OUR AVERAGE				
$5.8\pm0.6\pm0.4$		[466] AUBERT 05L	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$5.0\pm1.2\pm0.5$		[466] CHAO 04	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$4.6^{+1.8}_{-1.6}{}^{+0.6}_{-0.7}$		[466] BORNHEIM 03	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5.5^{+1.0}_{-1.9}\pm0.6$		[466] AUBERT 03L	BABR	Repl. by AUBERT 05L
$7.4^{+2.3}_{-2.2}\pm0.9$		[466] CASEY 02	BELL	Repl. by CHAO 04
$<$ 13.4	90	[466] ABE 01H	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 9.6	90	[466] AUBERT 01E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 12.7	90	[466] CRONIN-HEN...00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 20	90	GODANG 98	CLE2	Repl. by CRONIN-HENNESSY 00
$<$ 17	90	ASNER 96	CLE2	Repl. by GODANG 98
$<$ 240	90	[466] ALBRECHT 90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<$2300	90	[467] BEBEK 87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[466] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[467] BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$.

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma_{total}$ — Γ_{272}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$16.2\pm1.2\pm0.9$		[468] AUBERT,B 05G	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$10.9\pm3.3\pm1.6$		[468] AUBERT 03M	BABR	Repl. by AUBERT 05G
$<$130	90	[469] ADAM 96D	DLPH	$e^+e^- \rightarrow Z$
$<$220	90	[470] ABREU 95N	DLPH	Sup. by ADAM 96D
$<$450	90	[471] ALBRECHT 90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<$190	90	[472] BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[468] Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.
[469] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.
[470] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.
[471] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.
[472] BORTOLETTO 89 reports $< 1.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\rho^0\pi^+)/\Gamma_{total}$ — Γ_{273}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
8.7±1.1 OUR AVERAGE					
$8.8\pm1.0^{+0.6}_{-0.9}$		[473] AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$
$8.0^{+2.3}_{-2.0}\pm0.7$		[473] GORDON	02	BELL	$e^+e^- \to \Upsilon(\text{rS})$
$10.4^{+3.3}_{-3.4}\pm2.1$		[473] JESSOP	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$9.5\pm1.1\pm0.9$		[473] AUBERT	04Z	BABR	Repl. by AUBERT 05G
< 83	90	[474] ABE	00C	SLD	$e^+e^- \to Z$
<160	90	[475] ADAM	96D	DLPH	$e^+e^- \to Z$
< 43	90	ASNER	96	CLE2	Repl. by JESSOP 00
<260	90	[476] ABREU	95N	DLPH	Sup. by ADAM 96D
<150	90	[473] ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$
<170	90	[477] BORTOLETTO	89	CLEO	$e^+e^- \to \Upsilon(4S)$
<230	90	[477] BEBEK	87	CLEO	$e^+e^- \to \Upsilon(4S)$
<600	90	GILES	84	CLEO	Repl. by BEBEK 87

[473] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[474] ABE 00C assumes B($Z \to b\overline{b}$)=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.
[475] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.
[476] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.
[477] Papers assume the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\left[\Gamma(K^*(892)^0\pi^+)+\Gamma(\rho^0\pi^+)\right]/\Gamma_{total}$ — $(\Gamma_{202}+\Gamma_{273})/\Gamma$

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
$\mathbf{170^{+120}_{-80}\pm20}$	[478] ADAM	96D	DLPH	$e^+e^- \to Z$

[478] ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

$\Gamma(\pi^+ f_0(980)\times B(f_0(980)\to\pi^+\pi^-))/\Gamma_{total}$ — Γ_{274}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
< 3.0	90	[479] AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<140	90	[480] BORTOLETTO	89	CLEO	$e^+e^- \to \Upsilon(4S)$

[479] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[480] BORTOLETTO 89 reports $< 1.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ f_2(1270))/\Gamma_{total}$ — Γ_{275}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
8.2±2.1±1.4		[481,482] AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<240	90	[483] BORTOLETTO	89	CLEO	$e^+e^- \to \Upsilon(4S)$

[481] Reported $B(B^+ \to f_2(1270)\pi^+) \times B(f_2(1270) \to \pi^+\pi^-) = (2.3 \pm 0.6 \pm 0.4) \times 10^{-6}$ and rescaled using $B(f_2(1270) \to \pi^+\pi^-) = 0.28$.
[482] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[483] BORTOLETTO 89 reports $< 2.1 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\rho(1450)^0\pi^+)/\Gamma_{total}$ — Γ_{276}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<2.3	90	[484] AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$

[484] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_0(1370)\pi^+ \times B(f_0(1370) \to \pi^+\pi^-))/\Gamma_{total}$ — Γ_{277}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<3.0	90	[485] AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$

[485] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_0(600)\pi^+ \times B(f_0(600) \to \pi^+\pi^-))/\Gamma_{total}$ — Γ_{278}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<4.1	90	[486] AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$

[486] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+\pi^-\pi^+ \text{ nonresonant})/\Gamma_{total}$ — Γ_{279}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
< 4.6	90	[487] AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<41	90	BERGFELD	96B	CLE2	$e^+e^- \to \Upsilon(4S)$

[487] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+\pi^0\pi^0)/\Gamma_{total}$ — Γ_{280}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<8.9 \times 10^{-4}$	90	[488] ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$

[488] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^+\pi^0)/\Gamma_{total}$ — Γ_{281}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
12.0±1.9 OUR AVERAGE					
$13.2 \pm 2.3 ^{+1.4}_{-1.9}$		[489] ZHANG	05A	BELL	$e^+e^- \to \Upsilon(4S)$
10.9±1.9±1.9		[489] AUBERT	04Z	BABR	$e^+e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<$ 43	90	[489,490]	JESSOP	00	CLE2	$e^+e^- \to \Upsilon(4S)$
$<$ 77	90		ASNER	96	CLE2	Repl. by JESSOP 00
$<$550	90	[489]	ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$

[489] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[490] Assumes no nonresonant contributions of $B^+ \to \pi^+\pi^0\pi^0$.

$\Gamma(\pi^+\pi^-\pi^+\pi^0)/\Gamma_{total}$ — Γ_{282}/Γ

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
$<4.0 \times 10^{-3}$	90	[491]	ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$

[491] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^+\rho^0)/\Gamma_{total}$ — Γ_{283}/Γ

VALUE (units 10^{-6})	CL%		DOCUMENT ID		TECN	COMMENT
18 ±4 OUR AVERAGE			Error includes scale factor of 1.5.			
$16.8 \pm 2.2 \pm 2.3$		[492]	AUBERT,BE	06G	BABR	$e^+e^- \to \Upsilon(4S)$
$31.7 \pm 7.1 ^{+3.8}_{-6.7}$		[492,493]	ZHANG	03B	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$22.5 ^{+5.7}_{-5.4} \pm 5.8$		[492]	AUBERT	03V	BABR	Repl. by AUBERT,BE 06G
$<$1000	90	[492]	ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$

[492] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[493] The systematic error includes the error associated with the helicity-mix uncertainty.

$\Gamma(\rho^+ f_0(980) \times B(f_0(980) \to \pi^+\pi^-))/\Gamma_{total}$ — Γ_{284}/Γ

VALUE (units 10^{-6})	CL%		DOCUMENT ID		TECN	COMMENT
$<$1.9	90	[494]	AUBERT,BE	06G	BABR	$e^+e^- \to \Upsilon(4S)$

[494] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_1(1260)^+\pi^0)/\Gamma_{total}$ — Γ_{285}/Γ

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
$<1.7 \times 10^{-3}$	90	[495]	ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$

[495] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^0\pi^+)/\Gamma_{total}$ — Γ_{286}/Γ

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
$<9.0 \times 10^{-4}$	90	[496]	ALBRECHT	90B	ARG	$e^+e^- \to \Upsilon(4S)$

[496] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\omega\pi^+)/\Gamma_{total}$ — Γ_{287}/Γ

VALUE (units 10^{-6})	CL%		DOCUMENT ID		TECN	COMMENT
6.7±0.6 OUR AVERAGE						
$6.1 \pm 0.7 \pm 0.4$		[497]	AUBERT,B	06E	BABR	$e^+e^- \to \Upsilon(4S)$
$6.9 \pm 0.6 \pm 0.5$		[497]	JEN	06	BELL	$e^+e^- \to \Upsilon(4S)$
$11.3 ^{+3.3}_{-2.9} \pm 1.4$		[497]	JESSOP	00	CLE2	$e^+e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.5\pm0.9\pm0.5$		[497] AUBERT	04H	BABR	Repl. by AUBERT,B 06E
$5.7^{+1.4}_{-1.3}\pm0.6$		[497] WANG	04A	BELL	Repl. by JEN 06
$4.2^{+2.0}_{-1.8}\pm0.5$		[497] LU	02	BELL	Repl. by WANG 04A
$6.6^{+2.1}_{-1.8}\pm0.7$		[497] AUBERT	01G	BABR	Repl. by AUBERT 04H
$<$ 23	90	[497] BERGFELD	98	CLE2	Repl. by JESSOP 00
$<$400	90	[497] ALBRECHT	90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[497] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega\rho^+)/\Gamma_{total}$ — Γ_{288}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$10.6\pm2.1^{+1.6}_{-1.0}$		[498] AUBERT,B	06T	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$12.6^{+3.7}_{-3.3}\pm1.6$		[498] AUBERT	05O	BABR	Repl. by AUBERT,B 06T
$<$61	90	[498] BERGFELD	98	CLE2	

[498] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta\pi^+)/\Gamma_{total}$ — Γ_{289}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
4.9 ± 0.5 OUR AVERAGE					
$5.1\pm0.6\pm0.3$		[499] AUBERT,B	05K	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$4.8\pm0.7\pm0.3$		[499] CHANG	05A	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$1.2^{+2.8}_{-1.2}$		[499] RICHICHI	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$5.3\pm1.0\pm0.3$		[499] AUBERT	04H	BABR	Repl. by AUBERT,B 05K
$<$ 15	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00
$<$700	90	[499] ALBRECHT	90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[499] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta'\pi^+)/\Gamma_{total}$ — Γ_{290}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
2.6 ± 1.1 OUR AVERAGE Error includes scale factor of 2.0.					
$1.76^{+0.67}_{-0.62}{}^{+0.15}_{-0.14}$		[500] SCHUEMANN	06	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$4.0\pm0.8\pm0.4$		[500] AUBERT,B	05K	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<$ 4.5	90	[500] AUBERT	04H	BABR	Repl. by AUBERT,B 05K
$<$ 7.0	90	[500] ABE	01M	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$<$12	90	[500] AUBERT	01G	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$<$12	90	[500] RICHICHI	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$<$31	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

[500] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' \rho^+)/\Gamma_{total}$ — Γ_{291}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{8.7^{+3.1+2.3}_{-2.8-1.3}}$		[501] AUBERT	07E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<22	90	[501] AUBERT,B	04D	BABR	Repl. by AUBERT 07E
<33	90	[501] RICHICHI	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<47	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

[501] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta \rho^+)/\Gamma_{total}$ — Γ_{292}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
8.4±1.9±1.1		[502] AUBERT,B	05K	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<14	90	[502] AUBERT,B	04D	BABR	Repl. by AUBERT,B 05K
<15	90	[502] RICHICHI	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<32	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

[502] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi \pi^+)/\Gamma_{total}$ — Γ_{293}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
< 0.24	90	[503] AUBERT,B	06C	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.41	90	[503] AUBERT	04A	BABR	Repl. by AUBERT,B 06C
< 1.4	90	[503] AUBERT	01D	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
<153	90	[504] ABE	00C	SLD	$e^+e^- \rightarrow Z$
< 5	90	[503] BERGFELD	98	CLE2	

[503] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[504] ABE 00C assumes B($Z \rightarrow b\overline{b}$)=(21.7 ± 0.1)% and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})$% and $f_{B_s}=(10.5^{+1.8}_{-2.2})$%.

$\Gamma(\phi \rho^+)/\Gamma_{total}$ — Γ_{294}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN
<16	[505] BERGFELD	98	CLE2

[505] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_0^0 \pi^+)/\Gamma_{total}$ — Γ_{295}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<5.8	90	[506] AUBERT,BE	04	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[506] Assumes equal production of charged and neutral B mesons from $\Upsilon(4S)$ decays.

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-)/\Gamma_{total}$ — Γ_{296}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<8.6 \times 10^{-4}}$	90	[507] ALBRECHT	90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[507] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^0 a_1(1260)^+)/\Gamma_{total}$ Γ_{297}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<6.2 × 10⁻⁴	90	[508] BORTOLETTO	89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<6.0 \times 10^{-4}$	90	[509] ALBRECHT	90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<3.2 \times 10^{-3}$	90	[508] BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[508] BORTOLETTO 89 reports $< 5.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.
[509] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^0 a_2(1320)^+)/\Gamma_{total}$ Γ_{298}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<7.2 × 10⁻⁴	90	[510] BORTOLETTO	89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.6 \times 10^{-3}$	90	[511] BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[510] BORTOLETTO 89 reports $< 6.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.
[511] BEBEK 87reports $< 2.3 \times 10^{-3}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{total}$ Γ_{299}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<6.3 × 10⁻³	90	[512] ALBRECHT	90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[512] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^+ a_1(1260)^0)/\Gamma_{total}$ Γ_{300}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.3 × 10⁻²	90	[513] ALBRECHT	90B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[513] ALBRECHT 90B limit assumes equal production of $B^0\overline{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(h^+\pi^0)/\Gamma_{total}$ Γ_{301}/Γ

$h^+ = K^+$ or π^+

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
$\mathbf{16^{+6}_{-5} \pm 3.6}$	GODANG	98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\omega h^+)/\Gamma_{total}$ Γ_{302}/Γ

$h^+ = K^+$ or π^+

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
$\mathbf{13.8^{+2.7}_{-2.4}}$ **OUR AVERAGE**				
$13.4^{+3.3}_{-2.9} \pm 1.1$	[514] LU	02	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$14.3^{+3.6}_{-3.2} \pm 2.0$	[514] JESSOP	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$25\ ^{+8}_{-7} \pm 3$	[514] BERGFELD	98	CLE2	Repl. by JESSOP 00

[514] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(h^+ X^0 (\text{Familon}))/\Gamma_{\text{total}}$ — Γ_{303}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<49	90	[515] AMMAR	01B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[515] AMMAR 01B searched for the two-body decay of the B meson to a massless neutral feebly-interacting particle X^0 such as the familon, the Nambu-Goldstone boson associated with a spontaneously broken global family symmetry.

$\Gamma(p\overline{p}\pi^+)/\Gamma_{\text{total}}$ — Γ_{304}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{3.06^{+0.73}_{-0.62} \pm 0.37}$		[516,517] WANG	04	BELL	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 3.7	90	[516,518] ABE	02K	BELL	Repl. by WANG 04
<500	90	[519] ABREU	95N	DLPH	Sup. by ADAM 96D
<160	90	[520] BEBEK	89	CLEO	$e^+ e^- \to \Upsilon(4S)$
$570 \pm 150 \pm 210$		[521] ALBRECHT	88F	ARG	$e^+ e^- \to \Upsilon(4S)$

[516] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[517] The branching fraction for $M_{p\overline{p}} < 2.85$ GeV/c^2 is also reported.
[518] Explicitly vetoes resonant production of $p\overline{p}$ from Charmonium states.
[519] Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.
[520] BEBEK 89 reports $< 1.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \overline{B}^0$. We rescale to 50%.
[521] ALBRECHT 88F reports $(5.2 \pm 1.4 \pm 1.9) \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \overline{B}^0$. We rescale to 50%.

$\Gamma(p\overline{p}\pi^+ \text{nonresonant})/\Gamma_{\text{total}}$ — Γ_{305}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<53	90	BERGFELD	96B	CLE2	$e^+ e^- \to \Upsilon(4S)$

$\Gamma(p\overline{p}\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ — Γ_{306}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<5.2 \times 10^{-4}}$	90	[522] ALBRECHT	88F	ARG	$e^+ e^- \to \Upsilon(4S)$

[522] ALBRECHT 88F reports $< 4.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \overline{B}^0$. We rescale to 50%.

$\Gamma(p\overline{p}K^+)/\Gamma_{\text{total}}$ — Γ_{307}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
5.6 ±1.0 OUR AVERAGE Error includes scale factor of 2.4.				
$6.7 \pm 0.5 \pm 0.4$	[523,524,525] AUBERT,B	05L	BABR	$e^+ e^- \to \Upsilon(4S)$
$4.59^{+0.38}_{-0.34} \pm 0.50$	[523,524,525] WANG	05A	BELL	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5.66^{+0.67}_{-0.57} \pm 0.62$	[523,524,526] WANG	04	BELL	Repl. by WANG 05A
$4.3^{+1.1}_{-0.9} \pm 0.5$	[523,524] ABE	02K	BELL	Repl. by WANG 04

[523] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[524] Explicitly vetoes resonant production of $p\overline{p}$ from Charmonium states.
[525] Provides also results with $M_{p\overline{p}} < 2.85$ GeV/c^2 and angular asymmetry of $p\overline{p}$ system.
[526] The branching fraction for $M_{p\overline{p}} < 2.85$ GeV/c^2 is also reported.

$\Gamma(\Theta(1710)^{++}\overline{p}\times B(\Theta(1710)^{++}\to pK^+))/\Gamma_{total}$ — Γ_{308}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<0.091	90	[527] WANG	05A	BELL	$e^+e^-\to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.1	90	[527,528] AUBERT,B	05L	BABR	$e^+e^-\to \Upsilon(4S)$

[527] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[528] Provides upper limits depending on the pentaquark masses between 1.43 to 2.0 GeV/c^2.

$\Gamma(f_J(2220)K^+\times B(f_J(2220)\to p\overline{p}))/\Gamma_{total}$ — Γ_{309}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<0.41	90	[529] WANG	05A	BELL	$e^+e^-\to \Upsilon(4S)$

[529] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(p\overline{\Lambda}(1520))/\Gamma_{total}$ — Γ_{310}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<1.5	90	[530] AUBERT,B	05L	BABR	$e^+e^-\to \Upsilon(4S)$

[530] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(p\overline{p}K^+ \text{ nonresonant})/\Gamma_{total}$ — Γ_{311}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<89	90	BERGFELD	96B	CLE2	$e^+e^-\to \Upsilon(4S)$

$\Gamma(p\overline{p}K^*(892)^+)/\Gamma_{total}$ — Γ_{312}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
$\mathbf{10.3^{+3.6}_{-2.8}{}^{+1.3}_{-1.7}}$	[531,532] WANG	04	BELL	$e^+e^-\to \Upsilon(4S)$

[531] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[532] The branching fraction for $M_{p\overline{p}} < 2.85$ GeV/c^2 is also reported.

$\Gamma(p\overline{\Lambda})/\Gamma_{total}$ — Γ_{313}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
< 0.49	90	[533] CHANG	05	BELL	$e^+e^-\to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.5	90	[533] BORNHEIM	03	CLE2	$e^+e^-\to \Upsilon(4S)$
< 2.2	90	[533] ABE	02O	BELL	$e^+e^-\to \Upsilon(4S)$
< 2.6	90	[533] COAN	99	CLE2	$e^+e^-\to \Upsilon(4S)$
<60	90	[534] AVERY	89B	CLEO	$e^+e^-\to \Upsilon(4S)$
<93	90	[535] ALBRECHT	88F	ARG	$e^+e^-\to \Upsilon(4S)$

[533] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[534] AVERY 89B reports $< 5\times10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.
[535] ALBRECHT 88F reports $< 8.5\times10^{-5}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(p\overline{\Lambda}\gamma)/\Gamma_{\text{total}}$ — Γ_{314}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{2.16^{+0.58}_{-0.53}\pm0.20}$		[536] LEE	05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.9	90	[537] EDWARDS	03	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[536] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[537] Corresponds to $E_\gamma > 1.5$ GeV. The limit changes to 3.3×10^{-6} for $E_\gamma > 2.0$ GeV.

$\Gamma(p\overline{\Sigma}\gamma)/\Gamma_{\text{total}}$ — Γ_{315}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<4.6	90	[538] LEE	05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<7.9	90	[539] EDWARDS	03	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[538] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[539] Corresponds to $E_\gamma > 1.5$ GeV. The limit changes to 6.4×10^{-6} for $E_\gamma > 2.0$ GeV.

$\Gamma(p\overline{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ — Γ_{316}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<2.0 \times 10^{-4}}$	90	[540] ALBRECHT	88F	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

[540] ALBRECHT 88F reports $< 1.8 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\Lambda\overline{\Lambda}\pi^+)/\Gamma_{\text{total}}$ — Γ_{317}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<2.8	90	[541] LEE	04	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[541] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\Lambda\overline{\Lambda}K^+)/\Gamma_{\text{total}}$ — Γ_{318}/Γ

VALUE (units 10^{-6})	DOCUMENT ID		TECN	COMMENT
$\mathbf{2.91^{+0.9}_{-0.70}\pm0.38}$	[542] LEE	04	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[542] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Delta^0}p)/\Gamma_{\text{total}}$ — Γ_{319}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<3.8 \times 10^{-4}}$	90	[543] BORTOLETTO	89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[543] BORTOLETTO 89 reports $< 3.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(\Delta^{++}\overline{p})/\Gamma_{\text{total}}$ — Γ_{320}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<1.5 \times 10^{-4}}$	90	[544] BORTOLETTO	89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[544] BORTOLETTO 89 reports $< 1.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(D^+ p\overline{p})/\Gamma_{total}$ — Γ_{321}/Γ

VALUE	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<1.5 × 10^{-5}	90	[545] ABE	02W	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[545] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^+ p\overline{p})/\Gamma_{total}$ — Γ_{322}/Γ

VALUE	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<1.5 × 10^{-5}	90	[546] ABE	02W	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[546] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Lambda}_c^- p\pi^+)/\Gamma_{total}$ — Γ_{323}/Γ

VALUE (units 10^{-4})	*DOCUMENT ID*		*TECN*	*COMMENT*
2.1±0.6 OUR AVERAGE				
2.0±0.2±0.5	[547] GABYSHEV	06A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
2.4±0.6±0.6	[548] DYTMAN	02	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.9±0.5±0.5	[549] GABYSHEV	02	BELL	Repl. by GABYSHEV 06A
$6.2^{+2.3}_{-2.0}\pm1.6$	[550] FU	97	CLE2	Repl. by DYTMAN 02

[547] GABYSHEV 06A reports $(2.01 \pm 0.15 \pm 0.20) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[548] DYTMAN 02 reports $(2.4^{+0.63}_{-0.62}) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[549] GABYSHEV 02 reports $(1.87^{+0.51}_{-0.49}) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[550] FU 97 uses PDG 96 values of Λ_c branching fraction.

$\Gamma(\overline{\Lambda}_c^- \Delta(1232)^{++})/\Gamma_{total}$ — Γ_{324}/Γ

VALUE (units 10^{-5})	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<1.9	90	GABYSHEV	06A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\overline{\Lambda}_c^- \Delta_X(1600)^{++})/\Gamma_{total}$ — Γ_{325}/Γ

VALUE (units 10^{-5})	*DOCUMENT ID*		*TECN*	*COMMENT*
5.9±1.2±1.5	[551] GABYSHEV	06A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[551] GABYSHEV 06A reports $(5.9 \pm 1.0 \pm 0.6) \times 10^{-5}$ for $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\overline{\Lambda}_c^- \Delta_X(2420)^{++})/\Gamma_{total}$ — Γ_{326}/Γ

VALUE (units 10^{-5})	DOCUMENT ID		TECN	COMMENT
$\mathbf{4.7^{+1.1}_{-1.0}\pm1.2}$	[552] GABYSHEV	06A	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[552] GABYSHEV 06A reports $(4.7^{+1.0}_{-0.9} \pm 0.4) \times 10^{-5}$ for $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma((\overline{\Lambda}_c^- p)_s \pi^+)/\Gamma_{total}$ — Γ_{327}/Γ

$(\overline{\Lambda}_c^- p)_s$ denotes a low-mass enhancement near 3.35 GeV/c^2.

VALUE (units 10^{-5})	DOCUMENT ID		TECN	COMMENT
$\mathbf{3.9^{+0.9}_{-0.8}\pm1.0}$	[553] GABYSHEV	06A	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[553] GABYSHEV 06A reports $(3.9^{+0.8}_{-0.7} \pm 0.4) \times 10^{-5}$ for $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\overline{\Lambda}_c^- p\pi^+\pi^0)/\Gamma_{total}$ — Γ_{328}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{1.81\pm0.29^{+0.52}_{-0.50}}$		[554,555] DYTMAN	02	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.12	90	[556] FU	97	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[554] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[555] DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \overline{p}K^+\pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.
[556] FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\overline{\Lambda}_c^- p\pi^+\pi^+\pi^-)/\Gamma_{total}$ — Γ_{329}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{2.25\pm0.25^{+0.63}_{-0.61}}$		[557,558] DYTMAN	02	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.46	90	[559] FU	97	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[557] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[558] DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \overline{p}K^+\pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.
[559] FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\overline{\Lambda}_c^- p\pi^+\pi^+\pi^-\pi^0)/\Gamma_{total}$ — Γ_{330}/Γ

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{<1.34 \times 10^{-2}}$	90	[560] FU	97	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[560] FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\Lambda_c^+ \Lambda_c^- K^+)/\Gamma_{total}$ — Γ_{331}/Γ

VALUE (units 10^{-4})	*DOCUMENT ID*		*TECN*	*COMMENT*
$\mathbf{6.5^{+1.0}_{-0.9} \pm 3.6}$	[561] GABYSHEV	06	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[561] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and B($\Lambda_c^+ \rightarrow p K^- \pi^+$) = (5.0 ± 1.3)%.

$\Gamma(\overline{\Sigma}_c(2455)^0 p)/\Gamma_{total}$ — Γ_{332}/Γ

VALUE (units 10^{-5})	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
3.7±0.8±1.0		[562] GABYSHEV	06A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<8	90	[563,564] DYTMAN	02	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<9.3	90	[563,565] GABYSHEV	02	BELL	Repl. by GABYSHEV 06A

[562] GABYSHEV 06A reports $(3.7 \pm 0.7 \pm 0.4) \times 10^{-5}$ for B($\Lambda_c^+ \rightarrow p K^- \pi^+$) = 0.05. We rescale to our best value B($\Lambda_c^+ \rightarrow p K^- \pi^+$) = $(5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[563] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[564] DYTMAN 02 measurement uses B($\Lambda_c^- \rightarrow \overline{p} K^+ \pi^-$) = 5.0 ± 1.3%. The second error includes the systematic and the uncertainty of the branching ratio.

[565] Uses the value for $\Lambda_c \rightarrow p K^- \pi^+$ branching ratio (5.0 ± 1.3)%.

$\Gamma(\overline{\Sigma}_c(2520)^0 p)/\Gamma_{total}$ — Γ_{333}/Γ

VALUE (units 10^{-5})	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
<2.7	90	GABYSHEV	06A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<4.6	90	[566,567] GABYSHEV	02	BELL	Repl. by GABYSHEV 06A

[566] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[567] Uses the value for $\Lambda_c \rightarrow p K^- \pi^+$ branching ratio (5.0 ± 1.3)%.

$\Gamma(\overline{\Sigma}_c(2455)^0 p \pi^0)/\Gamma_{total}$ — Γ_{334}/Γ

VALUE (units 10^{-4})	*DOCUMENT ID*		*TECN*	*COMMENT*
4.4±1.4±1.1	[568,569] DYTMAN	02	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[568] DYTMAN 02 reports $(4.4 \pm 1.4) \times 10^{-4}$ for B($\Lambda_c^+ \rightarrow p K^- \pi^+$) = 0.05. We rescale to our best value B($\Lambda_c^+ \rightarrow p K^- \pi^+$) = $(5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[569] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Sigma}_c(2455)^0 p \pi^- \pi^+)/\Gamma_{total}$ — Γ_{335}/Γ

VALUE (units 10^{-4})	*DOCUMENT ID*		*TECN*	*COMMENT*
4.4±1.3±1.1	[570,571] DYTMAN	02	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

[570] DYTMAN 02 reports $(4.4 \pm 1.3) \times 10^{-4}$ for B($\Lambda_c^+ \rightarrow p K^- \pi^+$) = 0.05. We rescale to our best value B($\Lambda_c^+ \rightarrow p K^- \pi^+$) = $(5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

[571] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Sigma}_c(2455)^{--} p \pi^+ \pi^+)/\Gamma_{\text{total}}$ — Γ_{336}/Γ

VALUE (units 10^{-4})		DOCUMENT ID		TECN	COMMENT
$2.8 \pm 1.0 \pm 0.7$	[572,573]	DYTMAN	02	CLE2	$e^+ e^- \to \Upsilon(4S)$

[572] DYTMAN 02 reports $(2.8 \pm 1.0) \times 10^{-4}$ for $B(\Lambda_c^+ \to p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \to p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
[573] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Lambda}_c(2593)^- / \overline{\Lambda}_c(2625)^- p \pi^+)/\Gamma_{\text{total}}$ — Γ_{337}/Γ

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
$<1.9 \times 10^{-4}$	90	[574,575]	DYTMAN	02	CLE2	$e^+ e^- \to \Upsilon(4S)$

[574] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[575] DYTMAN 02 measurement uses $B(\Lambda_c^- \to \overline{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

$\Gamma(\overline{\Xi}_c^0 \Lambda_c^+ \times B(\overline{\Xi}_c^0 \to \overline{\Xi}^+ \pi^-))/\Gamma_{\text{total}}$ — Γ_{338}/Γ

VALUE (units 10^{-5})		DOCUMENT ID		TECN	COMMENT
$5.6^{+2.3}_{-1.9} \pm 1.5$	[576,577]	CHISTOV	06A	BELL	$e^+ e^- \to \Upsilon(4S)$

[576] CHISTOV 06A reports $(5.6^{+1.9}_{-1.5} \pm 1.9) \times 10^{-5}$ for $B(\Lambda_c^+ \to p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. We rescale to our best value $B(\Lambda_c^+ \to p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
[577] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Xi}_c^0 \Lambda_c^+ \times B(\overline{\Xi}_c^0 \to \overline{\Lambda} K^+ \pi^-))/\Gamma_{\text{total}}$ — Γ_{339}/Γ

VALUE (units 10^{-5})		DOCUMENT ID		TECN	COMMENT
$4.0^{+1.3}_{-1.2} \pm 1.0$	[578,579]	CHISTOV	06A	BELL	$e^+ e^- \to \Upsilon(4S)$

[578] CHISTOV 06A reports $(4.0^{+1.1}_{-0.9} \pm 1.3) \times 10^{-5}$ for $B(\Lambda_c^+ \to p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. We rescale to our best value $B(\Lambda_c^+ \to p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
[579] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+ e^+ e^-)/\Gamma_{\text{total}}$ — Γ_{340}/Γ

Test for ΔB=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
<0.0039	90	[580]	WEIR	90B	MRK2	$e^+ e^-$ 29 GeV

[580] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ — Γ_{341}/Γ

Test for ΔB=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%		DOCUMENT ID		TECN	COMMENT
<0.0091	90	[581]	WEIR	90B	MRK2	$e^+ e^-$ 29 GeV

[581] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+ \nu\overline{\nu})/\Gamma_{total}$ — Γ_{342}/Γ

Test for ΔB=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.0 \times 10^{-4}$	90	[582] AUBERT	05H	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[582] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ e^+ e^-)/\Gamma_{total}$ — Γ_{343}/Γ

Test for ΔB=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID		TECN	COMMENT
4.9±1.0 OUR AVERAGE					
$4.2^{+1.2}_{-1.1}\pm0.2$		[583] AUBERT,B	06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$6.3^{+1.9}_{-1.7}\pm0.3$		[584] ISHIKAWA	03	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$10.5^{+2.5}_{-2.2}\pm0.7$		[583] AUBERT	03U	BABR	Repl. by AUBERT,B 06J
$<$ 14	90	[583] ABE	02	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 9	90	[583] AUBERT	02L	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 24	90	[585] ANDERSON	01B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 990	90	[586] ALBRECHT	91E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<$68000	90	[587] WEIR	90B	MRK2	e^+e^- 29 GeV
$<$ 600	90	[588] AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 2500	90	[589] AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[583] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[584] Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.
[585] The result is for di-lepton masses above 0.5 GeV.
[586] ALBRECHT 91E reports $< 9.0 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.
[587] WEIR 90B assumes B^+ production cross section from LUND.
[588] AVERY 89B reports $< 5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.
[589] AVERY 87 reports $< 2.1 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{total}$ — Γ_{344}/Γ

Test for ΔB=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$0.39^{+0.10}_{-0.09}$ OUR AVERAGE					
$0.31^{+0.15}_{-0.12}\pm0.03$		[590] AUBERT,B	06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.45^{+0.14}_{-0.12}\pm0.03$		[591] ISHIKAWA	03	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.07^{+0.19}_{-0.11}\pm0.02$		[590] AUBERT	03U	BABR	Repl. by AUBERT,B 06J
$0.98^{+0.46}_{-0.36}\pm0.16$		[590] ABE	02	BELL	Repl. by ISHIKAWA 03
$<$ 1.2	90	[590] AUBERT	02L	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 3.68	90	[592] ANDERSON	01B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 5.2	90	[593] AFFOLDER	99B	CDF	$p\overline{p}$ at 1.8 TeV
$<$ 10	90	[594] ABE	96L	CDF	Repl. by AFFOLDER 99B
$<$ 240	90	[595] ALBRECHT	91E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<$6400	90	[596] WEIR	90B	MRK2	e^+e^- 29 GeV
$<$ 170	90	[597] AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<$ 380	90	[598] AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

[590] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[591] Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.

[592] The result is for di-lepton masses above 0.5 GeV.

[593] AFFOLDER 99B measured relative to $B^+ \rightarrow J/\psi(1S)K^+$.

[594] ABE 96L measured relative to $B^+ \rightarrow J/\psi(1S)K^+$ using PDG 94 branching ratios.

[595] ALBRECHT 91E reports $< 2.2\times10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\overline{B}^0$. We rescale to 50%.

[596] WEIR 90B assumes B^+ production cross section from LUND.

[597] AVERY 89B reports $< 1.5\times10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\overline{B}^0$. We rescale to 50%.

[598] AVERY 87 reports $< 3.2\times10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0\overline{B}^0$. We rescale to 50%.

$\Gamma(K^+\ell^+\ell^-)/\Gamma_{\text{total}}$ — Γ_{345}/Γ

Test for ΔB=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$4.4^{+0.8}_{-0.7}$ OUR AVERAGE Error includes scale factor of 1.1.			
$3.8^{+0.9}_{-0.8}\pm0.2$	[599] AUBERT,B 06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$5.3^{+1.1}_{-1.0}\pm0.3$	[599] ISHIKAWA 03	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

[599] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+\overline{\nu}\nu)/\Gamma_{\text{total}}$ — Γ_{346}/Γ

Test for ΔB=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2\times10^{-5}$	90	[600] AUBERT 05H	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<2.4\times10^{-4}$	90	[600] BROWDER 01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

[600] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+ e^+ e^-)/\Gamma_{total}$ — Γ_{347}/Γ

Test for ΔB=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.75^{+0.76}_{-0.65} \pm 0.38}$		[601] AUBERT,B	06J	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.20^{+1.34}_{-0.87} \pm 0.28$		[601] AUBERT	03U	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<$ 4.6	90	[602] ISHIKAWA	03	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$<$ 8.9	90	[601] ABE	02	BELL	Repl. by ISHIKAWA 03
$<$ 9.5	90	[601] AUBERT	02L	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<$690	90	[603] ALBRECHT	91E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

[601] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[602] Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.

[603] ALBRECHT 91E reports $< 6.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \overline{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^+ \mu^+ \mu^-)/\Gamma_{total}$ — Γ_{348}/Γ

Test for ΔB=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.8\ ^{+0.6}_{-0.4}}$ **OUR AVERAGE**					
$0.97^{+0.94}_{-0.69} \pm 0.14$		[604] AUBERT,B	06J	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.65^{+0.69}_{-0.53}{}^{+0.15}_{-0.16}$		[605] ISHIKAWA	03	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$3.07^{+2.58}_{-1.78} \pm 0.42$		[604] AUBERT	03U	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<$ 3.9	90	[604] ABE	02	BELL	Repl. by ISHIKAWA 03
$<$ 17.0	90	[604] AUBERT	02L	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<$1200	90	[606] ALBRECHT	91E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

[604] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

[605] Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence. The 90% C.L. upper limit is 2.2×10^{-6}.

[606] ALBRECHT 91E reports $< 1.1 \times 10^{-3}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \overline{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^+ \ell^+ \ell^-)/\Gamma_{total}$ — Γ_{349}/Γ

Test for ΔB=1 weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID		TECN	COMMENT
$\mathbf{7.3^{+5.0}_{-4.2} \pm 2.1}$		[607] AUBERT,B	06J	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<$22	90	[607] ISHIKAWA	03	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

[607] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{total}$ — Γ_{350}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0064	90	[608] WEIR	90B	MRK2	e^+e^- 29 GeV

[608] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+ e^- \mu^+)/\Gamma_{total}$ — Γ_{351}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.0064	90	[609] WEIR	90B	MRK2	e^+e^- 29 GeV

[609] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^+ \mu^-)/\Gamma_{total}$ — Γ_{352}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-7})	CL%	DOCUMENT ID		TECN	COMMENT
<0.91	90	[610] AUBERT,B	06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<8	90	[610] AUBERT	02L	BABR	Repl. by AUBERT,B 06J
$<6.4 \times 10^4$	90	[611] WEIR	90B	MRK2	e^+e^- 29 GeV

[610] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[611] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^- \mu^+)/\Gamma_{total}$ — Γ_{353}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-7})	CL%	DOCUMENT ID		TECN	COMMENT
<1.3	90	[612] AUBERT,B	06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<6.4 \times 10^4$	90	[613] WEIR	90B	MRK2	e^+e^- 29 GeV

[612] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[613] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^\pm \mu^\mp)/\Gamma_{total}$ — Γ_{354}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID		TECN	COMMENT
<0.91	90	[614] AUBERT,B	06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[614] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+ e^+ \mu^-)/\Gamma_{total}$ — Γ_{355}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID		TECN	COMMENT
<13	90	[615] AUBERT,B	06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[615] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+ e^- \mu^+)/\Gamma_{total}$ — Γ_{356}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID		TECN	COMMENT
<9.9	90	[616] AUBERT,B	06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

[616] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+ e^\pm \mu^\mp)/\Gamma_{total}$ — Γ_{357}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-7})	CL%	DOCUMENT ID		TECN	COMMENT
< 1.4	90	[617] AUBERT,B	06J	BABR	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<79	90	[617] AUBERT	02L	BABR	Repl. by AUBERT,B 06J

[617] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^- e^+ e^+)/\Gamma_{total}$ — Γ_{358}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.6 × 10^{-6}	90	[618] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0039	90	[619] WEIR	90B	MRK2	$e^+ e^-$ 29 GeV

[618] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[619] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{total}$ — Γ_{359}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.4 × 10^{-6}	90	[620] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0091	90	[621] WEIR	90B	MRK2	$e^+ e^-$ 29 GeV

[620] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[621] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{total}$ — Γ_{360}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.3 × 10^{-6}	90	[622] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0064	90	[623] WEIR	90B	MRK2	$e^+ e^-$ 29 GeV

[622] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[623] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\rho^- e^+ e^+)/\Gamma_{total}$ — Γ_{361}/Γ

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<2.6	90	[624] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[624] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho^- \mu^+ \mu^+)/\Gamma_{total}$ — Γ_{362}/Γ

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<5.0	90	[625] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[625] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho^- e^+ \mu^+)/\Gamma_{total}$ — Γ_{363}/Γ

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<3.3	90	[626] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[626] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^- e^+ e^+)/\Gamma_{total}$ — Γ_{364}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.0 × 10⁻⁶	90	[627] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0039	90	[628] WEIR	90B	MRK2	$e^+ e^-$ 29 GeV

[627] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[628] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^- \mu^+ \mu^+)/\Gamma_{total}$ — Γ_{365}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<1.8 × 10⁻⁶	90	[629] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0091	90	[630] WEIR	90B	MRK2	$e^+ e^-$ 29 GeV

[629] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[630] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^- e^+ \mu^+)/\Gamma_{total}$ — Γ_{366}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<2.0 × 10⁻⁶	90	[631] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0064	90	[632] WEIR	90B	MRK2	$e^+ e^-$ 29 GeV

[631] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
[632] WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^*(892)^- e^+ e^+)/\Gamma_{total}$ — Γ_{367}/Γ

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<2.8	90	[633] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[633] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^- \mu^+ \mu^+)/\Gamma_{total}$ — Γ_{368}/Γ

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<8.3	90	[634] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[634] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^- e^+ \mu^+)/\Gamma_{total}$ — Γ_{369}/Γ

Test of total lepton number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID		TECN	COMMENT
<4.4	90	[635] EDWARDS	02B	CLE2	$e^+ e^- \to \Upsilon(4S)$

[635] Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

POLARIZATION IN B^+ DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal (L) or both are transverse and parallel ($\|$) or perpendicular ($\perp$) to each other with the parameters Γ_L/Γ, $\Gamma_\perp/\Gamma$, and the relative phases $\phi_\|$ and $\phi_\perp$. See the definitions in the note on "Polarization in B Decays" review in the B^0 Particle Listings.

Γ_L/Γ in $B^+ \to \overline{D}^{*0} \rho^+$

VALUE	DOCUMENT ID		TECN	COMMENT
0.892±0.018±0.016	CSORNA	03	CLE2	$e^+ e^- \to \Upsilon(4S)$

Γ_L/Γ in $B^+ \to \overline{D}^{*0} K^{*+}$

VALUE	DOCUMENT ID		TECN	COMMENT
0.86±0.06±0.03	AUBERT	04K	BABR	$e^+ e^- \to \Upsilon(4S)$

Γ_L/Γ in $B^+ \to J/\psi K^{*+}$

VALUE	DOCUMENT ID		TECN	COMMENT
0.604±0.015±0.018	ITOH	05	BELL	$e^+ e^- \to \Upsilon(4S)$

$\Gamma_\perp/\Gamma$ in $B^+ \to J/\psi K^{*+}$

VALUE	DOCUMENT ID		TECN	COMMENT
0.180±0.014±0.010	ITOH	05	BELL	$e^+ e^- \to \Upsilon(4S)$

Γ_L/Γ in $B^+ \to \phi K^*(892)^+$

VALUE	DOCUMENT ID		TECN	COMMENT
0.50±0.07 OUR AVERAGE				
0.52±0.08±0.03	CHEN	05A	BELL	$e^+ e^- \to \Upsilon(4S)$
0.46±0.12±0.03	AUBERT	03V	BABR	$e^+ e^- \to \Upsilon(4S)$

$\Gamma_\perp/\Gamma$ in $B^+ \to \phi K^{*+}$

VALUE	DOCUMENT ID		TECN	COMMENT
0.19±0.08±0.02	CHEN	05A	BELL	$e^+ e^- \to \Upsilon(4S)$

$\phi_\|$ in $B^+ \to \phi K^{*+}$

VALUE (°)	DOCUMENT ID		TECN	COMMENT
2.10±0.28±0.04	CHEN	05A	BELL	$e^+ e^- \to \Upsilon(4S)$

$\phi_\perp$ in $B^+ \to \phi K^{*+}$

VALUE (°)	DOCUMENT ID		TECN	COMMENT
2.31±0.30±0.07	CHEN	05A	BELL	$e^+ e^- \to \Upsilon(4S)$

Γ_L/Γ in $B^+ \to \rho^0 K^*(892)^+$

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.96^{+0.04}_{-0.15}\pm0.04}$	AUBERT	03V	BABR	$e^+e^- \to \Upsilon(4S)$

$\Gamma_L/\Gamma(B^+ \to K^*(892)^0 \rho^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.48±0.08 OUR AVERAGE				
$0.52\pm0.10\pm0.04$	AUBERT,B	06G	BABR	$e^+e^- \to \Upsilon(4S)$
$0.43\pm0.11^{+0.05}_{-0.02}$	ZHANG	05D	BELL	$e^+e^- \to \Upsilon(4S)$

Γ_L/Γ in $B^+ \to \rho^+ \rho^0$

VALUE	DOCUMENT ID		TECN	COMMENT
0.91 ±0.04 OUR AVERAGE				
$0.905\pm0.042^{+0.023}_{-0.027}$	AUBERT,BE	06G	BABR	$e^+e^- \to \Upsilon(4S)$
$0.948\pm0.106\pm0.021$	ZHANG	03B	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.97\ ^{+0.03}_{-0.07}\ \pm0.04$	AUBERT	03V	BABR	Repl. by AUBERT,BE 06G

Γ_L/Γ in $B^+ \to \omega \rho^+$

VALUE	DOCUMENT ID		TECN	COMMENT
0.82±0.11±0.02	AUBERT,B	06T	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.88^{+0.12}_{-0.15}\pm0.03$	AUBERT	05O	BABR	Repl. by AUBERT,B 06T

CP VIOLATION

A_{CP} is defined as

$$\frac{B(B^- \to \overline{f}) - B(B^+ \to f)}{B(B^- \to \overline{f}) + B(B^+ \to f)},$$

the *CP*-violation charge asymmetry of exclusive B^- and B^+ decay.

$A_{CP}(B^+ \to J/\psi(1S) K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.015±0.017 OUR AVERAGE Error includes scale factor of 1.2.				
$0.030\pm0.014\pm0.010$	[636] AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$
$-0.026\pm0.022\pm0.017$	ABE	03B	BELL	$e^+e^- \to \Upsilon(4S)$
$0.018\pm0.043\pm0.004$	[637] BONVICINI	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.03\ \pm0.015\pm0.006$	AUBERT	04P	BABR	Repl. by AUBERT 05J
$0.003\pm0.030\pm0.004$	AUBERT	02F	BABR	Repl. by AUBERT 04P

[636] The result reported corresponds to $-A_{CP}$.

[637] A +0.3% correction is applied due to a slightly higher reconstruction efficiency for the positive kaons.

$A_{CP}(B^+ \to J/\psi(1S)\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.09 ±0.08 OUR AVERAGE				
0.123±0.085±0.004	AUBERT	04P	BABR	$e^+e^- \to \Upsilon(4S)$
−0.023±0.164±0.015	ABE	03B	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.01 ±0.22 ±0.01	AUBERT	02F	BABR	Repl. by AUBERT 04P

$A_{CP}(B^+ \to J/\psi K^*(892)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.048±0.029±0.016	[638] AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$

[638] The result reported corresponds to $-A_{CP}$.

$A_{CP}(B^+ \to \psi(2S)K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.025±0.024 OUR AVERAGE				
0.052±0.059±0.020	AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$
−0.042±0.020±0.017	ABE	03B	BELL	$e^+e^- \to \Upsilon(4S)$
0.02 ±0.091±0.01	[639] BONVICINI	00	CLE2	$e^+e^- \to \Upsilon(4S)$

[639] A +0.3% correction is applied due to a slightly higher reconstruction efficiency for the positive kaons.

$A_{CP}(B^+ \to \psi(2S)K^*(892)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.077±0.207±0.051	[640] AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$

[640] The result reported corresponds to $-A_{CP}$.

$A_{CP}(B^+ \to \chi_{c1}(1P)\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.07±0.18±0.02	KUMAR	06	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \chi_{c0}K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$-0.065\pm0.20^{+0.035}_{-0.024}$	GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \chi_{c1}K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.009±0.033 OUR AVERAGE				
−0.01 ±0.03 ±0.02	KUMAR	06	BELL	$e^+e^- \to \Upsilon(4S)$
−0.003±0.076±0.017	[641] AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$

[641] The result reported corresponds to $-A_{CP}$.

$A_{CP}(B^+ \to \chi_{c1}K^*(892)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.471±0.378±0.268	[642] AUBERT	05J	BABR	$e^+e^- \to \Upsilon(4S)$

[642] The result reported corresponds to $-A_{CP}$.

$A_{CP}(B^+ \to \overline{D}^0 \pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.008±0.008	ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D_{CP(+1)} \pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.035±0.024	ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D_{CP(-1)} \pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.017±0.026	ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \overline{D}^0 K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.066±0.036	ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.003±0.080±0.037	[643] ABE	03D	BELL	Repl. by SWAIN 03
0.04 ±0.06 ±0.03	[644] SWAIN	03	BELL	Repl. by ABE 06

[643] Corresponds to 90% confidence range $-0.15 < A_{CP} < 0.16$.
[644] Corresponds to 90% confidence range $-0.07 < A_{CP} < 0.15$.

$r_B(B^+ \to D^0 K^+)$

$r_B^{(*)}$ and $\delta_B^{(*)}$ are the amplitude ratios and relative strong phases between the amplitudes of $A_{CP}(B^+ \to D^{(*)0} K^+)$ and $A_{CP}(B^+ \to \overline{D}^{(*)0} K^+)$,

VALUE	DOCUMENT ID		TECN	COMMENT
0.14 ±0.06 OUR AVERAGE				
$0.159^{+0.054}_{-0.050}\pm0.050$	[645] POLUEKTOV	06	BELL	$e^+e^- \to \Upsilon(4S)$
0.12 ±0.08 ±0.05	[646] AUBERT,B	05Y	BABR	$e^+e^- \to \Upsilon(4S)$

[645] Uses a Dalitz plot analysis of the $\overline{D}^0 \to K^0_S \pi^+\pi^-$ decays; Combines the DK^+, $D^* K^+$ and DK^{*+} modes.
[646] Uses a Dalitz analysis of neutral D decays to $K^0_S \pi^+\pi^-$ in the processes $B^\pm \to D^{(*)} K^\pm$, $D^* \to D\pi^0$, $D\gamma$.

$\delta_B(B^+ \to D^0 K^+)$

VALUE (degrees)	DOCUMENT ID		TECN	COMMENT
135 ±26 OUR AVERAGE				
$145.7^{+19.0}_{-19.7}\pm23.1$	[647] POLUEKTOV	06	BELL	$e^+e^- \to \Upsilon(4S)$
$104 \pm45 \ ^{+23}_{-32}$	[648] AUBERT,B	05Y	BABR	$e^+e^- \to \Upsilon(4S)$

[647] Uses a Dalitz plot analysis of the $\overline{D}^0 \to K^0_S \pi^+\pi^-$ decays; Combines the DK^+, $D^* K^+$ and DK^{*+} modes.
[648] Uses a Dalitz analysis of neutral D decays to $K^0_S \pi^+\pi^-$ in the processes $B^\pm \to D^{(*)} K^\pm$, $D^* \to D\pi^0$, $D\gamma$.

$r_B(B^+ \to DK^{*+})$

r_B and δ_B are the amplitude ratios and relative strong phases between the amplitudes of $A_{CP}(B^+ \to DK^{*+})$ and $A_{CP}(B^+ \to \overline{D}K^{*+})$,

VALUE	DOCUMENT ID		TECN	COMMENT
$0.564^{+0.216}_{-0.155} \pm 0.093$	[649] POLUEKTOV	06	BELL	$e^+e^- \to \Upsilon(4S)$

[649] Uses a Dalitz plot analysis of the $\overline{D}^0 \to K^0_S \pi^+\pi^-$ decays; Combines the DK^+, D^*K^+ and DK^{*+} modes.

$\delta_B(B^+ \to DK^{*+})$

VALUE (degrees)	DOCUMENT ID		TECN	COMMENT
$242.6^{+20.2}_{-23.2} \pm 49.4$	[650] POLUEKTOV	06	BELL	$e^+e^- \to \Upsilon(4S)$

[650] Uses a Dalitz plot analysis of the $\overline{D}^0 \to K^0_S \pi^+\pi^-$ decays; Combines the DK^+, D^*K^+ and DK^{*+} modes.

$A_{CP}(B^+ \to [K^-\pi^+]_D K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$+0.88^{+0.77}_{-0.62} \pm 0.06$	SAIGO	05	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to [K^-\pi^+]_{\overline{D}} K^*(892)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$-0.22 \pm 0.61 \pm 0.17$	AUBERT,B	05V	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to [K^-\pi^+]_D \pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$+0.30^{+0.29}_{-0.25} \pm 0.06$	SAIGO	05	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to [\pi^+\pi^-\pi^0]_D K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$-0.02 \pm 0.16 \pm 0.03$	AUBERT,B	05T	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D_{CP(+1)} K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.22 ± 0.14 OUR AVERAGE	Error includes scale factor of 1.4.			
$0.06 \pm 0.14 \pm 0.05$	ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$
$0.35 \pm 0.13 \pm 0.04$	AUBERT	06J	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.07 \pm 0.17 \pm 0.06$	AUBERT	04N	BABR	Repl. by AUBERT 06J
$0.29 \pm 0.26 \pm 0.05$	[651] ABE	03D	BELL	Repl. by SWAIN 03
$0.06 \pm 0.19 \pm 0.04$	[652] SWAIN	03	BELL	Repl. by ABE 06

[651] Corresponds to 90% confidence range $-0.14 < A_{CP} < 0.73$.

[652] Corresponds to 90% confidence range $-0.26 < A_{CP} < 0.38$.

$A_{CP}(B^+ \to D_{CP(-1)} K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.09±0.10 OUR AVERAGE				
$-0.12\pm0.14\pm0.05$	ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$
$-0.06\pm0.13\pm0.04$	AUBERT	06J	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.22\pm0.24\pm0.04$	[653] ABE	03D	BELL	Repl. by SWAIN 03
$-0.19\pm0.17\pm0.05$	[654] SWAIN	03	BELL	Repl. by ABE 06

[653] Corresponds to 90% confidence range $-0.62 < A_{CP} < 0.18$.
[654] Corresponds to 90% confidence range $-0.47 < A_{CP} < 0.11$.

$A_{CP}(B^+ \to \overline{D}^{*0} \pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.014±0.015	ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to (D^*_{CP(+1)})^0 \pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.021±0.045	ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to (D^*_{CP(-1)})^0 \pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.090±0.051	ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D^{*0} K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.089±0.086	ABE	06	BELL	$e^+e^- \to \Upsilon(4S)$

$r^*_B(B^+ \to D^{*0} K^+)$

$r^{(*)}_B$ and $\delta^{(*)}_B$ are the amplitude ratios and relative strong phases between the amplitudes of $A_{CP}(B^+ \to D^{(*)0} K^+)$ and $A_{CP}(B^+ \to \overline{D}^{(*)0} K^+)$,

VALUE	DOCUMENT ID		TECN	COMMENT
0.17 ±0.08 OUR AVERAGE				
$0.175^{+0.108}_{-0.099}\pm0.050$	[655] POLUEKTOV	06	BELL	$e^+e^- \to \Upsilon(4S)$
$0.17\pm0.10\pm0.04$	[656] AUBERT,B	05Y	BABR	$e^+e^- \to \Upsilon(4S)$

[655] Uses a Dalitz plot analysis of the $\overline{D}^0 \to K^0_S \pi^+\pi^-$ decays; Combines the DK^+, D^*K^+ and DK^{*+} modes.
[656] Uses a Dalitz analysis of neutral D decays to $K^0_S \pi^+\pi^-$ in the processes $B^\pm \to D^{(*)}K^\pm$, $D^* \to D\pi^0$, $D\gamma$.

$\delta^*_B(B^+ \to D^{*0} K^+)$

VALUE (degrees)	DOCUMENT ID		TECN	COMMENT
299 ±31 OUR AVERAGE				
$302.0^{+33.8}_{-35.1}\pm23.7$	[657] POLUEKTOV	06	BELL	$e^+e^- \to \Upsilon(4S)$
$296\pm41\ ^{+20}_{-19}$	[658] AUBERT,B	05Y	BABR	$e^+e^- \to \Upsilon(4S)$

[657] Uses a Dalitz plot analysis of the $\overline{D}^0 \to K_S^0 \pi^+ \pi^-$ decays; Combines the $D K^+$, $D^* K^+$ and $D K^{*+}$ modes.
[658] Uses a Dalitz analysis of neutral D decays to $K_S^0 \pi^+ \pi^-$ in the processes $B^\pm \to D^{(*)} K^\pm$, $D^* \to D \pi^0$, $D \gamma$.

$A_{CP}(B^+ \to D^{*0}_{CP(+1)} K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.15±0.16 OUR AVERAGE				
$-0.20 \pm 0.22 \pm 0.04$	ABE	06	BELL	$e^+ e^- \to \Upsilon(4S)$
$-0.10 \pm 0.23 ^{+0.03}_{-0.04}$	AUBERT	05N	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D^{*}_{CP(-1)} K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.13±0.30±0.08	ABE	06	BELL	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D_{CP(+1)} K^*(892)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.08±0.19±0.08	AUBERT,B	05U	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D_{CP(-1)} K^*(892)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.26±0.40±0.12	AUBERT,B	05U	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D^{*+} \overline{D}^{*0})$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.15±0.11±0.02	AUBERT,B	06A	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D^{*+} \overline{D}^{0})$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.06±0.13±0.02	AUBERT,B	06A	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D^{+} \overline{D}^{*0})$

VALUE	DOCUMENT ID		TECN	COMMENT
0.13±0.18±0.04	AUBERT,B	06A	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to D^{+} \overline{D}^{0})$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.13±0.14±0.02	AUBERT,B	06A	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to K_S^0 \pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.00 ±0.04 OUR AVERAGE	Error includes scale factor of 1.2.			
$-0.029 \pm 0.039 \pm 0.010$	[659] AUBERT,BE	06C	BABR	$e^+ e^- \to \Upsilon(4S)$
$0.05 \pm 0.05 \pm 0.01$	[660] CHAO	05A	BELL	$e^+ e^- \to \Upsilon(4S)$
0.18 ± 0.24	[661] CHEN	00	CLE2	$e^+ e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.09 \pm 0.05 \pm 0.01$	[662] AUBERT,BE	05E	BABR	Repl. by AUBERT,BE 06C
$-0.05 \pm 0.08 \pm 0.01$	[663] AUBERT	04M	BABR	Repl. by AUBERT,BE 05E
$0.07 {}^{+0.09}_{-0.08} {}^{+0.01}_{-0.03}$	[664] UNNO	03	BELL	Repl. by CHAO 05A
$0.46 \pm 0.15 \pm 0.02$	[665] CASEY	02	BELL	Repl. by UNNO 03
$0.098 {}^{+0.430}_{-0.343} {}^{+0.020}_{-0.063}$	[666] ABE	01K	BELL	Repl. by CASEY 02
$-0.21 \pm 0.18 \pm 0.03$	[667] AUBERT	01E	BABR	Repl. by AUBERT 04M

[659] Corresponds to 90% confidence range $-0.092 < A_{CP} < 0.036$.
[660] Corresponds to a 90% CL interval of $-0.04 < A_{CP} < 0.13$.
[661] Corresponds to 90% confidence range $-0.22 < A_{CP} < 0.56$.
[662] Corresponds to 90% confidence range $-0.16 < A_{CP} < -0.02$.
[663] 90% CL interval $-0.18 < A_{CP} < 0.08$
[664] Corresponds to 90% confidence range $-0.10 < A_{CP} < +0.22$.
[665] Corresponds to 90% confidence range $+0.19 < A_{CP} < +0.72$.
[666] Corresponds to 90% confidence range $-0.53 < A_{CP} < 0.82$.
[667] Corresponds to 90% confidence range $-0.51 < A_{CP} < 0.09$.

$A_{CP}(B^+ \to K^+ \pi^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.04 ± 0.04 OUR AVERAGE			
$0.06 \pm 0.06 \pm 0.01$	[668] AUBERT 05L	BABR	$e^+ e^- \to \Upsilon(4S)$
$0.04 \pm 0.05 \pm 0.02$	[669] CHAO 04B	BELL	$e^+ e^- \to \Upsilon(4S)$
-0.29 ± 0.23	[670] CHEN 00	CLE2	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.06 \pm 0.06 \pm 0.02$	[668] CHAO 05A	BELL	Repl. by CHAO 04B
$-0.09 \pm 0.09 \pm 0.01$	[671] AUBERT 03L	BABR	Repl. by AUBERT 05L
$-0.02 \pm 0.19 \pm 0.02$	[672] CASEY 02	BELL	Repl. by CHAO 04B
$-0.059 {}^{+0.222}_{-0.196} {}^{+0.055}_{-0.017}$	[673] ABE 01K	BELL	Repl. by CASEY 02
$0.00 \pm 0.18 \pm 0.04$	[674] AUBERT 01E	BABR	Repl. by AUBERT 03L

[668] Corresponds to a 90% CL interval of $-0.06 < A_{CP} < 0.18$.
[669] Corresponds to 90% CL interval of $-0.05 < A_{CP} < 0.13$.
[670] Corresponds to 90% confidence range $-0.67 < A_{CP} < 0.09$.
[671] Corresponds to 90% confidence range $-0.24 < A_{CP} < 0.06$.
[672] Corresponds to 90% confidence range $-0.35 < A_{CP} < +0.30$.
[673] Corresponds to 90% confidence range $-0.40 < A_{CP} < 0.36$.
[674] Corresponds to 90% confidence range $-0.30 < A_{CP} < +0.30$.

$A_{CP}(B^+ \to K^+ \eta')$

VALUE	DOCUMENT ID	TECN	COMMENT
0.026 ± 0.021 OUR AVERAGE			
$0.028 \pm 0.028 \pm 0.021$	SCHUEMANN 06	BELL	$e^+ e^- \to \Upsilon(4S)$
$0.033 \pm 0.028 \pm 0.005$	[675] AUBERT 05M	BABR	$e^+ e^- \to \Upsilon(4S)$
$-0.11 \pm 0.11 \pm 0.02$	[676] AUBERT 02E	BABR	$e^+ e^- \to \Upsilon(4S)$
0.03 ± 0.12	[677] CHEN 00	CLE2	$e^+ e^- \to \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.037 \pm 0.045 \pm 0.011$	[678] AUBERT	03W	BABR	Repl. by AUBERT 05M
$-0.015 \pm 0.070 \pm 0.009$	[679] CHEN	02B	BELL	Repl. by SCHUEMANN 06
$0.06 \pm 0.15 \pm 0.01$	[680] ABE	01M	BELL	Repl. by CHEN 02B

[675] Corresponds to 90% confidence range $-0.012 < A_{CP} < 0.078$.
[676] Corresponds to 90% confidence range $-0.28 < A_{CP} < 0.07$.
[677] Corresponds to 90% confidence range $-0.17 < A_{CP} < 0.23$.
[678] Corresponds to 90% confidence range $-0.04 < A_{CP} < 0.11$.
[679] Corresponds to 90% confidence range $-0.13 < A_{CP} < 0.10$.
[680] Corresponds to 90% confidence range $-0.20 < A_{CP} < 0.32$.

$A_{CP}(B^+ \to \eta' K^*(892)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.30^{+0.33}_{-0.37} \pm 0.02}$	AUBERT	07E	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \eta K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.25±0.14 OUR AVERAGE				
$-0.20 \pm 0.15 \pm 0.01$	AUBERT,B	05K	BABR	$e^+ e^- \to \Upsilon(4S)$
$-0.49 \pm 0.31 \pm 0.07$	CHANG	05A	BELL	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.52 \pm 0.24 \pm 0.01$	AUBERT	04H	BABR	Repl. by AUBERT,B 05K

$A_{CP}(B^+ \to \eta K^*(892)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.01±0.08±0.02	AUBERT,B	06H	BABR	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.13 \pm 0.14 \pm 0.02$	AUBERT,B	04D	BABR	Repl. by AUBERT,B 06H

$A_{CP}(B^+ \to \eta K_0^*(1430)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.05±0.13±0.02	AUBERT,B	06H	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \eta K_2^*(1430)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.45±0.30±0.02	AUBERT,B	06H	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \omega K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.05±0.06 OUR AVERAGE				
$0.05 \pm 0.09 \pm 0.01$	AUBERT,B	06E	BABR	$e^+ e^- \to \Upsilon(4S)$
$0.05^{+0.08}_{-0.07} \pm 0.01$	JEN	06	BELL	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.09 \pm 0.17 \pm 0.01$	AUBERT	04H	BABR	Repl. by AUBERT,B 06E
$0.06^{+0.21}_{-0.18} \pm 0.01$	[681] WANG	04A	BELL	Repl. by JEN 06
$-0.21 \pm 0.28 \pm 0.03$	[682] LU	02	BELL	Repl. by WANG 04A

[681] Corresponds to 90% CL interval $0.15 < A_{CP} < 0.90$
[682] Corresponds to 90% confidence range $-0.70 < A_{CP} < +0.38$.

$A_{CP}(B^+ \to K^{*0}\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
-0.08 ± 0.10 OUR AVERAGE	Error includes scale factor of 1.8.			
$-0.149 \pm 0.064 \pm 0.022$	GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$0.068 \pm 0.078^{+0.070}_{-0.067}$	AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to K^+\pi^-\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.023 ± 0.031 OUR AVERAGE	Error includes scale factor of 1.2.			
$0.049 \pm 0.026 \pm 0.020$	GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$-0.013 \pm 0.037 \pm 0.011$	AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.01 \pm 0.07 \pm 0.03$	AUBERT	03M	BABR	Repl. by AUBERT,B 05N

$A_{CP}(B^+ \to f_0(980)K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$-0.04^{+0.08}_{-0.07}$ OUR AVERAGE	Error includes scale factor of 1.1.			
$-0.31 \pm 0.25 \pm 0.08$	[683] AUBERT	06O	BABR	$e^+e^- \to \Upsilon(4S)$
$-0.077 \pm 0.065^{+0.046}_{-0.026}$	GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$0.088 \pm 0.095^{+0.097}_{-0.056}$	AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$

[683] Measured in the $B^+ \to K^+K^-K^+$ decay.

$A_{CP}(B^+ \to f_2(1270)K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$-0.59 \pm 0.22 \pm 0.036$	GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to X_0(1550)K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$-0.04 \pm 0.07 \pm 0.02$	[684] AUBERT	06O	BABR	$e^+e^- \to \Upsilon(4S)$

[684] Measured in the $B^+ \to K^+K^-K^+$ decay.

$A_{CP}(B^+ \to \rho^0 K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$0.31^{+0.11}_{-0.09}$ OUR AVERAGE				
$0.30 \pm 0.11^{+0.11}_{-0.04}$	GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$0.32 \pm 0.13^{+0.10}_{-0.08}$	AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to K_0^*(1430)^0\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.00 ± 0.07 OUR AVERAGE	Error includes scale factor of 2.4.			
$0.076 \pm 0.038^{+0.028}_{-0.022}$	GARMASH	06	BELL	$e^+e^- \to \Upsilon(4S)$
$-0.064 \pm 0.032^{+0.023}_{-0.026}$	AUBERT,B	05N	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to K^*(892)^+ \pi^0)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.04±0.29±0.05	AUBERT	05X	BABR	$e^+e^- -> \Upsilon(4S)$

$A_{CP}(B^+ \to \rho^0 K^*(892)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{0.20^{+0.32}_{-0.29}\pm0.04}$	AUBERT	03V	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to K^*(892)^+ f_0(980))$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.34±0.21±0.03	AUBERT,B	06G	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to K^*(892)^0 \rho^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.01±0.16±0.02	AUBERT,B	06G	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to K^0 K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.10±0.26±0.03	[685] AUBERT,BE	06C	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.15±0.33±0.03	[686] AUBERT,BE	05E	BABR	Repl. by AUBERT,BE 06C

[685] Corresponds to 90% confidence range $-0.31 < A_{CP} < 0.54$.
[686] Corresponds to 90% confidence range $-0.43 < A_{CP} < 0.68$.

$A_{CP}(B^+ \to K^+ K^0_S K^0_S)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.04±0.11±0.02	[687] AUBERT,B	04V	BABR	$e^+e^- \to \Upsilon(4S)$

[687] Corresponds to 90% confidence range $-0.23 < A_{CP} < 0.15$.

$A_{CP}(B^+ \to K^+ K^- K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.017±0.026±0.015	AUBERT	06O	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.02 ±0.07 ±0.03	AUBERT	03M	BABR	Repl. by AUBERT 06O

$A_{CP}(B^+ \to K^{*+} K^+ K^-)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.11±0.08±0.03	AUBERT,B	06U	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to K^{*+} \pi^+ \pi^-)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.07±0.07±0.04	AUBERT,B	06U	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \phi K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
-0.01 ± 0.06 OUR AVERAGE				
$0.00 \pm 0.08 \pm 0.02$	AUBERT	06O	BABR	$e^+ e^- \to \Upsilon(4S)$
$-0.07 \pm 0.17 ^{+0.03}_{-0.02}$	ACOSTA	05J	CDF	$p\overline{p}$ at 1.96 TeV
$0.01 \pm 0.12 \pm 0.05$	[688] CHEN	03B	BELL	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.04 \pm 0.09 \pm 0.01$	[689] AUBERT	04A	BABR	Repl. by AUBERT 06O
$-0.05 \pm 0.20 \pm 0.03$	[690] AUBERT	02E	BABR	$e^+ e^- \to \Upsilon(4S)$

[688] Corresponds to 90% confidence range $-0.20 < A_{CP} < 0.22$.
[689] Corresponds to 90% confidence range $-0.10 < A_{CP} < 0.18$.
[690] Corresponds to 90% confidence range $-0.37 < A_{CP} < 0.28$.

$A_{CP}(B^+ \to \phi K^*(892)^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.05 ± 0.11 OUR AVERAGE				
$-0.02 \pm 0.14 \pm 0.03$	[691] CHEN	05A	BELL	$e^+ e^- \to \Upsilon(4S)$
$0.16 \pm 0.17 \pm 0.03$	AUBERT	03V	BABR	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.13 \pm 0.29 ^{+0.08}_{-0.11}$	[692] CHEN	03B	BELL	Repl. by CHEN 05A
$-0.43 ^{+0.36}_{-0.30} \pm 0.06$	[693] AUBERT	02E	BABR	Repl. by AUBERT 03V

[691] Corresponds to 90% confidence range $-0.25 < A_{CP} < 0.22$.
[692] Corresponds to 90% confidence range $-0.64 < A_{CP} < 0.36$.
[693] Corresponds to 90% confidence range $-0.88 < A_{CP} < 0.18$.

$A_{CP}(B^+ \to \phi K^+ \gamma)$

VALUE	DOCUMENT ID		TECN	COMMENT
$-0.26 \pm 0.14 \pm 0.05$	AUBERT	07Q	BABR	$e^+ e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \eta K^+ \gamma)$

VALUE	DOCUMENT ID		TECN	COMMENT
-0.13 ± 0.08 OUR AVERAGE				
$-0.09 \pm 0.12 \pm 0.01$	[694] AUBERT,B	06M	BABR	$e^+ e^- \to \Upsilon(4S)$
$-0.16 \pm 0.09 \pm 0.06$	[695] NISHIDA	05	BELL	$e^+ e^- \to \Upsilon(4S)$

[694] $m_{\eta K} < 3.25$ GeV/c^2.
[695] $m_{\eta K} < 2.4$ GeV/c^2

$A_{CP}(B^+ \to \pi^+ \pi^0)$

VALUE	DOCUMENT ID		TECN	COMMENT
-0.02 ± 0.07 OUR AVERAGE				
$-0.01 \pm 0.10 \pm 0.02$	[696] AUBERT	05L	BABR	$e^+ e^- \to \Upsilon(4S)$
$-0.02 \pm 0.10 \pm 0.01$	[697] CHAO	04B	BELL	$e^+ e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.00 \pm 0.10 \pm 0.02$	[698] CHAO	05A	BELL	Repl. by CHAO 04B
$-0.03 ^{+0.18}_{-0.17} \pm 0.02$	[699] AUBERT	03L	BABR	Repl. by AUBERT 05L
$0.30 \pm 0.30 ^{+0.06}_{-0.04}$	[700] CASEY	02	BELL	Repl. by CHAO 04B

[696] Corresponds to a 90% CL interval of $-0.19 < A_{CP} < 0.21$.
[697] This corresponds to 90% CL interval of $-0.18 < A_{CP} < 0.14$.
[698] Corresponds to a 90% CL interval of $-0.17 < A_{CP} < 0.16$.
[699] Corresponds to 90% confidence range $-0.32 < A_{CP} < 0.27$.
[700] Corresponds to 90% confidence range $-0.23 < A_{CP} < +0.86$.

$A_{CP}(B^+ \to \pi^+\pi^-\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{-0.007 \pm 0.077 \pm 0.025}$	AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.39 \pm 0.33 \pm 0.12$	AUBERT	03M	BABR	Repl. by AUBERT 05G

$A_{CP}(B^+ \to \rho^0\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{-0.074 \pm 0.120 ^{+0.035}_{-0.055}}$	AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.19 \pm 0.11 \pm 0.02$	AUBERT	04Z	BABR	Repl. by AUBERT,B 05G

$A_{CP}(B^+ \to f_2(1270)\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
$\mathbf{-0.004 \pm 0.247 ^{+0.028}_{-0.032}}$	AUBERT,B	05G	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \rho^+\pi^0)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.15±0.12 OUR AVERAGE				
$0.06 \pm 0.17 ^{+0.04}_{-0.05}$	ZHANG	05A	BELL	$e^+e^- \to \Upsilon(4S)$
$0.24 \pm 0.16 \pm 0.06$	AUBERT	04Z	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \rho^+\rho^0)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.08±0.13 OUR AVERAGE				
$-0.12 \pm 0.13 \pm 0.10$	AUBERT,BE	06G	BABR	$e^+e^- \to \Upsilon(4S)$
$0.00 \pm 0.22 \pm 0.03$	ZHANG	03B	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.19 \pm 0.23 \pm 0.03$	AUBERT	03V	BABR	Repl. by AUBERT,BE 06G

$A_{CP}(B^+ \to \omega\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.04±0.06 OUR AVERAGE				
$-0.01 \pm 0.10 \pm 0.01$	AUBERT,B	06E	BABR	$e^+e^- \to \Upsilon(4S)$
$-0.02 \pm 0.09 \pm 0.01$	JEN	06	BELL	$e^+e^- \to \Upsilon(4S)$
-0.34 ± 0.25	[701] CHEN	00	CLE2	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.03 \pm 0.16 \pm 0.01$	AUBERT	04H	BABR	Repl. by AUBERT,B 06E
$0.50 ^{+0.23}_{-0.20} \pm 0.02$	[702] WANG	04A	BELL	Repl. by JEN 06
$-0.01 ^{+0.29}_{-0.31} \pm 0.03$	[703] AUBERT	02E	BABR	Repl. by AUBERT 04H

[701] Corresponds to 90% confidence range $-0.75 < A_{CP} < 0.07$.
[702] Corresponds to 90% CL interval $-0.25 < A_{CP} < 0.41$
[703] Corresponds to 90% confidence range $-0.50 < A_{CP} < 0.46$.

$A_{CP}(B^+ \to \omega\rho^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.04±0.18±0.02	AUBERT,B	06T	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.05±0.26±0.02	AUBERT	05O	BABR	Repl. by AUBERT,B 06T

$A_{CP}(B^+ \to \eta\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.05±0.10 OUR AVERAGE				
−0.13±0.12±0.01	AUBERT,B	05K	BABR	$e^+e^- \to \Upsilon(4S)$
0.07±0.15±0.03	CHANG	05A	BELL	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
−0.44±0.18±0.01	AUBERT	04H	BABR	Repl. by AUBERT,B 05K

$A_{CP}(B^+ \to \eta'\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.15±0.15 OUR AVERAGE				
$0.20^{+0.37}_{-0.36}\pm0.04$	SCHUEMANN	06	BELL	$e^+e^- \to \Upsilon(4S)$
0.14±0.16±0.01	AUBERT,B	05K	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \eta\rho^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.02±0.18±0.02	AUBERT,B	05K	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to \eta'\rho^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.04±0.28±0.02	AUBERT	07E	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to p\overline{p}\pi^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.16±0.22±0.01	WANG	04	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to p\overline{p}K^+)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.05±0.11±0.01	WANG	04	BELL	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to K^+\ell^+\ell^-)$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.07±0.22±0.02	AUBERT,B	06J	BABR	$e^+e^- \to \Upsilon(4S)$

$A_{CP}(B^+ \to K^{*+}\ell^+\ell^-)$

VALUE	DOCUMENT ID		TECN	COMMENT
0.03±0.23±0.03	AUBERT,B	06J	BABR	$e^+e^- \to \Upsilon(4S)$

$\gamma(B^+ \to D^{(*)} K^{(*)+})$

For angle $\gamma(\phi_3)$ of the CKM unitarity triangle, see the review on "*CP* Violation" in the Reviews section.

VALUE (°)	DOCUMENT ID		TECN	COMMENT
57±17 OUR AVERAGE				
$53^{+15}_{-18}\pm10$	[704] POLUEKTOV	06	BELL	$e^+e^- \to \Upsilon(4S)$
$70\pm31^{+18}_{-15}$	[705] AUBERT,B	05Y	BABR	$e^+e^- \to \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$77^{+17}_{-19}\pm17$	[706] POLUEKTOV	04	BELL	Repl. by POLUEKTOV 06

[704] Uses a Dalitz plot analysis of the $\overline{D}^0 \to K^0_S \pi^+\pi^-$ decays; Combines the DK^+, D^*K^+ and DK^{*+} modes. The corresponding two standard deviations interval for gamma is $8^\circ < \gamma < 111^\circ$.

[705] Uses a Dalitz plot analysis of neutral $D \to K^0_S \pi^+\pi^-$ decays coming from $B^\pm \to DK^\pm$ and $B^\pm \to D^{*0}K^\pm$ followed by $D^{*0} \to D\pi^0$, $D\gamma$. The corresponding two standard deviations interval for gamma is $12^\circ < \gamma < 137^\circ$.

[706] Uses a Dalitz plot analysis of the 3-body $D \to K^0_S \pi^+\pi^-$ decays coming from $B^\pm \to DK^\pm$ and $B^\pm \to D^*K^\pm$ followed by $D^* \to D\pi^0$; here we use D to denote that the neutral D meson produced in the decay is an admixture of D^0 and $\overline{D}^0$. The corresponding two standard deviations interval for γ is $26^\circ < \gamma < 126^\circ$. POLUEKTOV 04 also reports the amplitude ratios and the strong phases.

$B^\pm$ REFERENCES

AUBERT	07E	PRL 98 051802	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	07H	PR D75 031101R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	07L	PRL 98 151802	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	07M	PRL 98 171801	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	07N	PR D75 072002	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	07Q	PR D75 051102R	B. Aubert *et al.*	(BABAR Collab.)
HOKUUE	07	PL B648 139	T. Hokuue *et al.*	(BELLE Collab.)
SATOYAMA	07	PL B647 67	N. Satoyama *et al.*	(BELLE Collab.)
URQUIJO	07	PR D75 032001	P. Urquijo *et al.*	(BELLE Collab.)
ABE	06	PR D73 051106R	K. Abe *et al.*	(BELLE Collab.)
ABULENCIA	06J	PRL 96 191801	A. Abulencia *et al.*	(CDF Collab.)
ACOSTA	06	PRL 96 202001	D. Acosta *et al.*	(CDF Collab.)
AUBERT	06	PR D73 011101R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	06F	PR D73 011103R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	06J	PR D73 051105R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	06K	PR D73 057101	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	06N	PR D74 031103R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	06O	PR D74 032003	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	06Z	PR D73 111104R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06A	PR D73 112004	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06C	PR D74 011102R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06E	PR D74 011106R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06G	PRL 97 201801	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06H	PRL 97 201802	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06J	PR D73 092001	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06M	PR D74 031102R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06P	PR D74 031105R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06T	PR D74 051102R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06U	PR D74 051104R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	06Y	PR D74 091105R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,BE	06A	PR D74 099903 (errat.)	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,BE	06C	PRL 97 171805	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,BE	06G	PRL 97 261801	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,BE	06H	PRL 97 261803	B. Aubert *et al.*	(BABAR Collab.)

AUBERT,BE	06J	PR D74 111102R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,BE	06M	PR D74 071101R	B. Aubert *et al.*	(BABAR Collab.)
CHISTOV	06A	PR D74 111105R	R. Chistov *et al.*	(BELLE Collab.)
FANG	06	PR D74 012007	F. Fang *et al.*	(BELLE Collab.)
GABYSHEV	06	PRL 97 202003	N. Gabyshev *et al.*	(BELLE Collab.)
GABYSHEV	06A	PRL 97 242001	N. Gabyshev *et al.*	(BELLE Collab.)
GARMASH	06	PRL 96 251803	A. Garmash *et al.*	(BELLE Collab.)
GOKHROO	06	PRL 97 162002	G. Gokhroo *et al.*	(BELLE Collab.)
IKADO	06	PRL 97 251802	K. Ikado *et al.*	(BELLE Collab.)
JEN	06	PR D74 111101R	C.-M. Jen *et al.*	(BELLE Collab.)
KUMAR	06	PR D74 051103R	R. Kumar *et al.*	(BELLE Collab.)
MOHAPATRA	06	PRL 96 221601	D. Mohapatra *et al.*	(BELLE Collab.)
POLUEKTOV	06	PR D73 112009	A. Poluektov *et al.*	(BELLE Collab.)
SCHUEMANN	06	PRL 97 061802	J. Schuemann *et al.*	(BELLE Collab.)
SONI	06	PL B634 155	N. Soni *et al.*	(BELLE Collab.)
ABE	05A	PRL 94 221805	K. Abe *et al.*	(BELLE Collab.)
ABE	05B	PR D71 072003	K. Abe *et al.*	(BELLE Collab.)
Also		PR D71 079903 (errat.)	K. Abe *et al.*	(BELLE Collab.)
ABE	05G	PRL 95 231802	K. Abe *et al.*	(BELLE Collab.)
ACOSTA	05J	PRL 95 031801	D. Acosta *et al.*	(CDF Collab.)
AUBERT	05	PRL 94 011801	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05B	PR D71 031501R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05G	PR D72 032004	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05H	PRL 94 101801	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05J	PRL 94 141801	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05K	PRL 94 171801	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05L	PRL 94 181802	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05M	PRL 94 191802	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05N	PR D71 031102R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05O	PR D71 031103R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05R	PR D71 071103R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05U	PR D71 091103R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	05X	PR D71 111101	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05B	PRL 95 041804	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05E	PR D72 011102R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05G	PR D72 052002	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05K	PRL 95 131803	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05L	PR D72 051101R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05N	PR D72 072003	B. Aubert *et al.*	(BABAR Collab.)
Also		PR D74 099903 (errat.)	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05O	PR D72 051102R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05T	PR D72 071102R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05U	PR D72 071103R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05V	PR D72 071104R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	05Y	PRL 95 121802	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,BE	05E	PRL 95 221801	B. Aubert *et al.*	(BABAR Collab.)
CHANG	05	PR D71 072007	M.-C. Chang *et al.*	(BELLE Collab.)
CHANG	05A	PR D71 091106R	P. Chang *et al.*	(BELLE Collab.)
CHAO	05A	PR D71 031502R	Y. Chao *et al.*	(BELLE Collab.)
CHEN	05A	PRL 94 221804	K.-F. Chen *et al.*	(BELLE Collab.)
GARMASH	05	PR D71 092003	A. Garmash *et al.*	(BELLE Collab.)
ITOH	05	PRL 95 091601	R. Itoh *et al.*	(BELLE Collab.)
LEE	05	PRL 95 061802	Y.-J. Lee *et al.*	(BELLE Collab.)
LIVENTSEV	05	PR D72 051109R	D. Liventsev *et al.*	(BELLE Collab.)
MAJUMDER	05	PRL 95 041803	G. Majumder *et al.*	(BELLE Collab.)
MOHAPATRA	05	PR D72 011101R	D. Mohapatra *et al.*	(BELLE Collab.)
NISHIDA	05	PL B610 23	S. Nishida *et al.*	(BELLE Collab.)
OKABE	05	PL B614 27	T. Okabe *et al.*	(BELLE Collab.)
SAIGO	05	PRL 94 091601	M. Saigo *et al.*	(BELLE Collab.)
WANG	05A	PL B617 141	M.-Z. Wang *et al.*	(BELLE Collab.)
XIE	05	PR D72 051105R	Q.L. Xie *et al.*	(BELLE Collab.)
YANG	05	PRL 94 111802	H. Yang *et al.*	(BELLE Collab.)
ZHANG	05A	PRL 94 031801	J. Zhang *et al.*	(BELLE Collab.)
ZHANG	05B	PR D71 091107R	L.M. Zhang *et al.*	(BELLE Collab.)
ZHANG	05D	PRL 95 141801	J. Zhang *et al.*	(BELLE Collab.)
ABDALLAH	04E	EPJ C33 307	J. Abdallah *et al.*	(DELPHI Collab.)
ABE	04D	PR D69 112002	K. Abe *et al.*	(BELLE Collab.)
AUBERT	04A	PR D69 011102	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	04C	PRL 92 111801	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	04H	PRL 92 061801	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	04K	PRL 92 141801	B. Aubert *et al.*	(BABAR Collab.)

AUBERT	04M	PRL 92 201802	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	04N	PRL 92 202002	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	04O	PRL 92 221803	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	04P	PRL 92 241802	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	04Q	PR D69 051101R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	04T	PR D69 071103R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT	04Y	PRL 93 041801	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	04Z	PRL 93 051802	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	04B	PR D70 011101R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	04D	PR D70 032006	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	04L	PRL 93 131804	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	04P	PR D70 092001	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	04S	PRL 93 181801	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	04U	PR D70 091105R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,B	04V	PRL 93 181805	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,BE	04	PR D70 111102R	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,BE	04A	PR D70 112006	B. Aubert *et al.*	(BABAR Collab.)
AUBERT,BE	04B	PR D70 091106	B. Aubert *et al.*	(BABAR Collab.)
CHAO	04	PR D69 111102R	Y. Chao *et al.*	(BELLE Collab.)
CHAO	04B	PRL 93 191802	Y. Chao *et al.*	(BELLE Collab.)
CHISTOV	04	PRL 93 051803	R. Chistov *et al.*	(BELLE Collab.)
DRUTSKOY	04	PRL 92 051801	A. Drutskoy *et al.*	(BELLE Collab.)
GARMASH	04	PR D69 012001	A. Garmash *et al.*	(BELLE Collab.)
LEE	04	PRL 93 211801	Y.-J. Lee *et al.*	(BELLE Collab.)
MAJUMDER	04	PR D70 111103R	G. Majumder *et al.*	(BELLE Collab.)
NAKAO	04	PR D69 112001	M. Nakao *et al.*	(BELLE Collab.)
POLUEKTOV	04	PR D70 072003	A. Poluektov *et al.*	(BELLE Collab.)
SCHWANDA	04	PRL 93 131803	C. Schwanda *et al.*	(BELLE Collab.)
WANG	04	PRL 92 131801	M.Z. Wang *et al.*	(BELLE Collab.)
WANG	04A	PR D70 012001	C.H. Wang *et al.*	(BELLE Collab.)
ZANG	04	PR D69 017101	S.L. Zang *et al.*	(BELLE Collab.)
ABE	03B	PR D67 032003	K. Abe *et al.*	(BELLE Collab.)
ABE	03D	PRL 90 131803	K. Abe *et al.*	(BELLE Collab.)
ADAM	03	PR D67 032001	N.E. Adam *et al.*	(CLEO Collab.)
ADAM	03B	PR D68 012004	N.E. Adam *et al.*	(CLEO Collab.)
ATHAR	03	PR D68 072003	S.B. Athar *et al.*	(CLEO Collab.)
AUBERT	03K	PRL 90 231801	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	03L	PRL 91 021801	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	03M	PRL 91 051801	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	03O	PRL 91 071801	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	03U	PRL 91 221802	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	03V	PRL 91 171802	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	03W	PRL 91 161801	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	03X	PR D68 092001	B. Aubert *et al.*	(BaBar Collab.)
BORNHEIM	03	PR D68 052002	A. Bornheim *et al.*	(CLEO Collab.)
CHEN	03B	PRL 91 201801	K.-F. Chen *et al.*	(BELLE Collab.)
CHOI	03	PRL 91 262001	S.-K. Choi *et al.*	(BELLE Collab.)
CSORNA	03	PR D67 112002	S.E. Csorna *et al.*	(CLEO Collab.)
EDWARDS	03	PR D68 011102R	K.W. Edwards *et al.*	(CLEO Collab.)
FANG	03	PRL 90 071801	F. Fang *et al.*	(BELLE Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang *et al.*	(BELLE Collab.)
ISHIKAWA	03	PRL 91 261601	A. Ishikawa *et al.*	(BELLE Collab.)
KROKOVNY	03B	PRL 91 262002	P. Krokovny *et al.*	(BELLE Collab.)
SWAIN	03	PR D68 051101R	S.K. Swain *et al.*	(BELLE Collab.)
UNNO	03	PR D68 011103R	Y. Unno *et al.*	(BELLE Collab.)
ZHANG	03B	PRL 91 221801	J. Zhang *et al.*	(BELLE Collab.)
ABE	02	PRL 88 021801	K. Abe *et al.*	(BELLE Collab.)
ABE	02B	PRL 88 031802	K. Abe *et al.*	(BELLE Collab.)
ABE	02H	PRL 88 171801	K. Abe *et al.*	(BELLE Collab.)
ABE	02K	PRL 88 181803	K. Abe *et al.*	(BELLE Collab.)
ABE	02N	PL B538 11	K. Abe *et al.*	(BELLE Collab.)
ABE	02O	PR D65 091103R	K. Abe *et al.*	(BELLE Collab.)
ABE	02W	PRL 89 151802	K. Abe *et al.*	(BELLE Collab.)
ACOSTA	02C	PR D65 092009	D. Acosta *et al.*	(CDF Collab.)
ACOSTA	02F	PR D66 052005	D. Acosta *et al.*	(CDF Collab.)
AHMED	02B	PR D66 031101R	S. Ahmed *et al.*	(CLEO Collab.)
AUBERT	02	PR D65 032001	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	02C	PRL 88 101805	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	02E	PR D65 051101R	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	02F	PR D65 091101R	B. Aubert *et al.*	(BaBar Collab.)
AUBERT	02L	PRL 88 241801	B. Aubert *et al.*	(BaBar Collab.)

BRIERE	02	PRL 89 081803	R. Briere et al.	(CLEO Collab.)
CASEY	02	PR D66 092002	B.C.K. Casey et al.	(BELLE Collab.)
CHEN	02B	PL B546 196	K.-F. Chen et al.	(BELLE Collab.)
DRUTSKOY	02	PL B542 171	A. Drutskoy et al.	(BELLE Collab.)
DYTMAN	02	PR D66 091101R	S.A. Dytman et al.	(CLEO Collab.)
ECKHART	02	PRL 89 251801	E. Eckhart et al.	(CLEO Collab.)
EDWARDS	02B	PR D65 111102R	K.W. Edwards et al.	(CLEO Collab.)
GABYSHEV	02	PR D66 091102R	N. Gabyshev et al.	(BELLE Collab.)
GARMASH	02	PR D65 092005	A. Garmash et al.	(BELLE Collab.)
GODANG	02	PRL 88 021802	R. Godang et al.	(CLEO Collab.)
GORDON	02	PL B542 183	A. Gordon et al.	(BELLE Collab.)
LU	02	PRL 89 191801	R.-S. Lu et al.	(BELLE Collab.)
MAHAPATRA	02	PRL 88 101803	R. Mahapatra et al.	(CLEO Collab.)
NISHIDA	02	PRL 89 231801	S. Nishida et al.	(BELLE Collab.)
ABE	01H	PRL 87 101801	K. Abe et al.	(BELLE Collab.)
ABE	01I	PRL 87 111801	K. Abe et al.	(BELLE Collab.)
ABE	01K	PR D64 071101	K. Abe et al.	(BELLE Collab.)
ABE	01L	PRL 87 161601	K. Abe et al.	(BELLE Collab.)
ABE	01M	PL B517 309	K. Abe et al.	(BELLE Collab.)
ALEXANDER	01B	PR D64 092001	J.P. Alexander et al.	(CLEO Collab.)
AMMAR	01B	PRL 87 271801	R. Ammar et al.	(CLEO Collab.)
ANDERSON	01B	PRL 87 181803	S. Anderson et al.	(CLEO Collab.)
AUBERT	01D	PRL 87 151801	B. Aubert et al.	(BaBar Collab.)
AUBERT	01E	PRL 87 151802	B. Aubert et al.	(BaBar Collab.)
AUBERT	01F	PRL 87 201803	B. Aubert et al.	(BaBar Collab.)
AUBERT	01G	PRL 87 221802	B. Aubert et al.	(BaBar Collab.)
BARATE	01E	EPJ C19 213	R. Barate et al.	(ALEPH Collab.)
BRIERE	01	PRL 86 3718	R.A. Biere et al.	(CLEO Collab.)
BROWDER	01	PRL 86 2950	T.E. Browder et al.	(CLEO Collab.)
EDWARDS	01	PRL 86 30	K.W. Edwards et al.	(CLEO Collab.)
GRITSAN	01	PR D64 077501	A. Gritsan et al.	(CLEO Collab.)
RICHICHI	01	PR D63 031103R	S.J. Richichi et al.	(CLEO Collab.)
ABBIENDI	00B	PL B476 233	G. Abbiendi et al.	(OPAL Collab.)
ABE	00C	PR D62 071101R	K. Abe et al.	(SLD Collab.)
AHMED	00B	PR D62 112003	S. Ahmed et al.	(CLEO Collab.)
ANASTASSOV	00	PRL 84 1393	A. Anastassov et al.	(CLEO Collab.)
BARATE	00R	PL B492 275	R. Barate et al.	(ALEPH Collab.)
BEHRENS	00	PR D61 052001	B.H. Behrens et al.	(CLEO Collab.)
BONVICINI	00	PRL 84 5940	G. Bonvicini et al.	(CLEO Collab.)
CHEN	00	PRL 85 525	S. Chen et al.	(CLEO Collab.)
COAN	00	PRL 84 5283	T.E. Coan et al.	(CLEO Collab.)
CRONIN-HEN...	00	PRL 85 515	D. Cronin-Hennessy et al.	(CLEO Collab.)
CSORNA	00	PR D61 111101	S.E. Csorna et al.	(CLEO Collab.)
JESSOP	00	PRL 85 2881	C.P. Jessop et al.	(CLEO Collab.)
RICHICHI	00	PRL 85 520	S.J. Richichi et al.	(CLEO Collab.)
ABBIENDI	99J	EPJ C12 609	G. Abbiendi et al.	(OPAL Collab.)
AFFOLDER	99B	PRL 83 3378	T. Affolder et al.	(CDF Collab.)
BARTELT	99	PRL 82 3746	J. Bartelt et al.	(CLEO Collab.)
COAN	99	PR D59 111101	T.E. Coan et al.	(CLEO Collab.)
ABE	98B	PR D57 5382	F. Abe et al.	(CDF Collab.)
ABE	98O	PR D58 072001	F. Abe et al.	(CDF Collab.)
ABE	98Q	PR D58 092002	F. Abe et al.	(CDF Collab.)
ACCIARRI	98S	PL B438 417	M. Acciarri et al.	(L3 Collab.)
ANASTASSOV	98	PRL 80 4127	A. Anastassov et al.	(CLEO Collab.)
ATHANAS	98	PRL 80 5493	M. Athanas et al.	(CLEO Collab.)
BARATE	98Q	EPJ C4 387	R. Barate et al.	(ALEPH Collab.)
BEHRENS	98	PRL 80 3710	B.H. Behrens et al.	(CLEO Collab.)
BERGFELD	98	PRL 81 272	T. Bergfeld et al.	(CLEO Collab.)
BRANDENB...	98	PRL 80 2762	G. Brandenbrug et al.	(CLEO Collab.)
GODANG	98	PRL 80 3456	R. Godang et al.	(CLEO Collab.)
ABE	97J	PRL 79 590	K. Abe et al.	(SLD Collab.)
ACCIARRI	97F	PL B396 327	M. Acciarri et al.	(L3 Collab.)
ARTUSO	97	PL B399 321	M. Artuso et al.	(CLEO Collab.)
ATHANAS	97	PRL 79 2208	M. Athanas et al.	(CLEO Collab.)
BROWDER	97	PR D56 11	T. Browder et al.	(CLEO Collab.)
FU	97	PRL 79 3125	X. Fu et al.	(CLEO Collab.)
JESSOP	97	PRL 79 4533	C.P. Jessop et al.	(CLEO Collab.)
ABE	96B	PR D53 3496	F. Abe et al.	(CDF Collab.)
ABE	96C	PRL 76 4462	F. Abe et al.	(CDF Collab.)
ABE	96H	PRL 76 2015	F. Abe et al.	(CDF Collab.)
ABE	96L	PRL 76 4675	F. Abe et al.	(CDF Collab.)

ABE	96Q	PR D54 6596	F. Abe *et al.*	(CDF Collab.)
ABE	96R	PRL 77 5176	F. Abe *et al.*	(CDF Collab.)
ADAM	96D	ZPHY C72 207	W. Adam *et al.*	(DELPHI Collab.)
ALEXANDER	96T	PRL 77 5000	J.P. Alexander *et al.*	(CLEO Collab.)
ASNER	96	PR D53 1039	D.M. Asner *et al.*	(CLEO Collab.)
BARISH	96B	PRL 76 1570	B.C. Barish *et al.*	(CLEO Collab.)
BERGFELD	96B	PRL 77 4503	T. Bergfeld *et al.*	(CLEO Collab.)
BISHAI	96	PL B369 186	M. Bishai *et al.*	(CLEO Collab.)
BUSKULIC	96J	ZPHY C71 31	D. Buskulic *et al.*	(ALEPH Collab.)
GIBAUT	96	PR D53 4734	D. Gibaut *et al.*	(CLEO Collab.)
PDG	96	PR D54 1	R. M. Barnett *et al.*	
ABREU	95N	PL B357 255	P. Abreu *et al.*	(DELPHI Collab.)
ABREU	95Q	ZPHY C68 13	P. Abreu *et al.*	(DELPHI Collab.)
ADAM	95	ZPHY C68 363	W. Adam *et al.*	(DELPHI Collab.)
AKERS	95T	ZPHY C67 379	R. Akers *et al.*	(OPAL Collab.)
ALBRECHT	95D	PL B353 554	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXANDER	95	PL B341 435	J. Alexander *et al.*	(CLEO Collab.)
Also		PL B347 469 (erratum)	J. Alexander *et al.*	(CLEO Collab.)
ARTUSO	95	PRL 75 785	M. Artuso *et al.*	(CLEO Collab.)
BARISH	95	PR D51 1014	B.C. Barish *et al.*	(CLEO Collab.)
BUSKULIC	95	PL B343 444	D. Buskulic *et al.*	(ALEPH Collab.)
ABE	94D	PRL 72 3456	F. Abe *et al.*	(CDF Collab.)
ALAM	94	PR D50 43	M.S. Alam *et al.*	(CLEO Collab.)
ALBRECHT	94D	PL B335 526	H. Albrecht *et al.*	(ARGUS Collab.)
ATHANAS	94	PRL 73 3503	M. Athanas *et al.*	(CLEO Collab.)
Also		PRL 74 3090 (erratum)	M. Athanas *et al.*	(CLEO Collab.)
PDG	94	PR D50 1173	L. Montanet *et al.*	(CERN, LBL, BOST+)
STONE	94	HEPSY 93-11	S. Stone	
Published in B Decays, 2nd Edition, World Scientific, Singapore				
ABREU	93D	ZPHY C57 181	P. Abreu *et al.*	(DELPHI Collab.)
ABREU	93G	PL B312 253	P. Abreu *et al.*	(DELPHI Collab.)
ACTON	93C	PL B307 247	P.D. Acton *et al.*	(OPAL Collab.)
ALBRECHT	93E	ZPHY C60 11	H. Albrecht *et al.*	(ARGUS Collab.)
ALEXANDER	93B	PL B319 365	J. Alexander *et al.*	(CLEO Collab.)
AMMAR	93	PRL 71 674	R. Ammar *et al.*	(CLEO Collab.)
BEAN	93B	PRL 70 2681	A. Bean *et al.*	(CLEO Collab.)
BUSKULIC	93D	PL B307 194	D. Buskulic *et al.*	(ALEPH Collab.)
Also		PL B325 537 (erratum)	D. Buskulic *et al.*	(ALEPH Collab.)
SANGHERA	93	PR D47 791	S. Sanghera *et al.*	(CLEO Collab.)
ALBRECHT	92C	PL B275 195	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	92E	PL B277 209	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	92G	ZPHY C54 1	H. Albrecht *et al.*	(ARGUS Collab.)
BORTOLETTO	92	PR D45 21	D. Bortoletto *et al.*	(CLEO Collab.)
BUSKULIC	92G	PL B295 396	D. Buskulic *et al.*	(ALEPH Collab.)
ALBRECHT	91B	PL B254 288	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	91C	PL B255 297	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	91E	PL B262 148	H. Albrecht *et al.*	(ARGUS Collab.)
BERKELMAN	91	ARNPS 41 1	K. Berkelman, S. Stone	(CORN, SYRA)
"Decays of *B* Mesons"				
FULTON	91	PR D43 651	R. Fulton *et al.*	(CLEO Collab.)
ALBRECHT	90B	PL B241 278	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	90J	ZPHY C48 543	H. Albrecht *et al.*	(ARGUS Collab.)
ANTREASYAN	90B	ZPHY C48 553	D. Antreasyan *et al.*	(Crystal Ball Collab.)
BORTOLETTO	90	PRL 64 2117	D. Bortoletto *et al.*	(CLEO Collab.)
Also		PR D45 21	D. Bortoletto *et al.*	(CLEO Collab.)
WEIR	90B	PR D41 1384	A.J. Weir *et al.*	(Mark II Collab.)
ALBRECHT	89G	PL B229 304	H. Albrecht *et al.*	(ARGUS Collab.)
AVERY	89B	PL B223 470	P. Avery *et al.*	(CLEO Collab.)
BEBEK	89	PRL 62 8	C. Bebek *et al.*	(CLEO Collab.)
BORTOLETTO	89	PRL 62 2436	D. Bortoletto *et al.*	(CLEO Collab.)
ALBRECHT	88F	PL B209 119	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	88K	PL B215 424	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	87C	PL B185 218	H. Albrecht *et al.*	(ARGUS Collab.)
ALBRECHT	87D	PL B199 451	H. Albrecht *et al.*	(ARGUS Collab.)
AVERY	87	PL B183 429	P. Avery *et al.*	(CLEO Collab.)
BEBEK	87	PR D36 1289	C. Bebek *et al.*	(CLEO Collab.)
ALAM	86	PR D34 3279	M.S. Alam *et al.*	(CLEO Collab.)
PDG	86	PL 170B 1	M. Aguilar-Benitez *et al.*	(CERN, CIT+)
GILES	84	PR D30 2279	R. Giles *et al.*	(CLEO Collab.)