

$N(2200) D_{15}$

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^-) \text{ Status: } **$$

OMITTED FROM SUMMARY TABLE

The mass is not well determined. A few early results have been omitted.

The latest GWU analysis (ARNDT 06) finds no evidence for this resonance.

 $N(2200)$ BREIT-WIGNER MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
≈ 2200 OUR ESTIMATE			
1900	BELL	83	DPWA $\pi^- p \rightarrow \Lambda K^0$
2180 ± 80	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1920	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
2228 ± 30	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2240 ± 65	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$

 $N(2200)$ BREIT-WIGNER WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
130	BELL	83	DPWA $\pi^- p \rightarrow \Lambda K^0$
400 ± 100	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
220	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
310 ± 50	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
761 ± 139	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$

 $N(2200)$ POLE POSITION**REAL PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2100 ± 60	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

-2×IMAGINARY PART

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
360 ± 80	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

 $N(2200)$ ELASTIC POLE RESIDUE**MODULUS $|r|$**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
20 ± 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

PHASE θ

<u>VALUE ($^\circ$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-90 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $N(2200)$ DECAY MODES

Mode
Γ_1 $N\pi$
Γ_2 $N\eta$
Γ_3 ΛK

 $N(2200)$ BRANCHING RATIOS

<u>$\Gamma(N\pi)/\Gamma_{\text{total}}$</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_1/Γ
0.10 ± 0.03	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.07 ± 0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.08 ± 0.04	BATINIC 95	DPWA	$\pi N \rightarrow N\pi, N\eta$	

<u>$\Gamma(N\eta)/\Gamma_{\text{total}}$</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.001 ± 0.01	BATINIC 95	DPWA	$\pi N \rightarrow N\pi, N\eta$	

<u>$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2200) \rightarrow N\eta$</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$(\Gamma_1 \Gamma_2)^{1/2}/\Gamma$
0.066	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$	

<u>$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2200) \rightarrow \Lambda K$</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$(\Gamma_1 \Gamma_3)^{1/2}/\Gamma$
-0.03	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
-0.05	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$	

 $N(2200)$ REFERENCES

ARNDT 06	PR C74 045205	R.A. Arndt <i>et al.</i>	(GWU)
BATINIC 95	PR C51 2310	M. Batinic <i>et al.</i>	(BOSK, UCLA)
Also	PR C57 1004 (erratum)	M. Batinic <i>et al.</i>	
BELL 83	NP B222 389	K.W. Bell <i>et al.</i>	(RL) IJP
CUTKOSKY 80	Toronto Conf. 19	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
Also	PR D20 2839	R.E. Cutkosky <i>et al.</i>	(CMU, LBL)
SAXON 80	NP B162 522	D.H. Saxon <i>et al.</i>	(RHEL, BRIS) IJP
BAKER 79	NP B156 93	R.D. Baker <i>et al.</i>	(RHEL) IJP
HOEHLER 79	PDAT 12-1	G. Hohler <i>et al.</i>	(KARLT) IJP
Also	Toronto Conf. 3	R. Koch	(KARLT) IJP