

B^0

$$I(J^P) = \frac{1}{2}(0^-)$$

Quantum numbers not measured. Values shown are quark-model predictions.

See also the B^\pm/B^0 ADMIXTURE and $B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE sections.

See the Note “Production and Decay of b -flavored Hadrons” at the beginning of the B^\pm Particle Listings and the Note on “ B^0 - \bar{B}^0 Mixing” near the end of the B^0 Particle Listings.

B^0 MASS

The fit uses m_{B^+} , ($m_{B^0} - m_{B^+}$), and m_{B^0} to determine m_{B^+} , m_{B^0} , and the mass difference.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------------|------|---------------------------------|
| 5279.53±0.33 OUR FIT | | | | |
| 5279.5 ±0.5 OUR AVERAGE | | | | |
| 5279.63±0.53±0.33 | | ¹ ACOSTA 06 | CDF | $p\bar{p}$ at 1.96 TeV |
| 5279.1 ±0.7 ±0.3 | 135 | ² CSORNA 00 | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| 5281.3 ±2.2 ±1.4 | 51 | ABE 96B | CDF | $p\bar{p}$ at 1.8 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 5279.2 ±0.54±2.0 | 340 | ALAM 94 | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| 5278.0 ±0.4 ±2.0 | | BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |
| 5279.6 ±0.7 ±2.0 | 40 | ³ ALBRECHT 90J | ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| 5278.2 ±1.0 ±3.0 | 40 | ALBRECHT 87C | ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| 5279.5 ±1.6 ±3.0 | 7 | ⁴ ALBRECHT 87D | ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| 5280.6 ±0.8 ±2.0 | | BEBEK 87 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Uses exclusively reconstructed final states containing a $J/\psi \rightarrow \mu^+\mu^-$ decays.

² CSORNA 00 uses fully reconstructed $135 B^0 \rightarrow J/\psi(\prime) K_S^0$ events and invariant masses without beam constraint.

³ ALBRECHT 90J assumes 10580 for $\gamma(4S)$ mass. Supersedes ALBRECHT 87C and ALBRECHT 87D.

⁴ Found using fully reconstructed decays with J/ψ . ALBRECHT 87D assume $m\gamma(4S) = 10577$ MeV.

$m_{B^0} - m_{B^+}$

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------------------------|------|---------------------------------|
| 0.37±0.24 OUR FIT | | | |
| 0.37±0.26 OUR AVERAGE | | | |
| 0.53±0.67±0.14 | ¹ ACOSTA 06 | CDF | $p\bar{p}$ at 1.96 TeV |
| 0.41±0.25±0.19 | ALAM 94 | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| -0.4 ±0.6 ±0.5 | BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |
| -0.9 ±1.2 ±0.5 | ALBRECHT 90J | ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| 2.0 ±1.1 ±0.3 | ² BEBEK 87 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Uses exclusively reconstructed final states containing a $J/\psi \rightarrow \mu^+ \mu^-$ decays.

² BEBEK 87 actually measure the difference between half of E_{cm} and the B^\pm or B^0 mass, so the $m_{B^0} - m_{B^\pm}$ is more accurate. Assume $m_{\Upsilon(4S)} = 10580$ MeV.

$$m_{B_H^0} - m_{B_L^0}$$

See the B^0 - \bar{B}^0 MIXING PARAMETERS section near the end of these B^0 Listings.

B^0 MEAN LIFE

See $B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE section for data on B -hadron mean life averaged over species of bottom particles.

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements and asymmetric lifetime errors.

| VALUE (10^{-12} s) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------|----------|------------------------------------|
| 1.530±0.009 OUR EVALUATION | | | | |
| 1.501 ^{+0.078} _{-0.074} ± 0.050 | | ¹ ABAZOV | 07S D0 | $p\bar{p}$ at 1.96 TeV |
| 1.524 ± 0.030 ± 0.016 | | ¹ ABULENCIA | 07A CDF | $p\bar{p}$ at 1.96 TeV |
| 1.504 ± 0.013 ^{+0.018} _{-0.013} | | ² AUBERT | 06G BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.530 ± 0.043 ± 0.023 | | ³ ABAZOV | 05W D0 | $p\bar{p}$ at 1.96 TeV |
| 1.534 ± 0.008 ± 0.010 | | ⁴ ABE | 05B BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.54 ± 0.05 ± 0.02 | | ⁵ ACOSTA | 05 CDF | $p\bar{p}$ at 1.96 TeV |
| 1.531 ± 0.021 ± 0.031 | | ⁶ ABDALLAH | 04E DLPH | $e^+ e^- \rightarrow Z$ |
| 1.533 ± 0.034 ± 0.038 | | ⁷ AUBERT | 03H BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.497 ± 0.073 ± 0.032 | | ⁸ ACOSTA | 02C CDF | $p\bar{p}$ at 1.8 TeV |
| 1.529 ± 0.012 ± 0.029 | | ⁹ AUBERT | 02H BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.546 ± 0.032 ± 0.022 | | ¹⁰ AUBERT | 01F BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.541 ± 0.028 ± 0.023 | | ⁹ ABBIENDI,G | 00B OPAL | $e^+ e^- \rightarrow Z$ |
| 1.518 ± 0.053 ± 0.034 | | ¹¹ BARATE | 00R ALEP | $e^+ e^- \rightarrow Z$ |
| 1.523 ± 0.057 ± 0.053 | | ¹² ABBIENDI | 99J OPAL | $e^+ e^- \rightarrow Z$ |
| 1.474 ± 0.039 ^{+0.052} _{-0.051} | | ¹¹ ABE | 98Q CDF | $p\bar{p}$ at 1.8 TeV |
| 1.52 ± 0.06 ± 0.04 | | ¹² ACCIARRI | 98S L3 | $e^+ e^- \rightarrow Z$ |
| 1.64 ± 0.08 ± 0.08 | | ¹² ABE | 97J SLD | $e^+ e^- \rightarrow Z$ |
| 1.532 ± 0.041 ± 0.040 | | ¹³ ABREU | 97F DLPH | $e^+ e^- \rightarrow Z$ |
| 1.25 ^{+0.15} _{-0.13} ± 0.05 | 121 | ⁸ BUSKULIC | 96J ALEP | $e^+ e^- \rightarrow Z$ |
| 1.49 ^{+0.17} _{-0.15} ± 0.08 | | ¹⁴ BUSKULIC | 96J ALEP | $e^+ e^- \rightarrow Z$ |
| 1.61 ^{+0.14} _{-0.13} ± 0.08 | | ^{11,15} ABREU | 95Q DLPH | $e^+ e^- \rightarrow Z$ |
| 1.63 ± 0.14 ± 0.13 | | ¹⁶ ADAM | 95 DLPH | $e^+ e^- \rightarrow Z$ |
| 1.53 ± 0.12 ± 0.08 | | ^{11,17} AKERS | 95T OPAL | $e^+ e^- \rightarrow Z$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--|------------------------|------------------------|--------------------------------|
| $1.473^{+0.052}_{-0.050} \pm 0.023$ | ³ ABAZOV | 05B D0 | Repl. by ABAZOV 05W |
| $1.40^{+0.11}_{-0.10} \pm 0.03$ | ¹ ABAZOV | 05C D0 | Repl. by ABAZOV 07S |
| $1.523^{+0.024}_{-0.023} \pm 0.022$ | ¹⁸ AUBERT | 03C BABR | Repl. by AUBERT 06G |
| $1.554 \pm 0.030 \pm 0.019$ | ¹⁰ ABE | 02H BELL | Repl. by ABE 05B |
| $1.58 \pm 0.09 \pm 0.02$ | ⁸ ABE | 98B CDF | Repl. by ACOSTA 02C |
| $1.54 \pm 0.08 \pm 0.06$ | ¹¹ ABE | 96C CDF | Repl. by ABE 98Q |
| $1.55 \pm 0.06 \pm 0.03$ | ¹⁹ BUSKULIC | 96J ALEP | $e^+ e^- \rightarrow Z$ |
| $1.61 \pm 0.07 \pm 0.04$ | ¹¹ BUSKULIC | 96J ALEP | Repl. by BARATE 00R |
| 1.62 ± 0.12 | ²⁰ ADAM | 95 DLPH | $e^+ e^- \rightarrow Z$ |
| $1.57 \pm 0.18 \pm 0.08$ | ¹²¹ ABE | 94D CDF | Repl. by ABE 98B |
| $1.17^{+0.29}_{-0.23} \pm 0.16$ | 96 | ¹¹ ABREU | 93D DLPH |
| $1.55 \pm 0.25 \pm 0.18$ | 76 | ¹⁶ ABREU | 93G DLPH |
| $1.51^{+0.24}_{-0.23}^{+0.12}_{-0.14}$ | 78 | ¹¹ ACTON | 93C OPAL |
| $1.52^{+0.20}_{-0.18}^{+0.07}_{-0.13}$ | 77 | ¹¹ BUSKULIC | 93D ALEP |
| $1.20^{+0.52}_{-0.36}^{+0.16}_{-0.14}$ | 15 | ²¹ WAGNER | 90 MRK2 $E_{cm}^{ee} = 29$ GeV |
| $0.82^{+0.57}_{-0.37} \pm 0.27$ | | ²² AVERILL | 89 HRS $E_{cm}^{ee} = 29$ GeV |

¹ Measured mean life using $B^0 \rightarrow J/\psi K_S$ decays.

² Measured using a simultaneous fit of the B^0 lifetime and $\bar{B}^0 B^0$ oscillation frequency Δm_d in the partially reconstructed $B^0 \rightarrow D^{*-} \ell \nu$ decays.

³ Measured mean life using $B^0 \rightarrow J/\psi K^{*0}$ decays.

⁴ Measurement performed using a combined fit of CP -violation, mixing and lifetimes.

⁵ Measured using the time-dependent angular analysis of $B_d^0 \rightarrow J/\psi K^{*0}$ decays.

⁶ Measurement performed using an inclusive reconstruction and B flavor identification technique.

⁷ Measurement performed with decays $B^0 \rightarrow D^{*-} \pi^+$ and $B^0 \rightarrow D^{*-} \rho^+$ using a partial reconstruction technique.

⁸ Measured mean life using fully reconstructed decays.

⁹ Data analyzed using partially reconstructed $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}$ decays.

¹⁰ Events are selected in which one B meson is fully reconstructed while the second B meson is reconstructed inclusively.

¹¹ Data analyzed using $D/D^* \ell X$ event vertices.

¹² Data analyzed using charge of secondary vertex.

¹³ Data analyzed using inclusive $D/D^* \ell X$.

¹⁴ Measured mean life using partially reconstructed $D^{*-} \pi^+ X$ vertices.

¹⁵ ABREU 95Q assumes $B(B^0 \rightarrow D^{**-} \ell^+ \nu_\ell) = 3.2 \pm 1.7\%$.

¹⁶ Data analyzed using vertex-charge technique to tag B charge.

¹⁷ AKERS 95T assumes $B(B_s^0 \rightarrow D_s^0 D^0) = 5.0 \pm 0.9\%$ to find B^+/B^0 yield.

¹⁸ AUBERT 03C uses a sample of approximately 14,000 exclusively reconstructed $B^0 \rightarrow D^*(2010)^- \ell \nu$ and simultaneously measures the lifetime and oscillation frequency.

¹⁹ Combined result of $D/D^* \ell X$ analysis, fully reconstructed B analysis, and partially reconstructed $D^{*-} \pi^+ X$ analysis.

²⁰ Combined ABREU 95Q and ADAM 95 result.

²¹ WAGNER 90 tagged B^0 mesons by their decays into $D^{*-} e^+ \nu$ and $D^{*-} \mu^+ \nu$ where the D^{*-} is tagged by its decay into $\pi^- \bar{D}^0$.

²² AVERILL 89 is an estimate of the B^0 mean lifetime assuming that $B^0 \rightarrow D^{*+} + X$ always.

MEAN LIFE RATIO τ_{B^+}/τ_{B^0}

τ_{B^+}/τ_{B^0} (direct measurements)

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements and asymmetric lifetime errors.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|-----------------------|----------------------------------|----------------------|
| 1.071 ± 0.009 OUR EVALUATION | | | | |
| 1.080 $\pm 0.016 \pm 0.014$ | ¹ ABAZOV | 05D D0 | $p\bar{p}$ at 1.96 TeV | |
| 1.066 $\pm 0.008 \pm 0.008$ | ² ABE | 05B BELL | $e^+ e^- \rightarrow \gamma(4S)$ | |
| 1.060 $\pm 0.021 \pm 0.024$ | ³ ABDALLAH | 04E DLPH | $e^+ e^- \rightarrow Z$ | |
| 1.093 $\pm 0.066 \pm 0.028$ | ⁴ ACOSTA | 02C CDF | $p\bar{p}$ at 1.8 TeV | |
| 1.082 $\pm 0.026 \pm 0.012$ | ⁵ AUBERT | 01F BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| 1.085 $\pm 0.059 \pm 0.018$ | ¹ BARATE | 00R ALEP | $e^+ e^- \rightarrow Z$ | |
| 1.079 $\pm 0.064 \pm 0.041$ | ⁶ ABBIENDI | 99J OPAL | $e^+ e^- \rightarrow Z$ | |
| 1.110 $\pm 0.056^{+0.033}_{-0.030}$ | ¹ ABE | 98Q CDF | $p\bar{p}$ at 1.8 TeV | |
| 1.09 $\pm 0.07 \pm 0.03$ | ⁶ ACCIARRI | 98S L3 | $e^+ e^- \rightarrow Z$ | |
| 1.01 $\pm 0.07 \pm 0.06$ | ⁶ ABE | 97J SLD | $e^+ e^- \rightarrow Z$ | |
| 1.27 $^{+0.23}_{-0.19} \pm 0.03$ | ⁴ BUSKULIC | 96J ALEP | $e^+ e^- \rightarrow Z$ | |
| 1.00 $^{+0.17}_{-0.15} \pm 0.10$ | ^{1,7} ABREU | 95Q DLPH | $e^+ e^- \rightarrow Z$ | |
| 1.06 $^{+0.13}_{-0.11} \pm 0.10$ | ⁸ ADAM | 95 DLPH | $e^+ e^- \rightarrow Z$ | |
| 0.99 $\pm 0.14 \pm 0.05$ | ^{1,9} AKERS | 95T OPAL | $e^+ e^- \rightarrow Z$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.091 $\pm 0.023 \pm 0.014$ | ⁵ ABE | 02H BELL | Repl. by ABE 05B | |
| 1.06 $\pm 0.07 \pm 0.02$ | ⁴ ABE | 98B CDF | Repl. by ACOSTA 02C | |
| 1.01 $\pm 0.11 \pm 0.02$ | ¹ ABE | 96C CDF | Repl. by ABE 98Q | |
| 1.03 $\pm 0.08 \pm 0.02$ | ¹⁰ BUSKULIC | 96J ALEP | $e^+ e^- \rightarrow Z$ | |
| 0.98 $\pm 0.08 \pm 0.03$ | ¹ BUSKULIC | 96J ALEP | Repl. by BARATE 00R | |
| 1.02 $\pm 0.16 \pm 0.05$ | 269 | ⁴ ABE | 94D CDF | Repl. by ABE 98B |
| 1.11 $^{+0.51}_{-0.39} \pm 0.11$ | 188 | ¹ ABREU | 93D DLPH | Sup. by ABREU 95Q |
| 1.01 $^{+0.29}_{-0.22} \pm 0.12$ | 253 | ⁸ ABREU | 93G DLPH | Sup. by ADAM 95 |
| 1.0 $^{+0.33}_{-0.25} \pm 0.08$ | 130 | ACTON | 93C OPAL | Sup. by AKERS 95T |
| 0.96 $^{+0.19}_{-0.15} \pm 0.18$ | 154 | ¹ BUSKULIC | 93D ALEP | Sup. by BUSKULIC 96J |

- ¹ Data analyzed using $D/D^*\mu X$ vertices.
- ² Measurement performed using a combined fit of CP -violation, mixing and lifetimes.
- ³ Measurement performed using an inclusive reconstruction and B flavor identification technique.
- ⁴ Measured using fully reconstructed decays.
- ⁵ Events are selected in which one B meson is fully reconstructed while the second B meson is reconstructed inclusively.
- ⁶ Data analyzed using charge of secondary vertex.
- ⁷ ABREU 95Q assumes $B(B^0 \rightarrow D^{**-} \ell^+ \nu_\ell) = 3.2 \pm 1.7\%$.
- ⁸ Data analyzed using vertex-charge technique to tag B charge.
- ⁹ AKERS 95T assumes $B(B^0 \rightarrow D_s^0 (*) D^0 (*)) = 5.0 \pm 0.9\%$ to find B^+/B^0 yield.
- ¹⁰ Combined result of $D/D^*\ell X$ analysis and fully reconstructed B analysis.

τ_{B^+}/τ_{B^0} (inferred from branching fractions)

These measurements are inferred from the branching fractions for semileptonic decay or other spectator-dominated decays by assuming that the rates for such decays are equal for B^0 and B^+ . We do not use measurements which assume equal production of B^0 and B^+ because of the large uncertainty in the production ratio.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------------------|------|---------------------------------------|
| 1.076±0.034 OUR EVALUATION | | | | | |
| 1.07 ±0.04 OUR AVERAGE | | | | | |
| 1.07 ± 0.04 ± 0.03 | | | URQUIJO | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| 1.067 $\pm 0.041 \pm 0.033$ | | | AUBERT,B | 06Y | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.95 $^{+0.117}_{-0.080} \pm 0.091$ | | | ¹ ARTUSO | 97 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| 1.15 ± 0.17 ± 0.06 | | | ² JESSOP | 97 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.93 ± 0.18 ± 0.12 | | | ³ ATHANAS | 94 | CLE2 Sup. by ARTUSO 97 |
| 0.91 ± 0.27 ± 0.21 | | | ⁴ ALBRECHT | 92C | ARG $e^+ e^- \rightarrow \gamma(4S)$ |
| 1.0 ± 0.4 | 29 | | ^{4,5} ALBRECHT | 92G | ARG $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.89 ± 0.19 ± 0.13 | | | ⁴ FULTON | 91 | CLEO $e^+ e^- \rightarrow \gamma(4S)$ |
| 1.00 ± 0.23 ± 0.14 | | | ⁴ ALBRECHT | 89L | ARG $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.49 to 2.3 | 90 | | ⁶ BEAN | 87B | CLEO $e^+ e^- \rightarrow \gamma(4S)$ |

¹ ARTUSO 97 uses partial reconstruction of $B \rightarrow D^* \ell \nu_\ell$ and independent of B^0 and B^+ production fraction.

² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

³ ATHANAS 94 uses events tagged by fully reconstructed B^- decays and partially or fully reconstructed B^0 decays.

⁴ Assumes equal production of B^0 and B^+ .

⁵ ALBRECHT 92G data analyzed using $B \rightarrow D_s \overline{D}$, $D_s \overline{D}^*$, $D_s^* \overline{D}$, $D_s^* \overline{D}^*$ events.

⁶ BEAN 87B assume the fraction of $B^0 \overline{B}^0$ events at the $\gamma(4S)$ is 0.41.

$$\text{sgn}(\text{Re}(\lambda_{CP})) \Delta\Gamma_{B_d^0} / \Gamma_{B_d^0}$$

$\Gamma_{B_d^0}$ and $\Delta\Gamma_{B_d^0}$ are the decay rate average and difference between two B_d^0 CP eigenstates (light – heavy). The λ_{CP} characterizes B^0 and \bar{B}^0 decays to states of charmonium plus K_L^0 , see the review on “CP Violation” in the reviews section.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|-----------------------|----------|----------------------------------|
| 0.009±0.037 OUR EVALUATION | | | |
| 0.008±0.037±0.018 | ¹ AUBERT,B | 04C BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Corresponds to 90% confidence range [−0.084, 0.068].

$$|\Delta\Gamma_{B_d^0}|/\Gamma_{B_d^0}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|---------|
| The data in this block is included in the average printed for a previous datablock. | | | | |

<0.18 95 ¹ABDALLAH 03B DLPH $e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.80 95 2,3 BEHRENS 00B CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ Using the measured $\tau_{B^0}=1.55 \pm 0.03$ ps.

² BEHRENS 00B uses high-momentum lepton tags and partially reconstructed $\bar{B}^0 \rightarrow D^{*+}\pi^-, \rho^-$ decays to determine the flavor of the B meson.

³ Assumes $\Delta_{md}=0.478 \pm 0.018 \text{ ps}^{-1}$ and $\tau_{B^0}=1.548 \pm 0.032$ ps.

B^0 DECAY MODES

\bar{B}^0 modes are charge conjugates of the modes below. Reactions indicate the weak decay vertex and do not include mixing. Modes which do not identify the charge state of the B are listed in the B^\pm/B^0 ADMIXTURE section.

The branching fractions listed below assume 50% $B^0\bar{B}^0$ and 50% B^+B^- production at the $\gamma(4S)$. We have attempted to bring older measurements up to date by rescaling their assumed $\gamma(4S)$ production ratio to 50:50 and their assumed D , D_s , D^* , and ψ branching ratios to current values whenever this would affect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

For inclusive branching fractions, e.g., $B \rightarrow D^\pm$ anything, the values usually are multiplicities, not branching fractions. They can be greater than one.

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|---|--|-----------------------------------|
| $\Gamma_1 \ell^+ \nu_\ell$ anything | [a] (10.33 ± 0.28) % | |
| $\Gamma_2 e^+ \nu_e X_c$ | (10.1 ± 0.4) % | |
| $\Gamma_3 D \ell^+ \nu_\ell$ anything | (9.6 ± 0.9) % | |
| $\Gamma_4 D^- \ell^+ \nu_\ell$ | [a] (2.17 ± 0.12) % | |
| $\Gamma_5 D^- \tau^+ \nu_\tau$ | (1.0 ± 0.4) % | |
| $\Gamma_6 D^*(2010)^- \ell^+ \nu_\ell$ | [a] (5.16 ± 0.11) % | |
| $\Gamma_7 D^*(2010)^- \tau^+ \nu_\tau$ | (1.6 ± 0.5) % | S=1.2 |
| $\Gamma_8 \bar{D}^0 \pi^- \ell^+ \nu_\ell$ | (4.3 ± 0.6) $\times 10^{-3}$ | |
| $\Gamma_9 D_0^*(2400)^- \ell^+ \nu_\ell \times B(D_0^{*-} \rightarrow \bar{D}^0 \pi^-)$ | (2.0 ± 0.9) $\times 10^{-3}$ | |
| $\Gamma_{10} D_2^*(2460)^- \ell^+ \nu_\ell \times B(D_2^{*-} \rightarrow \bar{D}^0 \pi^-)$ | (2.2 ± 0.6) $\times 10^{-3}$ | |
| $\Gamma_{11} \bar{D}^{(*)} n \pi \ell^+ \nu_\ell (n \geq 1)$ | (2.4 ± 0.5) % | |
| $\Gamma_{12} \bar{D}^{*0} \pi^- \ell^+ \nu_\ell$ | (4.9 ± 0.8) $\times 10^{-3}$ | |
| $\Gamma_{13} D_1(2420)^- \ell^+ \nu_\ell \times B(D_1^- \rightarrow \bar{D}^{*0} \pi^-)$ | (5.4 ± 2.1) $\times 10^{-3}$ | |
| $\Gamma_{14} D'_1(2430)^- \ell^+ \nu_\ell \times B(D'_1^- \rightarrow \bar{D}^{*0} \pi^-)$ | < 5.0 $\times 10^{-3}$ | CL=90% |
| $\Gamma_{15} D_2^*(2460)^- \ell^+ \nu_\ell \times B(D_2^{*-} \rightarrow \bar{D}^{*0} \pi^-)$ | < 3.0 $\times 10^{-3}$ | CL=90% |
| $\Gamma_{16} \rho^- \ell^+ \nu_\ell$ | [a] (2.47 ± 0.33) $\times 10^{-4}$ | |
| $\Gamma_{17} \pi^- \ell^+ \nu_\ell$ | [a] (1.36 ± 0.09) $\times 10^{-4}$ | |
| $\Gamma_{18} \pi^- \mu^+ \nu_\mu$ | | |

Inclusive modes

| | | |
|-----------------------------------|--|--------|
| $\Gamma_{19} K^\pm$ anything | (78 ± 8) % | |
| $\Gamma_{20} D^0 X$ | (8.1 ± 1.5) % | |
| $\Gamma_{21} \bar{D}^0 X$ | (47.4 ± 2.8) % | |
| $\Gamma_{22} D^+ X$ | < 3.9 % | CL=90% |
| $\Gamma_{23} D^- X$ | (36.9 ± 3.3) % | |
| $\Gamma_{24} D_s^+ X$ | ($10.3 \begin{array}{l} +2.1 \\ -1.8 \end{array}$) % | |
| $\Gamma_{25} D_s^- X$ | < 2.6 % | CL=90% |
| $\Gamma_{26} \Lambda_c^+ X$ | < 3.1 % | CL=90% |
| $\Gamma_{27} \bar{\Lambda}_c^- X$ | ($5.0 \begin{array}{l} +2.1 \\ -1.5 \end{array}$) % | |

| | | |
|---------------|-------------|----------------------|
| Γ_{28} | $\bar{c}X$ | (95 \pm 5) % |
| Γ_{29} | cX | (24.6 \pm 3.1) % |
| Γ_{30} | $\bar{c}cX$ | (119 \pm 6) % |

 D , D^* , or D_s modes

| | | |
|---------------|--|---|
| Γ_{31} | $D^- \pi^+$ | (2.68 \pm 0.13) $\times 10^{-3}$ |
| Γ_{32} | $D^- \rho^+$ | (7.7 \pm 1.3) $\times 10^{-3}$ |
| Γ_{33} | $D^- K^0 \pi^+$ | (4.9 \pm 0.9) $\times 10^{-4}$ |
| Γ_{34} | $D^- K^*(892)^+$ | (4.5 \pm 0.7) $\times 10^{-4}$ |
| Γ_{35} | $D^- \omega \pi^+$ | (2.8 \pm 0.6) $\times 10^{-3}$ |
| Γ_{36} | $D^- K^+$ | (2.0 \pm 0.6) $\times 10^{-4}$ |
| Γ_{37} | $D^- K^+ \bar{K}^0$ | < 3.1 $\times 10^{-4}$ |
| Γ_{38} | $D^- K^+ \bar{K}^*(892)^0$ | (8.8 \pm 1.9) $\times 10^{-4}$ |
| Γ_{39} | $\bar{D}^0 \pi^+ \pi^-$ | (8.4 \pm 0.9) $\times 10^{-4}$ |
| Γ_{40} | $D^*(2010)^- \pi^+$ | (2.76 \pm 0.13) $\times 10^{-3}$ |
| Γ_{41} | $D^- \pi^+ \pi^+ \pi^-$ | (8.0 \pm 2.5) $\times 10^{-3}$ |
| Γ_{42} | $(D^- \pi^+ \pi^+ \pi^-)$ nonresonant | (3.9 \pm 1.9) $\times 10^{-3}$ |
| Γ_{43} | $D^- \pi^+ \rho^0$ | (1.1 \pm 1.0) $\times 10^{-3}$ |
| Γ_{44} | $D^- a_1(1260)^+$ | (6.0 \pm 3.3) $\times 10^{-3}$ |
| Γ_{45} | $D^*(2010)^- \pi^+ \pi^0$ | (1.5 \pm 0.5) % |
| Γ_{46} | $D^*(2010)^- \rho^+$ | (6.8 \pm 0.9) $\times 10^{-3}$ |
| Γ_{47} | $D^*(2010)^- K^+$ | (2.14 \pm 0.16) $\times 10^{-4}$ |
| Γ_{48} | $D^*(2010)^- K^0 \pi^+$ | (3.0 \pm 0.8) $\times 10^{-4}$ |
| Γ_{49} | $D^*(2010)^- K^*(892)^+$ | (3.3 \pm 0.6) $\times 10^{-4}$ |
| Γ_{50} | $D^*(2010)^- K^+ \bar{K}^0$ | < 4.7 $\times 10^{-4}$ |
| Γ_{51} | $D^*(2010)^- K^+ \bar{K}^*(892)^0$ | (1.29 \pm 0.33) $\times 10^{-3}$ |
| Γ_{52} | $D^*(2010)^- \pi^+ \pi^+ \pi^-$ | (7.0 \pm 0.8) $\times 10^{-3}$ |
| Γ_{53} | $(D^*(2010)^- \pi^+ \pi^+ \pi^-)$ non-resonant | (0.0 \pm 2.5) $\times 10^{-3}$ |
| Γ_{54} | $D^*(2010)^- \pi^+ \rho^0$ | (5.7 \pm 3.2) $\times 10^{-3}$ |
| Γ_{55} | $D^*(2010)^- a_1(1260)^+$ | (1.30 \pm 0.27) % |
| Γ_{56} | $D^*(2010)^- \pi^+ \pi^+ \pi^- \pi^0$ | (1.76 \pm 0.27) % |
| Γ_{57} | $D^*- 3\pi^+ 2\pi^-$ | (4.7 \pm 0.9) $\times 10^{-3}$ |
| Γ_{58} | $D^*(2010)^- \rho \bar{\rho} \pi^+$ | (6.5 \pm 1.6) $\times 10^{-4}$ |
| Γ_{59} | $D^*(2010)^- \rho \bar{n}$ | (1.5 \pm 0.4) $\times 10^{-3}$ |
| Γ_{60} | $\bar{D}^*(2010)^- \omega \pi^+$ | (2.89 \pm 0.30) $\times 10^{-3}$ |
| Γ_{61} | $D_1(2430)^0 \omega \times$ $B(D_1(2430)^0 \rightarrow D^{*-} \pi^+)$ | (4.1 \pm 1.6) $\times 10^{-4}$ |
| Γ_{62} | $\bar{D}^{**-} \pi^+$ | [b] (2.1 \pm 1.0) $\times 10^{-3}$ |
| Γ_{63} | $D_1(2420)^- \pi^+ \times B(D_1^- \rightarrow D^- \pi^+ \pi^-)$ | (8.9 $^{+2.3}_{-3.5}$) $\times 10^{-5}$ |
| Γ_{64} | $D_1(2420)^- \pi^+ \times B(D_1^- \rightarrow D^{*-} \pi^+ \pi^-)$ | < 3.3 $\times 10^{-5}$ |
| | | CL=90% |

| | | | |
|---------------|--|-----------------------------------|--------|
| Γ_{65} | $\overline{D}_2^*(2460)^-\pi^+ \times$ $B(D_2^*(2460)^-\rightarrow D^0\pi^-)$ | $(-2.15 \pm 0.35) \times 10^{-4}$ | |
| Γ_{66} | $\overline{D}_0^*(2400)^-\pi^+ \times$ $B(D_0^*(2400)^-\rightarrow D^0\pi^-)$ | $(-6.0 \pm 3.0) \times 10^{-5}$ | |
| Γ_{67} | $D_2^*(2460)^-\pi^+ \times B((D_2^*)^-\rightarrow$ $D^{*-}\pi^+\pi^-)$ | $< 2.4 \times 10^{-5}$ | CL=90% |
| Γ_{68} | $\overline{D}_2^*(2460)^-\rho^+$ | $< 4.9 \times 10^{-3}$ | CL=90% |
| Γ_{69} | $D^0\overline{D}^0$ | $< 6 \times 10^{-5}$ | CL=90% |
| Γ_{70} | $D^{*0}\overline{D}^0$ | $< 2.9 \times 10^{-4}$ | CL=90% |
| Γ_{71} | D^-D^+ | $(-2.11 \pm 0.31) \times 10^{-4}$ | S=1.2 |
| Γ_{72} | $D^-D_s^+$ | $(-7.4 \pm 0.7) \times 10^{-3}$ | |
| Γ_{73} | $D^*(2010)^-D_s^+$ | $(-8.3 \pm 1.1) \times 10^{-3}$ | |
| Γ_{74} | $D^-D_s^{*+}$ | $(-7.6 \pm 1.6) \times 10^{-3}$ | |
| Γ_{75} | $D^*(2010)^-D_s^{*+}$ | $(-1.79 \pm 0.14) \%$ | |
| Γ_{76} | $D_{s0}(2317)^-K^+ \times$ $B(D_{s0}(2317)^-\rightarrow D_s^-\pi^0)$ | $(-4.4 \pm 1.4) \times 10^{-5}$ | |
| Γ_{77} | $D_{s0}(2317)^-\pi^+ \times$ $B(D_{s0}(2317)^-\rightarrow D_s^-\pi^0)$ | $< 2.5 \times 10^{-5}$ | CL=90% |
| Γ_{78} | $D_{sJ}(2457)^-K^+ \times$ $B(D_{sJ}(2457)^-\rightarrow D_s^-\pi^0)$ | $< 9.4 \times 10^{-6}$ | CL=90% |
| Γ_{79} | $D_{sJ}(2457)^-\pi^+ \times$ $B(D_{sJ}(2457)^-\rightarrow D_s^-\pi^0)$ | $< 4.0 \times 10^{-6}$ | CL=90% |
| Γ_{80} | $D_s^-D_s^+$ | $< 3.6 \times 10^{-5}$ | CL=90% |
| Γ_{81} | $D_s^{*-}D_s^+$ | $< 1.3 \times 10^{-4}$ | CL=90% |
| Γ_{82} | $D_s^{*-}D_s^{*+}$ | $< 2.4 \times 10^{-4}$ | CL=90% |
| Γ_{83} | $D_{s0}(2317)^+D^- \times$ $B(D_{s0}(2317)^+\rightarrow D_s^+\pi^0)$ | $(-9.9 \pm 4.2) \times 10^{-4}$ | S=1.5 |
| Γ_{84} | $D_{s0}(2317)^+D^- \times$ $B(D_{s0}(2317)^+\rightarrow D_s^{*+}\gamma)$ | $< 9.5 \times 10^{-4}$ | CL=90% |
| Γ_{85} | $D_{s0}(2317)^+D^*(2010)^- \times$ $B(D_{s0}(2317)^+\rightarrow D_s^+\pi^0)$ | $(-1.5 \pm 0.6) \times 10^{-3}$ | |
| Γ_{86} | $D_{sJ}(2457)^+D^-$ | $(-3.5 \pm 1.1) \times 10^{-3}$ | |
| Γ_{87} | $D_{sJ}(2457)^+D^- \times$ $B(D_{sJ}(2457)^+\rightarrow D_s^+\gamma)$ | $(-6.7 \pm 1.7) \times 10^{-4}$ | |
| Γ_{88} | $D_{sJ}(2457)^+D^- \times$ $B(D_{sJ}(2457)^+\rightarrow D_s^{*+}\gamma)$ | $< 6.0 \times 10^{-4}$ | CL=90% |
| Γ_{89} | $D_{sJ}(2457)^+D^- \times$ $B(D_{sJ}(2457)^+\rightarrow D_s^+\pi^+\pi^-)$ | $< 2.0 \times 10^{-4}$ | CL=90% |

| | | | | |
|----------------|---|---------------------|------------------|--------|
| Γ_{90} | $D_{sJ}(2457)^+ D^- \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^0)$ | < 3.6 | $\times 10^{-4}$ | CL=90% |
| Γ_{91} | $D^*(2010)^- D_{sJ}(2457)^+$ | (9.3 \pm 2.2) | $\times 10^{-3}$ | |
| Γ_{92} | $D_{sJ}(2457)^+ D^*(2010) \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)$ | (2.3 \pm 0.9) | $\times 10^{-3}$ | |
| Γ_{93} | $D^- D_{s1}(2536)^+ \times$ $B(D_{s1}(2536)^+ \rightarrow D^{*0} K^+)$ | (1.7 \pm 0.6) | $\times 10^{-4}$ | |
| Γ_{94} | $D^*(2010)^- D_{s1}(2536)^+ \times$ $B(D_{s1}(2536)^+ \rightarrow D^{*0} K^+)$ | (3.3 \pm 1.1) | $\times 10^{-4}$ | |
| Γ_{95} | $D^- D_{s1}(2536)^+ \times$ $B(D_{s1}(2536)^+ \rightarrow D^{*+} K^0)$ | (2.6 \pm 1.1) | $\times 10^{-4}$ | |
| Γ_{96} | $D^{*-} D_{s1}(2536)^+ \times$ $B(D_{s1}(2536)^+ \rightarrow D^{*+} K^0)$ | (5.0 \pm 1.7) | $\times 10^{-4}$ | |
| Γ_{97} | $D^- D_{sJ}(2573)^+ \times$ $B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)$ | < 1 | $\times 10^{-4}$ | CL=90% |
| Γ_{98} | $D^*(2010)^- D_{sJ}(2573)^+ \times$ $B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)$ | < 2 | $\times 10^{-4}$ | CL=90% |
| Γ_{99} | $D_s^+ \pi^-$ | (1.53 \pm 0.35) | $\times 10^{-5}$ | |
| Γ_{100} | $D_s^{*+} \pi^-$ | (3.0 \pm 0.7) | $\times 10^{-5}$ | |
| Γ_{101} | $D_s^+ \rho^-$ | < 6 | $\times 10^{-4}$ | CL=90% |
| Γ_{102} | $D_s^{*+} \rho^-$ | < 6 | $\times 10^{-4}$ | CL=90% |
| Γ_{103} | $D_s^+ a_0^-$ | < 1.9 | $\times 10^{-5}$ | CL=90% |
| Γ_{104} | $D_s^{*+} a_0^-$ | < 3.6 | $\times 10^{-5}$ | CL=90% |
| Γ_{105} | $D_s^+ a_1(1260)^-$ | < 2.2 | $\times 10^{-3}$ | CL=90% |
| Γ_{106} | $D_s^{*+} a_1(1260)^-$ | < 1.8 | $\times 10^{-3}$ | CL=90% |
| Γ_{107} | $D_s^+ a_2^-$ | < 1.9 | $\times 10^{-4}$ | CL=90% |
| Γ_{108} | $D_s^{*+} a_2^-$ | < 2.0 | $\times 10^{-4}$ | CL=90% |
| Γ_{109} | $D_s^- K^+$ | (2.9 \pm 0.5) | $\times 10^{-5}$ | |
| Γ_{110} | $D_s^{*-} K^+$ | (2.2 \pm 0.6) | $\times 10^{-5}$ | |
| Γ_{111} | $D_s^- K^*(892)^+$ | < 8 | $\times 10^{-4}$ | CL=90% |
| Γ_{112} | $D_s^{*-} K^*(892)^+$ | < 9 | $\times 10^{-4}$ | CL=90% |
| Γ_{113} | $D_s^- \pi^+ K^0$ | < 5 | $\times 10^{-3}$ | CL=90% |
| Γ_{114} | $D_s^{*-} \pi^+ K^0$ | < 2.6 | $\times 10^{-3}$ | CL=90% |
| Γ_{115} | $D_s^- \pi^+ K^*(892)^0$ | < 3.1 | $\times 10^{-3}$ | CL=90% |
| Γ_{116} | $D_s^{*-} \pi^+ K^*(892)^0$ | < 1.7 | $\times 10^{-3}$ | CL=90% |
| Γ_{117} | $\overline{D}^0 K^0$ | (5.2 \pm 0.7) | $\times 10^{-5}$ | |
| Γ_{118} | $\overline{D}^0 K^+ \pi^-$ | (8.8 \pm 1.7) | $\times 10^{-5}$ | |
| Γ_{119} | $\overline{D}^0 K^*(892)^0$ | (4.2 \pm 0.6) | $\times 10^{-5}$ | |
| Γ_{120} | $D_2^*(2460)^- K^+ \times$ $B(D_2^*(2460)^- \rightarrow \overline{D}^0 \pi^-)$ | (1.8 \pm 0.5) | $\times 10^{-5}$ | |
| Γ_{121} | $\overline{D}^0 K^+ \pi^- \text{ non-resonant}$ | < 3.7 | $\times 10^{-5}$ | CL=90% |

| | | | |
|----------------|---|----------------------------------|-----------|
| Γ_{122} | $\bar{D}^0 \pi^0$ | $(2.61 \pm 0.24) \times 10^{-4}$ | |
| Γ_{123} | $\bar{D}^0 \rho^0$ | $(3.2 \pm 0.5) \times 10^{-4}$ | |
| Γ_{124} | $\bar{D}^0 f_2$ | $(1.2 \pm 0.4) \times 10^{-4}$ | |
| Γ_{125} | $\bar{D}^0 \eta$ | $(2.02 \pm 0.35) \times 10^{-4}$ | $S=1.6$ |
| Γ_{126} | $\bar{D}^0 \eta'$ | $(1.25 \pm 0.23) \times 10^{-4}$ | $S=1.1$ |
| Γ_{127} | $\bar{D}^0 \omega$ | $(2.59 \pm 0.30) \times 10^{-4}$ | |
| Γ_{128} | $D^0 \phi$ | $< 1.16 \times 10^{-5}$ | $CL=90\%$ |
| Γ_{129} | $D^0 K^+ \pi^-$ | $< 1.9 \times 10^{-5}$ | $CL=90\%$ |
| Γ_{130} | $D^0 K^*(892)^0$ | $< 1.1 \times 10^{-5}$ | $CL=90\%$ |
| Γ_{131} | $\bar{D}^{*0} \gamma$ | $< 2.5 \times 10^{-5}$ | $CL=90\%$ |
| Γ_{132} | $\bar{D}^*(2007)^0 \pi^0$ | $(1.7 \pm 0.4) \times 10^{-4}$ | $S=1.5$ |
| Γ_{133} | $\bar{D}^*(2007)^0 \rho^0$ | $< 5.1 \times 10^{-4}$ | $CL=90\%$ |
| Γ_{134} | $\bar{D}^*(2007)^0 \eta$ | $(1.8 \pm 0.6) \times 10^{-4}$ | $S=1.8$ |
| Γ_{135} | $\bar{D}^*(2007)^0 \eta'$ | $(1.23 \pm 0.35) \times 10^{-4}$ | |
| Γ_{136} | $\bar{D}^*(2007)^0 \pi^+ \pi^-$ | $(6.2 \pm 2.2) \times 10^{-4}$ | |
| Γ_{137} | $\bar{D}^*(2007)^0 K^0$ | $(3.6 \pm 1.2) \times 10^{-5}$ | |
| Γ_{138} | $\bar{D}^*(2007)^0 K^*(892)^0$ | $< 6.9 \times 10^{-5}$ | $CL=90\%$ |
| Γ_{139} | $D^*(2007)^0 K^*(892)^0$ | $< 4.0 \times 10^{-5}$ | $CL=90\%$ |
| Γ_{140} | $D^*(2007)^0 \pi^+ \pi^+ \pi^- \pi^-$ | $(2.7 \pm 0.5) \times 10^{-3}$ | |
| Γ_{141} | $D^*(2010)^+ D^*(2010)^-$ | $(8.2 \pm 0.9) \times 10^{-4}$ | |
| Γ_{142} | $\bar{D}^*(2007)^0 \omega$ | $(2.7 \pm 0.8) \times 10^{-4}$ | $S=1.5$ |
| Γ_{143} | $D^*(2010)^+ D^-$ | $(6.1 \pm 1.5) \times 10^{-4}$ | $S=1.6$ |
| Γ_{144} | $D^*(2007)^0 \bar{D}^*(2007)^0$ | $< 9 \times 10^{-5}$ | $CL=90\%$ |
| Γ_{145} | $D^- D^0 K^+$ | $(1.7 \pm 0.4) \times 10^{-3}$ | |
| Γ_{146} | $D^- D^*(2007)^0 K^+$ | $(4.6 \pm 1.0) \times 10^{-3}$ | |
| Γ_{147} | $D^*(2010)^- D^0 K^+$ | $(3.1 \pm 0.6) \times 10^{-3}$ | |
| Γ_{148} | $D^*(2010)^- D^*(2007)^0 K^+$ | $(1.18 \pm 0.20) \%$ | |
| Γ_{149} | $D^- D^+ K^0$ | $< 1.7 \times 10^{-3}$ | $CL=90\%$ |
| Γ_{150} | $D^*(2010)^- D^+ K^0 + D^- D^*(2010)^+ K^0$ | $(6.5 \pm 1.6) \times 10^{-3}$ | |
| Γ_{151} | $D^*(2010)^- D^*(2010)^+ K^0$ | $(7.8 \pm 1.1) \times 10^{-3}$ | |
| Γ_{152} | $D^{*-} D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^{*+} K^0)$ | $(8.0 \pm 2.4) \times 10^{-4}$ | |
| Γ_{153} | $\bar{D}^0 D^0 K^0$ | $< 1.4 \times 10^{-3}$ | $CL=90\%$ |
| Γ_{154} | $\bar{D}^0 D^*(2007)^0 K^0 + \bar{D}^*(2007)^0 D^0 K^0$ | $< 3.7 \times 10^{-3}$ | $CL=90\%$ |
| Γ_{155} | $\bar{D}^*(2007)^0 D^*(2007)^0 K^0$ | $< 6.6 \times 10^{-3}$ | $CL=90\%$ |
| Γ_{156} | $(\bar{D} + \bar{D}^*)(D + D^*) K$ | $(4.3 \pm 0.7) \%$ | |

Charmonium modes

| | | | |
|----------------|---|-------------------------------------|--------|
| Γ_{157} | $\eta_c K^0$ | $(-8.9 \pm 1.6) \times 10^{-4}$ | |
| Γ_{158} | $\eta_c K^*(892)^0$ | $(-9.6 \pm 3.3) \times 10^{-4}$ | S=1.1 |
| Γ_{159} | $J/\psi(1S)K^0$ | $(-8.71 \pm 0.32) \times 10^{-4}$ | |
| Γ_{160} | $J/\psi(1S)K^+\pi^-$ | $(-1.2 \pm 0.6) \times 10^{-3}$ | |
| Γ_{161} | $J/\psi(1S)K^*(892)^0$ | $(-1.33 \pm 0.06) \times 10^{-3}$ | |
| Γ_{162} | $J/\psi(1S)\eta K_S^0$ | $(-8 \pm 4) \times 10^{-5}$ | |
| Γ_{163} | $J/\psi(1S)\eta' K_S^0$ | $< 2.5 \times 10^{-5}$ | CL=90% |
| Γ_{164} | $J/\psi(1S)\phi K^0$ | $(-9.4 \pm 2.6) \times 10^{-5}$ | |
| Γ_{165} | $J/\psi(1S)K(1270)^0$ | $(-1.3 \pm 0.5) \times 10^{-3}$ | |
| Γ_{166} | $J/\psi(1S)\pi^0$ | $(-2.05 \pm 0.24) \times 10^{-5}$ | |
| Γ_{167} | $J/\psi(1S)\eta$ | $(-9.5 \pm 1.9) \times 10^{-6}$ | |
| Γ_{168} | $J/\psi(1S)\pi^+\pi^-$ | $(-4.6 \pm 0.9) \times 10^{-5}$ | |
| Γ_{169} | $J/\psi(1S)\pi^+\pi^-$ nonresonant | $< 1.2 \times 10^{-5}$ | CL=90% |
| Γ_{170} | $J/\psi(1S)f_2$ | $< 4.6 \times 10^{-6}$ | CL=90% |
| Γ_{171} | $J/\psi(1S)\rho^0$ | $(-2.7 \pm 0.4) \times 10^{-5}$ | |
| Γ_{172} | $J/\psi(1S)\omega$ | $< 2.7 \times 10^{-4}$ | CL=90% |
| Γ_{173} | $J/\psi(1S)\phi$ | $< 9.2 \times 10^{-6}$ | CL=90% |
| Γ_{174} | $J/\psi(1S)\eta'(958)$ | $< 6.3 \times 10^{-5}$ | CL=90% |
| Γ_{175} | $J/\psi(1S)K^0\pi^+\pi^-$ | $(-1.0 \pm 0.4) \times 10^{-3}$ | |
| Γ_{176} | $J/\psi(1S)K^0\rho^0$ | $(-5.4 \pm 3.0) \times 10^{-4}$ | |
| Γ_{177} | $J/\psi(1S)K^*(892)^+\pi^-$ | $(-8 \pm 4) \times 10^{-4}$ | |
| Γ_{178} | $J/\psi(1S)K^*(892)^0\pi^+\pi^-$ | $(-6.6 \pm 2.2) \times 10^{-4}$ | |
| Γ_{179} | $X(3872)^-K^+$ | $< 5 \times 10^{-4}$ | CL=90% |
| Γ_{180} | $X(3872)^-K^+ \times B(X(3872)^- \rightarrow [c])$ | $< 5.4 \times 10^{-6}$ | CL=90% |
| | $J/\psi(1S)\pi^-\pi^0)$ | | |
| Γ_{181} | $X(3872)K^0 \times B(X \rightarrow J/\psi\pi^+\pi^-)$ | $< 1.03 \times 10^{-5}$ | CL=90% |
| Γ_{182} | $X(3872)K^0 \times B(X \rightarrow D^0\bar{D}^0\pi^0)$ | $(-1.7 \pm 0.8) \times 10^{-4}$ | |
| Γ_{183} | $X(3872)K^0 \times B(X \rightarrow \bar{D}^{*0}D^0)$ | $< 4.37 \times 10^{-4}$ | CL=90% |
| Γ_{184} | $J/\psi(1S)p\bar{p}$ | $< 8.3 \times 10^{-7}$ | CL=90% |
| Γ_{185} | $J/\psi(1S)\gamma$ | $< 1.6 \times 10^{-6}$ | CL=90% |
| Γ_{186} | $J/\psi(1S)\bar{D}^0$ | $< 1.3 \times 10^{-5}$ | CL=90% |
| Γ_{187} | $\psi(2S)K^0$ | $(-6.2 \pm 0.6) \times 10^{-4}$ | |
| Γ_{188} | $\psi(3770)K^0 \times B(\psi \rightarrow \bar{D}^0D^0)$ | $< 1.23 \times 10^{-4}$ | CL=90% |
| Γ_{189} | $\psi(3770)K^0 \times B(\psi \rightarrow D^-D^+)$ | $< 1.88 \times 10^{-4}$ | CL=90% |
| Γ_{190} | $\psi(2S)K^+\pi^-$ | $< 1 \times 10^{-3}$ | CL=90% |
| Γ_{191} | $\psi(2S)K^*(892)^0$ | $(-7.2 \pm 0.8) \times 10^{-4}$ | |
| Γ_{192} | $\chi_{c0}(1P)K^0$ | $< 1.13 \times 10^{-4}$ | CL=90% |
| Γ_{193} | $\chi_{c0}K^*(892)^0$ | $< 7.7 \times 10^{-4}$ | CL=90% |
| Γ_{194} | $\chi_{c2}K^0$ | $< 2.6 \times 10^{-5}$ | CL=90% |

| | | | | |
|----------------|----------------------------|-------------------|------------------|--------|
| Γ_{195} | $\chi_{c2} K^*(892)^0$ | < 3.6 | $\times 10^{-5}$ | CL=90% |
| Γ_{196} | $\chi_{c1}(1P) K^0$ | (3.9 \pm 0.4) | $\times 10^{-4}$ | |
| Γ_{197} | $\chi_{c1}(1P) K^*(892)^0$ | (3.2 \pm 0.6) | $\times 10^{-4}$ | |

K or K^* modes

| | | | | |
|----------------|--|-----------------------|------------------|--------|
| Γ_{198} | $K^+ \pi^-$ | (1.94 \pm 0.06) | $\times 10^{-5}$ | |
| Γ_{199} | $K^0 \pi^0$ | (9.8 \pm 0.6) | $\times 10^{-6}$ | |
| Γ_{200} | $\eta' K^0$ | (6.5 \pm 0.4) | $\times 10^{-5}$ | S=1.2 |
| Γ_{201} | $\eta' K^*(892)^0$ | (3.8 \pm 1.2) | $\times 10^{-6}$ | |
| Γ_{202} | ηK^0 | < 1.9 | $\times 10^{-6}$ | CL=90% |
| Γ_{203} | $\eta K^*(892)^0$ | (1.59 \pm 0.10) | $\times 10^{-5}$ | |
| Γ_{204} | $\eta K_0^*(1430)^0$ | (1.10 \pm 0.22) | $\times 10^{-5}$ | |
| Γ_{205} | $\eta K_2^*(1430)^0$ | (9.6 \pm 2.1) | $\times 10^{-6}$ | |
| Γ_{206} | ωK^0 | (5.0 \pm 0.6) | $\times 10^{-6}$ | |
| Γ_{207} | $a_0(980)^0 K^0 \times B(a_0(980)^0 \rightarrow \eta \pi^0)$ | < 7.8 | $\times 10^{-6}$ | CL=90% |
| Γ_{208} | $a_0(980)^{\pm} K^{\mp} \times B(a_0(980)^{\pm} \rightarrow \eta \pi^{\pm})$ | < 1.9 | $\times 10^{-6}$ | CL=90% |
| Γ_{209} | $a_0(1450)^{\pm} K^{\mp} \times B(a_0(1450)^{\pm} \rightarrow \eta \pi^{\pm})$ | < 3.1 | $\times 10^{-6}$ | CL=90% |
| Γ_{210} | $K_S^0 X^0$ (Familon) | < 5.3 | $\times 10^{-5}$ | CL=90% |
| Γ_{211} | $\omega K^*(892)^0$ | < 4.2 | $\times 10^{-6}$ | CL=90% |
| Γ_{212} | $K^+ K^-$ | < 4.1 | $\times 10^{-7}$ | CL=90% |
| Γ_{213} | $K^0 \bar{K}^0$ | (9.6 \pm 2.0) | $\times 10^{-7}$ | |
| Γ_{214} | $K_S^0 K_S^0 K_S^0$ | (6.2 \pm 1.2) | $\times 10^{-6}$ | S=1.3 |
| Γ_{215} | $K_S^0 K_S^0 K_L^0$ | < 1.6 | $\times 10^{-5}$ | CL=90% |
| Γ_{216} | $K^+ \pi^- \pi^0$ | (3.7 \pm 0.5) | $\times 10^{-5}$ | |
| Γ_{217} | $K^+ \rho^-$ | (8.5 \pm 2.8) | $\times 10^{-6}$ | S=1.7 |
| Γ_{218} | $(K^+ \pi^- \pi^0)$ non-resonant | < 9.4 | $\times 10^{-6}$ | CL=90% |
| Γ_{219} | $K_x^{*0} \pi^0$ | [d] (6.1 \pm 1.6) | $\times 10^{-6}$ | |
| Γ_{220} | $K^0 \pi^+ \pi^-$ charmless | (4.48 \pm 0.26) | $\times 10^{-5}$ | |
| Γ_{221} | $K^0 \pi^+ \pi^-$ non-resonant | (1.99 \pm 0.31) | $\times 10^{-5}$ | |
| Γ_{222} | $K^0 \rho^0$ | (5.4 \pm 0.9) | $\times 10^{-6}$ | |
| Γ_{223} | $K^0 f_0(980)$ | (5.5 \pm 0.9) | $\times 10^{-6}$ | |
| Γ_{224} | $K^*(892)^+ \pi^-$ | (9.8 \pm 1.3) | $\times 10^{-6}$ | S=1.2 |
| Γ_{225} | $K^*(1430)^+ \pi^-$ | (5.0 \pm 0.8) | $\times 10^{-5}$ | |
| Γ_{226} | $K_x^{*+} \pi^-$ | [d] (5.1 \pm 1.6) | $\times 10^{-6}$ | |
| Γ_{227} | $K^*(1410)^+ \pi^- \times B(K^*(1410)^+ \rightarrow K^0 \pi^+)$ | < 3.8 | $\times 10^{-6}$ | CL=90% |
| Γ_{228} | $K^*(1680)^+ \pi^- \times B(K^*(1680)^+ \rightarrow K^0 \pi^+)$ | < 2.6 | $\times 10^{-6}$ | CL=90% |

| | | | | |
|----------------|---|-----------------------------|------------------|--------|
| Γ_{229} | $K_2^*(1430)^+ \pi^- \times B(K_2^*(1430)^+ \rightarrow K^0 \pi^+)$ | < 2.1 | $\times 10^{-6}$ | CL=90% |
| Γ_{230} | $f_0(980) K^0 \times B(f_0(980) \rightarrow \pi^+ \pi^-)$ | (7.6 $^{+1.9}_{-2.1}$) | $\times 10^{-6}$ | |
| Γ_{231} | $f_2(1270) K^0 \times B(f_2(1270) \rightarrow \pi^+ \pi^-)$ | < 1.4 | $\times 10^{-6}$ | CL=90% |
| Γ_{232} | $K^*(892)^0 \pi^0$ | < 3.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{233} | $K_2^*(1430)^+ \pi^-$ | < 1.8 | $\times 10^{-5}$ | CL=90% |
| Γ_{234} | $K^0 K^- \pi^+$ | < 1.8 | $\times 10^{-5}$ | CL=90% |
| Γ_{235} | $\bar{K}^{*0} K^0 + K^{*0} \bar{K}^0$ | < 1.9 | $\times 10^{-6}$ | |
| Γ_{236} | $K^+ K^- \pi^0$ | < 1.9 | $\times 10^{-5}$ | CL=90% |
| Γ_{237} | $K^0 K^+ K^-$ | (2.47 ± 0.23) | $\times 10^{-5}$ | |
| Γ_{238} | $K^0 \phi$ | (8.6 $^{+1.3}_{-1.1}$) | $\times 10^{-6}$ | |
| Γ_{239} | $K^+ \pi^- \pi^+ \pi^-$ | [e] < 2.3 | $\times 10^{-4}$ | CL=90% |
| Γ_{240} | $K^*(892)^0 \pi^+ \pi^-$ | (5.4 ± 0.5) | $\times 10^{-5}$ | |
| Γ_{241} | $K^*(892)^0 \rho^0$ | (5.6 ± 1.6) | $\times 10^{-6}$ | |
| Γ_{242} | $K^*(892)^0 f_0(980)$ | < 4.3 | $\times 10^{-6}$ | CL=90% |
| Γ_{243} | $K_1(1400)^+ \pi^-$ | < 1.1 | $\times 10^{-3}$ | CL=90% |
| Γ_{244} | $a_1(1260)^- K^+$ | [e] (1.6 ± 0.4) | $\times 10^{-5}$ | |
| Γ_{245} | $b_1^- K^+ \times B(b_1^- \rightarrow \omega \pi^-)$ | (7.4 ± 1.4) | $\times 10^{-6}$ | |
| Γ_{246} | $K^*(892)^0 K^+ K^-$ | (2.75 ± 0.26) | $\times 10^{-5}$ | |
| Γ_{247} | $K^*(892)^0 \phi$ | (9.5 ± 0.8) | $\times 10^{-6}$ | |
| Γ_{248} | $K^*(892)^0 K^- \pi^+$ | (4.6 ± 1.4) | $\times 10^{-6}$ | |
| Γ_{249} | $K^*(892)^0 \bar{K}^*(892)^0$ | (1.28 $^{+0.37}_{-0.32}$) | $\times 10^{-6}$ | |
| Γ_{250} | $K^*(892)^0 K^+ \pi^-$ | < 2.2 | $\times 10^{-6}$ | CL=90% |
| Γ_{251} | $K^*(892)^0 K^*(892)^0$ | < 4.1 | $\times 10^{-7}$ | CL=90% |
| Γ_{252} | $K^*(892)^+ \rho^-$ | < 1.20 | $\times 10^{-5}$ | CL=90% |
| Γ_{253} | $K^*(892)^+ K^*(892)^-$ | < 1.41 | $\times 10^{-4}$ | CL=90% |
| Γ_{254} | $K_1(1400)^0 \rho^0$ | < 3.0 | $\times 10^{-3}$ | CL=90% |
| Γ_{255} | $K_1(1400)^0 \phi$ | < 5.0 | $\times 10^{-3}$ | CL=90% |
| Γ_{256} | $\phi(K\pi)_0^{*0}$ | (5.0 ± 0.9) | $\times 10^{-6}$ | |
| Γ_{257} | $\phi(K\pi)_0^{*0} (1.60 < m_{K\pi} < 2.15)$ | [f] < 1.7 | $\times 10^{-6}$ | CL=90% |
| Γ_{258} | $K_0^*(1430)^0 \phi$ | (4.6 ± 0.9) | $\times 10^{-6}$ | |
| Γ_{259} | $K^*(1680)^0 \phi$ | < 3.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{260} | $K^*(1780)^0 \phi$ | < 2.7 | $\times 10^{-6}$ | CL=90% |
| Γ_{261} | $K^*(2045)^0 \phi$ | < 1.53 | $\times 10^{-5}$ | CL=90% |
| Γ_{262} | $K_2^*(1430)^0 \rho^0$ | < 1.1 | $\times 10^{-3}$ | CL=90% |
| Γ_{263} | $K_2^*(1430)^0 \phi$ | (7.8 ± 1.3) | $\times 10^{-6}$ | |
| Γ_{264} | $K^0 \phi \phi$ | (4.1 $^{+1.7}_{-1.5}$) | $\times 10^{-6}$ | |
| Γ_{265} | $\eta' \eta' K^0$ | < 3.1 | $\times 10^{-5}$ | CL=90% |
| Γ_{266} | $K^*(892)^0 \gamma$ | (4.01 ± 0.20) | $\times 10^{-5}$ | |

| | | | |
|----------------|--------------------------------|--|--------|
| Γ_{267} | $\eta K^0 \gamma$ | $(-1.07^{+0.22}_{-0.15}) \times 10^{-5}$ | |
| Γ_{268} | $\eta' K^0 \gamma$ | $< 6.6 \times 10^{-6}$ | CL=90% |
| Γ_{269} | $K^0 \phi \gamma$ | $< 2.7 \times 10^{-6}$ | CL=90% |
| Γ_{270} | $K^+ \pi^- \gamma$ | $(4.6 \pm 1.4) \times 10^{-6}$ | |
| Γ_{271} | $K^*(1410) \gamma$ | $< 1.3 \times 10^{-4}$ | CL=90% |
| Γ_{272} | $K^+ \pi^- \gamma$ nonresonant | $< 2.6 \times 10^{-6}$ | CL=90% |
| Γ_{273} | $K^0 \pi^+ \pi^- \gamma$ | $(1.95 \pm 0.22) \times 10^{-5}$ | |
| Γ_{274} | $K^+ \pi^- \pi^0 \gamma$ | $(4.1 \pm 0.4) \times 10^{-5}$ | |
| Γ_{275} | $K_1(1270)^0 \gamma$ | $< 5.8 \times 10^{-5}$ | CL=90% |
| Γ_{276} | $K_1(1400)^0 \gamma$ | $< 1.5 \times 10^{-5}$ | CL=90% |
| Γ_{277} | $K_2^*(1430)^0 \gamma$ | $(1.24 \pm 0.24) \times 10^{-5}$ | |
| Γ_{278} | $K^*(1680)^0 \gamma$ | $< 2.0 \times 10^{-3}$ | CL=90% |
| Γ_{279} | $K_3^*(1780)^0 \gamma$ | $< 8.3 \times 10^{-5}$ | CL=90% |
| Γ_{280} | $K_4^*(2045)^0 \gamma$ | $< 4.3 \times 10^{-3}$ | CL=90% |

Light unflavored meson modes

| | | | |
|----------------|---|--------------------------------------|--------|
| Γ_{281} | $\rho^0 \gamma$ | $(9.3 \pm 2.1) \times 10^{-7}$ | S=1.1 |
| Γ_{282} | $\omega \gamma$ | $(4.6^{+2.0}_{-1.7}) \times 10^{-7}$ | |
| Γ_{283} | $\phi \gamma$ | $< 8.5 \times 10^{-7}$ | CL=90% |
| Γ_{284} | $\pi^+ \pi^-$ | $(5.13 \pm 0.24) \times 10^{-6}$ | |
| Γ_{285} | $\pi^0 \pi^0$ | $(1.62 \pm 0.31) \times 10^{-6}$ | S=1.3 |
| Γ_{286} | $\eta \pi^0$ | $< 1.3 \times 10^{-6}$ | CL=90% |
| Γ_{287} | $\eta \eta$ | $< 1.8 \times 10^{-6}$ | CL=90% |
| Γ_{288} | $\eta' \pi^0$ | $(1.5^{+1.0}_{-0.8}) \times 10^{-6}$ | S=1.5 |
| Γ_{289} | $\eta' \eta'$ | $< 2.4 \times 10^{-6}$ | CL=90% |
| Γ_{290} | $\eta' \eta$ | $< 1.7 \times 10^{-6}$ | CL=90% |
| Γ_{291} | $\eta' \rho^0$ | $< 1.3 \times 10^{-6}$ | CL=90% |
| Γ_{292} | $\eta' f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)$ | $< 1.5 \times 10^{-6}$ | CL=90% |
| Γ_{293} | $\eta \rho^0$ | $< 1.5 \times 10^{-6}$ | CL=90% |
| Γ_{294} | $\eta f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)$ | $< 4 \times 10^{-7}$ | CL=90% |
| Γ_{295} | $\omega \eta$ | $< 1.9 \times 10^{-6}$ | CL=90% |
| Γ_{296} | $\omega \eta'$ | $< 2.2 \times 10^{-6}$ | CL=90% |
| Γ_{297} | $\omega \rho^0$ | $< 1.5 \times 10^{-6}$ | CL=90% |
| Γ_{298} | $\omega f_0(980)$ | $< 1.5 \times 10^{-6}$ | CL=90% |
| Γ_{299} | $\omega \omega$ | $< 4.0 \times 10^{-6}$ | CL=90% |
| Γ_{300} | $\phi \pi^0$ | $< 2.8 \times 10^{-7}$ | CL=90% |
| Γ_{301} | $\phi \eta$ | $< 6 \times 10^{-7}$ | CL=90% |
| Γ_{302} | $\phi \eta'$ | $< 5 \times 10^{-7}$ | CL=90% |
| Γ_{303} | $\phi \rho^0$ | $< 1.3 \times 10^{-5}$ | CL=90% |
| Γ_{304} | $\phi \omega$ | $< 1.2 \times 10^{-6}$ | CL=90% |

| | | | | | |
|----------------|--|-----|---------------------|----------------------|--------|
| Γ_{305} | $\phi\phi$ | < | 1.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{306} | $a_0(980)^{\pm}\pi^{\mp} \times B(a_0(980)^{\pm} \rightarrow \eta\pi^{\pm})$ | < | 3.1 | $\times 10^{-6}$ | CL=90% |
| Γ_{307} | $a_0(1450)^{\pm}\pi^{\mp} \times B(a_0(1450)^{\pm} \rightarrow \eta\pi^{\pm})$ | < | 2.3 | $\times 10^{-6}$ | CL=90% |
| Γ_{308} | $\pi^{+}\pi^{-}\pi^0$ | < | 7.2 | $\times 10^{-4}$ | CL=90% |
| Γ_{309} | $\rho^0\pi^0$ | | (1.8 ± 0.5) | $\times 10^{-6}$ | |
| Γ_{310} | $\rho^{\mp}\pi^{\pm}$ | [g] | (2.28 ± 0.25) | $\times 10^{-5}$ | |
| Γ_{311} | $\pi^{+}\pi^{-}\pi^{+}\pi^{-}$ | < | 2.3 | $\times 10^{-4}$ | CL=90% |
| Γ_{312} | $\rho^0\rho^0$ | | (1.1 ± 0.4) | $\times 10^{-6}$ | |
| Γ_{313} | $\rho^0 f_0(980) \times B(f_0(980) \rightarrow \pi^{+}\pi^{-})$ | < | 5.3 | $\times 10^{-7}$ | CL=90% |
| Γ_{314} | $f_0(980)f_0(980) \times B(f_0(980) \rightarrow \pi^{+}\pi^{-})^2$ | < | 1.6 | $\times 10^{-7}$ | CL=90% |
| Γ_{315} | $a_1(1260)^{\mp}\pi^{\pm}$ | [g] | (3.3 ± 0.5) | $\times 10^{-5}$ | |
| Γ_{316} | $b_1^{\mp}\pi^{\pm} \times B(b_1^{\mp} \rightarrow \omega\pi^{\mp})$ | | (1.09 ± 0.15) | $\times 10^{-5}$ | |
| Γ_{317} | $a_2(1320)^{\mp}\pi^{\pm}$ | [g] | < | 3.0 $\times 10^{-4}$ | CL=90% |
| Γ_{318} | $\pi^{+}\pi^{-}\pi^0\pi^0$ | < | 3.1 | $\times 10^{-3}$ | CL=90% |
| Γ_{319} | $\rho^{+}\rho^{-}$ | | (2.42 ± 0.31) | $\times 10^{-5}$ | |
| Γ_{320} | $a_1(1260)^0\pi^0$ | < | 1.1 | $\times 10^{-3}$ | CL=90% |
| Γ_{321} | $\omega\pi^0$ | < | 1.2 | $\times 10^{-6}$ | CL=90% |
| Γ_{322} | $\pi^{+}\pi^{+}\pi^{-}\pi^{-}\pi^0$ | < | 9.0 | $\times 10^{-3}$ | CL=90% |
| Γ_{323} | $a_1(1260)^{+}\rho^{-}$ | < | 6.1 | $\times 10^{-5}$ | CL=90% |
| Γ_{324} | $a_1(1260)^0\rho^0$ | < | 2.4 | $\times 10^{-3}$ | CL=90% |
| Γ_{325} | $\pi^{+}\pi^{+}\pi^{+}\pi^{-}\pi^{-}\pi^{-}$ | < | 3.0 | $\times 10^{-3}$ | CL=90% |
| Γ_{326} | $a_1(1260)^{+}a_1(1260)^{-}$ | < | 2.8 | $\times 10^{-3}$ | CL=90% |
| Γ_{327} | $\pi^{+}\pi^{+}\pi^{+}\pi^{-}\pi^{-}\pi^{-}\pi^0$ | < | 1.1 | % | CL=90% |

Baryon modes

| | | | | | |
|----------------|---|-----|-------------------|--------------------|--------|
| Γ_{328} | $p\bar{p}$ | < | 1.1 | $\times 10^{-7}$ | CL=90% |
| Γ_{329} | $p\bar{p}\pi^{+}\pi^{-}$ | < | 2.5 | $\times 10^{-4}$ | CL=90% |
| Γ_{330} | $p\bar{p}K^0$ | | (2.7 ± 0.4) | $\times 10^{-6}$ | |
| Γ_{331} | $\Theta(1540)^{+}\bar{p} \times B(\Theta(1540)^{+} \rightarrow pK_S^0)$ | [h] | < | 5 $\times 10^{-8}$ | CL=90% |
| Γ_{332} | $f_J(2220)K^0 \times B(f_J(2220) \rightarrow p\bar{p})$ | < | 4.5 | $\times 10^{-7}$ | CL=90% |
| Γ_{333} | $p\bar{p}K^{*}(892)^0$ | | (1.5 ± 0.6) | $\times 10^{-6}$ | |
| Γ_{334} | $f_J(2220)K_0^{*} \times B(f_J(2220) \rightarrow p\bar{p})$ | < | 1.5 | $\times 10^{-7}$ | CL=90% |
| Γ_{335} | $p\bar{\Lambda}\pi^{-}$ | | (3.2 ± 0.4) | $\times 10^{-6}$ | |
| Γ_{336} | $p\bar{\Sigma}(1385)^{-}$ | < | 2.6 | $\times 10^{-7}$ | CL=90% |
| Γ_{337} | $\Delta^0\bar{\Lambda}$ | < | 9.3 | $\times 10^{-7}$ | CL=90% |
| Γ_{338} | $p\bar{\Lambda}K^{-}$ | < | 8.2 | $\times 10^{-7}$ | CL=90% |
| Γ_{339} | $p\bar{\Sigma}^0\pi^{-}$ | < | 3.8 | $\times 10^{-6}$ | CL=90% |

| | | | | | |
|----------------|---|---|-----------------|--------------------|--------|
| Γ_{340} | $\bar{\Lambda}\Lambda$ | < | 3.2 | $\times 10^{-7}$ | CL=90% |
| Γ_{341} | $\Delta^0 \bar{\Delta}^0$ | < | 1.5 | $\times 10^{-3}$ | CL=90% |
| Γ_{342} | $\Delta^{++} \bar{\Delta}^{--}$ | < | 1.1 | $\times 10^{-4}$ | CL=90% |
| Γ_{343} | $\bar{D}^0 p \bar{p}$ | (| 1.14 ± 0.09 | $) \times 10^{-4}$ | |
| Γ_{344} | $D_s^- \bar{\Lambda} p$ | (| 2.9 ± 0.9 | $) \times 10^{-5}$ | |
| Γ_{345} | $\bar{D}^*(2007)^0 p \bar{p}$ | (| 1.03 ± 0.13 | $) \times 10^{-4}$ | |
| Γ_{346} | $D^- p \bar{p} \pi^+$ | (| 3.38 ± 0.32 | $) \times 10^{-4}$ | |
| Γ_{347} | $D^{*-} p \bar{p} \pi^+$ | (| 4.8 ± 0.5 | $) \times 10^{-4}$ | |
| Γ_{348} | $\Theta_c \bar{p} \pi^+ \times B(\Theta_c \rightarrow D^- p)$ | < | 9 | $\times 10^{-6}$ | CL=90% |
| Γ_{349} | $\Theta_c \bar{p} \pi^+ \times B(\Theta_c \rightarrow D^{*-} p)$ | < | 1.4 | $\times 10^{-5}$ | CL=90% |
| Γ_{350} | $\bar{\Sigma}_c^{--} \Delta^{++}$ | < | 1.0 | $\times 10^{-3}$ | CL=90% |
| Γ_{351} | $\bar{\Lambda}_c^- p \pi^+ \pi^-$ | (| 1.3 ± 0.4 | $) \times 10^{-3}$ | |
| Γ_{352} | $\bar{\Lambda}_c^- p$ | (| 2.1 ± 0.7 | $) \times 10^{-5}$ | |
| Γ_{353} | $\bar{\Lambda}_c^- p \pi^0$ | < | 5.9 | $\times 10^{-4}$ | CL=90% |
| Γ_{354} | $\bar{\Lambda}_c^- p \pi^+ \pi^- \pi^0$ | < | 5.07 | $\times 10^{-3}$ | CL=90% |
| Γ_{355} | $\bar{\Lambda}_c^- p \pi^+ \pi^- \pi^+ \pi^-$ | < | 2.74 | $\times 10^{-3}$ | CL=90% |
| Γ_{356} | $\Lambda_c^+ \bar{p} \pi^+ \pi^-$ | (| 1.12 ± 0.32 | $) \times 10^{-3}$ | |
| Γ_{357} | $\Lambda_c^+ \bar{p} \pi^+ \pi^- \text{(nonresonant)}$ | (| 6.4 ± 1.9 | $) \times 10^{-4}$ | |
| Γ_{358} | $\bar{\Sigma}_c(2520)^{--} p \pi^+$ | (| 1.2 ± 0.4 | $) \times 10^{-4}$ | |
| Γ_{359} | $\bar{\Sigma}_c(2520)^0 p \pi^-$ | < | 3.8 | $\times 10^{-5}$ | CL=90% |
| Γ_{360} | $\bar{\Sigma}_c(2455)^0 p \pi^-$ | (| 1.5 ± 0.5 | $) \times 10^{-4}$ | |
| Γ_{361} | $\bar{\Sigma}_c(2455)^{--} p \pi^+$ | (| 2.2 ± 0.7 | $) \times 10^{-4}$ | |
| Γ_{362} | $\bar{\Lambda}_c^- \Lambda_c^+$ | < | 6.2 | $\times 10^{-5}$ | CL=90% |
| Γ_{363} | $\bar{\Lambda}_c(2593)^- / \bar{\Lambda}_c(2625)^- p$ | < | 1.1 | $\times 10^{-4}$ | CL=90% |
| Γ_{364} | $\Xi_c^- \Lambda_c^+ \times B(\Xi_c^- \rightarrow \Xi^+ \pi^- \pi^-)$ | (| 9 ± 5 | $) \times 10^{-5}$ | |
| Γ_{365} | $\Lambda_c^+ \Lambda_c^- K^0$ | (| 8 ± 5 | $) \times 10^{-4}$ | |

**Lepton Family number (*LF*) violating modes, or
 $\Delta B = 1$ weak neutral current (*B1*) modes**

| | | | | | | |
|----------------|------------------------|-----------|-----|-------------------|------------------|--------|
| Γ_{366} | $\gamma\gamma$ | <i>B1</i> | < | 6.2 | $\times 10^{-7}$ | CL=90% |
| Γ_{367} | $e^+ e^-$ | <i>B1</i> | < | 1.13 | $\times 10^{-7}$ | CL=90% |
| Γ_{368} | $e^+ e^- \gamma$ | <i>B1</i> | < | 1.2 | $\times 10^{-7}$ | CL=90% |
| Γ_{369} | $\mu^+ \mu^-$ | <i>B1</i> | < | 1.5 | $\times 10^{-8}$ | CL=90% |
| Γ_{370} | $\mu^+ \mu^- \gamma$ | <i>B1</i> | < | 1.6 | $\times 10^{-7}$ | CL=90% |
| Γ_{371} | $\tau^+ \tau^-$ | <i>B1</i> | < | 4.1 | $\times 10^{-3}$ | CL=90% |
| Γ_{372} | $\pi^0 \ell^+ \ell^-$ | <i>B1</i> | < | 1.2 | $\times 10^{-7}$ | CL=90% |
| Γ_{373} | $\pi^0 \nu \bar{\nu}$ | <i>B1</i> | < | 2.2 | $\times 10^{-4}$ | CL=90% |
| Γ_{374} | $\pi^0 e^+ e^-$ | <i>B1</i> | < | 1.4 | $\times 10^{-7}$ | CL=90% |
| Γ_{375} | $\pi^0 \mu^+ \mu^-$ | <i>B1</i> | < | 5.1 | $\times 10^{-7}$ | CL=90% |
| Γ_{376} | $K^0 \ell^+ \ell^-$ | <i>B1</i> | [a] | (2.9 ± 1.6) | $\times 10^{-7}$ | |
| Γ_{377} | $K^0 \nu \bar{\nu}$ | <i>B1</i> | < | 1.6 | $\times 10^{-4}$ | CL=90% |
| Γ_{378} | $\rho^0 \nu \bar{\nu}$ | <i>B1</i> | < | 4.4 | $\times 10^{-4}$ | CL=90% |

| | | | | |
|----------------|----------------------------|------|---|--------|
| Γ_{379} | $K^0 e^+ e^-$ | $B1$ | (-1.3 $^{+1.6}_{-1.1}$) $\times 10^{-7}$ | |
| Γ_{380} | $K^0 \mu^+ \mu^-$ | $B1$ | (-5.7 $^{+2.2}_{-1.8}$) $\times 10^{-7}$ | |
| Γ_{381} | $K^*(892)^0 \ell^+ \ell^-$ | $B1$ | [a] (-9.5 ± 1.8) $\times 10^{-7}$ | |
| Γ_{382} | $K^*(892)^0 e^+ e^-$ | $B1$ | (-1.04 $^{+0.35}_{-0.31}$) $\times 10^{-6}$ | |
| Γ_{383} | $K^*(892)^0 \mu^+ \mu^-$ | $B1$ | (-1.10 $^{+0.29}_{-0.26}$) $\times 10^{-6}$ | |
| Γ_{384} | $K^*(892)^0 \nu \bar{\nu}$ | $B1$ | < 3.4 $\times 10^{-4}$ | CL=90% |
| Γ_{385} | $\phi \nu \bar{\nu}$ | $B1$ | < 5.8 $\times 10^{-5}$ | CL=90% |
| Γ_{386} | $e^\pm \mu^\mp$ | LF | [g] < 9.2 $\times 10^{-8}$ | CL=90% |
| Γ_{387} | $\pi^0 e^\pm \mu^\mp$ | LF | < 1.4 $\times 10^{-7}$ | CL=90% |
| Γ_{388} | $K^0 e^\pm \mu^\mp$ | LF | < 2.7 $\times 10^{-7}$ | CL=90% |
| Γ_{389} | $K^*(892)^0 e^+ \mu^-$ | LF | < 5.3 $\times 10^{-7}$ | CL=90% |
| Γ_{390} | $K^*(892)^0 e^- \mu^+$ | LF | < 3.4 $\times 10^{-7}$ | CL=90% |
| Γ_{391} | $K^*(892)^0 e^\pm \mu^\mp$ | LF | < 5.8 $\times 10^{-7}$ | CL=90% |
| Γ_{392} | $e^\pm \tau^\mp$ | LF | [g] < 1.1 $\times 10^{-4}$ | CL=90% |
| Γ_{393} | $\mu^\pm \tau^\mp$ | LF | [g] < 3.8 $\times 10^{-5}$ | CL=90% |
| Γ_{394} | invisible | $B1$ | < 2.2 $\times 10^{-4}$ | CL=90% |
| Γ_{395} | $\nu \bar{\nu} \gamma$ | $B1$ | < 4.7 $\times 10^{-5}$ | CL=90% |

[a] An ℓ indicates an e or a μ mode, not a sum over these modes.

[b] \bar{D}^{**} represents an excited state with mass $2.2 < M < 2.8$ GeV/c².

[c] $X(3872)^+$ is a hypothetical charged partner of the $X(3872)$.

[d] Stands for the possible candidates of $K^*(1410)$, $K_0^*(1430)$ and $K_2^*(1430)$.

[e] B^0 and B_s^0 contributions not separated. Limit is on weighted average of the two decay rates.

[f] This decay refers to the coherent sum of resonant and nonresonant $J^P = 0^+$ $K\pi$ components with $1.60 < m_{K\pi} < 2.15$ GeV/c².

[g] The value is for the sum of the charge states or particle/antiparticle states indicated.

[h] $\Theta(1540)^+$ denotes a possible narrow pentaquark state.

B^0 BRANCHING RATIOS

For branching ratios in which the charge of the decaying B is not determined, see the B^\pm section.

$$\Gamma(\ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$$

$$\Gamma_1/\Gamma$$

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group

(HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

10.33±0.28 OUR EVALUATION

10.14±0.30 OUR AVERAGE

Error includes scale factor of 1.1.

| | | | |
|---|---------------------------|------|----------------------------------|
| $10.46 \pm 0.30 \pm 0.23$ | ¹ URQUIJO 07 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $9.64 \pm 0.27 \pm 0.33$ | ² AUBERT,B 06Y | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $10.78 \pm 0.60 \pm 0.69$ | ³ ARTUSO 97 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| $9.3 \pm 1.1 \pm 1.5$ | ALBRECHT 94 | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| $9.9 \pm 3.0 \pm 0.9$ | HENDERSON 92 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | |
| $10.32 \pm 0.36 \pm 0.35$ | ⁴ OKABE 05 | BELL | Repl. by URQUIJO 07 |
| $10.9 \pm 0.7 \pm 1.1$ | ATHANAS 94 | CLE2 | Sup. by ARTUSO 97 |

¹ URQUIJO 07 report a measurement of $(9.80 \pm 0.29 \pm 0.21)\%$ for the partial branching fraction of $B \rightarrow e\nu_e X_c$ decay with electron energy above 0.6 GeV. We converted the result to $B \rightarrow e\nu_e X$ branching fraction.

² The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame. The best precision on the ratio is achieved for a momentum threshold of 1.0 GeV: $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.074 \pm 0.041 \pm 0.026$.

³ ARTUSO 97 uses partial reconstruction of $B \rightarrow D^* \ell \nu_\ell$ and inclusive semileptonic branching ratio from BARISH 96B ($0.1049 \pm 0.0017 \pm 0.0043$).

⁴ The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame, and their ratio of $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.08 \pm 0.05 \pm 0.02$.

$\Gamma(e^+ \nu_e X_c)/\Gamma_{\text{total}}$

Γ_2/Γ

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------------------|------|----------------------------------|
| $10.08 \pm 0.30 \pm 0.22$ | ¹ URQUIJO 07 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Measure the independent B^+ and B^0 partial branching fractions with electron threshold energies of 0.4 GeV.

$\Gamma(D^- \ell^+ \nu_\ell)/\Gamma_{\text{total}}$

Γ_4/Γ

ℓ denotes e or μ , not the sum.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.0217±0.0012 OUR EVALUATION

0.0218±0.0012 OUR AVERAGE

| | | | |
|---|---------------------------|------|----------------------------------|
| $0.0221 \pm 0.0011 \pm 0.0012$ | ¹ AUBERT 08Q | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.0213 \pm 0.0012 \pm 0.0039$ | ABE 02E | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.0209 \pm 0.0013 \pm 0.0018$ | ² BARTEL 99 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.0235 \pm 0.0020 \pm 0.0044$ | ³ BUSKULIC 97 | ALEP | $e^+ e^- \rightarrow Z$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | |
| $0.0187 \pm 0.0015 \pm 0.0032$ | ⁴ ATHANAS 97 | CLE2 | Repl. by BARTEL 99 |
| $0.018 \pm 0.006 \pm 0.003$ | ⁵ FULTON 91 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.020 \pm 0.007 \pm 0.006$ | ⁶ ALBRECHT 89J | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses a fully reconstructed B meson as a tag on the recoil side.² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.³ BUSKULIC 97 assumes fraction (B^+) = fraction (B^0) = $(37.8 \pm 2.2)\%$ and PDG 96 values for B lifetime and branching ratio of D^* and D decays.⁴ ATHANAS 97 uses missing energy and missing momentum to reconstruct neutrino.⁵ FULTON 91 assumes assuming equal production of B^0 and B^+ at the $\gamma(4S)$ and uses Mark III D and D^* branching ratios.⁶ ALBRECHT 89J reports $0.018 \pm 0.006 \pm 0.005$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$. **$\Gamma(D^- \tau^+ \nu_\tau)/\Gamma_{\text{total}}$** **$\Gamma_5/\Gamma$**

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|----------|----------------------------------|
| $1.04 \pm 0.35 \pm 0.18$ | ¹ AUBERT | 08N BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses a fully reconstructed B meson as a tag on the recoil side. **$\Gamma(D^- \ell^+ \nu_\ell)/\Gamma(D \ell^+ \nu_\ell \text{anything})$** **$\Gamma_4/\Gamma_3$**

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|-----------|----------------------------------|
| $0.215 \pm 0.016 \pm 0.013$ | ¹ AUBERT | 07AN BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses a fully reconstructed B meson on the recoil side. **$\Gamma(D^*(2010)^- \ell^+ \nu_\ell)/\Gamma_{\text{total}}$** **$\Gamma_6/\Gamma$**

“OUR EVALUATION” is an average using rescaled values of the data listed below.

The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|---------|
| 0.0516 ± 0.0011 OUR EVALUATION | | | | |

 0.0523 ± 0.0022 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

| | | | | |
|---|-----|--------------------------|----------|----------------------------------|
| $0.0549 \pm 0.0016 \pm 0.0025$ | | ¹ AUBERT | 08Q BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.0469 \pm 0.0004 \pm 0.0034$ | | ² AUBERT | 08R BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.0590 \pm 0.0022 \pm 0.0050$ | | ³ ABDALLAH | 04D DLPH | $e^+ e^- \rightarrow Z^0$ |
| $0.0609 \pm 0.0019 \pm 0.0040$ | | ⁴ ADAM | 03 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.0459 \pm 0.0023 \pm 0.0040$ | | ⁵ ABE | 02F BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.0470 \pm 0.0013^{+0.0036}_{-0.0031}$ | | ⁶ ABREU | 01H DLPH | $e^+ e^- \rightarrow Z$ |
| $0.0526 \pm 0.0020 \pm 0.0046$ | | ⁷ ABBIENDI | 00Q OPAL | $e^+ e^- \rightarrow Z$ |
| $0.0553 \pm 0.0026 \pm 0.0052$ | | ⁸ BUSKULIC | 97 ALEP | $e^+ e^- \rightarrow Z$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $0.0490 \pm 0.0007^{+0.0036}_{-0.0035}$ | | ³ AUBERT | 05E BABR | Repl. by AUBERT 08R |
| $0.0539 \pm 0.0011 \pm 0.0034$ | | ⁹ ABDALLAH | 04D DLPH | $e^+ e^- \rightarrow Z^0$ |
| $0.0609 \pm 0.0019 \pm 0.0040$ | | ¹⁰ BRIERE | 02 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.0508 \pm 0.0021 \pm 0.0066$ | | ¹¹ ACKERSTAFF | 97G OPAL | Repl. by ABBIENDI 00Q |
| $0.0552 \pm 0.0017 \pm 0.0068$ | | ¹² ABREU | 96P DLPH | Repl. by ABREU 01H |
| $0.0449 \pm 0.0032 \pm 0.0039$ | 376 | ¹³ BARISH | 95 CLE2 | Repl. by ADAM 03 |
| $0.0518 \pm 0.0030 \pm 0.0062$ | 410 | ¹⁴ BUSKULIC | 95N ALEP | Sup. by BUSKULIC 97 |
| $0.045 \pm 0.003 \pm 0.004$ | | ¹⁵ ALBRECHT | 94 ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

| | | | | | |
|-----------------------|-----|---------------|-----|------|----------------------------------|
| 0.047 ± 0.005 ± 0.005 | 235 | 16 ALBRECHT | 93 | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| seen | 398 | 17 SANGHERA | 93 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.070 ± 0.018 ± 0.014 | | 18 ANTREASYAN | 90B | CBAL | $e^+ e^- \rightarrow \gamma(4S)$ |
| | | 19 ALBRECHT | 89C | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.060 ± 0.010 ± 0.014 | | 20 ALBRECHT | 89J | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.040 ± 0.004 ± 0.006 | | 21 BORTOLETTO | 89B | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.070 ± 0.012 ± 0.019 | 47 | 22 ALBRECHT | 87J | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² Measured using fully reconstructed D^* sample and a simultaneous fit to the Caprini-Lellouch-Neubert form factor parameters: $\rho^2 = 1.191 \pm 0.048 \pm 0.028$, $R_1(1) = 1.429 \pm 0.061 \pm 0.044$, and $R_2(1) = 0.827 \pm 0.038 \pm 0.022$.

³ Measured using fully reconstructed D^* sample.

⁴ Uses the combined fit of both $B^0 \rightarrow D^*(2010)^- \ell \nu$ and $B^+ \rightarrow \bar{D}(2007)^0 \ell \nu$ samples.

⁵ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

⁶ ABREU 01H measured using about 5000 partial reconstructed D^* sample.

⁷ ABBIENDI 00Q assumes the fraction $B(b \rightarrow B^0) = (39.7^{+1.8}_{-2.2})\%$. This result is an average of two methods using exclusive and partial D^* reconstruction.

⁸ BUSKULIC 97 assumes fraction (B^+) = fraction (B^0) = $(37.8 \pm 2.2)\%$ and PDG 96 values for B lifetime and D^* and D branching fractions.

⁹ Combines with previous partial reconstructed D^* measurement.

¹⁰ The results are based on the same analysis and data sample reported in ADAM 03.

¹¹ ACKERSTAFF 97G assumes fraction (B^+) = fraction (B^0) = $(37.8 \pm 2.2)\%$ and PDG 96 values for B lifetime and branching ratio of D^* and D decays.

¹² ABREU 96P result is the average of two methods using exclusive and partial D^* reconstruction.

¹³ BARISH 95 use $B(D^0 \rightarrow K^- \pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*+} \rightarrow D^0 \pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$.

¹⁴ BUSKULIC 95N assumes fraction (B^+) = fraction (B^0) = $38.2 \pm 1.3 \pm 2.2\%$ and $\tau_{B^0} = 1.58 \pm 0.06$ ps. $\Gamma(D^{*-} \ell^+ \nu_\ell)/\text{total} = [5.18 - 0.13(\text{fraction}(B^0) - 38.2) - 1.5(\tau_{B^0} - 1.58)]\%$.

¹⁵ ALBRECHT 94 assumes $B(D^{*+} \rightarrow D^0 \pi^+) = 68.1 \pm 1.0 \pm 1.3\%$. Uses partial reconstruction of D^{*+} and is independent of D^0 branching ratios.

¹⁶ ALBRECHT 93 reports $0.052 \pm 0.005 \pm 0.006$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$. We have taken their average e and μ value. They also obtain $\alpha = 2*\Gamma^0/(\Gamma^- + \Gamma^+) - 1 = 1.1 \pm 0.4 \pm 0.2$, $A_{AF} = 3/4*(\Gamma^- - \Gamma^+)/\Gamma = 0.2 \pm 0.08 \pm 0.06$ and a value of $|V_{cb}| = 0.036 - 0.045$ depending on model assumptions.

¹⁷ Combining $\bar{D}^{*0} \ell^+ \nu_\ell$ and $\bar{D}^{*-} \ell^+ \nu_\ell$ SANGHERA 93 test $V-A$ structure and fit the decay angular distributions to obtain $A_{FB} = 3/4*(\Gamma^- - \Gamma^+)/\Gamma = 0.14 \pm 0.06 \pm 0.03$. Assuming a value of V_{cb} , they measure V , A_1 , and A_2 , the three form factors for the $D^* \ell \nu_\ell$ decay, where results are slightly dependent on model assumptions.


¹⁸ ANTREASYAN 90B is average over B and $\bar{D}^*(2010)$ charge states.

¹⁹ The measurement of ALBRECHT 89C suggests a D^* polarization γ_L/γ_T of 0.85 ± 0.45 , or $\alpha = 0.7 \pm 0.9$.

²⁰ ALBRECHT 89J is ALBRECHT 87J value rescaled using $B(D^*(2010)^- \rightarrow D^0 \pi^-) = 0.57 \pm 0.04 \pm 0.04$. Superseded by ALBRECHT 93.

²¹ We have taken average of the the BORTOLETTO 89B values for electrons and muons, $0.046 \pm 0.005 \pm 0.007$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$. The measurement suggests a D^* polarization parameter value $\alpha = 0.65 \pm 0.66 \pm 0.25$.

²² ALBRECHT 87J assume μ - e universality, the $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.45$, the $B(D^0 \rightarrow K^- \pi^+) = (0.042 \pm 0.004 \pm 0.004)$, and the $B(D^*(2010)^- \rightarrow D^0 \pi^-) = 0.49 \pm 0.08$. Superseded by ALBRECHT 89J.



$\Gamma(D^*(2010)^- \ell^+ \nu_\ell) / \Gamma(D \ell^+ \nu_\ell \text{ anything})$

| VALUE | DOCUMENT ID | TECN | COMMENT | Γ_6/Γ_3 |
|--------------------------|---------------------|-----------|------------------------------------|---------------------|
| 0.537±0.031±0.036 | ¹ AUBERT | 07AN BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ | |

¹ Uses a fully reconstructed B meson on the recoil side.

$\Gamma(D^*(2010)^- \tau^+ \nu_\tau) / \Gamma_{\text{total}}$

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT | Γ_7/Γ |
|------------------------------|-------------------------------------|----------|------------------------------------|-------------------|
| 1.6 ± 0.5 OUR AVERAGE | Error includes scale factor of 1.2. | | | |
| 1.11±0.51±0.06 | ¹ AUBERT | 08N BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ | |

$2.02^{+0.40}_{-0.37} \pm 0.37$ ² MATYJA 07 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² Observed in the recoil of the accompanying B meson.

$\Gamma(D^0 \pi^- \ell^+ \nu_\ell) / \Gamma_{\text{total}}$

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT | Γ_8/Γ |
|---|--------------------------|----------|------------------------------------|-------------------|
| 4.3±0.6 OUR AVERAGE | | | | |
| 4.3±0.8±0.3 | ¹ AUBERT | 08Q BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ | |
| 4.3±0.9±0.2 | ^{1,2} LIVENTSEV | 08 BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 3.4±1.0±0.2 | ³ LIVENTSEV | 05 BELL | Repl. by LIVENTSEV 08 | |

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² LIVENTSEV 08 reports $(4.2 \pm 0.7 \pm 0.6) \times 10^{-3}$ for $B(B^0 \rightarrow D^- \ell^+ \nu_\ell) = (2.12 \pm 0.20) \times 10^{-2}$. We rescale to our best value $B(B^0 \rightarrow D^- \ell^+ \nu_\ell) = (2.17 \pm 0.12) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ LIVENTSEV 05 reports $[B(B^0 \rightarrow \bar{D}^0 \pi^- \ell^+ \nu_\ell)] / [B(B^+ \rightarrow \bar{D}^0 \ell^+ \nu_\ell)] = 0.15 \pm 0.03 \pm 0.03$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \ell^+ \nu_\ell) = (2.27 \pm 0.11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_0^*(2400)^- \ell^+ \nu_\ell \times B(D_0^{*-} \rightarrow \bar{D}^0 \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_9 / \Gamma$$

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------------|------|----------------------------------|
| $2.0 \pm 0.7 \pm 0.5$ | ¹ LIVENTSEV 08 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$$\Gamma(D_2^*(2460)^- \ell^+ \nu_\ell \times B(D_2^{*-} \rightarrow \bar{D}^0 \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{10} / \Gamma$$

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------------|------|----------------------------------|
| $2.2 \pm 0.4 \pm 0.4$ | ¹ LIVENTSEV 08 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$$\Gamma(\bar{D}^{*0} \pi^- \ell^+ \nu_\ell) / \Gamma_{\text{total}} \quad \Gamma_{12} / \Gamma$$

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---------|
| $4.9 \pm 0.8 \text{ OUR AVERAGE}$ | | | |

$4.8 \pm 0.8 \pm 0.4$

¹ AUBERT 08Q BABR $e^+ e^- \rightarrow \gamma(4S)$

$5.7 \pm 2.2 \pm 0.3$

^{1,2} LIVENTSEV 08 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.1 \pm 1.4 \pm 0.3$

^{3,4} LIVENTSEV 05 BELL Repl. by LIVENTSEV 08

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² LIVENTSEV 08 reports $(5.6 \pm 2.1 \pm 0.8) \times 10^{-3}$ for $B(B^0 \rightarrow D^- \ell^+ \nu_\ell) = (2.12 \pm 0.20) \times 10^{-2}$. We rescale to our best value $B(B^0 \rightarrow D^- \ell^+ \nu_\ell) = (2.17 \pm 0.12) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Excludes D^{*+} contribution to $D\pi$ modes.

⁴ LIVENTSEV 05 reports $[B(B^0 \rightarrow \bar{D}^{*0} \pi^- \ell^+ \nu_\ell)] / [B(B^+ \rightarrow \bar{D}^*(2007)^0 \ell^+ \nu_\ell)] = 0.10 \pm 0.02 \pm 0.01$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^*(2007)^0 \ell^+ \nu_\ell) = (6.07 \pm 0.29) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_1(2420)^- \ell^+ \nu_\ell \times B(D_1^- \rightarrow \bar{D}^{*0} \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{13} / \Gamma$$

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------------|------|----------------------------------|
| $5.4 \pm 1.9 \pm 0.9$ | ¹ LIVENTSEV 08 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$$\Gamma(D'_1(2430)^- \ell^+ \nu_\ell \times B(D'_1^- \rightarrow \bar{D}^{*0} \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{14} / \Gamma$$

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------------|------|----------------------------------|
| <5.0 | 90 | ¹ LIVENTSEV 08 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$\Gamma(D_2^*(2460)^-\ell^+\nu_\ell \times B(D_2^{*-} \rightarrow \bar{D}^{*0}\pi^-))/\Gamma_{\text{total}}$ Γ_{15}/Γ

| <u>VALUE</u> (units 10^{-3}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|------------------------|-------------|--------------------------------------|
| <3.0 | 90 | ¹ LIVENTSEV | 08 | BELL $e^+e^- \rightarrow \gamma(4S)$ |

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

 $\Gamma(\bar{D}^{(*)}n\pi\ell^+\nu_\ell(n \geq 1))/\Gamma(D\ell^+\nu_\ell \text{anything})$ Γ_{11}/Γ_3

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|---------------------|-------------|---------------------------------|
| 0.248±0.032±0.030 | ¹ AUBERT | 07AN BABR | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Uses a fully reconstructed B meson on the recoil side.

 $\Gamma(\rho^-\ell^+\nu_\ell)/\Gamma_{\text{total}}$ Γ_{16}/Γ

$\ell = e$ or μ , not sum over e and μ modes.

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
| 2.47±0.33 OUR AVERAGE | | | | |

| | | | | |
|--|----|------------------------|-----|--------------------------------------|
| 2.93±0.37±0.37 | | ¹ ADAM | 07 | CLE2 $e^+e^- \rightarrow \gamma(4S)$ |
| 2.17±0.54±0.32 | | ² HOKUUE | 07 | BELL $e^+e^- \rightarrow \gamma(4S)$ |
| 2.14±0.21±0.56 | | ³ AUBERT,B | 050 | BABR $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2.17±0.34 ^{+0.62} _{-0.68} | | ⁴ ATHAR | 03 | CLE2 Repl. by ADAM 07 |
| 3.29±0.42±0.72 | | ⁵ AUBERT | 03E | BABR Repl. by AUBERT,B 050 |
| 2.57±0.29 ^{+0.53} _{-0.62} | | ⁶ BEHRENS | 00 | CLE2 Repl. by ADAM 07 |
| 2.69±0.41 ^{+0.61} _{-0.64} | | ⁷ BEHRENS | 00 | CLE2 $e^+e^- \rightarrow \gamma(4S)$ |
| 2.5 ± 0.4 ^{+0.7} _{-0.9} | | ⁸ ALEXANDER | 96T | CLE2 Repl. by BEHRENS 00 |
| <4.1 | 90 | ⁹ BEAN | 93B | CLE2 $e^+e^- \rightarrow \gamma(4S)$ |

¹ The B^0 and B^+ results are combined assuming the isospin, B lifetimes, and relative charged/neutral B production at the $\gamma(4S)$.

² The signal events are tagged by a second B meson reconstructed in the semileptonic mode $B \rightarrow D^{(*)}\ell\nu_\ell$.

³ B^+ and B^0 decays combined assuming isospin symmetry. Systematic errors include both experimental and form-factor uncertainties.

⁴ ATHAR 03 reports systematic errors $+0.47$ _{-0.50} $\pm 0.41 \pm 0.01$, which are experimental systematic, systematic due to residual form-factor uncertainties in the signal, and systematic due to residual form-factor uncertainties in the cross-feed modes, respectively. We combine these in quadrature.

⁵ Uses isospin constraints and extrapolation to all electron energies according to five different form-factor calculations. The second error combines the systematic and theoretical uncertainties in quadrature.

⁶ Averaging with ALEXANDER 96T results including experimental and theoretical correlations considered, BEHRENS 00 reports systematic errors $+0.33$ _{-0.46} ± 0.41 , where the second error is theoretical model dependence. We combine these in quadrature.

⁷ BEHRENS 00 reports $+0.35$ _{-0.40} ± 0.50 , where the second error is the theoretical model dependence. We combine these in quadrature. B^+ and B^0 decays combined using isospin symmetry: $\Gamma(B^0 \rightarrow \rho^-\ell^+\nu) = 2\Gamma(B^+ \rightarrow \rho^0\ell^+\nu) \approx 2\Gamma(B^+ \rightarrow \omega\ell^+\nu)$. No evidence for $\omega\ell\nu$ is reported.

⁸ ALEXANDER 96T reports $+0.5$ _{-0.7} ± 0.5 where the second error is the theoretical model dependence. We combine these in quadrature. B^+ and B^0 decays combined using

isospin symmetry: $\Gamma(B^0 \rightarrow \rho^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \rho^0 \ell^+ \nu) \approx 2\Gamma(B^+ \rightarrow \omega \ell^+ \nu)$. No evidence for $\omega \ell \nu$ is reported.

⁹ BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\rho^0 \ell^+ \nu_\ell)$ and $\Gamma(\omega \ell^+ \nu_\ell)$ with this result, they obtain a limit $<(1.6\text{--}2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow (\omega \text{or } \rho^0) \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.08\text{--}0.13$ at 90% CL is derived as well.

$\Gamma(\pi^- \ell^+ \nu_\ell)/\Gamma_{\text{total}}$

Γ_{17}/Γ

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|------|---------------------------------------|
| $1.36 \pm 0.06 \pm 0.07$ OUR EVALUATION | | | |
| 1.41 ± 0.08 OUR AVERAGE | | | |
| $1.37 \pm 0.15 \pm 0.11$ | ^{1,2} ADAM | 07 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.46 \pm 0.07 \pm 0.08$ | ³ AUBERT | 07J | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.38 \pm 0.19 \pm 0.14$ | ⁴ HOKUE | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.33 \pm 0.17 \pm 0.11$ | ⁵ AUBERT,B | 06K | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $1.38 \pm 0.10 \pm 0.18$ | ⁶ AUBERT,B | 050 | BABR Repl. by AUBERT,B 06K |
| $1.33 \pm 0.18 \pm 0.13$ | ⁷ ATHAR | 03 | CLE2 Repl. by ADAM 07 |
| $1.8 \pm 0.4 \pm 0.4$ | ⁸ ALEXANDER | 96T | CLE2 Repl. by ATHAR 03 |

¹ The B^0 and B^+ results are combined assuming the isospin, B lifetimes, and relative charged/neutral B production at the $\gamma(4S)$.

² Also report the rate for $q^2 > 16 \text{ GeV}^2$ of $(0.41 \pm 0.08 \pm 0.04) \times 10^{-4}$ from which they obtain $|V_{ub}| = 3.6 \pm 0.4 \pm 0.2^{+0.6}_{-0.4}$ (last error is from theory).

³ The analysis uses events in which the signal B decays are reconstructed with an innovative loose neutrino reconstruction technique.

⁴ The signal events are tagged by a second B meson reconstructed in the semileptonic mode $B \rightarrow D^{(*)} \ell \nu_\ell$.

⁵ The signals are tagged by a second B meson reconstructed in a semileptonic or hadronic decay. The B^0 and B^+ results are combined assuming the isospin symmetry.

⁶ B^+ and B^0 decays combined assuming isospin symmetry. Systematic errors include both experimental and form-factor uncertainties.

⁷ ATHAR 03 reports systematic errors $0.11 \pm 0.01 \pm 0.07$, which are experimental systematic, systematic due to residual form-factor uncertainties in the signal, and systematic due to residual form-factor uncertainties in the cross-feed modes, respectively. We combine these in quadrature.

⁸ ALEXANDER 96T gives systematic errors $\pm 0.3 \pm 0.2$ where the second error reflects the estimated model dependence. We combine these in quadrature. Assumes isospin symmetry: $\Gamma(B^0 \rightarrow \pi^- \ell^+ \nu) = 2 \times \Gamma(B^+ \rightarrow \pi^0 \ell^+ \nu)$.

$\Gamma(\pi^- \mu^+ \nu_\mu)/\Gamma_{\text{total}}$

Γ_{18}/Γ

| VALUE | DOCUMENT ID | TECN |
|-------|-------------|------|
|-------|-------------|------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ¹ ALBRECHT 91C ARG

¹ In ALBRECHT 91C, one event is fully reconstructed providing evidence for the $b \rightarrow u$ transition.

$\Gamma(K^\pm \text{anything})/\Gamma_{\text{total}}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.78±0.08¹ ALBRECHT 96D ARG $e^+ e^- \rightarrow \gamma(4S)$ ¹ Average multiplicity. Γ_{19}/Γ $\Gamma(D^0 X)/\Gamma_{\text{total}}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.081±0.014±0.005¹ AUBERT 07N BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.063 \pm 0.019 \pm 0.005$ ¹ AUBERT,BE 04B BABR Repl. by AUBERT 07N¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties. Γ_{20}/Γ $\Gamma(\bar{D}^0 X)/\Gamma_{\text{total}}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.474±0.020^{+0.020}_{-0.019}¹ AUBERT 07N BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.511 \pm 0.031 \pm 0.028$ ¹ AUBERT,BE 04B BABR Repl. by AUBERT 07N¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties. Γ_{21}/Γ $\Gamma(D^0 X)/[\Gamma(D^0 X) + \Gamma(\bar{D}^0 X)]$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.146±0.022±0.006¹ AUBERT 07N BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.110 \pm 0.031 \pm 0.008$

AUBERT,BE 04B BABR Repl. by AUBERT 07N

 $\Gamma_{20}/(\Gamma_{20}+\Gamma_{21})$ $\Gamma(D^+ X)/\Gamma_{\text{total}}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

<0.039

¹ AUBERT 07N BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.051

¹ AUBERT,BE 04B BABR Repl. by AUBERT 07N¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties. Γ_{22}/Γ $\Gamma(D^- X)/\Gamma_{\text{total}}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.369±0.016^{+0.030}_{-0.027}¹ AUBERT 07N BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.397 \pm 0.030 \pm 0.040$ ¹ AUBERT,BE 04B BABR Repl. by AUBERT 07N Γ_{23}/Γ ¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D^+ X)/[\Gamma(D^+ X) + \Gamma(D^- X)]$ $\Gamma_{22}/(\Gamma_{22} + \Gamma_{23})$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|--------------------|-------------|----------------------------------|--|
| 0.058±0.028±0.006 | AUBERT | 07N BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.055±0.040±0.006 | AUBERT,BE | 04B BABR | Repl. by AUBERT 07N | |

 $\Gamma(D_s^+ X)/\Gamma_{\text{total}}$ Γ_{24}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|------------------------|-------------|----------------------------------|--|
| 0.103±0.012^{+0.017}_{-0.014} | ¹ AUBERT | 07N BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.109±0.021 ^{+0.039} _{-0.024} | ¹ AUBERT,BE | 04B BABR | Repl. by AUBERT 07N | |

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(D_s^- X)/\Gamma_{\text{total}}$ Γ_{25}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|------------|------------------------|-------------|----------------------------------|--|
| <0.026 | 90 | ¹ AUBERT | 07N BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <0.087 | 90 | ¹ AUBERT,BE | 04B BABR | Repl. by AUBERT 07N | |

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(D_s^+ X)/[\Gamma(D_s^+ X) + \Gamma(D_s^- X)]$ $\Gamma_{24}/(\Gamma_{24} + \Gamma_{25})$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|--------------------|-------------|----------------------------------|--|
| 0.879±0.066±0.005 | AUBERT | 07N BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.733±0.092±0.010 | AUBERT,BE | 04B BABR | Repl. by AUBERT 07N | |

 $\Gamma(\Lambda_c^+ X)/\Gamma_{\text{total}}$ Γ_{26}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|------------|------------------------|-------------|----------------------------------|--|
| <0.031 | 90 | ¹ AUBERT | 07N BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <0.038 | 90 | ¹ AUBERT,BE | 04B BABR | Repl. by AUBERT 07N | |

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(\bar{\Lambda}_c^- X)/\Gamma_{\text{total}}$ Γ_{27}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|------------------------|-------------|----------------------------------|--|
| 0.05 ±0.010^{+0.019}_{-0.011} | ¹ AUBERT | 07N BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.049±0.017 ^{+0.018} _{-0.011} | ¹ AUBERT,BE | 04B BABR | Repl. by AUBERT 07N | |

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\Lambda_c^+ X)/[\Gamma(\Lambda_c^+ X) + \Gamma(\bar{\Lambda}_c^- X)]$ VALUE**0.243^{+0.119}_{-0.121} ± 0.003**DOCUMENT ID

AUBERT

TECN

07N

COMMENTBABR $e^+ e^- \rightarrow \gamma(4S)$ $\Gamma_{26}/(\Gamma_{26}+\Gamma_{27})$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.286 ± 0.142 ± 0.007

AUBERT,BE 04B BABR Repl. by AUBERT 07N

 $\Gamma(\bar{c}X)/\Gamma_{\text{total}}$ VALUE**0.947^{+0.030}_{-0.040} ± 0.045**DOCUMENT ID¹ AUBERTTECN

07N

COMMENTBABR $e^+ e^- \rightarrow \gamma(4S)$ Γ_{28}/Γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.039 ± 0.051^{+0.063}_{-0.058}¹ AUBERT,BE 04B BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(cX)/\Gamma_{\text{total}}$ VALUE**0.246^{+0.024}_{-0.017} ± 0.021**DOCUMENT ID¹ AUBERTTECN

07N

COMMENTBABR $e^+ e^- \rightarrow \gamma(4S)$ Γ_{29}/Γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.237 ± 0.036^{+0.041}_{-0.027}¹ AUBERT,BE 04B BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(\bar{c}cX)/\Gamma_{\text{total}}$ VALUE**1.193^{+0.030}_{-0.049} ± 0.053**DOCUMENT ID¹ AUBERTTECN

07N

COMMENTBABR $e^+ e^- \rightarrow \gamma(4S)$ Γ_{30}/Γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.276 ± 0.062^{+0.088}_{-0.074}¹ AUBERT,BE 04B BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(D^- \pi^+)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})**2.68^{+0.13}_{-0.13} OUR AVERAGE**EVTSDOCUMENT IDTECNCOMMENT Γ_{31}/Γ

2.55 ± 0.05 ± 0.16

¹ AUBERT 07H BABR $e^+ e^- \rightarrow \gamma(4S)$

3.03 ± 0.23 ± 0.23

² AUBERT,BE 06J BABR $e^+ e^- \rightarrow \gamma(4S)$

2.68 ± 0.12 ± 0.24

^{1,3} AHMED 02B CLE2 $e^+ e^- \rightarrow \gamma(4S)$

2.7 ± 0.6 ± 0.5

⁴ BORTOLETTO92 CLEO $e^+ e^- \rightarrow \gamma(4S)$

4.8 ± 1.1 ± 1.1

⁵ ALBRECHT 90J ARG $e^+ e^- \rightarrow \gamma(4S)$

5.1 ± 2.8 ± 1.3

⁶ BEBEK 87 CLEO $e^+ e^- \rightarrow \gamma(4S)$

5.1 -2.5 -1.2

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------------------|----------------|-----------------------|-----|------|------------------------------------|
| $2.94 \pm 0.21 \pm 0.09$ | ^{1,7} | AUBERT,B | 040 | BABR | Repl. by AUBERT 07H |
| $2.9 \pm 0.4 \pm 0.1$ | ⁸ | ALAM | 94 | CLE2 | Repl. by AHMED 02B |
| $3.1 \pm 1.3 \pm 1.0$ | ⁷ | ⁵ ALBRECHT | 88K | ARG | $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.

³ AHMED 02B reports an additional uncertainty on the branching ratios to account for 4.5% uncertainty on relative production of B^0 and B^+ , which is not included here.

⁴ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁵ ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ production ratio is 45:55. Superseded by ALBRECHT 90J which assumes 50:50.

⁶ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

⁷ AUBERT,B 040 reports $[B(B^0 \rightarrow D^- \pi^+)] \times [B(D^+ \rightarrow K_S^0 \pi^+)] = (42.7 \pm 2.1 \pm 2.2) \times 10^{-6}$. We divide by our best value $B(D^+ \rightarrow K_S^0 \pi^+) = (1.45 \pm 0.04) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁸ ALAM 94 reports $[B(B^0 \rightarrow D^- \pi^+)] \times [B(D^+ \rightarrow K^- \pi^+ \pi^+)] = (0.265 \pm 0.032 \pm 0.023) \times 10^{-3}$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.22 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- \rho^+)/\Gamma_{\text{total}}$

Γ_{32}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|------|-----------------------|------|---|
| 0.0077 ± 0.0013 OUR AVERAGE | | | | |
| $0.0076 \pm 0.0013 \pm 0.0002$ | 79 | ¹ ALAM | 94 | CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $0.009 \pm 0.005 \pm 0.003$ | 9 | ² ALBRECHT | 90J | ARG $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------------|---|-----------------------|-----|--|
| $0.022 \pm 0.012 \pm 0.009$ | 6 | ² ALBRECHT | 88K | ARG $e^+ e^- \rightarrow \Upsilon(4S)$ |
|-----------------------------|---|-----------------------|-----|--|

¹ ALAM 94 reports $[B(B^0 \rightarrow D^- \rho^+)] \times [B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.000704 \pm 0.000096 \pm 0.000070$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.22 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ production ratio is 45:55. Superseded by ALBRECHT 90J which assumes 50:50.

$\Gamma(D^- K^0 \pi^+)/\Gamma_{\text{total}}$

Γ_{33}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|------|---|
| $4.9 \pm 0.7 \pm 0.5$ | ¹ AUBERT,BE | 05B | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- K^*(892)^+)/\Gamma_{\text{total}}$

Γ_{34}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|------|---|
| $4.5 \pm 0.7 \pm 0.5$ OUR AVERAGE | ¹ AUBERT,BE | 05B | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $4.6 \pm 0.6 \pm 0.5$ | ¹ MAHAPATRA | 02 | CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^-\omega\pi^+)/\Gamma_{\text{total}}$ Γ_{35}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|----------------------------|------|---------------------------------|
| 0.0028±0.0005±0.0004 | ¹ ALEXANDER 01B | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. The signal is consistent with all observed $\omega\pi^+$ having proceeded through the $\rho'/^+$ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

 $\Gamma(D^-\bar{K}^+)/\Gamma_{\text{total}}$ Γ_{36}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|------|---------------------------------|
| (2.04±0.50±0.27) × 10⁻⁴ | ¹ ABE 01I | BELL | $e^+e^- \rightarrow \gamma(4S)$ |

¹ ABE 01I reports $B(B^0 \rightarrow D^-\bar{K}^+)/B(B^0 \rightarrow D^-\pi^+) = 0.068 \pm 0.015 \pm 0.007$. We multiply by our best value $B(B^0 \rightarrow D^-\pi^+) = (3.0 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and the second error is systematic error from using our best value.

 $\Gamma(D^-\bar{K}^+\bar{K}^0)/\Gamma_{\text{total}}$ Γ_{37}/Γ

| VALUE (units 10 ⁻⁴) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-----|--------------------------|------|---------------------------------|
| <3.1 | 90 | ¹ DRUTSKOY 02 | BELL | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^-\bar{K}^+(\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{38}/Γ

| VALUE (units 10 ⁻⁴) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|--------------------------|------|---------------------------------|
| 8.8±1.1±1.5 | ¹ DRUTSKOY 02 | BELL | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{39}/Γ

| VALUE (units 10 ⁻⁴) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-----|------|------------------------|------|---------------------------------|
| 8.4±0.4±0.8 | | | ¹ KUZMIN 07 | BELL | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------|----|---------------------------|------|---------------------------------|
| 8.0±0.6±1.5 | | 1,2 SATPATHY 03 | BELL | Repl. by KUZMIN 07 |
| < 16 | 90 | ¹ ALAM 94 | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| < 70 | 90 | ³ BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |
| <340 | 90 | ⁴ BEBEK 87 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |
| 700 ± 500 | 5 | ⁵ BEHRENDS 83 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² No assumption about the intermediate mechanism is made in the analysis.

³ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D . The product branching fraction into $D_0^*(2340)\pi$ followed by $D_0^*(2340) \rightarrow D^0\pi$ is < 0.0001 at 90% CL and into $D_2^*(2460)$ followed by $D_2^*(2460) \rightarrow D^0\pi$ is < 0.0004 at 90% CL.

⁴ BEBEK 87 assume the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%. $B(D^0 \rightarrow K^-\pi^+) = (4.2 \pm 0.4 \pm 0.4)\%$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-) = (9.1 \pm 0.8 \pm 0.8)\%$ were used.

⁵ Corrected by us using assumptions: $B(D^0 \rightarrow K^-\pi^+) = (0.042 \pm 0.006)$ and $B(\gamma(4S) \rightarrow B^0\bar{B}^0) = 50\%$. The product branching ratio is $B(B^0 \rightarrow \bar{D}^0\pi^+\pi^-)B(\bar{D}^0 \rightarrow K^+\pi^-) = (0.39 \pm 0.26) \times 10^{-2}$.

| $\Gamma(D^*(2010)^-\pi^+)/\Gamma_{\text{total}}$ | Γ_{40}/Γ | | | |
|--|----------------------|-------------------|------|---------------------------------|
| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
| 2.76±0.13 OUR AVERAGE | | | | |
| 2.79±0.08±0.17 | | 1 AUBERT 07H | BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| 2.7 ± 0.4 ± 0.1 | | 2,3 AUBERT,BE 06J | BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| 2.81±0.24±0.05 | | 4 BRANDENB... 98 | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| 2.6 ± 0.3 ± 0.4 | 82 | 5 ALAM 94 | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| 3.37±0.96±0.02 | | 6 BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |
| 2.36±0.88±0.02 | 12 | 7 ALBRECHT 90J | ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| 2.36 ^{+1.50} _{-1.10} ±0.02 | 5 | 8 BEBEK 87 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------|----|-----------------|------|---------------------------------|
| 10 ± 4 ± 1 | 8 | 9 AKERS 94J | OPAL | $e^+e^- \rightarrow Z$ |
| 2.7 ± 1.4 ± 1.0 | 5 | 10 ALBRECHT 87C | ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| 3.5 ± 2 ± 2 | | 11 ALBRECHT 86F | ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| 17 ± 5 ± 5 | 41 | 12 GILES 84 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² AUBERT,BE 06J reports $[B(B^0 \rightarrow D^*(2010)^-\pi^+)] / [B(B^0 \rightarrow D^-\pi^+)] = 0.99 \pm 0.11 \pm 0.08$. We multiply by our best value $B(B^0 \rightarrow D^-\pi^+) = (2.68 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.

⁴ BRANDENBURG 98 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the D^* reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of $B(D^* \rightarrow D\pi)$.

⁵ ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0\pi^+)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

⁶ BORTOLETTO 92 reports $(4.0 \pm 1.0 \pm 0.7) \times 10^{-3}$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

⁷ ALBRECHT 90J reports $(2.8 \pm 0.9 \pm 0.6) \times 10^{-3}$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

⁸ BEBEK 87 reports $(2.8^{+1.5}_{-1.2}{}^{+1.0}_{-0.6}) \times 10^{-3}$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$.

We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92 and ALBRECHT 90J.

⁹ Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and 38% B_d production fraction.

¹⁰ ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\gamma(4S) \rightarrow B^+B^-) = 55\%$ and $B(\gamma(4S) \rightarrow B^0\bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

¹¹ ALBRECHT 86F uses pseudomass that is independent of D^0 and D^+ branching ratios.

¹² Assumes $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.60^{+0.08}_{-0.15}$. Assumes $B(\gamma(4S) \rightarrow B^0\bar{B}^0) = 0.40 \pm 0.02$ Does not depend on D branching ratios.

$\Gamma(D^-\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{41}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------|---------------------------|-------------|---------------------------------|
| 0.0080±0.0021±0.0014 | ¹ BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

 $\Gamma((D^-\pi^+\pi^+\pi^-)\text{nonresonant})/\Gamma_{\text{total}}$ Γ_{42}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------|---------------------------|-------------|---------------------------------|
| 0.0039±0.0014±0.0013 | ¹ BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

 $\Gamma(D^-\pi^+\rho^0)/\Gamma_{\text{total}}$ Γ_{43}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------|---------------------------|-------------|---------------------------------|
| 0.0011±0.0009±0.0004 | ¹ BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

 $\Gamma(D^-\alpha_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{44}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------|---------------------------|-------------|---------------------------------|
| 0.0060±0.0022±0.0024 | ¹ BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

 $\Gamma(D^*(2010)^-\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{45}/Γ

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------|-------------|-----------------------|-------------|---------------------------------|
| 0.0152±0.0052±0.0001 | 51 | ¹ ALBRECHT | 90J ARG | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|---|-----------------------|---------|---------------------------------|
| 0.015 ± 0.008 ± 0.008 | 8 | ² ALBRECHT | 87C ARG | $e^+e^- \rightarrow \gamma(4S)$ |
|-----------------------|---|-----------------------|---------|---------------------------------|

¹ ALBRECHT 90J reports $0.018 \pm 0.004 \pm 0.005$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

² ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\gamma(4S) \rightarrow B^+B^-) = 55\%$ and $B(\gamma(4S) \rightarrow B^0\bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

 $\Gamma(D^*(2010)^-\rho^+)/\Gamma_{\text{total}}$ Γ_{46}/Γ

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------------|-------------|--------------------|-------------|----------------|
| 0.0068 ± 0.0009 OUR AVERAGE | | | | |

| | | | | |
|--------------------------|--|---------------------|----|--------------------------------------|
| 0.0068 ± 0.0003 ± 0.0009 | | ¹ CSORNA | 03 | CLE2 $e^+e^- \rightarrow \gamma(4S)$ |
|--------------------------|--|---------------------|----|--------------------------------------|

| | | | | |
|--------------------------|--|---------------------------|------|---------------------------------|
| 0.0160 ± 0.0113 ± 0.0001 | | ² BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |
|--------------------------|--|---------------------------|------|---------------------------------|

| | | | | |
|-----------------------------|----|-----------------------|---------|---------------------------------|
| 0.00589 ± 0.00352 ± 0.00004 | 19 | ³ ALBRECHT | 90J ARG | $e^+e^- \rightarrow \gamma(4S)$ |
|-----------------------------|----|-----------------------|---------|---------------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------------|----|---------------------|----|--------------------------------------|
| 0.0074 ± 0.0010 ± 0.0014 | 76 | ^{4,5} ALAM | 94 | CLE2 $e^+e^- \rightarrow \gamma(4S)$ |
|--------------------------|----|---------------------|----|--------------------------------------|

| | | | | |
|-----------------------------|----|-------------------|----|--------------------------------------|
| 0.081 ± 0.029 +0.059 -0.024 | 19 | ⁶ CHEN | 85 | CLEO $e^+e^- \rightarrow \gamma(4S)$ |
|-----------------------------|----|-------------------|----|--------------------------------------|

- ¹ Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$ resonance. The second error combines the systematic and theoretical uncertainties in quadrature. CSORNA 03 includes data used in ALAM 94. A full angular fit to three complex helicity amplitudes is performed.
- ² BORTOLETTO 92 reports $0.019 \pm 0.008 \pm 0.011$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .
- ³ ALBRECHT 90J reports $0.007 \pm 0.003 \pm 0.003$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .
- ⁴ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0\pi^+)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.
- ⁵ This decay is nearly completely longitudinally polarized, $\Gamma_L/\Gamma = (93 \pm 5 \pm 5)\%$, as expected from the factorization hypothesis (ROSNER 90). The nonresonant $\pi^+\pi^0$ contribution under the ρ^+ is less than 9% at 90% CL.
- ⁶ Uses $B(D^* \rightarrow D^0\pi^+) = 0.6 \pm 0.15$ and $B(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 0.4$. Does not depend on D branching ratios.

 $\Gamma(D^*(2010)^- K^+)/\Gamma_{\text{total}}$ Γ_{47}/Γ VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

2.14±0.16 OUR AVERAGE $2.14 \pm 0.12 \pm 0.10$ ¹ AUBERT 06A BABR $e^+e^- \rightarrow \Upsilon(4S)$ $2.0 \pm 0.4 \pm 0.1$ ² ABE 01I BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 06A reports $[B(B^0 \rightarrow D^*(2010)^- K^+)] / [B(B^0 \rightarrow D^*(2010)^- \pi^+)] = 0.0776 \pm 0.0034 \pm 0.0029$. We multiply by our best value $B(B^0 \rightarrow D^*(2010)^- \pi^+) = (2.76 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABE 01I reports $[B(B^0 \rightarrow D^*(2010)^- K^+)] / [B(B^0 \rightarrow D^*(2010)^- \pi^+)] = 0.074 \pm 0.015 \pm 0.006$. We multiply by our best value $B(B^0 \rightarrow D^*(2010)^- \pi^+) = (2.76 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(D^*(2010)^- K^0\pi^+)/\Gamma_{\text{total}}$ Γ_{48}/Γ VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

3.0±0.7±0.3¹ AUBERT,BE 05B BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D^*(2010)^- K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{49}/Γ VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

3.3±0.6 OUR AVERAGE $3.2 \pm 0.6 \pm 0.3$ ¹ AUBERT,BE 05B BABR $e^+e^- \rightarrow \Upsilon(4S)$ $3.8 \pm 1.3 \pm 0.8$ ² MAHAPATRA 02 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and an unpolarized final state.

$\Gamma(D^*(2010)^- K^+ \bar{K}^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|-----------------------|-------------|---------------------------------------|
| <4.7 | 90 | ¹ DRUTSKOY | 02 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^*(2010)^- K^+ \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-----------------------|-------------|---------------------------------------|
| 12.9 ± 2.2 ± 2.5 | ¹ DRUTSKOY | 02 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{52}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------------|------------|---|-------------|----------------|
| 0.0070 ± 0.0008 OUR AVERAGE | | Error includes scale factor of 1.3. See the ideogram below. | | |

| | | | | |
|-----------------------------|---------------------------|------|----------------------------------|----------------------------------|
| 0.00681 ± 0.00023 ± 0.00072 | ¹ MAJUMDER | 04 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.0063 ± 0.0010 ± 0.0011 | ^{2,3} ALAM | 94 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.0134 ± 0.0036 ± 0.0001 | ⁴ BORTOLETTO92 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ | |
| 0.0101 ± 0.0041 ± 0.0001 | ⁵ ALBRECHT | 90J | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|-----------------------|--------------------|-----|---------------------------------------|
| 0.033 ± 0.009 ± 0.016 | ⁶ ALBRECHT | 87C | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| <0.042 | 90 | ⁷ BEBEK | 87 | CLEO $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.


³ The three pion mass is required to be between 1.0 and 1.6 GeV consistent with an a_1 meson. (If this channel is dominated by a_1^+ , the branching ratio for $\bar{D}^* - a_1^+$ is twice that for $\bar{D}^* - \pi^+ \pi^+ \pi^-$.)

⁴ BORTOLETTO 92 reports $0.0159 \pm 0.0028 \pm 0.0037$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

⁵ ALBRECHT 90J reports $0.012 \pm 0.003 \pm 0.004$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

⁶ ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\gamma(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\gamma(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

⁷ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.



$\Gamma((D^*(2010)^- \pi^+ \pi^+ \pi^-) \text{ nonresonant}) / \Gamma_{\text{total}}$

Γ_{53}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|---------------------------|------|----------------------------------|
| 0.0000±0.0019±0.0016 | ¹ BORTOLETTO92 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

$\Gamma(D^*(2010)^- \pi^+ \rho^0) / \Gamma_{\text{total}}$

Γ_{54}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|---------------------------|------|----------------------------------|
| 0.00573±0.00317±0.00004 | ¹ BORTOLETTO92 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ BORTOLETTO 92 reports $0.0068 \pm 0.0032 \pm 0.0021$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(D^*(2010)^- a_1(1260)^+)/\Gamma_{\text{total}}$

Γ_{55}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|---------------------------|------|----------------------------------|
| 0.0130±0.0027 OUR AVERAGE | | | |
| 0.0126±0.0020±0.0022 | ^{1,2} ALAM 94 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.0152±0.0070±0.0001 | ³ BORTOLETTO92 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ ALAM 94 value is twice their $\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^-) / \Gamma_{\text{total}}$ value based on their observation that the three pions are dominantly in the $a_1(1260)$ mass range 1.0 to 1.6 GeV.

² ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0) / B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-) / B(D^0 \rightarrow K^- \pi^+)$.

³ BORTOLETTO 92 reports $0.018 \pm 0.006 \pm 0.006$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(D^*(2010)^-\pi^+\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{56}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------------|------|-----------------------------------|
| 0.0176 ± 0.0027 OUR AVERAGE | | | | |
| $0.0172 \pm 0.0014 \pm 0.0024$ | | ¹ ALEXANDER 01B | CLE2 | $e^+e^- \rightarrow \Upsilon(4S)$ |
| $0.0345 \pm 0.0181 \pm 0.0003$ | 28 | ² ALBRECHT 90J | ARG | $e^+e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega\pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

² ALBRECHT 90J reports $0.041 \pm 0.015 \pm 0.016$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(D^*-3\pi^+2\pi^-)/\Gamma_{\text{total}}$ Γ_{57}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--|--------------------------|------|-----------------------------------|
| $4.72 \pm 0.59 \pm 0.71$ | ¹ MAJUMDER 04 | BELL | $e^+e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^-p\bar{p}\pi^+)/\Gamma_{\text{total}}$ Γ_{58}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|--------------------------|------|-----------------------------------|
| $6.5^{+1.3}_{-1.2} \pm 1.0$ | ¹ ANDERSON 01 | CLE2 | $e^+e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^-p\bar{n})/\Gamma_{\text{total}}$ Γ_{59}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--|--------------------------|------|-----------------------------------|
| $14.5^{+3.4}_{-3.0} \pm 2.7$ | ¹ ANDERSON 01 | CLE2 | $e^+e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2010)^-\omega\pi^+)/\Gamma_{\text{total}}$ Γ_{60}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT | |
|---|------------------------------|------|-----------------------------------|--|
| 2.89 ± 0.30 OUR AVERAGE | | | | |
| $2.88 \pm 0.21 \pm 0.31$ | ¹ AUBERT 06L | BABR | $e^+e^- \rightarrow \Upsilon(4S)$ | |
| $2.9 \pm 0.3 \pm 0.4$ | ^{1,2} ALEXANDER 01B | CLE2 | $e^+e^- \rightarrow \Upsilon(4S)$ | |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² The signal is consistent with all observed $\omega\pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

$\Gamma(D_1(2430)^0 \omega \times B(D_1(2430)^0 \rightarrow D^{*-} \pi^+)) / \Gamma_{\text{total}}$ Γ_{61}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|----------|----------------------------------|
| 4.1±1.2±1.1 | 1 AUBERT | 06L BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by fitting the events with $\cos \theta_{D^*} < 0.5$ and scaling up the result by a factor of 4/3. No interference effects between $B^0 \rightarrow D'_1 \omega$ and $D^* \omega \pi$ are assumed.

 $\Gamma(\bar{D}^{**-} \pi^+) / \Gamma_{\text{total}}$ Γ_{62}/Γ

D^{**-} represents an excited state with mass $2.2 < M < 2.8 \text{ GeV}/c^2$.

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|---------------|----------|----------------------------------|
| 2.1±1.0±0.1 | 1,2 AUBERT,BE | 06J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ AUBERT,BE 06J reports $[B(B^0 \rightarrow \bar{D}^{**-} \pi^+)] / [B(B^0 \rightarrow D^- \pi^+)] = 0.77 \pm 0.22 \pm 0.29$. We multiply by our best value $B(B^0 \rightarrow D^- \pi^+) = (2.68 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.

 $\Gamma(D_1(2420)^- \pi^+ \times B(D_1^- \rightarrow D^- \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{63}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|----------|----------------------------------|
| 0.89±0.15±0.17 | 1 ABE | 05A BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D_1(2420)^- \pi^+ \times B(D_1^- \rightarrow D^{*-} \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{64}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|----------------------------------|
| <0.33 | 90 | 1 ABE | 05A BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}_2^*(2460)^- \pi^+ \times B(D_2^*(2460)^- \rightarrow D^0 \pi^-)) / \Gamma_{\text{total}}$ Γ_{65}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------------|-------------|----------------------------------|---------|
| 2.15±0.17±0.31 | 1,2 KUZMIN | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|--------|---------|----------------------------------|
| <14.7 | 90 | 1 ALAM | 94 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
|-------|----|--------|---------|----------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Our second uncertainty combines systematics and model errors quoted in the paper.

 $\Gamma(\bar{D}_0^*(2400)^- \pi^+ \times B(D_0^*(2400)^- \rightarrow D^0 \pi^-)) / \Gamma_{\text{total}}$ Γ_{66}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|---------|----------------------------------|
| 0.60±0.13±0.27 | 1,2 KUZMIN | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Our second uncertainty combines systematics and model errors quoted in the paper.

 $\Gamma(D_2^*(2460)^- \pi^+ \times B((D_2^*)^- \rightarrow D^{*-} \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{67}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|----------------------------------|
| <0.24 | 90 | 1 ABE | 05A BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

| $\Gamma(\overline{D}_2^*(2460)^-\rho^+)/\Gamma_{\text{total}}$ | | | | Γ_{68}/Γ |
|--|-----|-------------------|------|--------------------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
| <0.0049 | 90 | ¹ ALAM | 94 | CLE2 $e^+e^- \rightarrow \gamma(4S)$ |

¹ ALAM 94 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^-\pi^+)$ and $B(D_2^*(2460)^+\rightarrow D^0\pi^+) = 30\%$.

| $\Gamma(D^0\overline{D}^0)/\Gamma_{\text{total}}$ | | | | Γ_{69}/Γ |
|---|-----|-----------------------|------|--------------------------------------|
| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
| <0.6 | 90 | ¹ AUBERT,B | 06A | BABR $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

| $\Gamma(D^{*0}\overline{D}^0)/\Gamma_{\text{total}}$ | | | | Γ_{70}/Γ |
|--|-----|-----------------------|------|--------------------------------------|
| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
| <2.9 | 90 | ¹ AUBERT,B | 06A | BABR $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

| $\Gamma(D^-D^+)/\Gamma_{\text{total}}$ | | | | Γ_{71}/Γ |
|---|-----|-------------------------------------|------|--------------------------------------|
| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
| 2.11±0.31 OUR AVERAGE | | Error includes scale factor of 1.2. | | |
| 1.97±0.20±0.20 | | ¹ FRATINA | 07 | BELL $e^+e^- \rightarrow \gamma(4S)$ |
| 2.8 ± 0.4 ± 0.5 | | ¹ AUBERT,B | 06A | BABR $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.91±0.51±0.30 | | ¹ MAJUMDER | 05 | BELL Repl. by FRATINA 07 |
| < 9.4 | 90 | ¹ LIPELES | 00 | CLE2 $e^+e^- \rightarrow \gamma(4S)$ |
| <59 | 90 | BARATE | 98Q | ALEP $e^+e^- \rightarrow Z$ |
| <12 | 90 | ASNER | 97 | CLE2 $e^+e^- \rightarrow \gamma(4S)$ |
| 1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. | | | | |

| $\Gamma(D^-D_s^+)/\Gamma_{\text{total}}$ | | | | Γ_{72}/Γ |
|---|------|---------------------------|------|--------------------------------------|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.0074±0.0007 OUR AVERAGE | | | | |
| 0.0075±0.0005±0.0006 | | ¹ ZUPANC | 07 | BELL $e^+e^- \rightarrow \gamma(4S)$ |
| 0.0068±0.0015±0.0005 | | ² AUBERT | 06N | BABR $e^+e^- \rightarrow \gamma(4S)$ |
| 0.0070±0.0025±0.0006 | | ³ GIBAUT | 96 | CLE2 $e^+e^- \rightarrow \gamma(4S)$ |
| 0.010 ± 0.009 ± 0.001 | | ⁴ ALBRECHT | 92G | ARG $e^+e^- \rightarrow \gamma(4S)$ |
| 0.0055±0.0031±0.0004 | | ⁵ BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.012 ± 0.007 | 3 | ⁶ BORTOLETTO90 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

¹ ZUPANC 07 reports $(7.5 \pm 0.2 \pm 1.1) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$.

We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AUBERT 06N reports $(0.64 \pm 0.13 \pm 0.10) \times 10^{-2}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.0462 \pm 0.0062$.

We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ GIBAUT 96 reports $0.0087 \pm 0.0024 \pm 0.0020$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ALBRECHT 92G reports $0.017 \pm 0.013 \pm 0.006$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ branching ratios, e.g., $B(D^+ \rightarrow K^-\pi^+\pi^+) = 7.7 \pm 1.0\%$.

⁵ BORTOLETTO 92 reports $0.0080 \pm 0.0045 \pm 0.0030$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.030 \pm 0.011$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁶ BORTOLETTO 90 assume $B(D_s \rightarrow \phi\pi^+) = 2\%$. Superseded by BORTOLETTO 92.

$$\Gamma(D^*(2010)^- D_s^+)/\Gamma_{\text{total}} \quad \Gamma_{73}/\Gamma$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------------|------|-----------------------------------|
| 0.0083±0.0011 OUR AVERAGE | | | | |
| 0.0075±0.0013±0.0006 | | ¹ AUBERT 06N | BABR | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.0085±0.0016±0.0007 | | ² AUBERT 03I | BABR | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.0090±0.0017±0.0007 | | ³ AHMED 00B | CLE2 | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.009 ± 0.006 ± 0.001 | | ⁴ ALBRECHT 92G | ARG | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.011 ± 0.006 ± 0.001 | | ⁵ BORTOLETTO92 | CLEO | $e^+e^- \rightarrow \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.0074±0.0022±0.0006 | | ⁶ GIBAUT 96 | CLE2 | Repl. by AHMED 00B |
| 0.024 ± 0.014 | 3 | ⁷ BORTOLETTO90 | CLEO | $e^+e^- \rightarrow \Upsilon(4S)$ |

¹ AUBERT 06N reports $(0.71 \pm 0.13 \pm 0.09) \times 10^{-2}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.0462 \pm 0.0062$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AUBERT 03I reports $0.0103 \pm 0.0014 \pm 0.0013$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AHMED 00B reports $0.0110 \pm 0.0018 \pm 0.0011$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ALBRECHT 92G reports $0.014 \pm 0.010 \pm 0.003$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ and $D^*(2010)^+$ branching ratios, e.g., $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$, $B(D^+ \rightarrow K^-\pi^+\pi^+) = 7.1 \pm 1.0\%$, and $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 55 \pm 4\%$.

⁵ BORTOLETTO 92 reports $0.016 \pm 0.009 \pm 0.006$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.030 \pm 0.011$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our

best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

⁶GIBAUT 96 reports $0.0093 \pm 0.0023 \pm 0.0016$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
⁷BORTOLETTO 90 assume $B(D_s \rightarrow \phi\pi^+) = 2\%$. Superseded by BORTOLETTO 92.

$\Gamma(D^- D_s^{*+})/\Gamma_{\text{total}}$

Γ_{74}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|---------------------------|------|-----------------------------------|
| 0.0076±0.0016 OUR AVERAGE | | | |
| 0.0073±0.0017±0.0006 | ¹ AUBERT 06N | BABR | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.0080±0.0033±0.0006 | ² GIBAUT 96 | CLE2 | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.017 ± 0.012 ± 0.001 | ³ ALBRECHT 92G | ARG | $e^+e^- \rightarrow \Upsilon(4S)$ |

¹AUBERT 06N reports $(0.69 \pm 0.16 \pm 0.09) \times 10^{-2}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.0462 \pm 0.0062$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²GIBAUT 96 reports $0.0100 \pm 0.0035 \pm 0.0022$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ALBRECHT 92G reports $0.027 \pm 0.017 \pm 0.009$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ branching ratios, e.g., $B(D^+ \rightarrow K^-\pi^+\pi^+) = 7.7 \pm 1.0\%$.

$[\Gamma(D^*(2010)^- D_s^+) + \Gamma(D^*(2010)^- D_s^{*+})]/\Gamma_{\text{total}}$

$(\Gamma_{73} + \Gamma_{75})/\Gamma$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|---------|
| 2.6±0.4 OUR AVERAGE | | | | |

¹AUBERT 03I reports $(3.00 \pm 0.19 \pm 0.39) \times 10^{-2}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
²BORTOLETTO 90 reports $(7.5 \pm 2.0) \times 10^{-2}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.02$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^*(2010)^- D_s^{*+})/\Gamma_{\text{total}}$

Γ_{75}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-------------|------|---------|
| 0.0179±0.0014 OUR AVERAGE | | | |

¹AUBERT 06N
²AUBERT 05V
³AUBERT 03I
⁴AHMED 00B
⁵ALBRECHT 92G
• • • We do not use the following data for averages, fits, limits, etc. • • •
⁶GIBAUT 96 CLE2 Repl. by AHMED 00B

¹ AUBERT 06N reports $(1.68 \pm 0.21 \pm 0.19) \times 10^{-2}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.0462 \pm 0.0062$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² A partial reconstruction technique is used and the result is independent of the particle decay rate of D_s^+ meson. It also provides a model-independent determination of $B(D_s^+ \rightarrow \phi\pi^+) = (4.81 \pm 0.52 \pm 0.38)\%$.

³ AUBERT 03I reports $0.0197 \pm 0.0015 \pm 0.0030$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ AHMED 00B reports $0.0182 \pm 0.0037 \pm 0.0025$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ ALBRECHT 92G reports $0.026 \pm 0.014 \pm 0.006$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ and $D^*(2010)^+$ branching ratios, e.g., $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$, $B(D^+ \rightarrow K^-\pi^+\pi^+) = 7.1 \pm 1.0\%$, and $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 55 \pm 4\%$.

⁶ GIBAUT 96 reports $0.0203 \pm 0.0050 \pm 0.0036$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_{s0}(2317)^- K^+ \times B(D_{s0}(2317)^- \rightarrow D_s^- \pi^0)) / \Gamma_{\text{total}} \quad \Gamma_{76}/\Gamma$$

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---|--------------------------|------|---------------------------------|
| $4.4^{+1.4}_{-1.3} \pm 0.4$ | ¹ DRUTSKOY 05 | BELL | $e^+e^- \rightarrow \gamma(4S)$ |

¹ DRUTSKOY 05 reports $(5.3^{+1.5}_{-1.3} \pm 1.6) \times 10^{-5}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$.

We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_{s0}(2317)^- \pi^+ \times B(D_{s0}(2317)^- \rightarrow D_s^- \pi^0)) / \Gamma_{\text{total}} \quad \Gamma_{77}/\Gamma$$

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|--------------------------|------|---------------------------------|
| <2.5 | 90 | ¹ DRUTSKOY 05 | BELL | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(D_{sJ}(2457)^- K^+ \times B(D_{sJ}(2457)^- \rightarrow D_s^- \pi^0)) / \Gamma_{\text{total}} \quad \Gamma_{78}/\Gamma$$

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|--------------------------|------|---------------------------------|
| <0.94 | 90 | ¹ DRUTSKOY 05 | BELL | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(D_{sJ}(2457)^-\pi^+ \times B(D_{sJ}(2457)^- \rightarrow D_s^- \pi^0)) / \Gamma_{\text{total}}$ Γ_{79}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|-----------------------|-------------|---------------------------------------|
| <0.40 | 90 | ¹ DRUTSKOY | 05 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D_s^- D_s^+)/\Gamma_{\text{total}}$ Γ_{80}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------------|------------|---------------------|-------------|---------------------------------------|
| < 3.6 × 10⁻⁵ | 90 | ¹ ZUPANC | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----|------------------------|-----|---------------------------------------|
| $< 10 \times 10^{-5}$ | 90 | ¹ AUBERT,BE | 05F | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
|-----------------------|----|------------------------|-----|---------------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D_s^{*-} D_s^+)/\Gamma_{\text{total}}$ Γ_{81}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------------|------------|------------------------|-------------|---------------------------------------|
| < 1.3 × 10⁻⁴ | 90 | ¹ AUBERT,BE | 05F | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D_s^{*-} D_s^{*+})/\Gamma_{\text{total}}$ Γ_{82}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------------|------------|------------------------|-------------|---------------------------------------|
| < 2.4 × 10⁻⁴ | 90 | ¹ AUBERT,BE | 05F | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D_{s0}(2317)^+ D^- \times B(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0)) / \Gamma_{\text{total}}$ Γ_{83}/Γ

| <u>VALUE</u> (units 10^{-3}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|--------------------|-------------|----------------|
|---------------------------------|--------------------|-------------|----------------|

0.99^{+0.42}_{-0.34} OUR AVERAGE Error includes scale factor of 1.5.

| | | | |
|---|-------------------------|-----|---------------------------------------|
| $1.5 \begin{array}{l} +0.5 \\ -0.4 \end{array} \pm 0.1$ | ^{1,2} AUBERT,B | 04S | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
|---|-------------------------|-----|---------------------------------------|

| | | | |
|---|-------------------------|-----|---------------------------------------|
| $0.71 \begin{array}{l} +0.30 \\ -0.25 \end{array} \pm 0.06$ | ^{1,3} KROKOVNY | 03B | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
|---|-------------------------|-----|---------------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² AUBERT,B 04S reports $(1.8 \pm 0.4 \begin{array}{l} +0.7 \\ -0.5 \end{array}) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$.

We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ KROKOVNY 03B reports $(0.86 \begin{array}{l} +0.33 \\ -0.26 \end{array} \pm 0.26) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$.

We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(D_{s0}(2317)^+ D^- \times B(D_{s0}(2317)^+ \rightarrow D_s^{*+} \gamma)) / \Gamma_{\text{total}}$ Γ_{84}/Γ

| <u>VALUE</u> (units 10^{-3}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|-----------------------|-------------|---------------------------------------|
| <0.95 | 90 | ¹ KROKOVNY | 03B | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(D_{s0}(2317)^+ D^*(2010)^- \times B(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0)) / \Gamma_{\text{total}} \quad \Gamma_{85}/\Gamma$$

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|----------------------------------|
| $1.5 \pm 0.4^{+0.5}_{-0.4}$ | ¹ AUBERT,B | 04S BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(D_{sJ}(2457)^+ D^-) / \Gamma_{\text{total}} \quad \Gamma_{86}/\Gamma$$

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------|-------------|----------------------------------|
| 3.5 ± 1.1 OUR AVERAGE | | | |
| $2.6 \pm 1.5 \pm 0.7$ | ¹ AUBERT | 06N BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $4.8^{+2.2}_{-1.6} \pm 1.1$ | ^{2,3} AUBERT,B | 04S BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $3.9^{+1.5}_{-1.3} \pm 0.9$ | ^{2,4} KROKOVNY | 03B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses a missing-mass method in the events that one of the B mesons is fully reconstructed. |

² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

³ AUBERT,B 04S reports $[B(B^0 \rightarrow D_{sJ}(2457)^+ D^-)] \times [B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0)] = (2.3^{+1.0}_{-0.7} \pm 0.3) \times 10^{-3}$. We divide by our best value $B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ KROKOVNY 03B reports $[B(B^0 \rightarrow D_{sJ}(2457)^+ D^-)] \times [B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0)] = (1.9^{+0.7}_{-0.6} \pm 0.2) \times 10^{-3}$. We divide by our best value $B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_{sJ}(2457)^+ D^- \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)) / \Gamma_{\text{total}} \quad \Gamma_{87}/\Gamma$$

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------------------|-------------|----------------------------------|
| $0.67^{+0.17}_{-0.14}$ OUR AVERAGE | | | |
| $0.66^{+0.25}_{-0.16} \pm 0.05$ | ^{1,2} AUBERT,B | 04S BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.67^{+0.22}_{-0.20} \pm 0.05$ | ^{1,3} KROKOVNY | 03B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² AUBERT,B 04S reports $(0.8 \pm 0.2^{+0.3}_{-0.2}) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$.

We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ KROKOVNY 03B reports $(0.82^{+0.22}_{-0.19} \pm 0.25) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_{sJ}(2457)^+ D^- \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \gamma)) / \Gamma_{\text{total}} \quad \Gamma_{88}/\Gamma$$

| <u>VALUE (units 10^{-3})</u> | <u>CL %</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-----------------------|-------------|----------------------------------|
| <0.60 | 90 | ¹ KROKOVNY | 03B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(D_{sJ}(2457)^+ D^- \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{89}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-----------------------|-------------|----------------------------------|
| <0.20 | 90 | ¹ KROKOVNY | 03B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D_{sJ}(2457)^+ D^- \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^0))/\Gamma_{\text{total}}$ Γ_{90}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-----------------------|-------------|----------------------------------|
| <0.36 | 90 | ¹ KROKOVNY | 03B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^*(2010)^- D_{sJ}(2457)^+)/\Gamma_{\text{total}}$ Γ_{91}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| 9.3±2.2 OUR AVERAGE | | | |

| | | | |
|--|---------------------|----------|----------------------------------|
| 8.8±2.0±1.4 | ¹ AUBERT | 06N BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 11 $\begin{array}{l} +5 \\ -4 \end{array}$ ± 3 | 2,3 AUBERT,B | 04S BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses a missing-mass method in the events that one of the B mesons is fully reconstructed.

² AUBERT,B 04S reports $[B(B^0 \rightarrow D^*(2010)^- D_{sJ}(2457)^+)] \times [B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0)] = (5.5 \pm 1.2^{+2.2}_{-1.6}) \times 10^{-3}$. We divide by our best value $B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D_{sJ}(2457)^+ D^*(2010)^- \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma))/\Gamma_{\text{total}}$ Γ_{92}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-----------------------|-------------|----------------------------------|
| 2.3±0.3^{+0.9}_{-0.6} | ¹ AUBERT,B | 04S BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^- D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^{*0} K^+))/\Gamma_{\text{total}}$ Γ_{93}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------|-------------|----------------------------------|
| 1.71±0.48±0.32 | | ¹ AUBERT | 08B BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5 90 AUBERT 03X BABR Repl. by AUBERT 08B

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^*(2010)^- D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^{*0} K^+))/\Gamma_{\text{total}}$ Γ_{94}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------|-------------|----------------------------------|
| 3.32±0.88±0.66 | | ¹ AUBERT | 08B BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7 90 AUBERT 03X BABR Repl. by AUBERT 08B

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(D^- D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^{*+} K^0)) / \Gamma_{\text{total}}$ Γ_{95}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------|---------------------|----------|----------------------------------|
| 2.61 ± 1.03 ± 0.31 | ¹ AUBERT | 08B BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^{*-} D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^{*+} K^0)) / \Gamma_{\text{total}}$ Γ_{96}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------|---------------------|----------|----------------------------------|
| 5.00 ± 1.51 ± 0.67 | ¹ AUBERT | 08B BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^- D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)) / \Gamma_{\text{total}}$ Γ_{97}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|----------------------------------|
| <1 | 90 | AUBERT | 03X BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

 $\Gamma(D^*(2010)^- D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)) / \Gamma_{\text{total}}$ Γ_{98}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|----------------------------------|
| <2 | 90 | AUBERT | 03X BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

 $\Gamma(D_s^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{99}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-----|-------------|------|---------|
| 15.3 ± 3.5 OUR AVERAGE | | | | |

14 ± 4 ± 1

20 $^{+9}_{-7}$ ± 2

¹ AUBERT 07K BABR $e^+ e^- \rightarrow \gamma(4S)$

² KROKOVNY 02 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

26 ± 9 ± 2

< 230 90

< 1300 90

³ AUBERT 03D BABR Repl. by AUBERT 07K

⁴ ALEXANDER 93B CLE2 $e^+ e^- \rightarrow \gamma(4S)$

⁵ BORTOLETTO 90 CLEO $e^+ e^- \rightarrow \gamma(4S)$

¹ AUBERT 07K reports $[B(B^0 \rightarrow D_s^+ \pi^-)] \times [B(D_s^+ \rightarrow \phi \pi^+)] = (0.63 \pm 0.15 \pm 0.05) \times 10^{-6}$. We divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² KROKOVNY 02 reports $[B(B^0 \rightarrow D_s^+ \pi^-)] \times [B(D_s^+ \rightarrow \phi \pi^+)] = (0.86^{+0.37}_{-0.30} \pm 0.11) \times 10^{-6}$. We divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AUBERT 03D reports $[B(B^0 \rightarrow D_s^+ \pi^-)] \times [B(D_s^+ \rightarrow \phi \pi^+)] = (1.13 \pm 0.33 \pm 0.21) \times 10^{-6}$. We divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ALEXANDER 93B reports < 270×10^{-6} for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.0438$.

⁵ BORTOLETTO 90 assume $B(D_s \rightarrow \phi \pi^+) = 2\%$.

| $[\Gamma(D_s^+ \pi^-) + \Gamma(D_s^- K^+)/\Gamma_{\text{total}}]$ | $(\Gamma_{99} + \Gamma_{109})/\Gamma$ | | | |
|--|---------------------------------------|--------------------|-------------|----------------------------------|
| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| <0.0010 | 90 | 1 ALBRECHT | 93E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| ¹ ALBRECHT 93E reports $< 1.7 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$. | | | | |

| $\Gamma(D_s^{*+} \pi^-)/\Gamma_{\text{total}}$ | Γ_{100}/Γ | | | |
|---|-----------------------|--------------------|----------------------------------|----------------------------------|
| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| $3.0 \pm 0.7 \pm 0.2$ | 1 AUBERT | 07K BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 4.1 | 90 | AUBERT | 03D BABR | Repl. by AUBERT 07K |
| <40 | 90 | 2 ALEXANDER | 93B CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| ¹ AUBERT 07K reports $[B(B^0 \rightarrow D_s^{*+} \pi^-)] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (1.32 \pm 0.27 \pm 0.15) \times 10^{-6}$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |
| ² ALEXANDER 93B reports $< 44 \times 10^{-5}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$. | | | | |

| $[\Gamma(D_s^{*+} \pi^-) + \Gamma(D_s^{*-} K^+)/\Gamma_{\text{total}}]$ | $(\Gamma_{100} + \Gamma_{110})/\Gamma$ | | | |
|--|--|--------------------|-------------|----------------------------------|
| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| <0.0007 | 90 | 1 ALBRECHT | 93E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| ¹ ALBRECHT 93E reports $< 1.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$. | | | | |

| $\Gamma(D_s^+ \rho^-)/\Gamma_{\text{total}}$ | Γ_{101}/Γ | | | |
|---|-----------------------|--------------------|-------------|----------------------------------|
| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| <0.0006 | 90 | 1 ALEXANDER | 93B CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.0014 | 90 | 2 ALBRECHT | 93E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| ¹ ALEXANDER 93B reports $< 6.6 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$. | | | | |
| ² ALBRECHT 93E reports $< 2.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$. | | | | |

| $\Gamma(D_s^{*+} \rho^-)/\Gamma_{\text{total}}$ | Γ_{102}/Γ | | | |
|---|-----------------------|--------------------|-------------|----------------------------------|
| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| <0.0006 | 90 | 1 ALEXANDER | 93B CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.0015 | 90 | 2 ALBRECHT | 93E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| ¹ ALEXANDER 93B reports $< 7.4 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$. | | | | |
| ² ALBRECHT 93E reports $< 2.5 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$. | | | | |

$\Gamma(D_s^+ a_0^-)/\Gamma_{\text{total}}$ Γ_{103}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|---------------------|-------------|----------------------------------|
| <1.9 | 90 | ¹ AUBERT | 06x BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D_s^{*+} a_0^-)/\Gamma_{\text{total}}$ Γ_{104}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|---------------------|-------------|----------------------------------|
| <3.6 | 90 | ¹ AUBERT | 06x BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D_s^+ a_1(1260)^-)/\Gamma_{\text{total}}$ Γ_{105}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------|------------|-----------------------|-------------|----------------------------------|
| <0.0022 | 90 | ¹ ALBRECHT | 93E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ ALBRECHT 93E reports $< 3.5 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$. $\Gamma(D_s^{*+} a_1(1260)^-)/\Gamma_{\text{total}}$ Γ_{106}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------|------------|-----------------------|-------------|----------------------------------|
| <0.0018 | 90 | ¹ ALBRECHT | 93E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ ALBRECHT 93E reports $< 2.9 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$. $\Gamma(D_s^+ a_2^-)/\Gamma_{\text{total}}$ Γ_{107}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|---------------------|-------------|----------------------------------|
| <19 | 90 | ¹ AUBERT | 06x BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D_s^{*+} a_2^-)/\Gamma_{\text{total}}$ Γ_{108}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|---------------------|-------------|----------------------------------|
| <20 | 90 | ¹ AUBERT | 06x BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D_s^- K^+)/\Gamma_{\text{total}}$ Γ_{109}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|----------------|
| 29 ± 5 OUR AVERAGE | | | | |

 $28 \pm 5 \pm 2$ $37^{+11}_{-10} \pm 3$ ¹AUBERT 07K BABR $e^+ e^- \rightarrow \gamma(4S)$ ²KROKOVNY 02 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $26 \pm 10 \pm 2$ < 190 < 1300 ³AUBERT 03D BABR Repl. by AUBERT 07K⁴ALEXANDER 93B CLE2 $e^+ e^- \rightarrow \gamma(4S)$ ⁵BORTOLETTO90 CLEO $e^+ e^- \rightarrow \gamma(4S)$

¹AUBERT 07K reports $[B(B^0 \rightarrow D_s^- K^+)] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (1.21 \pm 0.17 \pm 0.11) \times 10^{-6}$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²KROKOVNY 02 reports $[B(B^0 \rightarrow D_s^- K^+)] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (1.61^{+0.45}_{-0.38} \pm 0.21) \times 10^{-6}$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³AUBERT 03D reports $[B(B^0 \rightarrow D_s^- K^+)] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (1.16 \pm 0.36 \pm 0.24) \times 10^{-6}$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ALEXANDER 93B reports $< 230 \times 10^{-6}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$.

⁵BORTOLETTO 90 assume $B(D_s \rightarrow \phi\pi^+) = 2\%$.

$\Gamma(D_s^{*-} K^+)/\Gamma_{\text{total}}$

Γ_{110}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------------|----------|----------------------------------|
| 2.2±0.6±0.2 | | ¹ AUBERT | 07K BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 2.5 | 90 | AUBERT | 03D BABR | Repl. by AUBERT 07K |
| <14 | 90 | ² ALEXANDER | 93B CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹AUBERT 07K reports $[B(B^0 \rightarrow D_s^{*-} K^+)] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (0.97 \pm 0.24 \pm 0.12) \times 10^{-6}$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²ALEXANDER 93B reports $< 17 \times 10^{-5}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$.

$\Gamma(D_s^- K^*(892)^+)/\Gamma_{\text{total}}$

Γ_{111}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------------|----------|----------------------------------|
| <0.0008 | 90 | ¹ ALEXANDER | 93B CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.0028 | 90 | ² ALBRECHT | 93E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ALEXANDER 93B reports $< 9.7 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$.

²ALBRECHT 93E reports $< 4.6 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$.

$\Gamma(D_s^{*-} K^*(892)^+)/\Gamma_{\text{total}}$

Γ_{112}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------------|----------|----------------------------------|
| <0.0009 | 90 | ¹ ALEXANDER | 93B CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.004 | 90 | ² ALBRECHT | 93E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ ALEXANDER 93B reports $< 11.0 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$.

² ALBRECHT 93E reports $< 5.8 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$.

$\Gamma(D_s^- \pi^+ K^0)/\Gamma_{\text{total}}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|-----------------------|------|----------------------------------|
| <0.005 | 90 | ¹ ALBRECHT | 93E | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ ALBRECHT 93E reports $< 7.3 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$.

Γ_{113}/Γ

$\Gamma(D_s^{*-} \pi^+ K^0)/\Gamma_{\text{total}}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------|-----|-----------------------|------|----------------------------------|
| <0.0026 | 90 | ¹ ALBRECHT | 93E | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ ALBRECHT 93E reports $< 4.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$.

Γ_{114}/Γ

$\Gamma(D_s^- \pi^+ K^*(892)^0)/\Gamma_{\text{total}}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------|-----|-----------------------|------|----------------------------------|
| <0.0031 | 90 | ¹ ALBRECHT | 93E | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ ALBRECHT 93E reports $< 5.0 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$.

Γ_{115}/Γ

$\Gamma(D_s^{*-} \pi^+ K^*(892)^0)/\Gamma_{\text{total}}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------|-----|-----------------------|------|----------------------------------|
| <0.0017 | 90 | ¹ ALBRECHT | 93E | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ ALBRECHT 93E reports $< 2.7 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.0438$.

Γ_{116}/Γ

$\Gamma(\bar{D}^0 K^0)/\Gamma_{\text{total}}$

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-----------------------|----------|----------------------------------|
| 5.2±0.7 OUR AVERAGE | | | |
| $5.3 \pm 0.7 \pm 0.3$ | ¹ AUBERT,B | 06L BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

$5.0^{+1.3}_{-1.2} \pm 0.6$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

Γ_{117}/Γ

$\Gamma(\bar{D}^0 K^+ \pi^-)/\Gamma_{\text{total}}$

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|---------------------|----------|----------------------------------|
| 88±15±9 | ¹ AUBERT | 06A BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

Γ_{118}/Γ

$\Gamma(\overline{D}^0 K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{119}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|-----------------------|-------------|----------------|----------------------------------|
| 4.2±0.6 OUR AVERAGE | | | | |
| $4.0 \pm 0.7 \pm 0.3$ | ¹ AUBERT,B | 06L | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $4.8^{+1.1}_{-1.0} \pm 0.5$ | ¹ KROKOVNY | 03 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $5.7 \pm 0.9 \pm 0.6$ | ¹ AUBERT | 06A | BABR | Repl. by AUBERT,B 06L |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D_2^*(2460)^- K^+ \times B(D_2^*(2460)^- \rightarrow \overline{D}^0 \pi^-))/\Gamma_{\text{total}}$ Γ_{120}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|--|---------------------|-------------|----------------|----------------------------------|
| $18.3 \pm 4.0 \pm 3.1$ | ¹ AUBERT | 06A | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. | | | | |

 $\Gamma(\overline{D}^0 K^+ \pi^- \text{ non-resonant})/\Gamma_{\text{total}}$ Γ_{121}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|------------|---------------------|-------------|----------------|----------------------------------|
| <37 | 90 | ¹ AUBERT | 06A | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\overline{D}^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{122}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|------------|---------------------|-------------|----------------|----------------------------------|
| 2.61±0.24 OUR AVERAGE | | | | | |
| $2.25 \pm 0.14 \pm 0.35$ | | ¹ BLYTH | 06 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $2.9 \pm 0.2 \pm 0.3$ | | ¹ AUBERT | 04B | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $2.74^{+0.36}_{-0.32} \pm 0.55$ | | ¹ COAN | 02 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $3.1 \pm 0.4 \pm 0.5$ ¹ ABE 02J BELL Repl. by BLYTH 06 <1.2 90 ² NEMATI 98 CLE2 Repl. by COAN 02 <4.8 90 ³ ALAM 94 CLE2 Repl. by NEMATI 98¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² NEMATI 98 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.³ ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$. $\Gamma(\overline{D}^0 \rho^0)/\Gamma_{\text{total}}$ Γ_{123}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|--|------------|-----------------------|-------------|----------------|----------------------------------|
| $3.19 \pm 0.20 \pm 0.45$ | | ^{1,2} KUZMIN | 07 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $2.9 \pm 1.0 \pm 0.4$ ¹ SATPATHY 03 BELL Repl. by KUZMIN 07 < 3.9 90 ³ NEMATI 98 CLE2 $e^+ e^- \rightarrow \gamma(4S)$ < 5.5 90 ⁴ ALAM 94 CLE2 Repl. by NEMATI 98 < 6.0 90 ⁵ BORTOLETTO92 CLEO $e^+ e^- \rightarrow \gamma(4S)$ < 27.0 90 ⁶ ALBRECHT 88K ARG $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Our second uncertainty combines systematics and model errors quoted in the paper.

³ NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

⁴ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

⁵ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁶ ALBRECHT 88K reports < 0.003 assuming $B^0 \bar{B}^0 : B^+ B^-$ production ratio is 45:55. We rescale to 50%.

$\Gamma(\bar{D}^0 f_2)/\Gamma_{\text{total}}$

Γ_{124}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--|---------------|------|------------------------------------|
| $1.20 \pm 0.18 \pm 0.38$ | 1,2 KUZMIN 07 | BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Our second uncertainty combines systematics and model errors quoted in the paper.

$\Gamma(\bar{D}^0 \eta)/\Gamma_{\text{total}}$

Γ_{125}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------------------------|------|------------------------------------|
| 2.02 ± 0.35 OUR AVERAGE | | Error includes scale factor of 1.6. | | |
| 1.77 $\pm 0.16 \pm 0.21$ | | 1 BLYTH 06 | BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 2.5 $\pm 0.2 \pm 0.3$ | | 1 AUBERT 04B | BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.4 $\begin{array}{l} +0.5 \\ -0.4 \end{array} \pm 0.3$ | | 1 ABE 02J | BELL | Repl. by BLYTH 06 |
| <1.3 | 90 | 2 NEMATI 98 | CLE2 | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| <6.8 | 90 | 3 ALAM 94 | CLE2 | Repl. by NEMATI 98 |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(\bar{D}^0 \eta')/\Gamma_{\text{total}}$

Γ_{126}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------------------------|------|------------------------------------|
| 1.25 ± 0.23 OUR AVERAGE | | Error includes scale factor of 1.1. | | |
| 1.14 $\pm 0.20 \begin{array}{l} +0.10 \\ -0.13 \end{array}$ | | 1 SCHUMANN 05 | BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.7 $\pm 0.4 \pm 0.2$ | | 1 AUBERT 04B | BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <9.4 | 90 | 2 NEMATI 98 | CLE2 | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| <8.6 | 90 | 3 ALAM 94 | CLE2 | Repl. by NEMATI 98 |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(\bar{D}^0\eta')/\Gamma(\bar{D}^0\eta)$

| VALUE | CL% |
|---------------------|-----|
| 0.7±0.2 ±0.1 | |

 $\Gamma_{126}/\Gamma_{125}$

| DOCUMENT ID | TECN | COMMENT |
|-------------|----------|----------------------------------|
| AUBERT | 04B BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

 $\Gamma(\bar{D}^0\omega)/\Gamma_{\text{total}}$

| VALUE (units 10^{-4}) | CL% |
|------------------------------|-----|
| 2.59±0.30 OUR AVERAGE | |

| 1 | BLYTH | 06 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
|--|-----------|------------------|---------|----------------------------------|
| ¹ | AUBERT | 04B | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.8 | ± 0.5 | $+0.4$ -0.3 | 1 ABE | 02J BELL Repl. by BLYTH 06 |
| <5.1 | 90 | 2 NEMATI | 98 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <6.3 | 90 | 3 ALAM | 94 CLE2 | Repl. by NEMATI 98 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² NEMATI 98 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.³ ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$. $\Gamma(D^0\phi)/\Gamma_{\text{total}}$

| VALUE (units 10^{-6}) | CL% |
|--------------------------|-----|
| <11.6 | 90 |

| DOCUMENT ID | TECN | COMMENT |
|---------------------|-----------|----------------------------------|
| ¹ AUBERT | 07AO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. Γ_{128}/Γ

| VALUE (units 10^{-5}) | CL% |
|--------------------------|-----|
| <1.1 | 90 |

| DOCUMENT ID | TECN | COMMENT |
|-----------------------|----------|----------------------------------|
| ¹ AUBERT,B | 06L BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|-----------------------|---------|----------------------------------|
| <1.8 | 90 | ¹ KROKOVNY | 03 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
|------|----|-----------------------|---------|----------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. Γ_{130}/Γ

| VALUE (units 10^{-6}) | CL% |
|--------------------------|-----|
| <19 | 90 |

| DOCUMENT ID | TECN | COMMENT |
|---------------------|----------|----------------------------------|
| ¹ AUBERT | 06A BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D^0K^+\pi^-)/\Gamma_{\text{total}}$ Γ_{129}/Γ

| VALUE | CL% |
|----------------------------------|-----|
| <2.5 × 10⁻⁵ | 90 |

| DOCUMENT ID | TECN | COMMENT |
|-----------------------|----------|----------------------------------|
| ¹ AUBERT,B | 05Q BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------------|----|---------------------|---------|----------------------------------|
| <5.0 × 10 ⁻⁵ | 90 | ¹ ARTUSO | 00 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
|-------------------------|----|---------------------|---------|----------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. Γ_{131}/Γ

| VALUE | CL% |
|----------------------------------|-----|
| <2.5 × 10⁻⁵ | 90 |

| DOCUMENT ID | TECN | COMMENT |
|-----------------------|----------|----------------------------------|
| ¹ AUBERT,B | 05Q BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

$\Gamma(\bar{D}^*(2007)^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{132}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------------|------|---|
| 1.7 ± 0.4 OUR AVERAGE | | | | Error includes scale factor of 1.5. See the ideogram below. |
| 1.39 ± 0.18 ± 0.26 | | ¹ BLYTH 06 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 2.9 ± 0.4 ± 0.5 | | ¹ AUBERT 04B | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 2.20 ^{+0.59} _{-0.52} ± 0.79 | | ¹ COAN 02 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •


| | | | | |
|---|----|------------------------|------|--------------------|
| 2.7 ^{+0.8} _{-0.7} ± 0.5 | | ¹ ABE 02J | BELL | Repl. by BLYTH 06 |
| <4.4 | 90 | ² NEMATI 98 | CLE2 | Repl. by COAN 02 |
| <9.7 | 90 | ³ ALAM 94 | CLE2 | Repl. by NEMATI 98 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² NEMATI 98 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+ \pi^-)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

WEIGHTED AVERAGE
1.7±0.4 (Error scaled by 1.5)

 $\Gamma(\bar{D}^0 \pi^0)/\Gamma(\bar{D}^*(2007)^0 \pi^0)$ $\Gamma_{122}/\Gamma_{132}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------|-------------------------|-------------------------------------|
| 1.14 ± 0.26 OUR AVERAGE | | | Error includes scale factor of 1.3. |
| 1.62 ± 0.23 ± 0.35 | | ¹ BLYTH 06 | $e^+ e^- \rightarrow \gamma(4S)$ |
| 1.0 ± 0.1 ± 0.2 | | ¹ AUBERT 04B | $e^+ e^- \rightarrow \gamma(4S)$ |

$\Gamma(\bar{D}^*(2007)^0 \rho^0)/\Gamma_{\text{total}}$ Γ_{133}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|--------------------------|------|----------------------------------|
| $<5.1 \times 10^{-4}$ | 90 | ¹ SATPATHY 03 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <0.00056 | 90 | ² NEMATI 98 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <0.00117 | 90 | ³ ALAM 94 | CLE2 | Repl. by NEMATI 98 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² NEMATI 98 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+ \pi^-)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

 $\Gamma(\bar{D}^*(2007)^0 \eta)/\Gamma_{\text{total}}$ Γ_{134}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------------------------|------|----------------------------------|
| 1.8 ± 0.6 OUR AVERAGE | | Error includes scale factor of 1.8. | | |
| 1.40 $\pm 0.28 \pm 0.26$ | | ¹ BLYTH 06 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 2.6 $\pm 0.4 \pm 0.4$ | | ¹ AUBERT 04B | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <4.6 | 90 | ¹ ABE 02J | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| <2.6 | 90 | ² NEMATI 98 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <6.9 | 90 | ³ ALAM 94 | CLE2 | Repl. by NEMATI 98 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² NEMATI 98 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+ \pi^-)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

 $\Gamma(\bar{D}^0 \eta)/\Gamma(\bar{D}^*(2007)^0 \eta)$ $\Gamma_{125}/\Gamma_{134}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|----------------------------------|
| 0.99 ± 0.19 OUR AVERAGE | | | |
| 1.27 $\pm 0.29 \pm 0.25$ | BLYTH 06 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.9 $\pm 0.2 \pm 0.1$ | AUBERT 04B | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

 $\Gamma(\bar{D}^*(2007)^0 \eta')/\Gamma(\bar{D}^*(2007)^0 \eta)$ $\Gamma_{135}/\Gamma_{134}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|----------------------------------|
| $0.5 \pm 0.3 \pm 0.1$ | AUBERT 04B | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

 $\Gamma(\bar{D}^*(2007)^0 \eta')/\Gamma_{\text{total}}$ Γ_{135}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|---------------------------|------|----------------------------------|
| 1.23 ± 0.35 OUR AVERAGE | | | | |
| 1.21 $\pm 0.34 \pm 0.22$ | | ¹ SCHUMANN 05 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 1.3 $\pm 0.7 \pm 0.2$ | | ^{1,2} AUBERT 04B | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <14 | 90 | BRANDENB... 98 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <19 | 90 | ³ NEMATI 98 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <27 | 90 | ⁴ ALAM 94 | CLE2 | Repl. by NEMATI 98 |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Reports an upper limit $< 2.6 \times 10^{-4}$ at 90% CL.

³ NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

⁴ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(\bar{D}^0 \eta')/\Gamma(\bar{D}^*(2007)^0 \eta')$

$\Gamma_{126}/\Gamma_{135}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|------------------------------------|
| $1.3 \pm 0.8 \pm 0.2$ | AUBERT | 04B | $e^+ e^- \rightarrow \Upsilon(4S)$ |

$\Gamma(\bar{D}^0 \omega)/\Gamma(\bar{D}^*(2007)^0 \omega)$

$\Gamma_{127}/\Gamma_{142}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|------|---|
| 0.78 ± 0.14 OUR AVERAGE | Error includes scale factor of 1.1. | | |
| $1.04 \pm 0.20 \pm 0.17$ | BLYTH | 06 | BELL $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $0.7 \pm 0.1 \pm 0.1$ | AUBERT | 04B | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |

$\Gamma(\bar{D}^*(2007)^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{136}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|--------------|------|---|
| $(6.2 \pm 1.2 \pm 1.8) \times 10^{-4}$ | 1,2 SATPATHY | 03 | BELL $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² No assumption about the intermediate mechanism is made in the analysis.

$\Gamma(\bar{D}^*(2007)^0 K^0)/\Gamma_{\text{total}}$

Γ_{137}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|---|
| $3.6 \pm 1.2 \pm 0.3$ | 1 | AUBERT,B | 06L | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------|----|-----------------------|----|---|
| < 6.6 | 90 | ¹ KROKOVNY | 03 | BELL $e^+ e^- \rightarrow \Upsilon(4S)$ |
|---------|----|-----------------------|----|---|

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 K^*(892)^0)/\Gamma_{\text{total}}$

Γ_{138}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-----------------------|------|---|
| $< 6.9 \times 10^{-5}$ | 90 | ¹ KROKOVNY | 03 | BELL $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 K^*(892)^0)/\Gamma_{\text{total}}$

Γ_{139}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-----------------------|------|---|
| $< 4.0 \times 10^{-5}$ | 90 | ¹ KROKOVNY | 03 | BELL $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 \pi^+ \pi^+ \pi^- \pi^-)/\Gamma_{\text{total}}$

Γ_{140}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---------|
| 2.7 ± 0.5 OUR AVERAGE | | | |

| | | | |
|--------------------------|-----------------------|----|---|
| $2.60 \pm 0.47 \pm 0.37$ | ¹ MAJUMDER | 04 | BELL $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $3.0 \pm 0.7 \pm 0.6$ | ¹ EDWARDS | 02 | CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

| $\Gamma(D^*(2007)^0 \pi^+ \pi^+ \pi^- \pi^-)/\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^- \pi^0)$ | Γ_{140}/Γ_{56} | | |
|---|----------------------------|------|----------------------------------|
| VALUE | DOCUMENT ID | TECN | COMMENT |
| 0.17±0.04±0.02 | ¹ EDWARDS 02 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

| $\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$ | Γ_{141}/Γ | | | |
|---|-----------------------|-------------|------|---------|
| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |

| 8.2±0.9 OUR AVERAGE | | | | |
|--|--|---------------------------|------|----------------------------------|
| 8.1±0.6±1.0 | | ¹ AUBERT,B 06A | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 8.1±0.8±1.1 | | ¹ MIYAKE 05 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 9.9 ^{+4.2} _{-3.3} ±1.2 | | ¹ LIPELES 00 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--|----|---------------------------|------|-------------------------|
| 8.3±1.6±1.2 | | ^{1,2} AUBERT 02M | BABR | Repl. by AUBERT,B 06B |
| 6.2 ^{+4.0} _{-2.9} ±1.0 | | ³ ARTUSO 99 | CLE2 | Repl. by LIPELES 00 |
| <61 | 90 | ⁴ BARATE 98Q | ALEP | $e^+ e^- \rightarrow Z$ |
| <22 | 90 | ⁵ ASNER 97 | CLE2 | Repl. by ARTUSO 99 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² AUBERT 02M also assumes the measured CP -odd fraction of the final states is $0.22 \pm 0.18 \pm 0.03$.

³ ARTUSO 99 uses $B(\gamma(4S) \rightarrow B^0 \bar{B}^0) = (48 \pm 4)\%$.

⁴ BARATE 98Q (ALEPH) observes 2 events with an expected background of 0.10 ± 0.03 which corresponds to a branching ratio of $(2.3^{+1.9}_{-1.2} \pm 0.4) \times 10^{-3}$.

⁵ ASNER 97 at CLEO observes 1 event with an expected background of 0.022 ± 0.011 . This corresponds to a branching ratio of $(5.3^{+7.1}_{-3.7} \pm 1.0) \times 10^{-4}$.

| $\Gamma(\bar{D}^*(2007)^0 \omega)/\Gamma_{\text{total}}$ | Γ_{142}/Γ | | | |
|--|-----------------------|-------------|------|---------|
| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |

| 2.7 ±0.8 OUR AVERAGE | | | | |
|-----------------------------|----|-------------------------|------|----------------------------------|
| 2.29±0.39±0.40 | | ¹ BLYTH 06 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 4.2 ±0.7 ±0.9 | 90 | ¹ AUBERT 04B | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|------------------------|------|----------------------------------|
| < 7.9 | 90 | ¹ ABE 02J | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 7.4 | 90 | ² NEMATI 98 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <21 | 90 | ³ ALAM 94 | CLE2 | Repl. by NEMATI 98 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² NEMATI 98 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+ \pi^-)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(D^*(2010)^+ D^-)/\Gamma_{\text{total}}$ Γ_{143}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

6.1±1.5 OUR AVERAGE

Error includes scale factor of 1.6.

5.7±0.7±0.7 ¹AUBERT,B 06A BABR $e^+ e^- \rightarrow \gamma(4S)$ 11.7±2.6^{+2.2}_{-2.5} 1,2 ABE 02Q BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.8±1.0±1.3 1 AUBERT 03J BABR Repl. by AUBERT,B 06B

14.8±3.8^{+2.8}_{-3.1} 1,3 ABE 02Q BELL $e^+ e^- \rightarrow \gamma(4S)$ < 6.3 90 ¹LIPELES 00 CLE2 $e^+ e^- \rightarrow \gamma(4S)$ <56 90 BARATE 98Q ALEP $e^+ e^- \rightarrow Z$ <18 90 ASNER 97 CLE2 $e^+ e^- \rightarrow \gamma(4S)$ ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² The measurement is performed using fully reconstructed D^* and D^+ decays.³ The measurement is performed using a partial reconstruction technique for the D^* and fully reconstructed D^+ decays as a cross check. $\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$ Γ_{144}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

< 0.9 90 ¹AUBERT,B 06A BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<270 90 BARATE 98Q ALEP $e^+ e^- \rightarrow Z$ ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D^- D^0 K^+)/\Gamma_{\text{total}}$ Γ_{145}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

1.7±0.3±0.3 ¹AUBERT 03X BABR $e^+ e^- \rightarrow \gamma(4S)$ ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D^- D^*(2007)^0 K^+)/\Gamma_{\text{total}}$ Γ_{146}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

4.6±0.7±0.7 ¹AUBERT 03X BABR $e^+ e^- \rightarrow \gamma(4S)$ ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D^*(2010)^- D^0 K^+)/\Gamma_{\text{total}}$ Γ_{147}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

3.1^{+0.4}_{-0.3}±0.4 ¹AUBERT 03X BABR $e^+ e^- \rightarrow \gamma(4S)$ ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D^*(2010)^- D^*(2007)^0 K^+)/\Gamma_{\text{total}}$ Γ_{148}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

11.8±1.0±1.7 ¹AUBERT 03X BABR $e^+ e^- \rightarrow \gamma(4S)$ ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(D^- D^+ K^0)/\Gamma_{\text{total}}$ Γ_{149}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| <1.7 | 90 | 1 AUBERT | 03X BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $[\Gamma(D^*(2010)^- D^+ K^0) + \Gamma(D^- D^*(2010)^+ K^0)]/\Gamma_{\text{total}}$ Γ_{150}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------------------------|
| $6.5 \pm 1.2 \pm 1.0$ | 1 AUBERT | 03X BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D^*(2010)^- D^*(2010)^+ K^0)/\Gamma_{\text{total}}$ Γ_{151}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| 7.8 ± 1.1 OUR AVERAGE | | | |

| | | | |
|-----------------------|--------------|----------|----------------------------------|
| $6.8 \pm 0.8 \pm 1.4$ | 1,2 DALSENO | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $8.8 \pm 0.8 \pm 1.4$ | 1,2 AUBERT,B | 06Q BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----------------------------|----------|----------|-----------------------|
| $8.8^{+1.5}_{-1.4} \pm 1.3$ | 1 AUBERT | 03X BABR | Repl. by AUBERT,B 06Q |
|-----------------------------|----------|----------|-----------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² The result is rescaled by a factor of 2 to convert from K_S^0 to K^0 . $\Gamma(D^{*-} D_{s1}(2536)^+ \times B(D_{s1}(2536)^+ \rightarrow D^+ K^0))/\Gamma_{\text{total}}$ Γ_{152}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| 8.0 ± 2.4 OUR AVERAGE | | | |

| | | | |
|-----------------------------|--------------|----------|----------------------------------|
| $7.6^{+4.8+1.6}_{-4.2-1.4}$ | 1,2 DALSENO | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $8.2 \pm 2.6 \pm 1.2$ | 1,2 AUBERT,B | 06Q BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² The result is rescaled by a factor of 2 to convert from K_S^0 to K^0 . $\Gamma(\bar{D}^0 D^0 K^0)/\Gamma_{\text{total}}$ Γ_{153}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| <1.4 | 90 | 1 AUBERT | 03X BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $[\Gamma(\bar{D}^0 D^*(2007)^0 K^0) + \Gamma(\bar{D}^*(2007)^0 D^0 K^0)]/\Gamma_{\text{total}}$ Γ_{154}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| <3.7 | 90 | 1 AUBERT | 03X BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\bar{D}^*(2007)^0 D^*(2007)^0 K^0)/\Gamma_{\text{total}}$ Γ_{155}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| <6.6 | 90 | 1 AUBERT | 03X BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma((\overline{D} + \overline{D}^*)(D + D^*)K)/\Gamma_{\text{total}}$ Γ_{156}/Γ

| <u>VALUE (units 10^{-2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|----------------------------------|
| 4.3±0.3±0.6 | ¹ AUBERT | 03X BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\eta_c K^0)/\Gamma_{\text{total}}$ Γ_{157}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| 0.89±0.16 OUR AVERAGE | | | |

| | | | |
|---------------------------------|-------------------------|-----------|----------------------------------|
| $0.64^{+0.22}_{-0.20} \pm 0.20$ | ^{1,2} AUBERT | 07AV BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.93 \pm 0.16 \pm 0.16$ | ^{1,3} AUBERT,B | 04B BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.23 \pm 0.23^{+0.40}_{-0.41}$ | ¹ FANG | 03 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.09^{+0.55}_{-0.42} \pm 0.33$ | ⁴ EDWARDS | 01 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² AUBERT 07AV reports $[B(B^0 \rightarrow \eta_c K^0)] \times [B(\eta_c(1S) \rightarrow p\bar{p})] = (0.83^{+0.28}_{-0.26} \pm 0.05) \times 10^{-6}$. We divide by our best value $B(\eta_c(1S) \rightarrow p\bar{p}) = (1.3 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AUBERT,B 04B reports $[B(B^0 \rightarrow \eta_c K^0)] \times [B(\eta_c(1S) \rightarrow K\bar{K}\pi)] = (0.0648 \pm 0.0085 \pm 0.0071) \times 10^{-3}$. We divide by our best value $B(\eta_c(1S) \rightarrow K\bar{K}\pi) = (7.0 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ EDWARDS 01 assumes equal production of B^0 and B^+ at the $\gamma(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma\eta_c)$ in those modes have been accounted for.

 $\Gamma(\eta_c K^0)/\Gamma(J/\psi(1S)K^0)$ $\Gamma_{157}/\Gamma_{159}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-----------------------|-------------|----------------------------------|
| 1.39±0.20±0.45 | ¹ AUBERT,B | 04B BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses BABAR measurement of $B(B^0 \rightarrow J/\psi K^0) = (8.5 \pm 0.5 \pm 0.6) \times 10^{-4}$. $\Gamma(\eta_c K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{158}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-------------------------------------|
| 0.96±0.33 OUR AVERAGE | | | Error includes scale factor of 1.1. |

| | | | |
|---------------------------------|-----------------------|-----------|----------------------------------|
| $0.79^{+0.25}_{-0.23} \pm 0.24$ | ^{1,2} AUBERT | 07AV BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.62 \pm 0.32^{+0.55}_{-0.60}$ | ² FANG | 03 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ AUBERT 07AV reports $[B(B^0 \rightarrow \eta_c K^*(892)^0)] \times [B(\eta_c(1S) \rightarrow p\bar{p})] = (1.03^{+0.27}_{-0.24} \pm 0.17) \times 10^{-6}$. We divide by our best value $B(\eta_c(1S) \rightarrow p\bar{p}) = (1.3 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\eta_c K^*(892)^0)/\Gamma(\eta_c K^0)$ $\Gamma_{158}/\Gamma_{157}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|----------------------------------|
| 1.33±0.36^{+0.24}_{-0.33} | FANG | 03 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

$\Gamma(J/\psi(1S)K^0)/\Gamma_{\text{total}}$ Γ_{159}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-------------|--------------------|-------------|----------------------------------|
| 8.71 ± 0.32 OUR AVERAGE | | | | | |
| 8.6 ± 1.3 | ± 0.3 | | 1,2 AUBERT | 07AV BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 8.69 ± 0.22 | ± 0.30 | | 2 AUBERT | 05J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 7.9 ± 0.4 | ± 0.9 | | 2 ABE | 03B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 9.5 ± 0.8 | ± 0.6 | | 2 AVERY | 00 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| 11.5 ± 2.3 | ± 1.7 | | 3 ABE | 96H CDF | $p\bar{p}$ at 1.8 TeV |
| 7.0 ± 4.1 | ± 0.1 | | 4 BORTOLETTO92 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |
| 9.3 ± 7.2 | ± 0.1 | 2 | 5 ALBRECHT | 90J ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 8.3 ± 0.4 | ± 0.5 | | 2 AUBERT | 02 BABR | Repl. by AUBERT 05J |
| 8.5 ± 1.4 | ± 0.6 | | 2 JESSOP | 97 CLE2 | Repl. by AVERY 00 |
| 7.5 ± 2.4 | ± 0.8 | 10 | 4 ALAM | 94 CLE2 | Sup. by JESSOP 97 |
| <50 | 90 | | ALAM | 86 CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ AUBERT 07AV reports $[B(B^0 \rightarrow J/\psi(1S)K^0)] \times [B(J/\psi(1S) \rightarrow p\bar{p})] = (1.87^{+0.28}_{-0.26} \pm 0.07) \times 10^{-6}$. We divide by our best value $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

³ ABE 96H assumes that $B(B^+ \rightarrow J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$.

⁴ BORTOLETTO 92 reports $(6 \pm 3 \pm 2) \times 10^{-4}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

⁵ ALBRECHT 90J reports $(8 \pm 6 \pm 2) \times 10^{-4}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)K^+\pi^-)/\Gamma_{\text{total}}$ Γ_{160}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-------------|--------------------|-------------|----------------------------------|
| $1.16 \pm 0.56 \pm 0.01$ | | | 1 BORTOLETTO92 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3 90 2 ALBRECHT 87D ARG $e^+ e^- \rightarrow \gamma(4S)$

<6.3 90 2 GILES 84 CLEO $e^+ e^- \rightarrow \gamma(4S)$

¹ BORTOLETTO 92 reports $(1.0 \pm 0.4 \pm 0.3) \times 10^{-3}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ALBRECHT 87D assume $B^+ B^- / B^0 \bar{B}^0$ ratio is 55/45. $K\pi$ system is specifically selected as nonresonant.

$\Gamma(J/\psi(1S)K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{161}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| 1.33 ± 0.06 OUR AVERAGE | | | | |
| 1.30 $+0.22$ -0.21 | ± 0.04 | 1,2 AUBERT | 07AV BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.309 ± 0.026 ± 0.077 | | 2 AUBERT | 05J BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.29 ± 0.05 ± 0.13 | | 2 ABE | 02N BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.74 ± 0.20 ± 0.18 | | 3 ABE | 980 CDF | $p\bar{p} 1.8 \text{ TeV}$ |
| 1.32 ± 0.17 ± 0.17 | | 4 JESSOP | 97 CLE2 | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.28 ± 0.66 ± 0.01 | 6 | 5 BORTOLETTO92 | CLEO | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1.28 ± 0.60 ± 0.01 | 6 | 6 ALBRECHT | 90J ARG | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 4.07 ± 1.82 ± 0.04 | 5 | 7 BEBEK | 87 CLEO | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.24 ± 0.05 ± 0.09 | | 2 AUBERT | 02 BABR | Repl. by AUBERT 05J |
| 1.36 ± 0.27 ± 0.22 | | 8 ABE | 96H CDF | Sup. by ABE 980 |
| 1.69 ± 0.31 ± 0.18 | 29 | 9 ALAM | 94 CLE2 | Sup. by JESSOP 97 |
| | | 10 ALBRECHT | 94G ARG | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 4.0 ± 0.30 | | 11 ALBAJAR | 91E UA1 | $E_{\text{cm}}^{p\bar{p}} = 630 \text{ GeV}$ |
| 3.3 ± 0.18 | 5 | 12 ALBRECHT | 87D ARG | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 4.1 ± 0.18 | 5 | 13 ALAM | 86 CLEO | Repl. by BEBEK 87 |

¹AUBERT 07AV reports $[B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] \times [B(J/\psi(1S) \rightarrow p\bar{p})] = (2.82^{+0.30+0.36}_{-0.28-0.35}) \times 10^{-6}$. We divide by our best value $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ ABE 980 reports $[B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)]/[B(B^+ \rightarrow J/\psi(1S)K^+)] = 1.76 \pm 0.14 \pm 0.15$. We multiply by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (9.9 \pm 1.0) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁵ BORTOLETTO 92 reports $(1.1 \pm 0.5 \pm 0.3) \times 10^{-3}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁶ ALBRECHT 90J reports $(1.1 \pm 0.5 \pm 0.2) \times 10^{-3}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁷ BEBEK 87 reports $(3.5 \pm 1.6 \pm 0.3) \times 10^{-3}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Updated in BORTOLETTO 92 to use the same assumptions.

⁸ ABE 96H assumes that $B(B^+ \rightarrow J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$.

⁹ The neutral and charged B events together are predominantly longitudinally polarized, $\Gamma_L/\Gamma = 0.080 \pm 0.08 \pm 0.05$. This can be compared with a prediction using HQET, 0.73 (KRAMER 92). This polarization indicates that the $B \rightarrow \psi K^*$ decay is dominated by the $CP = -1$ CP eigenstate. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹⁰ ALBRECHT 94G measures the polarization in the vector-vector decay to be predominantly longitudinal, $\Gamma_T/\Gamma = 0.03 \pm 0.16 \pm 0.15$ making the neutral decay a CP eigenstate when the K^{*0} decays through $K_S^0 \pi^0$.

¹¹ ALBAJAR 91E assumes B_d^0 production fraction of 36%.

¹² ALBRECHT 87D assume $B^+ B^- / B^0 \bar{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.

¹³ ALAM 86 assumes B^\pm / B^0 ratio is 60/40. The observation of the decay $B^+ \rightarrow J/\psi K^*(892)^+$ (HAAS 85) has been retracted in this paper.

$\Gamma(J/\psi(1S)K^*(892)^0)/\Gamma(J/\psi(1S)K^0)$

$\Gamma_{161}/\Gamma_{159}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|---------------------------------------|
| 1.50±0.09 OUR AVERAGE | | | |
| 1.51±0.05±0.08 | AUBERT | 05J | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| 1.39±0.36±0.10 | ABE | 96Q | CDF $p\bar{p}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1.49±0.10±0.08 | ¹ AUBERT | 02 | BABR Repl. by AUBERT 05J |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)\eta K_S^0)/\Gamma_{\text{total}}$

Γ_{162}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|----------------------------------|
| 8.4±2.6±2.7 | | | |
| ¹ AUBERT | 04Y | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)\eta' K_S^0)/\Gamma_{\text{total}}$

Γ_{163}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------------------|------|---------------------------------------|
| <2.5 | 90 | ¹ XIE | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)\phi K^0)/\Gamma_{\text{total}}$

Γ_{164}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------|------|---------------------------------------|
| (9.4±2.6) $\times 10^{-5}$ OUR AVERAGE | | | |
| (10.2±3.8±1.0) $\times 10^{-5}$ | ¹ AUBERT | 03O | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| (8.8 $^{+3.5}_{-3.0}$ ±1.3) $\times 10^{-5}$ | ² ANASTASSOV | 00 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ANASTASSOV 00 finds 10 events on a background of 0.5 ± 0.2 . Assumes equal production of B^0 and B^+ at the $\gamma(4S)$, a uniform Dalitz plot distribution, isotropic $J/\psi(1S)$ and ϕ decays, and $B(B^+ \rightarrow J/\psi(1S)\phi K^+) = B(B^0 \rightarrow J/\psi(1S)\phi K^0)$.

$\Gamma(J/\psi(1S)K(1270)^0)/\Gamma_{\text{total}}$

Γ_{165}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|----------------------------------|
| 1.30±0.34±0.32 | | | |
| ¹ ABE | 01L | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses the PDG value of $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.00 \pm 0.10) \times 10^{-3}$.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$

Γ_{166}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----------------------|-------------|------|----------------------------------|
| 2.05±0.24 OUR AVERAGE | | | | |
| 1.94±0.22±0.17 | ¹ AUBERT,B | 06B | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 2.3 ± 0.5 ± 0.2 | ¹ ABE | 03B | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 2.5 $^{+1.1}_{-0.9}$ ± 0.2 | ¹ AVERY | 00 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------|---------------------|------------------------|------|------------------------|
| 2.0 ± 0.6 ± 0.2 | ¹ AUBERT | 02 | BABR | Repl. by AUBERT, B 06B |
| < 32 | 90 | ² ACCIARRI | 97C | L3 |
| < 5.8 | 90 | BISHAI | 96 | CLE2 Sup. by AVERY 00 |
| <690 | 90 | ¹ ALEXANDER | 95 | CLE2 Sup. by BISHAI 96 |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ACCIARRI 97C assumes B^0 production fraction ($39.5 \pm 4.0\%$) and B_s ($12.0 \pm 3.0\%$).

$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$

Γ_{167}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|--------------------|------|---|
| 9.5±1.7±0.8 | | ¹ CHANG | 07A | BELL $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|-----------------------|-----|---|
| < 27 | 90 | ¹ AUBERT | 030 | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |
| <1200 | 90 | ² ACCIARRI | 97C | L3 |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ACCIARRI 97C assumes B^0 production fraction ($39.5 \pm 4.0\%$) and B_s ($12.0 \pm 3.0\%$).

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{168}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|------|---|
| (4.6±0.7±0.6) × 10⁻⁵ | ¹ AUBERT | 03B | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\pi^+\pi^- \text{ nonresonant})/\Gamma_{\text{total}}$

Γ_{169}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|---|
| <1.2 | 90 | ¹ AUBERT | 07AC | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)f_2)/\Gamma_{\text{total}}$

Γ_{170}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|---|
| <0.46 | 90 | ¹ AUBERT | 07AC | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\rho^0)/\Gamma_{\text{total}}$

Γ_{171}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|---|
| 2.7±0.3±0.2 | | ¹ AUBERT | 07AC | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------|---------------------|--------|------|---|
| 1.6±0.6±0.4 | ¹ AUBERT | 03B | BABR | Repl. by AUBERT 07AC |
| <25 | 90 | BISHAI | 96 | CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\omega)/\Gamma_{\text{total}}$

Γ_{172}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-------------|------|---|
| <2.7 × 10⁻⁴ | 90 | BISHAI | 96 | CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ |

$\Gamma(J/\psi(1S)\phi)/\Gamma_{\text{total}}$ Γ_{173}/Γ

| <i>VALUE</i> (units 10^{-6}) | <i>CL%</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|------------|---------------------|-------------|----------------------------------|
| <9.2 | 90 | ¹ AUBERT | 030 BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(J/\psi(1S)\eta'(958))/\Gamma_{\text{total}}$ Γ_{174}/Γ

| <i>VALUE</i> (units 10^{-5}) | <i>CL%</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|------------|---------------------|-------------|----------------------------------|
| <6.3 | 90 | ¹ AUBERT | 030 BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(J/\psi(1S)K^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{175}/Γ

| <i>VALUE</i> (units 10^{-4}) | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|-----------------------|-------------|--------------------|
| 10.3±3.3±1.5 | ¹ AFFOLDER | 02B CDF | $p\bar{p}$ 1.8 TeV |

¹ Uses $B^0 \rightarrow J/\psi(1S)K_S^0$ decay as a reference and $B(B^0 \rightarrow J/\psi(1S)K^0) = 8.3 \times 10^{-4}$.

 $\Gamma(J/\psi(1S)K^0\rho^0)/\Gamma_{\text{total}}$ Γ_{176}/Γ

| <i>VALUE</i> (units 10^{-4}) | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|-----------------------|-------------|--------------------|
| 5.4±2.9±0.9 | ¹ AFFOLDER | 02B CDF | $p\bar{p}$ 1.8 TeV |

¹ Uses $B^0 \rightarrow J/\psi(1S)K_S^0$ decay as a reference and $B(B^0 \rightarrow J/\psi(1S)K^0) = 8.3 \times 10^{-4}$.

 $\Gamma(J/\psi(1S)K^*(892)^+\pi^-)/\Gamma_{\text{total}}$ Γ_{177}/Γ

| <i>VALUE</i> (units 10^{-4}) | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|-----------------------|-------------|--------------------|
| 7.7±4.1±1.3 | ¹ AFFOLDER | 02B CDF | $p\bar{p}$ 1.8 TeV |

¹ Uses $B^0 \rightarrow J/\psi(1S)K_S^0$ decay as a reference and $B(B^0 \rightarrow J/\psi(1S)K^0) = 8.3 \times 10^{-4}$.

 $\Gamma(J/\psi(1S)K^*(892)^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{178}/Γ

| <i>VALUE</i> (units 10^{-4}) | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|-----------------------|-------------|--------------------|
| 6.6±1.9±1.1 | ¹ AFFOLDER | 02B CDF | $p\bar{p}$ 1.8 TeV |

¹ Uses $B^0 \rightarrow J/\psi(1S)K^{*(892)}{}^0$ decay as a reference and $B(B^0 \rightarrow J/\psi(1S)K^0) = 12.4 \times 10^{-4}$.

 $\Gamma(X(3872)^-K^+)/\Gamma_{\text{total}}$ Γ_{179}/Γ

| <i>VALUE</i> | <i>CL%</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|--------------------------------|------------|---------------------|-------------|----------------------------------|
| <5 × 10⁻⁴ | 90 | ¹ AUBERT | 06E BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Perform measurements of absolute branching fractions using a missing mass technique.

 $\Gamma(X(3872)^-K^+ \times B(X(3872)^- \rightarrow J/\psi(1S)\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{180}/Γ

| <i>VALUE</i> (units 10^{-6}) | <i>CL%</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|------------|---------------------|-------------|----------------------------------|
| <5.4 | 90 | ¹ AUBERT | 05B BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. The isovector-X hypothesis is excluded with a likelihood test at 1×10^{-4} level.

$\Gamma(X(3872)K^0 \times B(X \rightarrow J/\psi \pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{181}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------------------------|
| <10.3 | 90 | 1,2 AUBERT | 06 BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ The lower limit is also given to be 1.34×10^{-6} at 90% CL.

² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(X(3872)K^0 \times B(X \rightarrow D^0 \bar{D}^0 \pi^0))/\Gamma_{\text{total}}$ Γ_{182}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|----------------------------------|
| 1.66 ± 0.70 +0.32 -0.37 | 1 GOKHROO | 06 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Measure the near-threshold enhancements in the $(D^0 \bar{D}^0 \pi^0)$ system at a mass $3875.2 \pm 0.7^{+0.3}_{-1.6} \pm 0.8$ MeV/c².

 $\Gamma(X(3872)K^0 \times B(X \rightarrow \bar{D}^{*0} D^0))/\Gamma_{\text{total}}$ Γ_{183}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------------------------|
| <4.37 | 90 | 1 AUBERT | 08B BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(J/\psi(1S)p\bar{p})/\Gamma_{\text{total}}$ Γ_{184}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------|--------------------|-------------|----------------------------------|
| <8.3 × 10⁻⁷ | 90 | 1 XIE | 05 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------|--------------------|-------------|----------------------------------|
| <1.9 × 10⁻⁶ | 90 | 1 AUBERT | 03K BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(J/\psi(1S)\gamma)/\Gamma_{\text{total}}$ Γ_{185}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------------------------|
| <1.6 | 90 | 1 AUBERT,B | 04T BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(J/\psi(1S)\bar{D}^0)/\Gamma_{\text{total}}$ Γ_{186}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------------------------|
| <1.3 | 90 | 1 AUBERT | 05U BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------------------------|
| <2.0 | 90 | 1 ZHANG | 05B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\psi(2S)K^0)/\Gamma_{\text{total}}$ Γ_{187}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------------------------|
| 6.2 ± 0.6 OUR AVERAGE | | | | |
| 6.46 ± 0.65 ± 0.51 | | 1 AUBERT | 05J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 6.7 ± 1.1 | | 1 ABE | 03B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 5.0 ± 1.1 ± 0.6 | | 1 RICHICHI | 01 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|-----|-----------|-----------|---------------------------|------|------|----------------------------------|
| 6.9 | ± 1.1 | ± 1.1 | ¹ AUBERT | 02 | BABR | Repl. by AUBERT 05J |
| < 8 | | 90 | ¹ ALAM | 94 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <15 | | 90 | ¹ BORTOLETTO92 | CLEO | | $e^+ e^- \rightarrow \gamma(4S)$ |
| <28 | | 90 | ¹ ALBRECHT | 90J | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\psi(3770)K^0 \times B(\psi \rightarrow \bar{D}^0 D^0))/\Gamma_{\text{total}}$ Γ_{188}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|---------------------------------------|
| <1.23 | 90 | ¹ AUBERT | 08B | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\psi(3770)K^0 \times B(\psi \rightarrow D^- D^+))/\Gamma_{\text{total}}$ Γ_{189}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|---------------------------------------|
| <1.88 | 90 | ¹ AUBERT | 08B | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\psi(2S)K^0)/\Gamma(J/\psi(1S)K^0)$ $\Gamma_{187}/\Gamma_{159}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|---------------------|------|---------------------------------------|
| 0.82 ± 0.13 ± 0.12 | ¹ AUBERT | 02 | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\psi(2S)K^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{190}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|-----------------------|------|--------------------------------------|
| <0.001 | 90 | ¹ ALBRECHT | 90J | ARG $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\psi(2S)K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{191}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----|-----------------------|------|---------------------------------------|
| 7.2 ± 0.8 OUR AVERAGE | | | | |
| 6.49 ± 0.59 ± 0.97 | | ¹ AUBERT | 05J | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| 7.6 ± 1.1 ± 1.0 | | ¹ RICHICHI | 01 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| 9.0 ± 2.2 ± 0.9 | | ² ABE | 980 | CDF $p\bar{p}$ 1.8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------------|----|---------------------------|------|------|----------------------------------|
| <19 | 90 | ¹ ALAM | 94 | CLE2 | Repl. by RICHICHI 01 |
| 14 ± 8 ± 4 | | ¹ BORTOLETTO92 | CLEO | | $e^+ e^- \rightarrow \gamma(4S)$ |
| <23 | 90 | ¹ ALBRECHT | 90J | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ABE 980 reports $[B(B^0 \rightarrow \psi(2S)K^*(892)^0)]/[B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.908 \pm 0.194 \pm 0.10$. We multiply by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (9.9 \pm 1.0) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\psi(2S)K^*(892)^0)/\Gamma(\psi(2S)K^0)$ $\Gamma_{191}/\Gamma_{187}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|------|---------------------------------------|
| 1.00 ± 0.14 ± 0.09 | AUBERT | 05J | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

$\Gamma(\chi_{c0}(1P)K^0)/\Gamma_{\text{total}}$ Γ_{192}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|----------------------|-------------|---------------------------------------|
| < 1.13 | 90 | ¹ GARMASH | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <12.4 | 90 | ² AUBERT | 05K | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| < 5.0 | 90 | ³ EDWARDS | 01 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.³ EDWARDS 01 assumes equal production of B^0 and B^+ at the $\gamma(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma \eta_c)$ in those modes have been accounted for. $\Gamma(\chi_{c0}K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{193}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|------------|---------------------|-------------|---------------------------------------|
| < 7.7 × 10⁻⁴ | 90 | ¹ AUBERT | 05K | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\chi_{c2}K^0)/\Gamma_{\text{total}}$ Γ_{194}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------|-------------|---------------------------------------|
| < 2.6 × 10⁻⁵ | 90 | ¹ SONI | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <4.1 × 10 ⁻⁵ | 90 | ¹ AUBERT | 05K | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\chi_{c2}K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{195}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------|-------------|---------------------------------------|
| < 3.6 × 10⁻⁵ | 90 | ¹ AUBERT | 05K | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <7.1 × 10 ⁻⁵ | 90 | ¹ SONI | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\chi_{c1}(1P)K^0)/\Gamma_{\text{total}}$ Γ_{196}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------|-------------|---------------------------------------|
| 3.9 ± 0.4 OUR AVERAGE | | | | |
| 3.51 ± 0.33 ± 0.45 | | ¹ SONI | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| 4.53 ± 0.41 ± 0.51 | | ¹ AUBERT | 05J | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| 3.0 $\begin{array}{l} +1.5 \\ -1.0 \end{array}$ ± 0.2 | | ² AVERY | 00 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 4.1 ± 1.3 ± 0.2 | | ³ AUBERT | 02 | BABR Repl. by AUBERT 05J |
| <27 | 90 | ¹ ALAM | 94 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² AVERY 00 reports $(3.9 \begin{array}{l} +1.9 \\ -1.3 \end{array} \pm 0.4) \times 10^{-4}$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$.We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (36.0 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.³ AUBERT 02 reports $(5.4 \pm 1.4 \pm 1.1) \times 10^{-4}$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$.We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (36.0 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\chi_{c1}(1P) K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{197}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------|-------------|---------------------------------------|
| 3.2 ± 0.6 OUR AVERAGE | | | | |
| 3.14 ± 0.34 ± 0.72 | | ¹ SONI | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| 3.27 ± 0.42 ± 0.64 | | ¹ AUBERT | 05J | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 3.6 ± 1.2 ± 0.2 | | ² AUBERT | 02 | BABR Repl. by AUBERT 05J |
| <21 | 90 | ³ ALAM | 94 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² AUBERT 02 reports $(4.8 \pm 1.4 \pm 0.9) \times 10^{-4}$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (36.0 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.³ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\chi_{c1}(1P) K^0)/\Gamma(J/\psi(1S) K^0)$ $\Gamma_{196}/\Gamma_{159}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|---------------------------------------|
| 0.50±0.15±0.03 | ¹ AUBERT | 02 | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1 AUBERT 02 reports $0.66 \pm 0.11 \pm 0.17$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (36.0 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. | | | |

 $\Gamma(\chi_{c1}(1P) K^*(892)^0)/\Gamma(\chi_{c1}(1P) K^0)$ $\Gamma_{197}/\Gamma_{196}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|---------------------------------------|
| 0.72±0.11±0.12 | AUBERT | 05J | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.89 ± 0.34 ± 0.17 | ¹ AUBERT | 02 | BABR Repl. by AUBERT 05J |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. | | | |

 $\Gamma(K^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{198}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-----------------------------|-------------|---------------------------------------|
| 19.4± 0.6 OUR AVERAGE | | | | |
| 19.1 ± 0.6 ± 0.6 | | ¹ AUBERT | 07B | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| 19.9 ± 0.4 ± 0.8 | | ¹ LIN | 07A | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| 18.0 ± 2.3 ± 1.2 | | ¹ BORNHEIM | 03 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 18.5 ± 1.0 ± 0.7 | | ¹ CHAO | 04 | BELL Repl. by LIN 07A |
| 17.9 ± 0.9 ± 0.7 | | ¹ AUBERT | 02Q | BABR Repl. by AUBERT 07B |
| 22.5 ± 1.9 ± 1.8 | | ¹ CASEY | 02 | BELL Repl. by CHAO 04 |
| 19.3 ± 3.4 ± 1.5 | | ¹ ABE | 01H | BELL Repl. by CASEY 02 |
| 16.7 ± 1.6 ± 1.3 | | ¹ AUBERT | 01E | BABR Repl. by AUBERT 02Q |
| < 66 | 90 | ² ABE | 00C | SLD $e^+ e^- \rightarrow Z$ |
| 17.2 ± 2.5 ± 1.2 | | ¹ CRONIN-HEN..00 | CLE2 | Repl. by BORNHEIM 03 |
| 15 ± 5 ± 1.4 | | GODANG | 98 | CLE2 Repl. by CRONIN-HENNESSY 00 |

| | | | | | |
|-------|-------------------------|-----------------------|-----|------|----------------------------------|
| 24 | $\frac{+17}{-11} \pm 2$ | ³ ADAM | 96D | DLPH | $e^+ e^- \rightarrow Z$ |
| < 17 | 90 | ASNER | 96 | CLE2 | Sup. by ADAM 96D |
| < 30 | 90 | ⁴ BUSKULIC | 96V | ALEP | $e^+ e^- \rightarrow Z$ |
| < 90 | 90 | ⁵ ABREU | 95N | DLPH | Sup. by ADAM 96D |
| < 81 | 90 | ⁶ AKERS | 94L | OPAL | $e^+ e^- \rightarrow Z$ |
| < 26 | 90 | ⁷ BATTLE | 93 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 180 | 90 | ALBRECHT | 91B | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 90 | 90 | ⁸ AVERY | 89B | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 320 | 90 | AVERY | 87 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

³ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

⁴ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

⁵ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B_s^0 decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

⁶ Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and B_d^0 (B_s^0) fraction 39.5% (12%).

⁷ BATTLE 93 assumes equal production of $B^0\bar{B}^0$ and $B^+\bar{B}^-$ at $\gamma(4S)$.

⁸ Assumes the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$.

$\Gamma(K^+\pi^-)/\Gamma(K^0\pi^0)$

$\Gamma_{198}/\Gamma_{199}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|------------------|------|---------------------------------------|
| 2.16 ± 0.16 ± 0.16 | LIN | 07A | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1.20 $^{+0.50 + 0.22}_{-0.58 - 0.32}$ | ¹ ABE | 01H | BELL Repl. by LIN 07A |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$[\Gamma(K^+\pi^-) + \Gamma(\pi^+\pi^-)]/\Gamma_{\text{total}}$

$(\Gamma_{198} + \Gamma_{284})/\Gamma$

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|---------------------|------|---------------------------------------|
| 19 ± 6 OUR AVERAGE | | | | |
| 28 $^{+15}_{-10} \pm 20$ | | ¹ ADAM | 96D | DLPH $e^+ e^- \rightarrow Z$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 18 $^{+6 + 3}_{-5 - 4}$ | 17.2 | ASNER | 96 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| 24 $^{+8 \pm 2}_{-7}$ | | ² BATTLE | 93 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |

¹ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

² BATTLE 93 assumes equal production of $B^0\bar{B}^0$ and $B^+\bar{B}^-$ at $\gamma(4S)$.

$\Gamma(K^0\pi^0)/\Gamma_{\text{total}}$ Γ_{199}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-----------------------------|-------------|----------------------------------|
| 9.8 ± 0.6 OUR AVERAGE | | | | |
| 10.3 $\pm 0.7 \pm 0.6$ | | ¹ AUBERT 08E | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 9.2 $\pm 0.7 \pm 0.6$ | | ¹ LIN 07A | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $12.8^{+4.0}_{-3.3}{}^{+1.7}_{-1.4}$ | | ¹ BORNHEIM 03 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 11.4 $\pm 0.9 \pm 0.6$ | | ¹ AUBERT 05Y | BABR | Repl. by AUBERT 08E |
| 11.4 $\pm 1.7 \pm 0.8$ | | ¹ AUBERT 04M | BABR | Repl. by AUBERT 05Y |
| $11.7^{+2.3}_{-1.3}{}^{+1.2}_{-1.3}$ | | ¹ CHAO 04 | BELL | Repl. by LIN 07A |
| $8.0^{+3.3}_{-3.1}{}^{+1.6}_{-1.6}$ | | ¹ CASEY 02 | BELL | Repl. by CHAO 04 |
| $16.0^{+7.2}_{-5.9}{}^{+2.5}_{-2.7}$ | | ¹ ABE 01H | BELL | Repl. by CASEY 02 |
| $8.2^{+3.1}_{-2.7}{}^{+1.2}_{-1.2}$ | | ¹ AUBERT 01E | BABR | Repl. by AUBERT 04M |
| $14.6^{+5.9}_{-5.1}{}^{+2.4}_{-3.3}$ | | ¹ CRONIN-HEN..00 | CLE2 | Repl. by BORNHEIM 03 |
| <41 | 90 | GODANG 98 | CLE2 | Repl. by CRONIN-HENNESSY 00 |
| <40 | 90 | ASNER 96 | CLE2 | Rep. by GODANG 98 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\eta' K^0)/\Gamma_{\text{total}}$ Γ_{200}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------------|-------------|----------------------------------|
| 65 ± 4 OUR AVERAGE | | | |
| Error includes scale factor of 1.2. | | | |
| 66.6 $\pm 2.6 \pm 2.8$ | ¹ AUBERT 07AE | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $58.9^{+3.6}_{-3.5} \pm 4.3$ | ¹ SCHUEMANN 06 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $89^{+18}_{-16} \pm 9$ | ¹ RICHICHI 00 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 67.4 $\pm 3.3 \pm 3.2$ | ¹ AUBERT 05M | BABR | AUBERT 07AE |
| 60.6 $\pm 5.6 \pm 4.6$ | ¹ AUBERT 03W | BABR | Repl. by AUBERT 05M |
| $55^{+19}_{-16} \pm 8$ | ¹ ABE 01M | BELL | Repl. by SCHUEMANN 06 |
| $42^{+13}_{-11} \pm 4$ | ¹ AUBERT 01G | BABR | Repl. by AUBERT 03W |
| $47^{+27}_{-20} \pm 9$ | BEHRENS 98 | CLE2 | Repl. by RICHICHI 00 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\eta' K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{201}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------------|-------------|---------------------------------------|
| $3.8 \pm 1.1 \pm 0.5$ | | | | |
| ¹ AUBERT 07E | | | | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 2.6 | 90 | ¹ SCHUEMANN 07 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 7.6 | 90 | ¹ AUBERT,B 04D | BABR | Repl. by AUBERT 07E |
| <24 | 90 | ¹ RICHICHI 00 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <39 | 90 | BEHRENS 98 | CLE2 | Repl. by RICHICHI 00 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\eta K^0)/\Gamma_{\text{total}}$ Γ_{202}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-----------------------|-------------|---------------------------------------|
| < 1.9 | 90 | ¹ CHANG | 07B | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 2.9 | 90 | ¹ AUBERT,B | 06V | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| < 2.5 | 90 | ¹ AUBERT,B | 05K | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| < 2.0 | 90 | ¹ CHANG | 05A | BELL Repl. by CHANG 07B |
| < 5.2 | 90 | ¹ AUBERT | 04H | BABR Repl. by AUBERT,B 05K |
| < 9.3 | 90 | ¹ RICHICHI | 00 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| <33 | 90 | BEHRENS | 98 | CLE2 Repl. by RICHICHI 00 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\eta K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{203}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-----------------------|-------------|---------------------------------------|
| 15.9 ± 1.0 OUR AVERAGE | | | | |
| 15.2 $\pm 1.2 \pm 1.0$ | | ¹ WANG | 07B | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| 16.5 $\pm 1.1 \pm 0.8$ | | ¹ AUBERT,B | 06H | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| $13.8^{+5.5}_{-4.6} \pm 1.6$ | | ¹ RICHICHI | 00 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 18.6 $\pm 2.3 \pm 1.2$ | | ¹ AUBERT,B | 04D | BABR Repl. by AUBERT,B 06H |
| <30 | 90 | BEHRENS | 98 | CLE2 Repl. by RICHICHI 00 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\eta K_0^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{204}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|---------------------------------------|
| $11.0 \pm 1.6 \pm 1.5$ | ¹ AUBERT,B | 06H | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\eta K_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{205}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|---------------------------------------|
| $9.6 \pm 1.8 \pm 1.1$ | ¹ AUBERT,B | 06H | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\omega K^0)/\Gamma_{\text{total}}$ Γ_{206}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-----------------------|-------------|---------------------------------------|
| 5.0 ± 0.6 OUR AVERAGE | | | | |
| 5.4 $\pm 0.8 \pm 0.3$ | | ¹ AUBERT | 07AE | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| $4.4^{+0.8}_{-0.7} \pm 0.4$ | | ¹ JEN | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| $10.0^{+5.4}_{-4.2} \pm 1.4$ | | ¹ JESSOP | 00 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 6.2 $\pm 1.0 \pm 0.4$ | | ¹ AUBERT,B | 06E | BABR Repl. by AUBERT 07AE |
| $5.9^{+1.6}_{-1.3} \pm 0.5$ | | ¹ AUBERT | 04H | BABR Repl. by AUBERT,B 06E |
| $4.0^{+1.9}_{-1.6} \pm 0.5$ | | ¹ WANG | 04A | BELL Repl. by JEN 06 |
| <13 | 90 | ¹ AUBERT | 01G | BABR Repl. by AUBERT 04H |
| <57 | 90 | ¹ BERGFELD | 98 | CLE2 Repl. by JESSOP 00 |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_0(980)^{\pm} K^{\mp} \times B(a_0(980)^{\pm} \rightarrow \eta\pi^{\pm})) / \Gamma_{\text{total}}$ Γ_{208}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|--|
| <1.9 | 90 | ¹ AUBERT | 07Y | BABR $e^+e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|------------------------|----|--------------------------|
| <2.1 | 90 | ¹ AUBERT,BE | 04 | BABR Repl. by AUBERT 07Y |
|------|----|------------------------|----|--------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_0(1450)^{\pm} K^{\mp} \times B(a_0(1450)^{\pm} \rightarrow \eta\pi^{\pm})) / \Gamma_{\text{total}}$ Γ_{209}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|--|
| <3.1 | 90 | ¹ AUBERT | 07Y | BABR $e^+e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_0(980)^0 K^0 \times B(a_0(980)^0 \rightarrow \eta\pi^0)) / \Gamma_{\text{total}}$ Γ_{207}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------------------------|------|--|
| <7.8 | 90 | ¹ AUBERT,BE | 04 | BABR $e^+e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of charged and neutral B mesons from $\Upsilon(4S)$ decays.

$\Gamma(K_S^0 X^0 (\text{Familon})) / \Gamma_{\text{total}}$ Γ_{210}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|--------------------|------|--|
| <53 | 90 | ¹ AMMAR | 01B | CLE2 $e^+e^- \rightarrow \Upsilon(4S)$ |

¹ AMMAR 01B searched for the two-body decay of the B meson to a massless neutral feebly-interacting particle X^0 such as the familon, the Nambu-Goldstone boson associated with a spontaneously broken global family symmetry.

$\Gamma(\omega K^*(892)^0) / \Gamma_{\text{total}}$ Γ_{211}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-----------------------|------|--|
| < 4.2 | 90 | ¹ AUBERT,B | 06T | BABR $e^+e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|-----------------------|-----|----------------------------|
| < 6.0 | 90 | ¹ AUBERT | 050 | BABR Repl. by AUBERT,B 06T |
| <23 | 90 | ¹ BERGFELD | 98 | CLE2 |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ K^-) / \Gamma_{\text{total}}$ Γ_{212}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------------------|------|--|
| < 0.41 | 90 | ¹ LIN | 07 | BELL $e^+e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|--------------------------|-----|--|
| < 0.5 | 90 | ¹ AUBERT | 07B | BABR $e^+e^- \rightarrow \Upsilon(4S)$ |
| < 1.8 | 90 | ² ABULENCIA,A | 06D | CDF $p\bar{p}$ at 1.96 TeV |
| < 0.37 | 90 | ABE | 05G | BELL Repl. by LIN 07 |
| < 0.7 | 90 | CHAO | 04 | BELL $e^+e^- \rightarrow \Upsilon(4S)$ |
| < 0.8 | 90 | ¹ BORNHEIM | 03 | CLE2 $e^+e^- \rightarrow \Upsilon(4S)$ |
| < 0.6 | 90 | ¹ AUBERT | 02Q | BABR $e^+e^- \rightarrow \Upsilon(4S)$ |
| < 0.9 | 90 | ¹ CASEY | 02 | BELL $e^+e^- \rightarrow \Upsilon(4S)$ |
| < 2.7 | 90 | ¹ ABE | 01H | BELL $e^+e^- \rightarrow \Upsilon(4S)$ |

| | | | | | |
|-------|----|--------------------------|-----|------|----------------------------------|
| < 2.5 | 90 | ¹ AUBERT | 01E | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 66 | 90 | ³ ABE | 00C | SLD | $e^+ e^- \rightarrow Z$ |
| < 1.9 | 90 | ¹ CRONIN-HEN. | .00 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 4.3 | 90 | GODANG | 98 | CLE2 | Repl. by CRONIN-HENNESSY 00 |
| < 46 | | ⁴ ADAM | 96D | DLPH | $e^+ e^- \rightarrow Z$ |
| < 4 | 90 | ASNER | 96 | CLE2 | Repl. by GODANG 98 |
| < 18 | 90 | ⁵ BUSKULIC | 96V | ALEP | $e^+ e^- \rightarrow Z$ |
| <120 | 90 | ⁶ ABREU | 95N | DLPH | Sup. by ADAM 96D |
| < 7 | 90 | ¹ BATTLE | 93 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ABULENCIA,A 06D obtains this from $\Gamma(K^+ K^-)/\Gamma(K^+ \pi^-) < 0.10$ at 90% CL, assuming $B(B^0 \rightarrow K^+ \pi^-) = (18.9 \pm 0.7) \times 10^{-6}$.

³ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

⁴ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

⁵ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

⁶ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B_s^0 decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

$\Gamma(K^0 \bar{K}^0)/\Gamma_{\text{total}}$

Γ_{213}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------|
|---|------------|--------------------|-------------|----------------|

$0.96^{+0.20}_{-0.18}$ OUR AVERAGE

| | | | | | |
|--|----|------------------------|-----|------|----------------------------------|
| $0.87^{+0.25}_{-0.20} \pm 0.09$ | | ¹ LIN | 07 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.08 \pm 0.28 \pm 0.11$ | | ¹ AUBERT,BE | 06C | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.8 $\pm 0.3 \pm 0.9$ | | ¹ ABE | 05G | BELL | Repl. by LIN 07 |
| $1.19^{+0.40}_{-0.35} \pm 0.13$ | | ¹ AUBERT,BE | 05E | BABR | Repl. by AUBERT,BE 06C |
| < 1.8 | 90 | ¹ AUBERT | 04M | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 1.5 | 90 | ¹ CHAO | 04 | BELL | Repl. by ABE 05G |
| < 3.3 | 90 | ¹ BORNHEIM | 03 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 4.1 | 90 | ¹ CASEY | 02 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| <17 | 90 | GODANG | 98 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{214}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
|---|--------------------|-------------|----------------|

$6.2^{+1.2}_{-1.1}$ OUR AVERAGE Error includes scale factor of 1.3.

| | | | | | |
|-----------------------------|--|-----------------------|----|------|----------------------------------|
| $6.9^{+0.9}_{-0.8} \pm 0.6$ | | ¹ AUBERT,B | 05 | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $4.2^{+1.6}_{-1.3} \pm 0.8$ | | ¹ GARMASH | 04 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K_S^0 K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{215}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-----------------------|-------------|----------------------------------|
| <16 | 90 | ¹ AUBERT,B | 06R BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{216}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|----------------------------------|
| $36.6^{+4.2}_{-4.3} \pm 3.0$ | | ¹ CHANG | 04 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----|----|----------------------|---------|----------------------------------|
| <40 | 90 | ¹ ECKHART | 02 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
|-----|----|----------------------|---------|----------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K^+ \rho^-)/\Gamma_{\text{total}}$ Γ_{217}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|-------------------------------------|
| 8.5 ± 2.8 OUR AVERAGE | | | | Error includes scale factor of 1.7. |

$15.1^{+3.4+2.4}_{-3.3-2.6}$ ¹CHANG 04 BELL $e^+ e^- \rightarrow \gamma(4S)$

$7.3^{+1.3}_{-1.2} \pm 1.3$ ¹AUBERT 03T BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----|----|---------------------|---------|----------------------------------|
| <32 | 90 | ¹ JESSOP | 00 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <35 | 90 | ASNER | 96 CLE2 | Repl. by JESSOP 00 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma((K^+ \pi^- \pi^0) \text{ non-resonant})/\Gamma_{\text{total}}$ Γ_{218}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| <9.4 | 90 | ¹ CHANG | 04 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K_x^{*0} \pi^0)/\Gamma_{\text{total}}$ Γ_{219}/Γ

K_x^{*0} stands for the possible candidates of $K^*(1410)$, $K_0^*(1430)$ and $K_2^*(1430)$.

| <u>VALUE (units 10^{-6})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------------------------|
| $6.1^{+1.6+0.5}_{-1.5-0.6}$ | ¹ CHANG | 04 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K^0 \pi^+ \pi^- \text{ charmless})/\Gamma_{\text{total}}$ Γ_{220}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|----------------|
| 44.8 ± 2.6 OUR AVERAGE | | | | |

$47.5 \pm 2.4 \pm 3.7$ ¹GARMASH 07 BELL $e^+ e^- \rightarrow \gamma(4S)$

$43.0 \pm 2.3 \pm 2.3$ ²AUBERT 06I BABR $e^+ e^- \rightarrow \gamma(4S)$

$50 \pm 10 \pm 7$ ²ECKHART 02 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$43.7 \pm 3.8 \pm 3.4$ ²AUBERT,B 04O BABR Repl. by AUBERT 06I

$45.4 \pm 5.2 \pm 5.9$ ²GARMASH 04 BELL Repl. by GARMASH 07

| | | | | |
|------|----|----------|---------|----------------------------------|
| <440 | 90 | ALBRECHT | 91E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
|------|----|----------|---------|----------------------------------|

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(K^0 \pi^+ \pi^- \text{ non-resonant})/\Gamma_{\text{total}}$ Γ_{221}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------------------|-------------|------------------------------------|
| $19.9 \pm 2.5^{+1.7}_{-2.0}$ | ¹ GARMASH 07 | BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays. $\Gamma(K^0 \rho^0)/\Gamma_{\text{total}}$ Γ_{222}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------------------|-------------|------------------------------------|
| 5.4 ± 0.9 OUR AVERAGE | | | | |
| $4.9 \pm 0.8 \pm 0.9$ | 90 | ¹ AUBERT 07F | BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $6.1 \pm 1.0^{+1.1}_{-1.2}$ | 90 | ² GARMASH 07 | BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---------|----|--------------------|-----|------|------------------------------------|
| < 39 | 90 | ASNER | 96 | CLEO | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| < 320 | 90 | ALBRECHT | 91B | ARG | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| < 500 | 90 | ³ AVERY | 89B | CLEO | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| < 64000 | 90 | ⁴ AVERY | 87 | CLEO | $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.² Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.³ Avery 89B reports $< 5.8 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.⁴ Avery 87 reports < 0.08 assuming the $\Upsilon(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%. $\Gamma(K^0 f_0(980))/\Gamma_{\text{total}}$ Γ_{223}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------------------|-------------|------------------------------------|
| $5.5 \pm 0.7 \pm 0.6$ | 90 | ¹ AUBERT 06I | BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|------------------------|------|------------------------------------|
| < 360 | 90 | ² AVERY 89B | CLEO | $e^+ e^- \rightarrow \Upsilon(4S)$ |
|-------|----|------------------------|------|------------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.² Avery 89B reports $< 4.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%. $\Gamma(K^*(892)^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{224}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-------------------------------------|-------------|------------------------------------|
| 9.8 ± 1.3 OUR AVERAGE | | Error includes scale factor of 1.2. | | |
| $8.4 \pm 1.1^{+1.0}_{-0.9}$ | | ¹ GARMASH 07 | BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $11.0 \pm 1.5 \pm 0.71$ | | ² AUBERT 06I | BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $16 \begin{array}{l} +6 \\ -5 \end{array} \pm 2$ | | ² ECKHART 02 | CLE2 | $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------------------------|-----------------------|--------------------|------|---------------------------------------|
| $12.9 \pm 2.4 \pm 1.4$ | ² AUBERT,B | 040 | BABR | Repl. by AUBERT 06I |
| $14.8^{+4.6}_{-4.4}{}^{+2.8}_{-1.3}$ | ² CHANG | 04 | BELL | Repl. by GARMASH 07 |
| < 72 | 90 | ASNER | 96 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| <620 | 90 | ALBRECHT | 91B | ARG $e^+ e^- \rightarrow \gamma(4S)$ |
| <380 | 90 | ³ Avery | 89B | CLEO $e^+ e^- \rightarrow \gamma(4S)$ |
| <560 | 90 | ⁴ Avery | 87 | CLEO $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

³ Avery 89B reports $< 4.4 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

⁴ Avery 87 reports $< 7 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(1430)^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{225}/Γ

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT |
|------------------------------|----------------------|------|---------------------------------------|
| $49.7 \pm 3.8^{+6.8}_{-8.2}$ | ¹ GARMASH | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

$\Gamma(K_x^* \pi^-)/\Gamma_{\text{total}}$

Γ_{226}/Γ

K_x^{*+} stands for the possible candidates of $K^*(1410)$, $K_0^*(1430)$ and $K_2^*(1430)$.

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|--------------------|------|---------------------------------------|
| $5.1 \pm 1.5^{+0.6}_{-0.7}$ | ¹ CHANG | 04 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(1410)^+ \pi^- \times B(K^*(1410)^+ \rightarrow K^0 \pi^+))/\Gamma_{\text{total}}$

Γ_{227}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|----------------------|------|---------------------------------------|
| <3.8 | 90 | ¹ GARMASH | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

$\Gamma(K^*(1680)^+ \pi^- \times B(K^*(1680)^+ \rightarrow K^0 \pi^+))/\Gamma_{\text{total}}$

Γ_{228}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|----------------------|------|---------------------------------------|
| <2.6 | 90 | ¹ GARMASH | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

$\Gamma(K_2^*(1430)^+ \pi^- \times B(K_2^*(1430)^+ \rightarrow K^0 \pi^+))/\Gamma_{\text{total}}$

Γ_{229}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|----------------------|------|---------------------------------------|
| <2.1 | 90 | ¹ GARMASH | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

$\Gamma(f_0(980) K^0 \times B(f_0(980) \rightarrow \pi^+ \pi^-))/\Gamma_{\text{total}}$

Γ_{230}/Γ

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|----------------------|------|---------------------------------------|
| $7.6 \pm 1.7^{+0.9}_{-1.3}$ | ¹ GARMASH | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

$\Gamma(f_2(1270)K^0 \times B(f_2(1270) \rightarrow \pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{231}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|-------------------------|-------------|----------------------------------|
| <1.4 | 90 | ¹ GARMASH 07 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

 $\Gamma(K^*(892)^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{232}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------|-----------------------|-------------|----------------------------------|
| <3.5 × 10⁻⁶ | 90 | ¹ CHANG 04 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----|-----------|------|----------------------------------|
| $<3.6 \times 10^{-6}$ | 90 | JESSOP 00 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| $<2.8 \times 10^{-5}$ | 90 | ASNER 96 | CLE2 | Repl. by JESSOP 00 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K_2^*(1430)^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{233}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|-------------------------|-------------|----------------------------------|
| < 18 | 90 | ¹ GARMASH 04 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|--------------|-----|----------------------------------|
| <2600 | 90 | ALBRECHT 91B | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
|-------|----|--------------|-----|----------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K^0 K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{234}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|-------------------------|-------------|----------------------------------|
| <18 × 10⁻⁶ | 90 | ¹ GARMASH 04 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------|----|-------------------------|------|----------------------------------|
| <21 × 10 ⁻⁶ | 90 | ¹ ECKHART 02 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
|------------------------|----|-------------------------|------|----------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $[\Gamma(\bar{K}^{*0} K^0) + \Gamma(K^{*0} \bar{K}^0)]/\Gamma_{\text{total}}$ Γ_{235}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|----------------------------|-------------|----------------------------------|
| <1.9 | ¹ AUBERT,BE 06N | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{236}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|-------------------------|-------------|----------------------------------|
| <19 × 10⁻⁶ | 90 | ¹ ECKHART 02 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K^0 K^+ K^-)/\Gamma_{\text{total}}$ Γ_{237}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
| 24.7 ± 2.3 OUR AVERAGE | | | | |

$23.8 \pm 2.0 \pm 1.6$

$28.3 \pm 3.3 \pm 4.0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|--------------|-----|----------------------------------|
| <1300 | 90 | ALBRECHT 91E | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
|-------|----|--------------|-----|----------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^0\phi)/\Gamma_{\text{total}}$ Γ_{238}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
|---------------------------------|------------|--------------------|-------------|----------------|

8.6 \pm 1.3 OUR AVERAGE $8.4^{+1.5}_{-1.3} \pm 0.5$ $9.0^{+2.2}_{-1.8} \pm 0.7$ ¹ AUBERT 04A BABR $e^+ e^- \rightarrow \gamma(4S)$ ¹ CHEN 03B BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $8.1^{+3.1}_{-2.5} \pm 0.8$ ¹ AUBERT 01D BABR $e^+ e^- \rightarrow \gamma(4S)$ < 12.3 ¹ BRIERE 01 CLE2 $e^+ e^- \rightarrow \gamma(4S)$ < 31 ¹ BERGFELD 98 CLE2 < 88 ASNER 96 CLE2 $e^+ e^- \rightarrow \gamma(4S)$ < 720 ALBRECHT 91B ARG $e^+ e^- \rightarrow \gamma(4S)$ < 420 ² AVERY 89B CLEO $e^+ e^- \rightarrow \gamma(4S)$ < 1000 ³ AVERY 87 CLEO $e^+ e^- \rightarrow \gamma(4S)$ ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² Avery 89B reports $< 4.9 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.³ Avery 87 reports $< 1.3 \times 10^{-3}$ assuming the $\gamma(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%. $\Gamma(K^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{239}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|----------------|
|--------------|------------|--------------------|-------------|----------------|

 $< 2.3 \times 10^{-4}$ ¹ ADAM 96D DLPH $e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $< 2.1 \times 10^{-4}$ ² ABREU 95N DLPH Sup. by ADAM 96D¹ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.² Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B_s^0 decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons. $\Gamma(K^*(892)^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{240}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
|---------------------------------|------------|--------------------|-------------|----------------|

54.5 \pm 2.9 \pm 4.3¹ AUBERT 07AS BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 < 1400 ALBRECHT 91E ARG $e^+ e^- \rightarrow \gamma(4S)$ ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^*(892)^0\rho^0)/\Gamma_{\text{total}}$ Γ_{241}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
|---------------------------------|------------|--------------------|-------------|----------------|

5.6 \pm 0.9 \pm 1.3¹ AUBERT,B 06G BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------|----|---------------------|-----|------|----------------------------------|
| < 34 | 90 | ² GODANG | 02 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <286 | 90 | ³ ABE | 00C | SLD | $e^+ e^- \rightarrow Z$ |
| <460 | 90 | ALBRECHT | 91B | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| <580 | 90 | ⁴ Avery | 89B | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |
| <960 | 90 | ⁵ Avery | 87 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 2.4×10^{-5} .

³ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

⁴ Avery 89B reports $< 6.7 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

⁵ Avery 87 reports $< 1.2 \times 10^{-3}$ assuming the $\gamma(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^0 f_0(980))/\Gamma_{\text{total}}$

Γ_{242}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-----------------------|------|---------------------------------------|
| < 4.3 | 90 | ¹ AUBERT,B | 06G | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|--------------------|-----|---------------------------------------|
| <170 | 90 | ² Avery | 89B | CLEO $e^+ e^- \rightarrow \gamma(4S)$ |
|------|----|--------------------|-----|---------------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Avery 89B reports $< 2.0 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_1(1400)^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{243}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------|------|--------------------------------------|
| $< 1.1 \times 10^{-3}$ | 90 | ALBRECHT | 91B | ARG $e^+ e^- \rightarrow \gamma(4S)$ |

$\Gamma(a_1(1260)^- K^+)/\Gamma_{\text{total}}$

Γ_{244}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-----------------------|------|---------------------------------------|
| $16.3 \pm 2.9 \pm 2.3$ | | ^{1,2} AUBERT | 08F | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|--------------------|-----|------------------------------|
| <230 | 90 | ³ ADAM | 96D | DLPH $e^+ e^- \rightarrow Z$ |
| <390 | 90 | ⁴ ABREU | 95N | DLPH Sup. by ADAM 96D |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes a_1^\pm decays only to 3π and $B(a_1^\pm \rightarrow \pi^\pm \pi^\mp \pi^\pm) = 0.5$.

³ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

⁴ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B_s^0 decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

$\Gamma(b_1^- K^+ \times B(b_1^- \rightarrow \omega \pi^-))/\Gamma_{\text{total}}$

Γ_{245}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|---------------------------------------|
| $7.4 \pm 1.0 \pm 1.0$ | | ¹ AUBERT | 07BI | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^0 K^+ K^-)/\Gamma_{\text{total}}$ Γ_{246}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------|-------------|------------------------------------|
| $27.5 \pm 1.3 \pm 2.2$ | | ¹ AUBERT | 07AS BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<610 90 ALBRECHT 91E ARG $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K^*(892)^0 \phi)/\Gamma_{\text{total}}$ Γ_{247}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------|
| 9.5 ± 0.8 OUR AVERAGE | | | | |

$9.2 \pm 0.7 \pm 0.6$ ¹ AUBERT 07D BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$10.0^{+1.6}_{-1.5}{}^{+0.7}_{-0.8}$ ¹ CHEN 03B BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$11.5^{+4.5}_{-3.7}{}^{+1.8}_{-1.7}$ ¹ BRIERE 01 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.2 \pm 0.9 \pm 0.5$ ¹ AUBERT,B 04W BABR Repl. by AUBERT 07D

$11.2 \pm 1.3 \pm 0.8$ ¹ AUBERT 03V BABR Repl. by AUBERT,B 04W

$8.7^{+2.5}_{-2.1} \pm 1.1$ ¹ AUBERT 01D BABR Repl. by AUBERT 03V

<384 90 ² ABE 00C SLD $e^+ e^- \rightarrow Z$

< 21 90 ¹ BERGFELD 98 CLE2

< 43 90 ASNER 96 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

<320 90 ALBRECHT 91B ARG $e^+ e^- \rightarrow \Upsilon(4S)$

<380 90 ³ AVERY 89B CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

<380 90 ⁴ AVERY 87 CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

³ Avery 89B reports $< 4.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

⁴ Avery 87 reports $< 4.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.

 $\Gamma(K^*(892)^0 K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{248}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|------------------------------------|
| $4.6 \pm 1.1 \pm 0.8$ | ¹ AUBERT | 07AS BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{249}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------------|-------------|------------------------------------|
| $1.28^{+0.35}_{-0.30} \pm 0.11$ | | ¹ AUBERT | 08I BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 22 90 ² GODANG 02 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

<469 90 ³ ABE 00C SLD $e^+ e^- \rightarrow Z$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 1.9×10^{-5} .

³ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

$\Gamma(K^*(892)^0 K^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{250}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|-----------|----------------------------------|
| <2.2 | 90 | ¹ AUBERT | 07AS BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^0 K^*(892)^0)/\Gamma_{\text{total}}$

Γ_{251}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|----------|----------------------------------|
| < 0.41 | 90 | ¹ AUBERT | 08I BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|---------|----------------------------------|
| <37 | 90 | ² GODANG | 02 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 2.9×10^{-5} .

$\Gamma(K^*(892)^+ \rho^-)/\Gamma_{\text{total}}$

Γ_{252}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-----------------------|----------|----------------------------------|
| <12.0 | 90 | ¹ AUBERT,B | 06G BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^+ K^*(892)^-)/\Gamma_{\text{total}}$

Γ_{253}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|---------|----------------------------------|
| <141 | 90 | ¹ GODANG | 02 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 8.9×10^{-5} .

$\Gamma(K_1(1400)^0 \rho^0)/\Gamma_{\text{total}}$

Γ_{254}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|----------------------------------|
| $<3.0 \times 10^{-3}$ | 90 | ALBRECHT | 91B ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

$\Gamma(\phi(K\pi)^{*0}_0)/\Gamma_{\text{total}}$

Γ_{256}/Γ

This decay refers to the coherent sum of resonant and nonresonant $J^P = 0^+$ $K\pi$ components with $1.13 < m_{K\pi} < 1.53 \text{ GeV}/c^2$.

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|----------|----------------------------------|
| $5.0 \pm 0.8 \pm 0.3$ | 90 | ¹ AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\phi(K\pi)^{*0} (1.60 < m_{K\pi} < 2.15)) / \Gamma_{\text{total}}$ Γ_{257}/Γ

This decay refers to the coherent sum of resonant and nonresonant $J^P = 0^+$ $K\pi$ components with $1.60 < m_{K\pi} < 2.15$ GeV/c².

| VALUE (units 10 ⁻⁶) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-----|---------------------|-----------|----------------------------------|
| <1.7 | 90 | ¹ AUBERT | 07AO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K_1(1400)^0 \phi) / \Gamma_{\text{total}}$ Γ_{255}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|----------------------------------|
| $<5.0 \times 10^{-3}$ | 90 | ALBRECHT | 91B ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

 $\Gamma(K_0^*(1430)^0 \phi) / \Gamma_{\text{total}}$ Γ_{258}/Γ

| VALUE (units 10 ⁻⁶) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|---------------------|----------|----------------------------------|
| 4.6 ± 0.7 ± 0.6 | ¹ AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ²AUBERT,B 04W BABR Repl. by AUBERT 07D

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Observed 181 ± 17 events with statistical significance greater than 10 σ .

 $\Gamma(K^*(1680)^0 \phi) / \Gamma_{\text{total}}$ Γ_{259}/Γ

| VALUE (units 10 ⁻⁶) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-----|---------------------|-----------|----------------------------------|
| <3.5 | 90 | ¹ AUBERT | 07AO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K^*(1780)^0 \phi) / \Gamma_{\text{total}}$ Γ_{260}/Γ

| VALUE (units 10 ⁻⁶) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-----|---------------------|-----------|----------------------------------|
| <2.7 | 90 | ¹ AUBERT | 07AO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K^*(2045)^0 \phi) / \Gamma_{\text{total}}$ Γ_{261}/Γ

| VALUE (units 10 ⁻⁶) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-----|---------------------|-----------|----------------------------------|
| <15.3 | 90 | ¹ AUBERT | 07AO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K_2^*(1430)^0 \rho^0) / \Gamma_{\text{total}}$ Γ_{262}/Γ

| VALUE (units 10 ⁻⁶) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-----|-------------|---------|----------------------------------|
| $<1.1 \times 10^3$ | 90 | ALBRECHT | 91B ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

 $\Gamma(K_2^*(1430)^0 \phi) / \Gamma_{\text{total}}$ Γ_{263}/Γ

| VALUE (units 10 ⁻⁶) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-----|---------------------|----------|----------------------------------|
| 7.8 ± 1.1 ± 0.6 | | ¹ AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ²AUBERT,B 04W BABR Repl. by AUBERT 07D
 ALBRECHT 91B ARG $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² The angular distribution of $B \rightarrow \phi K^*(1430)$ provides evidence with statistical significance of 3.2σ .

$\Gamma(K^0\phi\phi)/\Gamma_{\text{total}}$

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT | Γ_{264}/Γ |
|---|------------------------|----------|-----------------------------------|-----------------------|
| $4.1^{+1.7}_{-1.4} \pm 0.4$ | ¹ AUBERT,BE | 06H BABR | $e^+e^- \rightarrow \Upsilon(4S)$ | |

¹ Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$ and for a $\phi\phi$ invariant mass below $2.85\text{ GeV}/c^2$.

$\Gamma(\eta'\eta'K^0)/\Gamma_{\text{total}}$

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT | Γ_{265}/Γ |
|--------------------------|-----|-----------------------|----------|-----------------------------------|-----------------------|
| <31 | 90 | ¹ AUBERT,B | 06P BABR | $e^+e^- \rightarrow \Upsilon(4S)$ | |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^0\gamma)/\Gamma_{\text{total}}$

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT | Γ_{266}/Γ |
|--|-----|------------------------|----------|-----------------------------------|-----------------------|
| 40.1 ± 2.0 OUR AVERAGE | | | | | |
| $39.2 \pm 2.0 \pm 2.4$ | | ¹ AUBERT,BE | 04A BABR | $e^+e^- \rightarrow \Upsilon(4S)$ | |
| $40.1 \pm 2.1 \pm 1.7$ | | ² NAKAO | 04 BELL | $e^+e^- \rightarrow \Upsilon(4S)$ | |
| $45.5^{+7.2}_{-6.8} \pm 3.4$ | | ³ COAN | 00 CLE2 | $e^+e^- \rightarrow \Upsilon(4S)$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------|----|---------------------|----------|-----------------------------------|
| < 110 | 90 | ACOSTA | 02G CDF | $p\bar{p}$ at 1.8 TeV |
| $42.3 \pm 4.0 \pm 2.2$ | | ² AUBERT | 02C BABR | Repl. by AUBERT,BE 04A |
| < 210 | 90 | ⁴ ADAM | 96D DLPH | $e^+e^- \rightarrow Z$ |
| $40 \pm 17 \pm 8$ | | ⁵ AMMAR | 93 CLE2 | Repl. by COAN 00 |
| < 420 | 90 | ALBRECHT | 89G ARG | $e^+e^- \rightarrow \Upsilon(4S)$ |
| < 240 | 90 | ⁶ AVERY | 89B CLEO | $e^+e^- \rightarrow \Upsilon(4S)$ |
| < 2100 | 90 | AVERY | 87 CLEO | $e^+e^- \rightarrow \Upsilon(4S)$ |

¹ Uses the production ratio of charged and neutral B from $\Upsilon(4S)$ decays $R^+/0 = 1.006 \pm 0.048$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. No evidence for a nonresonant $K\pi\gamma$ contamination was seen; the central value assumes no contamination.

⁴ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

⁵ AMMAR 93 observed 6.6 ± 2.8 events above background.

⁶ AVERY 89B reports $< 2.8 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\eta K^0\gamma)/\Gamma_{\text{total}}$

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT | Γ_{267}/Γ |
|--|-------------------------|----------|-----------------------------------|-----------------------|
| $10.7^{+2.2}_{-1.5}$ OUR AVERAGE | | | | |
| $11.3^{+2.8}_{-1.6} \pm 0.6$ | ^{1,2} AUBERT,B | 06M BABR | $e^+e^- \rightarrow \Upsilon(4S)$ | |
| $8.7^{+3.1}_{-2.7} \pm 1.6$ | ^{2,3} NISHIDA | 05 BELL | $e^+e^- \rightarrow \Upsilon(4S)$ | |

$^1 m_{\eta' K} < 3.25 \text{ GeV}/c^2$. 2 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $^3 m_{\eta K} < 2.4 \text{ GeV}/c^2$ $\Gamma(\eta' K^0 \gamma)/\Gamma_{\text{total}}$ Γ_{268}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| <6.6 | 90 | 1,2 AUBERT,B | 06M BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

 $^1 m_{\eta' K} < 3.25 \text{ GeV}/c^2$. 2 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^0 \phi \gamma)/\Gamma_{\text{total}}$ Γ_{269}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| <2.7 | 90 | 1 AUBERT | 07Q BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

| | | | | |
|--------|----|------------|---------|----------------------------------|
| <8.3 | 90 | 1 DRUTSKOY | 04 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
|--------|----|------------|---------|----------------------------------|

 1 Assumes equal production of B^+ and B^0 at $\gamma(4S)$. $\Gamma(K^+ \pi^- \gamma)/\Gamma_{\text{total}}$ Γ_{270}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|----------------------------------|
| (4.6\pm1.3\pm0.5) $\times 10^{-6}$ | 1,2 NISHIDA | 02 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

 1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $^2 1.25 \text{ GeV}/c^2 < M_{K\pi} < 1.6 \text{ GeV}/c^2$ $\Gamma(K^*(1410)\gamma)/\Gamma_{\text{total}}$ Γ_{271}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|----------------------------------|
| <1.3 $\times 10^{-4}$ | 90 | 1 NISHIDA | 02 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

 1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^+ \pi^- \gamma \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{272}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|----------------------------------|
| <2.6 $\times 10^{-6}$ | 90 | 1,2 NISHIDA | 02 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

 1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $^2 1.25 \text{ GeV}/c^2 < M_{K\pi} < 1.6 \text{ GeV}/c^2$ $\Gamma(K^0 \pi^+ \pi^- \gamma)/\Gamma_{\text{total}}$ Γ_{273}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| 1.95\pm0.22 OUR AVERAGE | | | |

| | | | |
|----------------------------|------------|----------|----------------------------------|
| 1.85 \pm 0.21 \pm 0.12 | 1,2 AUBERT | 07R BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 2.40 \pm 0.4 \pm 0.3 | 2,3 YANG | 05 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

 $^1 M_{K\pi\pi} < 1.8 \text{ GeV}/c^2$. 2 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $^3 M_{K\pi\pi} < 2.0 \text{ GeV}/c^2$.

| $\Gamma(K^+\pi^-\pi^0\gamma)/\Gamma_{\text{total}}$ | Γ_{274}/Γ |
|--|---|
| <u>VALUE (units 10^{-5})</u> | <u>DOCUMENT ID</u> |
| $4.07 \pm 0.22 \pm 0.31$ | 1,2 AUBERT 07R BABR $e^+e^- \rightarrow \gamma(4S)$ |
| ¹ $M_{K\pi\pi} < 1.8 \text{ GeV}/c^2$. | |
| ² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. | |

| $\Gamma(K_1(1270)^0\gamma)/\Gamma_{\text{total}}$ | Γ_{275}/Γ |
|--|--|
| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> |
| < 5.8 | 90 |
| 1 YANG 05 BELL $e^+e^- \rightarrow \gamma(4S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | |
| <700 90 | 2 ALBRECHT 89G ARG $e^+e^- \rightarrow \gamma(4S)$ |
| ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. | |
| ² ALBRECHT 89G reports < 0.0078 assuming the $\gamma(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%. | |

| $\Gamma(K_1(1400)^0\gamma)/\Gamma_{\text{total}}$ | Γ_{276}/Γ |
|--|--|
| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> |
| < 1.5 | 90 |
| 1 YANG 05 BELL $e^+e^- \rightarrow \gamma(4S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | |
| <430 90 | 2 ALBRECHT 89G ARG $e^+e^- \rightarrow \gamma(4S)$ |
| ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. | |
| ² ALBRECHT 89G reports < 0.0048 assuming the $\gamma(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%. | |

| $\Gamma(K_2^*(1430)^0\gamma)/\Gamma_{\text{total}}$ | Γ_{277}/Γ |
|--|---|
| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> |
| 1.24 ± 0.24 OUR AVERAGE | |
| 1.22 $\pm 0.25 \pm 0.10$ | 1 AUBERT,B 04U BABR $e^+e^- \rightarrow \gamma(4S)$ |
| 1.3 $\pm 0.5 \pm 0.1$ | 1 NISHIDA 02 BELL $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | |
| <40 90 | 2 ALBRECHT 89G ARG $e^+e^- \rightarrow \gamma(4S)$ |
| ¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. | |
| ² ALBRECHT 89G reports < 4.4×10^{-4} assuming the $\gamma(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%. | |

| $\Gamma(K^*(1680)^0\gamma)/\Gamma_{\text{total}}$ | Γ_{278}/Γ |
|--|-----------------------|
| <u>VALUE</u> | <u>CL%</u> |
| <0.0020 | 90 |
| 1 ALBRECHT 89G ARG $e^+e^- \rightarrow \gamma(4S)$ | |
| ¹ ALBRECHT 89G reports < 0.0022 assuming the $\gamma(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%. | |

| $\Gamma(K_3^*(1780)^0 \gamma)/\Gamma_{\text{total}}$ | | | | Γ_{279}/Γ |
|---|------------|--------------------|-------------|------------------------------------|
| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| < 83 | 90 | 1,2 NISHIDA | 05 BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <10000 | 90 | 3 ALBRECHT | 89G ARG | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. | | | | |
| 2 Uses $B(K_3^*(1780) \rightarrow \eta K) = 0.11^{+0.05}_{-0.04}$. | | | | |
| 3 ALBRECHT 89G reports < 0.011 assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%. | | | | |

| $\Gamma(K_4^*(2045)^0 \gamma)/\Gamma_{\text{total}}$ | | | | Γ_{280}/Γ |
|--|------------|--------------------|-------------|------------------------------------|
| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| <0.0043 | 90 | 1 ALBRECHT | 89G ARG | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1 ALBRECHT 89G reports < 0.0048 assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%. | | | | |

| $\Gamma(\rho^0 \gamma)/\Gamma_{\text{total}}$ | | | | Γ_{281}/Γ |
|---|------------|---------------------|------------------------------------|-------------------------------------|
| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.93\pm0.21 OUR AVERAGE | | | | Error includes scale factor of 1.1. |
| 0.79 $^{+0.22}_{-0.20}\pm 0.06$ | | | | |
| 1 AUBERT 07L BABR $e^+ e^- \rightarrow \Upsilon(4S)$ | | | | |
| 1.25 $^{+0.37}_{-0.33}\pm 0.07$ | | | | |
| 1 MOHAPATRA 06 BELL $e^+ e^- \rightarrow \Upsilon(4S)$ | | | | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| 0.0 \pm 0.2 \pm 0.1 | 90 | 1 AUBERT 05 BABR | Repl. by AUBERT 07L | |
| < 0.8 | 90 | 1 MOHAPATRA 05 BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ | |
| < 1.2 | 90 | 1 AUBERT 04C BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ | |
| <17 | 90 | 1 COAN 00 CLE2 | $e^+ e^- \rightarrow \Upsilon(4S)$ | |
| 1 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. | | | | |

| $\Gamma(\omega \gamma)/\Gamma_{\text{total}}$ | | | | Γ_{282}/Γ |
|---|------------|---------------------|------------------------------------|-----------------------|
| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.46\pm0.20 OUR AVERAGE | | | | |
| 0.40 $^{+0.24}_{-0.20}\pm 0.05$ | | | | |
| 1 AUBERT 07L BABR $e^+ e^- \rightarrow \Upsilon(4S)$ | | | | |
| 0.56 $^{+0.34}_{-0.27}\pm 0.05$ | | | | |
| 1 MOHAPATRA 06 BELL $e^+ e^- \rightarrow \Upsilon(4S)$ | | | | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <1.0 | 90 | 1 AUBERT 05 BABR | Repl. by AUBERT 07L | |
| <0.8 | 90 | 1 MOHAPATRA 05 BELL | Repl. by MOHAPATRA 06 | |
| <1.0 | 90 | 1 AUBERT 04C BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ | |
| <9.2 | 90 | 1 COAN 00 CLE2 | $e^+ e^- \rightarrow \Upsilon(4S)$ | |
| 1 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. | | | | |

| $\Gamma(\phi\gamma)/\Gamma_{\text{total}}$ | | Γ_{283}/Γ | | | |
|---|-----|-----------------------|------|----------------------------------|--|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<8.5 \times 10^{-7}$ | 90 | 1 AUBERT,BE 05C | BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| $<0.33 \times 10^{-5}$ | 90 | 1 COAN 00 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ | |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

| $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ | | Γ_{284}/Γ | | | |
|---|-----|-----------------------|------|----------------------------------|--|
| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT | |
| 5.13±0.24 OUR AVERAGE | | | | | |
| 5.5 ± 0.4 ± 0.3 | | 1 AUBERT 07B | BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| 5.1 ± 0.2 ± 0.2 | | 1 LIN 07A | BELL | $e^+ e^- \rightarrow \gamma(4S)$ | |
| 4.1 ± 1.1 ± 0.1 | | 2 ABULENCIA,A 06D | CDF | $p\bar{p}$ at 1.96 TeV | |
| 4.5 +1.4 +0.5 -1.2 -0.4 | | 1 BORNHEIM 03 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| 4.4 ± 0.6 ± 0.3 | | 1 CHAO 04 | BELL | Repl. by LIN 07A | |
| 4.7 ± 0.6 ± 0.2 | | 1 AUBERT 02Q | BABR | Repl. by AUBERT 07B | |
| 5.4 ± 1.2 ± 0.5 | | 1 CASEY 02 | BELL | Repl. by CHAO 04 | |
| 5.6 +2.3 +0.4 -2.0 -0.5 | | 1 ABE 01H | BELL | Repl. by CASEY 02 | |
| 4.1 ± 1.0 ± 0.7 | | 1 AUBERT 01E | BABR | Repl. by AUBERT 02Q | |
| < 67 | 90 | 3 ABE 00C | SLD | $e^+ e^- \rightarrow Z$ | |
| 4.3 +1.6 -1.4 ± 0.5 | | 1 CRONIN-HEN..00 | CLE2 | Repl. by BORNHEIM 03 | |
| < 15 | 90 | GODANG 98 | CLE2 | Repl. by CRONIN-HENNESSY 00 | |
| < 45 | 90 | 4 ADAM 96D | DLPH | $e^+ e^- \rightarrow Z$ | |
| < 20 | 90 | ASNER 96 | CLE2 | Repl. by GODANG 98 | |
| < 41 | 90 | 5 BUSKULIC 96V | ALEP | $e^+ e^- \rightarrow Z$ | |
| < 55 | 90 | 6 ABREU 95N | DLPH | Sup. by ADAM 96D | |
| < 47 | 90 | 7 AKERS 94L | OPAL | $e^+ e^- \rightarrow Z$ | |
| < 29 | 90 | 1 BATTLE 93 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ | |
| < 130 | 90 | 1 ALBRECHT 90B | ARG | $e^+ e^- \rightarrow \gamma(4S)$ | |
| < 77 | 90 | 8 BORTOLETTO89 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ | |
| < 260 | 90 | 8 BEBEK 87 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ | |
| < 500 | 90 | GILES 84 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ | |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ABULENCIA,A 06D reports $[B(B^0 \rightarrow \pi^+\pi^-)] / [B(B^0 \rightarrow K^+\pi^-)] = 0.21 \pm 0.05 \pm 0.03$. We multiply by our best value $B(B^0 \rightarrow K^+\pi^-) = (1.94 \pm 0.06) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7 \pm 1.8)\%$ and $f_{B_s} = (10.5 \pm 1.8)\%$.

⁴ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

⁵ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

⁶ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

⁷ Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and B_d^0 (B_s^0) fraction 39.5% (12%).

⁸ Paper assumes the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+\pi^-)/\Gamma(K^+\pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------|------|---------|
| 0.257±0.014 OUR AVERAGE | | | |

| | | | | |
|---|-------------|-----|------|---------------------------------|
| 0.26 ± 0.01 ± 0.01 | LIN | 07A | BELL | $e^+e^- \rightarrow \gamma(4S)$ |
| 0.21 ± 0.05 ± 0.03 | ABULENCIA,A | 06D | CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.29 +0.13 +0.01 -0.12 -0.02 | ABE | 01H | BELL | Repl. by LIN 07A |

 $\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----|-------------|------|---------|
| 1.62±0.31 OUR AVERAGE | | | | |

| | | | | |
|----------------------------|----------|------|------|---------------------------------|
| 1.47 ± 0.25 ± 0.12 | 1 AUBERT | 07BC | BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| 2.3 +0.4 +0.2 -0.5 -0.3 | 1 CHAO | 05 | BELL | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------|------------|-----|------|---------------------------------|
| 1.17 ± 0.32 ± 0.10 | 1 AUBERT | 05L | BABR | Repl. by AUBERT 07BC |
| < 3.6 | 1 AUBERT | 03L | BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| 2.1 ± 0.6 ± 0.3 | 1 AUBERT | 03S | BABR | Repl. by AUBERT 05L |
| < 4.4 | 1 BORNHEIM | 03 | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| 1.7 ± 0.6 ± 0.2 | 1 LEE | 03 | BELL | Repl. by CHAO 05 |
| < 5.7 | 1 ASNER | 02 | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| < 6.4 | 1 CASEY | 02 | BELL | $e^+e^- \rightarrow \gamma(4S)$ |
| < 9.3 | GODANG | 98 | CLE2 | Repl. by ASNER 02 |
| < 9.1 | ASNER | 96 | CLE2 | Repl. by GODANG 98 |
| <60 | 2 ACCIARRI | 95H | L3 | $e^+e^- \rightarrow Z$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

 $\Gamma(\eta\pi^0)/\Gamma_{\text{total}}$

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
| < 1.3 | | | | |

| | | | | | |
|---|----|------------|-----|------|---------------------------------|
| < 1.3 | 90 | 1 AUBERT | 06W | BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 2.5 | 90 | 1 CHANG | 05A | BELL | $e^+e^- \rightarrow \gamma(4S)$ |
| < 2.5 | 90 | 1 AUBERT,B | 04D | BABR | Repl. by AUBERT 06W |
| < 2.9 | 90 | 1 RICHICHI | 00 | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| < 8 | 90 | BEHRENS | 98 | CLE2 | Repl. by RICHICHI 00 |
| < 250 | 90 | 2 ACCIARRI | 95H | L3 | $e^+e^- \rightarrow Z$ |
| <1800 | 90 | 1 ALBRECHT | 90B | ARG | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
| < 1.8 | | | | |

| | | | | | |
|-------|----|------------|-----|------|---------------------------------|
| < 1.8 | 90 | 1 AUBERT,B | 06V | BABR | $e^+e^- \rightarrow \gamma(4S)$ |
|-------|----|------------|-----|------|---------------------------------|

 $\Gamma_{284}/\Gamma_{198}$ Γ_{285}/Γ Γ_{286}/Γ Γ_{287}/Γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|-----------------------|-----|------|------------------------------------|
| < 2.0 | 90 | ¹ CHANG | 05A | BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| < 2.8 | 90 | ¹ AUBERT,B | 04X | BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| < 18 | 90 | BEHRENS | 98 | CLE2 | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| <410 | 90 | ² ACCIARRI | 95H | L3 | $e^+ e^- \rightarrow Z$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

$\Gamma(\eta' \pi^0)/\Gamma_{\text{total}}$

Γ_{288}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

1.5^{+1.0}_{-0.8} OUR AVERAGE Error includes scale factor of 1.5.

$0.8^{+0.8}_{-0.6} \pm 0.1$ ¹ AUBERT 06W BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$2.8 \pm 1.0 \pm 0.3$ ¹ SCHUEMANN 06 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.0^{+1.4}_{-1.0} \pm 0.8$ 90 ¹ AUBERT,B 04D BABR Repl. by AUBERT 06W

< 5.7 90 ¹ RICHICHI 00 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

<11 90 BEHRENS 98 CLE2 Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' \eta')/\Gamma_{\text{total}}$

Γ_{289}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

< 2.4 90 ¹ AUBERT,B 06V BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.5 90 ¹ SCHUEMANN 07 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

<10 90 ¹ AUBERT,B 04X BABR Repl. by AUBERT,B 06V

<47 90 BEHRENS 98 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' \eta)/\Gamma_{\text{total}}$

Γ_{290}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

< 1.7 90 ¹ AUBERT 06W BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.5 90 ¹ SCHUEMANN 07 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

< 4.6 90 ¹ AUBERT,B 04X BABR $e^+ e^- \rightarrow \Upsilon(4S)$

<27 90 BEHRENS 98 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' \rho^0)/\Gamma_{\text{total}}$

Γ_{291}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

< 1.3 90 ¹ SCHUEMANN 07 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|-----------------------|-----|------|----------------------------------|
| < 3.7 | 90 | AUBERT | 07E | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 4.3 | 90 | ¹ AUBERT,B | 04D | BABR | Repl. by AUBERT 07E |
| <12 | 90 | ¹ RICHICHI | 00 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <23 | 90 | BEHRENS | 98 | CLE2 | Repl. by RICHICHI 00 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\eta' f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{292}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|----------------------------------|
| <1.5 | 90 | AUBERT | 07E | $e^+ e^- \rightarrow \gamma(4S)$ |

$\Gamma(\eta \rho^0) / \Gamma_{\text{total}}$ Γ_{293}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|---------------------------------------|
| < 1.5 | 90 | ¹ AUBERT | 07Y | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|-----------------------|-----|------|----------------------------------|
| < 1.9 | 90 | ¹ WANG | 07B | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 1.5 | 90 | ¹ AUBERT,B | 04D | BABR | Repl. by AUBERT 07Y |
| <10 | 90 | ¹ RICHICHI | 00 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <13 | 90 | BEHRENS | 98 | CLE2 | Repl. by RICHICHI 00 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\eta f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{294}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|---------------------------------------|
| <0.4 | 90 | ¹ AUBERT | 07Y | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\omega \eta) / \Gamma_{\text{total}}$ Γ_{295}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-----------------------|------|---------------------------------------|
| < 1.9 | 90 | ¹ AUBERT,B | 05K | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------------------|----|-----------------------|-----|------|-----------------------|
| $4.0^{+1.3}_{-1.2} \pm 0.4$ | | ¹ AUBERT,B | 04X | BABR | Repl. by AUBERT,B 05K |
| <12 | 90 | ¹ BERGFELD | 98 | CLE2 | |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\omega \eta') / \Gamma_{\text{total}}$ Γ_{296}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------------------------|------|---------------------------------------|
| < 2.2 | 90 | ¹ SCHUEMANN | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|-----------------------|-----|------|----------------------------------|
| < 2.8 | 90 | ¹ AUBERT,B | 04X | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| <60 | 90 | ¹ BERGFELD | 98 | CLE2 | |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\omega\rho^0)/\Gamma_{\text{total}}$ Γ_{297}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| < 1.5 | 90 | 1 AUBERT,B | 06T BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|------------|----------|-----------------------|
| < 3.3 | 90 | 1 AUBERT | 050 BABR | Repl. by AUBERT,B 06T |
| <11 | 90 | 1 BERGFELD | 98 CLE2 | |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$ Γ_{298}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| <1.5 | 90 | 1 AUBERT,B | 06T BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_{299}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| < 4.0 | 90 | 1 AUBERT,B | 06T BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----|----|------------|---------|--|
| <19 | 90 | 1 BERGFELD | 98 CLE2 | |
|-----|----|------------|---------|--|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ Γ_{300}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| <0.28 | 90 | 1 AUBERT,B | 06C BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|------------|----------|-----------------------|
| <1.0 | 90 | 1 AUBERT,B | 04D BABR | Repl. by AUBERT,B 06C |
| <5 | 90 | 1 BERGFELD | 98 CLE2 | |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$ Γ_{301}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| <0.6 | 90 | 1 AUBERT,B | 06V BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|------------|----------|-----------------------|
| <1.0 | 90 | 1 AUBERT,B | 04X BABR | Repl. by AUBERT,B 06V |
| <9 | 90 | 1 BERGFELD | 98 CLE2 | |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\phi\eta')/\Gamma_{\text{total}}$ Γ_{302}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------------------------|
| < 0.5 | 90 | 1 SCHUEMANN | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|------------|----------|----------------------------------|
| < 1.0 | 90 | 1 AUBERT,B | 06V BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 4.5 | 90 | 1 AUBERT,B | 04X BABR | Repl. by AUBERT,B 06V |
| <31 | 90 | 1 BERGFELD | 98 CLE2 | |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\phi\rho^0)/\Gamma_{\text{total}}$ Γ_{303}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|-----------------------------|
| < 13 | 90 | 1 BERGFELD | 98 | CLE2 |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <156 | 90 | 2 ABE | 00C | SLD $e^+ e^- \rightarrow Z$ |
| 1 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. | | | | |
| 2 ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$. | | | | |

 $\Gamma(\phi\omega)/\Gamma_{\text{total}}$ Γ_{304}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| < 1.2 | 90 | 1 AUBERT,B | 06T | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <21 | 90 | 1 BERGFELD | 98 | CLE2 |
| 1 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. | | | | |

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_{305}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|---|
| $< 1.5 \times 10^{-6}$ | 90 | 1 AUBERT,B | 04X | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| < 3.21×10^{-4} | 90 | 2 ABE | 00C | SLD $e^+ e^- \rightarrow Z$ |
| < 1.2×10^{-5} | 90 | 1 BERGFELD | 98 | CLE2 |
| < 3.9×10^{-5} | 90 | ASNER | 96 | CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. | | | | |
| 2 ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$. | | | | |

 $\Gamma(a_0(980)^{\pm}\pi^{\mp} \times B(a_0(980)^{\pm} \rightarrow \eta\pi^{\pm}))/\Gamma_{\text{total}}$ Γ_{306}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| <3.1 | 90 | 1 AUBERT | 07Y | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <5.1 | 90 | 1 AUBERT,BE | 04 | BABR Repl. by AUBERT 07Y |
| 1 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. | | | | |

 $\Gamma(a_0(1450)^{\pm}\pi^{\mp} \times B(a_0(1450)^{\pm} \rightarrow \eta\pi^{\pm}))/\Gamma_{\text{total}}$ Γ_{307}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| <2.3 | 90 | 1 AUBERT | 07Y | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. | | | | |

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{308}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|--|
| $< 7.2 \times 10^{-4}$ | 90 | 1 ALBRECHT | 90B | ARG $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 1 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$. | | | | |

$\Gamma(\rho^0\pi^0)/\Gamma_{\text{total}}$ Γ_{309}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------|-------------------------|-------------|---------------------------------|
| 1.8 ±0.5 OUR AVERAGE | | | | |
| $3.12^{+0.88+0.60}_{-0.82-0.76}$ | | ¹ DRAGIC 06 | BELL | $e^+e^- \rightarrow \gamma(4S)$ |
| $1.4 \pm 0.6 \pm 0.3$ | | ¹ AUBERT 04Z | BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| $1.6^{+2.0}_{-1.4} \pm 0.8$ | | ¹ JESSOP 00 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----|---------------------------|------|---------------------------------|
| $5.1 \pm 1.6 \pm 0.9$ | | DRAGIC 04 | BELL | Repl. by DRAGIC 06 |
| < 5.3 | 90 | ¹ GORDON 02 | BELL | Repl. by DRAGIC 04 |
| < 24 | 90 | ASNER 96 | CLEO | Repl. by JESSOP 00 |
| < 400 | 90 | ¹ ALBRECHT 90B | ARG | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\rho^\mp\pi^\pm)/\Gamma_{\text{total}}$ Γ_{310}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|-------------------------|-------------|---------------------------------|
| 22.8±2.5 OUR AVERAGE | | | | |
| $22.6 \pm 1.8 \pm 2.2$ | | ¹ AUBERT 03T | BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| $20.8^{+6.0+2.8}_{-6.3-3.1}$ | | ¹ GORDON 02 | BELL | $e^+e^- \rightarrow \gamma(rS)$ |
| $27.6^{+8.4}_{-7.4} \pm 4.2$ | | ¹ JESSOP 00 | CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|----------|----|---------------------------|------|---------------------------------|
| < 88 | 90 | ASNER 96 | CLE2 | Repl. by JESSOP 00 |
| < 520 | 90 | ¹ ALBRECHT 90B | ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| < 5200 | 90 | ² BEBEK 87 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² BEBEK 87 reports $< 6.1 \times 10^{-3}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{311}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|------------|-----------------------|-------------|------------------------|
| $< 2.3 \times 10^{-4}$ | 90 | ¹ ADAM 96D | DLPH | $e^+e^- \rightarrow Z$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------|----|---------------------------|------|---------------------------------|
| $< 2.8 \times 10^{-4}$ | 90 | ² ABREU 95N | DLPH | Sup. by ADAM 96D |
| $< 6.7 \times 10^{-4}$ | 90 | ³ ALBRECHT 90B | ARG | $e^+e^- \rightarrow \gamma(4S)$ |

¹ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

² Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

³ ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$.

 $\Gamma(\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{312}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-------------------------|-------------|---------------------------------|
| $1.07 \pm 0.33 \pm 0.19$ | | ¹ AUBERT 07G | BABR | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|---------------------------|------|----------------------------------|----------------------------------|
| < 1.1 | 90 | ¹ AUBERT | 05I | BABR | Repl. by AUBERT 07G |
| < 2.1 | 90 | ¹ AUBERT | 03V | BABR | Repl. by AUBERT 05I |
| < 18 | 90 | ² GODANG | 02 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <136 | 90 | ³ ABE | 00C | SLD | $e^+ e^- \rightarrow Z$ |
| <280 | 90 | ¹ ALBRECHT | 90B | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| <290 | 90 | ⁴ BORTOLETTO89 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ | |
| <430 | 90 | ⁴ BEBEK | 87 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 1.4×10^{-5} .

³ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

⁴ Paper assumes the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho^0 f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$

Γ_{313}/Γ

| VALUE (units 10^{-6}) | CL % | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------|------|---------------------------------------|
| <0.53 | 90 | ¹ AUBERT | 07G | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(f_0(980) f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)^2) / \Gamma_{\text{total}}$

Γ_{314}/Γ

| VALUE (units 10^{-6}) | CL % | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------|------|---------------------------------------|
| <0.16 | 90 | ¹ AUBERT | 07G | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(a_1(1260)^\mp \pi^\pm) / \Gamma_{\text{total}}$

Γ_{315}/Γ

| VALUE (units 10^{-6}) | CL % | DOCUMENT ID | TECN | COMMENT |
|--|------|-----------------------|------|---------------------------------------|
| $33.2 \pm 3.8 \pm 3.0$ | | ^{1,2} AUBERT | 06V | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|---------------------------|------|----------------------------------|----------------------------------|
| < 630 | 90 | ¹ ALBRECHT | 90B | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 490 | 90 | ³ BORTOLETTO89 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ | |
| <1000 | 90 | ³ BEBEK | 87 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes $a_1(1260)$ decays only to 3π and $B(a_1^\pm \rightarrow \pi^\pm \pi^\mp \pi^\pm) = 0.5$.

³ Paper assumes the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(b_1^\mp \pi^\pm \times B(b_1^\mp \rightarrow \omega \pi^\mp)) / \Gamma_{\text{total}}$

Γ_{316}/Γ

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|------|---------------------------------------|
| $10.9 \pm 1.2 \pm 0.9$ | ¹ AUBERT | 07BI | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

| $\Gamma(a_2(1320)^{\mp}\pi^{\pm})/\Gamma_{\text{total}}$ | | | | Γ_{317}/Γ | |
|---|-----|---------------------------|------|---------------------------------|---------------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<3.0 \times 10^{-4}$ | 90 | ¹ BORTOLETTO89 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| $<1.4 \times 10^{-3}$ | 90 | ¹ BEBEK | 87 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |
| ¹ Paper assumes the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%. | | | | | |

| $\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ | | | | Γ_{318}/Γ | |
|---|-----|-----------------------|------|-----------------------|---------------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<3.1 \times 10^{-3}$ | 90 | ¹ ALBRECHT | 90B | ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| ¹ ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$. | | | | | |

| $\Gamma(\rho^+\rho^-)/\Gamma_{\text{total}}$ | | | | Γ_{319}/Γ | |
|---|------------------|-----------------------|--------------------------|---------------------------------|----------------------|
| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT | |
| 24.2 ± 3.1 OUR AVERAGE | | | | | |
| 25.5 ± 2.1 | $+3.6$ -3.9 | ¹ AUBERT | 07BF BABR | $e^+e^- \rightarrow \gamma(4S)$ | |
| 22.8 ± 3.8 | $+2.3$ -2.6 | ¹ SOMOV | 06 BELL | $e^+e^- \rightarrow \gamma(4S)$ | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| 25 | $+7$ -6 | ¹ AUBERT | 04G BABR | Repl. by AUBERT, B 04R | |
| 30 | ± 4 | ± 5 | ^{1,2} AUBERT, B | 04R BABR | Repl. by AUBERT 07BF |
| <2200 | 90 | ¹ ALBRECHT | 90B ARG | $e^+e^- \rightarrow \gamma(4S)$ | |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² The quoted result is obtained after combining with AUBERT 04G result by AUBERT 04R alone gives $(33 \pm 4 \pm 5) \times 10^{-6}$.

| $\Gamma(a_1(1260)^0\pi^0)/\Gamma_{\text{total}}$ | | | | Γ_{320}/Γ | |
|---|-----|-----------------------|------|-----------------------|---------------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<1.1 \times 10^{-3}$ | 90 | ¹ ALBRECHT | 90B | ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| ¹ ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$. | | | | | |

| $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ | | | | Γ_{321}/Γ | |
|---|-----|------------------------|----------|---------------------------------|--|
| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT | |
| < 1.2 | 90 | ¹ AUBERT, B | 04D BABR | $e^+e^- \rightarrow \gamma(4S)$ | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| < 2.0 | 90 | ¹ JEN | 06 BELL | $e^+e^- \rightarrow \gamma(4S)$ | |
| < 1.9 | 90 | ¹ WANG | 04A BELL | $e^+e^- \rightarrow \gamma(4S)$ | |
| < 3 | 90 | ¹ AUBERT | 01G BABR | $e^+e^- \rightarrow \gamma(4S)$ | |
| < 5.5 | 90 | ¹ JESSOP | 00 CLE2 | $e^+e^- \rightarrow \gamma(4S)$ | |
| < 14 | 90 | ¹ BERGFELD | 98 CLE2 | Repl. by JESSOP 00 | |
| <460 | 90 | ² ALBRECHT | 90B ARG | $e^+e^- \rightarrow \gamma(4S)$ | |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$.

$\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{322}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|---------------------------------|
| $<9.0 \times 10^{-3}$ | 90 | 1 ALBRECHT | 90B ARG | $e^+e^- \rightarrow \gamma(4S)$ |

1 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$. $\Gamma(a_1(1260)^+\rho^-)/\Gamma_{\text{total}}$ Γ_{323}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------------------|------|---------------------------------|
| < 61 | 90 | 1,2 AUBERT,B 060 | BABR | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|------------|---------|---------------------------------|
| <3400 | 90 | 1 ALBRECHT | 90B ARG | $e^+e^- \rightarrow \gamma(4S)$ |
|-------|----|------------|---------|---------------------------------|

1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.2 Assumes $a_1(1260)$ decays only to 3π and $B(a_1^\pm \rightarrow \pi^\pm\pi^\mp\pi^\pm) = 0.5$. $\Gamma(a_1(1260)^0\rho^0)/\Gamma_{\text{total}}$ Γ_{324}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|---------------------------------|
| $<2.4 \times 10^{-3}$ | 90 | 1 ALBRECHT | 90B ARG | $e^+e^- \rightarrow \gamma(4S)$ |

1 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$. $\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^-\pi^+)/\Gamma_{\text{total}}$ Γ_{325}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|---------------------------------|
| $<3.0 \times 10^{-3}$ | 90 | 1 ALBRECHT | 90B ARG | $e^+e^- \rightarrow \gamma(4S)$ |

1 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$. $\Gamma(a_1(1260)^+a_1(1260)^-)/\Gamma_{\text{total}}$ Γ_{326}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|----------------|------|---------------------------------|
| $<2.8 \times 10^{-3}$ | 90 | 1 BORTOLETTO89 | CLEO | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----|------------|---------|---------------------------------|
| $<6.0 \times 10^{-3}$ | 90 | 2 ALBRECHT | 90B ARG | $e^+e^- \rightarrow \gamma(4S)$ |
|-----------------------|----|------------|---------|---------------------------------|

1 BORTOLETTO 89 reports $< 3.2 \times 10^{-3}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$.

We rescale to 50%.

2 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$. $\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{327}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|---------------------------------|
| $<1.1 \times 10^{-2}$ | 90 | 1 ALBRECHT | 90B ARG | $e^+e^- \rightarrow \gamma(4S)$ |

1 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$. $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{328}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|--------------------------------------|
| < 0.11 | 90 | 1 TSAI | 07 | BELL $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------|----|---------------------------|------|------|------------------------------------|
| < 0.41 | 90 | ¹ CHANG | 05 | BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| < 0.27 | 90 | ¹ AUBERT | 04U | BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| < 1.4 | 90 | ¹ BORNHEIM | 03 | CLE2 | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| < 1.2 | 90 | ¹ ABE | 020 | BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| < 7.0 | 90 | ¹ COAN | 99 | CLE2 | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| < 18 | 90 | ² BUSKULIC | 96V | ALEP | $e^+ e^- \rightarrow Z$ |
| <350 | 90 | ³ ABREU | 95N | DLPH | Sup. by ADAM 96D |
| < 34 | 90 | ⁴ BORTOLETTO89 | CLEO | | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| <120 | 90 | ⁵ ALBRECHT | 88F | ARG | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| <170 | 90 | ⁴ BEBEK | 87 | CLEO | $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

³ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

⁴ Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

⁵ ALBRECHT 88F reports $< 1.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{329}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|--------------------|------|---|
| <2.5 | 90 | ¹ BEBEK | 89 | CLEO $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------------|----|-----------------------|-----|------|------------------------------------|
| <9.5 | 90 | ² ABREU | 95N | DLPH | Sup. by ADAM 96D |
| $5.4 \pm 1.8 \pm 2.0$ | | ³ ALBRECHT | 88F | ARG | $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ BEBEK 89 reports $< 2.9 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

² Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

³ ALBRECHT 88F reports $6.0 \pm 2.0 \pm 2.2$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(p\bar{p}K^0)/\Gamma_{\text{total}}$

Γ_{330}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-----|---------------------|------|---|
| 2.7 ±0.4 OUR AVERAGE | | | | |
| $3.0 \pm 0.5 \pm 0.3$ | | ¹ AUBERT | 07AV | BABR $e^+ e^- \rightarrow \Upsilon(4S)$ |
| $2.40^{+0.64}_{-0.44} \pm 0.28$ | | 1,2,3 WANG | 05A | BELL $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.88^{+0.77}_{-0.60} \pm 0.23$ 1,2,4 WANG 04 BELL Repl. by WANG 05A

<7.2 90 1,2 ABE 02K BELL Repl. by WANG 04

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Explicitly vetoes resonant production of $p\bar{p}$ from charmonium states and pK^0 production from Λ_c .

³ Provides also results with $M_{p\bar{p}} < 2.85 \text{ GeV}/c^2$ and angular asymmetry of $p\bar{p}$ system.

⁴ The branching fraction for $M_{p\bar{p}} < 2.85$ is also reported.

$\Gamma(\Theta(1540)^+ \bar{p} \times B(\Theta(1540)^+ \rightarrow p K_S^0)) / \Gamma_{\text{total}}$ Γ_{331}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|-----------|----------------------------------|
| <0.05 | 90 | ¹ AUBERT | 07AV BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.23 90 ¹WANG 05A BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(f_J(2220) K^0 \times B(f_J(2220) \rightarrow p \bar{p})) / \Gamma_{\text{total}}$ Γ_{332}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|-----------|----------------------------------|
| <0.45 | 90 | ¹ AUBERT | 07AV BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(p \bar{p} K^*(892)^0) / \Gamma_{\text{total}}$ Γ_{333}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----|---------------------|-----------|----------------------------------|
| 1.47 ± 0.45 ± 0.40 | 90 | ¹ AUBERT | 07AV BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7.6 90 ¹WANG 04 BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(f_J(2220) K_0^* \times B(f_J(2220) \rightarrow p \bar{p})) / \Gamma_{\text{total}}$ Γ_{334}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|-----------|----------------------------------|
| <0.15 | 90 | ¹ AUBERT | 07AV BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(p \bar{\Lambda} \pi^-) / \Gamma_{\text{total}}$ Γ_{335}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----|-------------------|----------|----------------------------------|
| 3.23 ± 0.33 ± 0.29 | 90 | ¹ WANG | 07C BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.62^{+0.44}_{-0.40} \pm 0.31$ 1,2 WANG 05A BELL Repl. by WANG 07C

$3.97^{+1.00}_{-0.80} \pm 0.56$ 1 WANG 03 BELL Repl. by WANG 05A

< 13 90 ¹COAN 99 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

<180 90 ³ALBRECHT 88F ARG $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Provides also results with $M_{p\bar{p}} < 2.85 \text{ GeV}/c^2$ and angular asymmetry of $p\bar{\Lambda}$ system.

³ ALBRECHT 88F reports $< 2.0 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

 $\Gamma(p \bar{\Sigma}(1385)^-) / \Gamma_{\text{total}}$ Γ_{336}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------------|----------|----------------------------------|
| <0.26 | 90 | ¹ WANG | 07C BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\Delta^0 \bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{337}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------------------------|
| <0.93 | 90 | 1 WANG | 07C BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(p \bar{\Lambda} K^-)/\Gamma_{\text{total}}$ Γ_{338}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------------------------|
| <0.82 | 90 | 1 WANG | 03 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(p \bar{\Sigma}^0 \pi^-)/\Gamma_{\text{total}}$ Γ_{339}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------|--------------------|-------------|----------------------------------|
| <3.8 × 10⁻⁶ | 90 | 1 WANG | 03 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{\Lambda} \Lambda)/\Gamma_{\text{total}}$ Γ_{340}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------------------------|
| <0.32 | 90 | 1 TSAI | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|------------|----------|----------------------------------|
| <0.69 | 90 | 1 CHANG | 05 BELL | Repl. by TSAI 2007 |
| <1.2 | 90 | 1 BORNHEIM | 03 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <1.0 | 90 | 1 ABE | 020 BELL | Repl. by CHANG 05 |
| <3.9 | 90 | 1 COAN | 99 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\Delta^0 \bar{\Delta}^0)/\Gamma_{\text{total}}$ Γ_{341}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------|------------|--------------------|-------------|----------------------------------|
| <0.0015 | 90 | 1 BORTOLETTO89 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ BORTOLETTO 89 reports < 0.0018 assuming $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

 $\Gamma(\Delta^{++} \bar{\Delta}^{--})/\Gamma_{\text{total}}$ Γ_{342}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------|--------------------|-------------|----------------------------------|
| <1.1 × 10⁻⁴ | 90 | 1 BORTOLETTO89 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ BORTOLETTO 89 reports $< 1.3 \times 10^{-4}$ assuming $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

 $\Gamma(\bar{D}^0 p \bar{p})/\Gamma_{\text{total}}$ Γ_{343}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|--------------------|-------------|----------------|
| 1.14 ± 0.09 OUR AVERAGE | | | |

1.13 ± 0.06 ± 0.08 1 AUBERT,B 06S BABR $e^+ e^- \rightarrow \gamma(4S)$

1.18 ± 0.15 ± 0.16 1 ABE 02W BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(D_s^-\bar{\Lambda}p)/\Gamma_{\text{total}}$ Γ_{344}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---|------------------|------|----------------------------------|
| $2.9 \pm 0.9 \pm 0.2$ | 1,2 MEDVEDEVA 07 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² MEDVEDEVA 07 reports $(2.9 \pm 0.7 \pm 0.5 \pm 0.4) \times 10^{-5}$ for $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.38 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{D}^*(2007)^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{345}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|----------------------------------|
| 1.03 ± 0.13 OUR AVERAGE | | | |
| $1.01 \pm 0.10 \pm 0.09$ | 1 AUBERT,B 06S BABR | | $e^+ e^- \rightarrow \gamma(4S)$ |

| | | | |
|---------------------------------|----------------|--|----------------------------------|
| $1.20^{+0.33}_{-0.29} \pm 0.21$ | 1 ABE 02W BELL | | $e^+ e^- \rightarrow \gamma(4S)$ |
|---------------------------------|----------------|--|----------------------------------|

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^- p\bar{p}\pi^+)/\Gamma_{\text{total}}$ Γ_{346}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|------|----------------------------------|
| $3.38 \pm 0.14 \pm 0.29$ | 1 AUBERT,B 06S BABR | | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^{*-} p\bar{p}\pi^+)/\Gamma_{\text{total}}$ Γ_{347}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|------|----------------------------------|
| $4.81 \pm 0.22 \pm 0.44$ | 1 AUBERT,B 06S BABR | | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\Theta_c \bar{p}\pi^+ \times B(\Theta_c \rightarrow D^- p))/\Gamma_{\text{total}}$ Γ_{348}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|----------------------------------|
| <9 | 90 | 1 AUBERT,B 06S BABR | | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\Theta_c \bar{p}\pi^+ \times B(\Theta_c \rightarrow D^{*-} p))/\Gamma_{\text{total}}$ Γ_{349}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------------|------|----------------------------------|
| <14 | 90 | 1 AUBERT,B 06S BABR | | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{\Sigma}_c^{--} \Delta^{++})/\Gamma_{\text{total}}$ Γ_{350}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------|-----|---------------|------|----------------------------------|
| <0.0010 | 90 | 1 PROCARIO 94 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ PROCARIO 94 reports < 0.0012 for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.043$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.050$.

| $\Gamma(\bar{\Lambda}_c^- p \pi^+ \pi^-)/\Gamma_{\text{total}}$ | Γ_{351}/Γ | | | |
|---|-----------------------|-------------|----------------|----------------------------------|
| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 1.3 ± 0.4 OUR AVERAGE | | | | |
| 1.7 $^{+0.3}_{-0.2} \pm 0.4$ | ¹ DYTMAN | 02 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.10 \pm 0.20 \pm 0.29$ | ² GABYSHEV | 02 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $1.33^{+0.46}_{-0.42} \pm 0.37$ | ³ FU | 97 | CLE2 | Repl. by DYTMAN 02 |

¹ DYTMAN 02 reports $(1.67^{+0.27}_{-0.25}) \times 10^{-3}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² GABYSHEV 02 reports $(1.1 \pm 0.2) \times 10^{-3}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ FU 97 uses PDG 96 values of Λ_c branching fraction.

| $\Gamma(\bar{\Lambda}_c^- p)/\Gamma_{\text{total}}$ | Γ_{352}/Γ | | | |
|---|-----------------------|-------------------------|-------------|---------------------------------------|
| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 2.1 ± 0.7 OUR AVERAGE | | | | |
| $2.10^{+0.67}_{-0.55} {}^{+0.77}_{-0.46}$ | | ¹ AUBERT | 07AV BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $2.19^{+0.56}_{-0.49} \pm 0.65$ | | ^{1,2} GABYSHEV | 03 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 9 | 90 | ^{1,3} DYTMAN | 02 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| < 3.1 | 90 | ^{1,4} GABYSHEV | 02 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| < 21 | 90 | ⁵ FU | 97 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² The second error for GABYSHEV 03 includes the systematic and the error of $\Lambda_c \rightarrow \bar{p} K^+ \pi^-$ decay branching fraction.

³ DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

⁴ Uses the value for $\Lambda_c \rightarrow p K^- \pi^+$ branching ratio ($5.0 \pm 1.3\%$).

⁵ FU 97 uses PDG 96 values of Λ_c branching ratio.

| $\Gamma(\bar{\Lambda}_c^- p \pi^0)/\Gamma_{\text{total}}$ | Γ_{353}/Γ | | | |
|---|-----------------------|--------------------|-------------|---------------------------------------|
| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| $< 5.9 \times 10^{-4}$ | 90 | ¹ FU | 97 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |

¹ FU 97 uses PDG 96 values of Λ_c branching ratio.

| $\Gamma(\bar{\Lambda}_c^- p \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ | Γ_{354}/Γ | | | |
|---|-----------------------|--------------------|-------------|---------------------------------------|
| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| $< 5.07 \times 10^{-3}$ | 90 | ¹ FU | 97 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |

¹ FU 97 uses PDG 96 values of Λ_c branching ratio.

| $\Gamma(\bar{\Lambda}_c^- p \pi^+ \pi^- \pi^+ \pi^-) / \Gamma_{\text{total}}$ | Γ_{355} / Γ |
|---|-------------------------|
| $<2.74 \times 10^{-3}$ | 90 |

¹ FU 97 uses PDG 96 values of Λ_c branching ratio.

| $\Gamma(\Lambda_c^+ \bar{p} \pi^+ \pi^-)/\Gamma_{\text{total}}$ | Γ_{356}/Γ | | |
|---|-----------------------|---------|----------------------------------|
| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
| 11.2 \pm 1.4 \pm 2.9 | 1,2 PARK | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² PARK 07 reports $(11.2 \pm 0.5 \pm 3.2) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$.

We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(\Lambda_c^+ \bar{p} \pi^+ \pi^- \text{(nonresonant)}) / \Gamma_{\text{total}}$ | Γ_{357} / Γ | | |
|--|-------------------------|---------|----------------------------------|
| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
| 6.4 \pm 1.0 \pm 1.7 | 1,2 PARK | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² PARK 07 reports $(6.4 \pm 0.4 \pm 1.9) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$.

We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(\Sigma_c(2520)^--\rho\pi^+)/\Gamma_{\text{total}}$ | Γ_{358}/Γ | | |
|--|-----------------------|------|---------------------------------|
| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
| 1.20±0.27±0.31 | 1, ² PARK | 07 | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

³ GABYSHEV 02 BELL Repl. by PARK 07

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(1S)$.

²PARK 07 reports $(1.2 \pm 0.1 \pm 0.4) \times 10^{-4}$ for $B(A^+ \rightarrow p\pi^+$

We rescale to our best value $B(A^+ \rightarrow \mu K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error

³ GABYSHEV 02 reports $(1.63^{+0.64}_{-0.50}) \times 10^{-4}$ for $B(\Lambda^+ \rightarrow p K^- \pi^+) = 0.05$. We

rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(\overline{\Sigma}_c(2520)^0 p\pi^-)/\Gamma_{\text{total}}$ | Γ_{359}/Γ | | | |
|--|-----------------------|-------------|------|----------------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
| $<0.38 \times 10^{-4}$ | 90 | 1 PARK 07 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.21 \times 10^{-4}$ 90 1,2 GABYSHEV 02 BELL Repl. by PARK 07

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses the value for $\Lambda_c \rightarrow p K^- \pi^+$ branching ratio (5.0

$\Gamma(\bar{\Sigma}_c(2455)^0 p\pi^-)/\Gamma_{\text{total}}$ Γ_{360}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---------------------------------------|
| 1.5 ± 0.5 OUR AVERAGE | | | | |
| $1.4 \pm 0.3 \pm 0.4$ | | 1,2 PARK | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| $2.2 \pm 0.7 \pm 0.6$ | | 3 DYTMAN | 02 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $0.5^{+0.5}_{-0.4} \pm 0.1$ | 90 | 4 GABYSHEV | 02 | BELL Repl. by PARK 07 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² PARK 07 reports $(1.4 \pm 0.2 \pm 0.4) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$.We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.³ DYTMAN 02 reports $(2.2 \pm 0.7) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.⁴ GABYSHEV 02 reports $(0.48^{+0.46}_{-0.41}) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma(\bar{\Sigma}_c(2455)^--p\pi^+)/\Gamma_{\text{total}}$ Γ_{361}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|---------------------------------------|
| 2.2 ± 0.7 OUR AVERAGE | | | |
| $2.1 \pm 0.3 \pm 0.5$ | 1,2 PARK | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| $3.7 \pm 1.1 \pm 1.0$ | 3 DYTMAN | 02 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | |
| $2.4 \pm 0.7 \pm 0.6$ | 4 GABYSHEV | 02 | BELL Repl. by PARK 07 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² PARK 07 reports $(2.1 \pm 0.2 \pm 0.6) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$.We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.³ DYTMAN 02 reports $(3.7 \pm 1.1) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.⁴ GABYSHEV 02 reports $(2.38^{+0.75}_{-0.69}) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma(\bar{\Lambda}_c^- \Lambda_c^+)/\Gamma_{\text{total}}$ Γ_{362}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---------------------------------------|
| <6.2 | 90 | 1 UCHIDA | 08 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\bar{\Lambda}_c(2593)^- / \bar{\Lambda}_c(2625)^- p)/\Gamma_{\text{total}}$ Γ_{363}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|------|---------------------------------------|
| $<1.1 \times 10^{-4}$ | 90 | 1,2 DYTMAN | 02 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² DYTMAN 02 measurement uses $B(\bar{\Lambda}_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

 $\Gamma(\Xi_c^- \Lambda_c^+ \times B(\Xi_c^- \rightarrow \Xi^+ \pi^- \pi^-))/\Gamma_{\text{total}}$ Γ_{364}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|------|---------------------------------------|
| $9.3^{+4.2}_{-3.4} \pm 2.4$ | 1,2 CHISTOV | 06A | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ CHISTOV 06A reports $(9.3^{+3.7}_{-2.8} \pm 3.1) \times 10^{-5}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$.

We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\Lambda_c^+ \Lambda_c^- K^0)/\Gamma_{\text{total}}$ Γ_{365}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|------|---------------------------------------|
| $7.9^{+2.9}_{-2.3} \pm 4.3$ | 1 GABYSHEV | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3)\%$.

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{366}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|---------------------------------------|
| $<6.2 \times 10^{-7}$ | 90 | 1 VILLA | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $<1.7 \times 10^{-6}$ | 90 | 1 AUBERT | 01I | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| $<3.9 \times 10^{-5}$ | 90 | 2 ACCIARRI | 95I | L3 $e^+ e^- \rightarrow Z$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ACCIARRI 95I assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_{367}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|---------------------------------------|
| $<11.3 \times 10^{-8}$ | 90 | 1 AUBERT | 08P | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $< 6.1 \times 10^{-8}$ | 90 | 1 AUBERT | 05W | BABR Repl. by AUBERT 08P |
| $< 1.9 \times 10^{-7}$ | 90 | 1 CHANG | 03 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 8.3 \times 10^{-7}$ | 90 | 1 BERGFELD | 00B | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 1.4 \times 10^{-5}$ | 90 | 2 ACCIARRI | 97B | L3 $e^+ e^- \rightarrow Z$ |
| $< 5.9 \times 10^{-6}$ | 90 | AMMAR | 94 | CLE2 Repl. by BERGFELD 00B |
| $< 2.6 \times 10^{-5}$ | 90 | 3 AVERY | 89B | CLEO $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 7.6 \times 10^{-5}$ | 90 | 4 ALBRECHT | 87D | ARG $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 6.4 \times 10^{-5}$ | 90 | 5 AVERY | 87 | CLEO $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 3 \times 10^{-4}$ | 90 | GILES | 84 | CLEO Repl. by AVERY 87 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_s , and Λ_b .

³ Avery 89B reports $< 3 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

⁴ ALBRECHT 87D reports $< 8.5 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

⁵ Avery 87 reports $< 8 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(e^+e^-\gamma)/\Gamma_{\text{total}}$

Γ_{368}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------|----------|---------------------------------|
| $< 1.2 \times 10^{-7}$ | 90 | AUBERT | 08C BABR | $e^+e^- \rightarrow \gamma(4S)$ |

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

Γ_{369}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|--|
| $< 1.5 \times 10^{-8}$ | 90 | 1 AALTONEN | 08I CDF | $p\bar{p}$ at 1.96 TeV |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 5.2 \times 10^{-8}$ | 90 | 2 AUBERT | 08P BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| $< 3.9 \times 10^{-8}$ | 90 | 3 ABULENCIA | 05 CDF | Repl. by AALTONEN 08I |
| $< 8.3 \times 10^{-8}$ | 90 | 2 AUBERT | 05W BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| $< 1.5 \times 10^{-7}$ | 90 | 4 ACOSTA | 04D CDF | $p\bar{p}$ at 1.96 TeV |
| $< 1.6 \times 10^{-7}$ | 90 | 2 CHANG | 03 BELL | $e^+e^- \rightarrow \gamma(4S)$ |
| $< 6.1 \times 10^{-7}$ | 90 | 2 BERGFELD | 00B CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| $< 4.0 \times 10^{-5}$ | 90 | ABBOTT | 98B D0 | $p\bar{p}$ 1.8 TeV |
| $< 6.8 \times 10^{-7}$ | 90 | 5 ABE | 98 CDF | $p\bar{p}$ at 1.8 TeV |
| $< 1.0 \times 10^{-5}$ | 90 | 6 ACCIARRI | 97B L3 | $e^+e^- \rightarrow Z$ |
| $< 1.6 \times 10^{-6}$ | 90 | 7 ABE | 96L CDF | Repl. by ABE 98 |
| $< 5.9 \times 10^{-6}$ | 90 | AMMAR | 94 CLE2 | $e^+e^- \rightarrow \gamma(4S)$ |
| $< 8.3 \times 10^{-6}$ | 90 | 8 ALBAJAR | 91C UA1 | $E_{\text{cm}}^{p\bar{p}} = 630 \text{ GeV}$ |
| $< 1.2 \times 10^{-5}$ | 90 | 9 ALBAJAR | 91C UA1 | $E_{\text{cm}}^{p\bar{p}} = 630 \text{ GeV}$ |
| $< 4.3 \times 10^{-5}$ | 90 | 10 Avery | 89B CLEO | $e^+e^- \rightarrow \gamma(4S)$ |
| $< 4.5 \times 10^{-5}$ | 90 | 11 ALBRECHT | 87D ARG | $e^+e^- \rightarrow \gamma(4S)$ |
| $< 7.7 \times 10^{-5}$ | 90 | 12 Avery | 87 CLEO | $e^+e^- \rightarrow \gamma(4S)$ |
| $< 2 \times 10^{-4}$ | 90 | GILES | 84 CLEO | Repl. by Avery 87 |

¹ Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.86 \pm 0.59$, and the number of $B^+ \rightarrow J/\psi K^+$ decays.

² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

³ Assumes production cross section $\sigma(B^+)/\sigma(B_s) = 3.71 \pm 0.41$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+\mu^- K^+) = (5.88 \pm 0.26) \times 10^{-5}$.

⁴ Assumes production cross-section $\sigma(B_s)/\sigma(B^+) = 0.100/0.391$ and the CDF measured value of $\sigma(B^+) = 3.6 \pm 0.6 \mu\text{b}$.

⁵ ABE 98 assumes production of $\sigma(B^0) = \sigma(B^+)$ and $\sigma(B_s)/\sigma(B^0) = 1/3$. They normalize to their measured $\sigma(B^0, p_T(B) > 6, |y| < 1.0) = 2.39 \pm 0.32 \pm 0.44 \mu\text{b}$.

⁶ ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_s , and Λ_b .

⁷ ABE 96L assumes equal B^0 and B^+ production. They normalize to their measured $\sigma(B^+, p_T(B) > 6 \text{ GeV}/c, |y| < 1) = 2.39 \pm 0.54 \mu\text{b}$.

⁸ B^0 and B_s^0 are not separated.

- ⁹ Obtained from unseparated B^0 and B_s^0 measurement by assuming a $B^0:B_s^0$ ratio 2:1.
¹⁰ Avery 89B reports $< 5 \times 10^{-3}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.
¹¹ ALBRECHT 87D reports $< 5 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.
¹² Avery 87 reports $< 9 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(\mu^+\mu^-\gamma)/\Gamma_{\text{total}}$ Γ_{370}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------|----------|---------------------------------|
| $< 1.6 \times 10^{-7}$ | 90 | AUBERT | 08C BABR | $e^+e^- \rightarrow \gamma(4S)$ |

 $\Gamma(\tau^+\tau^-\gamma)/\Gamma_{\text{total}}$ Γ_{371}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|---------------------|----------|---------------------------------|
| $< 4.1 \times 10^{-3}$ | 90 | ¹ AUBERT | 06S BABR | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{374}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|---------------------|-----------|---------------------------------|
| $< 1.4 \times 10^{-7}$ | 90 | ¹ AUBERT | 07AG BABR | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\pi^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{375}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|---------------------|-----------|---------------------------------|
| $< 5.1 \times 10^{-7}$ | 90 | ¹ AUBERT | 07AG BABR | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\pi^0 \ell^+ \ell^-)/\Gamma_{\text{total}}$ Γ_{372}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|---------------------|-----------|---------------------------------|
| $< 1.2 \times 10^{-7}$ | 90 | ¹ AUBERT | 07AG BABR | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\pi^0 \nu\bar{\nu})/\Gamma_{\text{total}}$ Γ_{373}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interaction.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------------|----------|---------------------------------|
| $< 2.2 \times 10^{-4}$ | 90 | ¹ CHEN | 07D BELL | $e^+e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{379}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE (units 10^{-7}) | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-----|-----------------------|----------|---------------------------------|
| $1.3^{+1.6}_{-1.1} \pm 0.2$ | | ¹ AUBERT,B | 06J BABR | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|---|-----------------------------|---------------------|-----------------------|------|----------------------------------|----------------------------------|
| - | $2.1^{+2.3}_{-1.6} \pm 0.8$ | ¹ AUBERT | 03U | BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| < | 5.4 | 90 | ² ISHIKAWA | 03 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| < | 27 | 90 | ¹ ABE | 02 | BELL | Repl. by ISHIKAWA 03 |
| < | 38 | 90 | ¹ AUBERT | 02L | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| < | 84.5 | 90 | ³ ANDERSON | 01B | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| < | 3000 | 90 | ALBRECHT | 91E | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| < | 5200 | 90 | ⁴ AVERY | 87 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes equal production of B^0 and B^+ at $\gamma(4S)$.

³ The result is for di-lepton masses above 0.5 GeV.

⁴ AVERY 87 reports $< 6.5 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$

Γ_{380}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE (units 10^{-7}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

5.7 $^{+2.2}_{-1.8}$ OUR AVERAGE

| | | | | | |
|------------------------------|-----------------------|-----|------|----------------------------------|--|
| 5.9 $^{+3.3}_{-2.6} \pm 0.7$ | ¹ AUBERT,B | 06J | BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |
| 5.6 $^{+2.9}_{-2.3} \pm 0.5$ | ² ISHIKAWA | 03 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|----------------------------------|---------------------|-----------------------|------|-----------------------|----------------------------------|
| 1.63 $^{+0.82}_{-0.63} \pm 0.14$ | ¹ AUBERT | 03U | BABR | Repl. by AUBERT,B 06J | |
| <33 | 90 | ¹ ABE | 02 | BELL | Repl. by ISHIKAWA 03 |
| <36 | 90 | AUBERT | 02L | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| <66.4 | 90 | ³ ANDERSON | 01B | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| <5200 | 90 | ALBRECHT | 91E | ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| <3600 | 90 | ⁴ AVERY | 87 | CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes equal production of B^0 and B^+ at $\gamma(4S)$. The second error is a total of systematic uncertainties including model dependence.

³ The result is for di-lepton masses above 0.5 GeV.

⁴ AVERY 87 reports $< 4.5 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K^0 \ell^+ \ell^-)/\Gamma_{\text{total}}$

Γ_{376}/Γ

| VALUE (units 10^{-7}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-----------------------|------|---------------------------------------|
| 2.9 $^{+1.6}_{-1.3} \pm 0.3$ | | ¹ AUBERT,B | 06J | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------|----|-----------------------|----|------|----------------------------------|
| <6.8 | 90 | ¹ ISHIKAWA | 03 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
|------|----|-----------------------|----|------|----------------------------------|

¹ Assumes equal production of B^0 and B^+ at $\gamma(4S)$.

$\Gamma(K^0 \nu \bar{\nu})/\Gamma_{\text{total}}$ Γ_{377}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|------------|--------------------|-------------|----------------------------------|
| $< 1.6 \times 10^{-4}$ | 90 | ¹ CHEN | 07D BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\rho^0 \nu \bar{\nu})/\Gamma_{\text{total}}$ Γ_{378}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|------------|--------------------|-------------|----------------------------------|
| $< 4.4 \times 10^{-4}$ | 90 | ¹ CHEN | 07D BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^*(892)^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{382}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|-----------------------|-------------|----------------------------------|
| $1.04^{+0.33}_{-0.29} \pm 0.11$ | | ¹ AUBERT,B | 06J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------------------------------|----|-----------------------|----------|----------------------------------|
| $1.11^{+0.56}_{-0.47} \pm 0.11$ | | ¹ AUBERT | 03U BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 2.4 | 90 | ² ISHIKAWA | 03 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 6.4 | 90 | ¹ ABE | 02 BELL | Repl. by ISHIKAWA 03 |
| < 6.7 | 90 | ¹ AUBERT | 02L BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| <290 | 90 | ALBRECHT | 91E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² Assumes equal production of B^0 and B^+ at $\gamma(4S)$. $\Gamma(K^*(892)^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{383}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| <u>VALUE</u> (units 10^{-6}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------------|------------|--------------------|-------------|----------------|
| $1.10^{+0.29}_{-0.26}$ OUR AVERAGE | | | | |

 $0.87^{+0.38}_{-0.33} \pm 0.12$ ¹ AUBERT,B 06J BABR $e^+ e^- \rightarrow \gamma(4S)$ $1.33^{+0.42}_{-0.37} \pm 0.11$ ² ISHIKAWA 03 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------------------------------|----|-----------------------|----------|----------------------------------|
| $0.86^{+0.79}_{-0.58} \pm 0.11$ | | ¹ AUBERT | 03U BABR | Repl. by AUBERT,B 06J |
| < 4.2 | 90 | ¹ ABE | 02 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 3.3 | 90 | AUBERT | 02L BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| < 4.0 | 90 | ³ AFFOLDER | 99B CDF | $p\bar{p}$ at 1.8 TeV |
| < 25 | 90 | ⁴ ABE | 96L CDF | Repl. by AFFOLDER 99B |
| < 23 | 90 | ⁵ ALBAJAR | 91C UA1 | $E_{cm}^{pp} = 630$ GeV |
| <340 | 90 | ALBRECHT | 91E ARG | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² Assumes equal production of B^0 and B^+ at $\gamma(4S)$. The second error is a total of systematic uncertainties including model dependence.³ AFFOLDER 99B measured relative to $B^0 \rightarrow J/\psi(1S) K^*(892)^0$.⁴ ABE 96L measured relative to $B^0 \rightarrow J/\psi(1S) K^*(892)^0$ using PDG 94 branching ratios.⁵ ALBAJAR 91C assumes 36% of \bar{b} quarks give B^0 mesons.

$\Gamma(K^*(892)^0 \ell^+ \ell^-)/\Gamma_{\text{total}}$ Γ_{381}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE (units 10^{-7}) | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----------------------|----------|----------------------------------|
| 9.5±1.8 OUR AVERAGE | | | |
| $8.1^{+2.1}_{-1.9} \pm 0.9$ | ¹ AUBERT,B | 06J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $11.7^{+3.0}_{-2.7} \pm 0.9$ | ¹ ISHIKAWA | 03 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Assumes equal production of B^0 and B^+ at $\gamma(4S)$. $\Gamma(K^*(892)^0 \nu \bar{\nu})/\Gamma_{\text{total}}$ Γ_{384}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------------|----------|----------------------------------|
| $< 3.4 \times 10^{-4}$ | 90 | ¹ CHEN | 07D BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $< 1.0 \times 10^{-3}$ | 90 | ² ADAM | 96D DLPH | $e^+ e^- \rightarrow Z$ |
| 1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. | | | | |
| 2 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. | | | | |

 $\Gamma(\phi \nu \bar{\nu})/\Gamma_{\text{total}}$ Γ_{385}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------------|----------|----------------------------------|
| $< 5.8 \times 10^{-5}$ | 90 | ¹ CHEN | 07D BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. | | | | |

 $\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{386}/Γ

Test of lepton family number conservation. Allowed by higher-order electroweak interactions.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-----------------------|----------|----------------------------------|
| $< 9.2 \times 10^{-8}$ | 90 | ¹ AUBERT | 08P BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $< 1.8 \times 10^{-7}$ | 90 | ¹ AUBERT | 05W BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 1.7 \times 10^{-7}$ | 90 | ¹ CHANG | 03 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 15 \times 10^{-7}$ | 90 | ¹ BERGFELD | 00B CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 3.5 \times 10^{-6}$ | 90 | ABE | 98V CDF | $p\bar{p}$ at 1.8 TeV |
| $< 1.6 \times 10^{-5}$ | 90 | ² ACCIARRI | 97B L3 | $e^+ e^- \rightarrow Z$ |
| $< 5.9 \times 10^{-6}$ | 90 | AMMAR | 94 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 3.4 \times 10^{-5}$ | 90 | ³ AVERY | 89B CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 4.5 \times 10^{-5}$ | 90 | ⁴ ALBRECHT | 87D ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 7.7 \times 10^{-5}$ | 90 | ⁵ AVERY | 87 CLEO | $e^+ e^- \rightarrow \gamma(4S)$ |
| $< 3 \times 10^{-4}$ | 90 | GILES | 84 CLEO | Repl. by AVERY 87 |

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_s , and Λ_b .³ Paper assumes the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.⁴ ALBRECHT 87D reports $< 5 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.⁵ AVERY 87 reports $< 9 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^0 e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{387}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|-----------|----------------------------------|
| $<1.4 \times 10^{-7}$ | 90 | 1 AUBERT | 07AG BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^0 e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{388}/Γ

Test of lepton family number conservation.

| VALUE (units 10^{-7}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|----------------------------------|
| < 2.7 | 90 | 1 AUBERT,B | 06J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<40 90 1 AUBERT 02L BABR Repl. by AUBERT,B 06J

1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^*(892)^0 e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{389}/Γ

| VALUE (units 10^{-7}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|----------------------------------|
| < 5.3 | 90 | 1 AUBERT,B | 06J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

1 Assumes equal production of B^0 and B^+ at $\gamma(4S)$. $\Gamma(K^*(892)^0 e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{390}/Γ

| VALUE (units 10^{-7}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|----------------------------------|
| < 3.4 | 90 | 1 AUBERT,B | 06J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

1 Assumes equal production of B^0 and B^+ at $\gamma(4S)$. $\Gamma(K^*(892)^0 e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{391}/Γ

Test of lepton family number conservation.

| VALUE (units 10^{-7}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|----------------------------------|
| < 5.8 | 90 | 1 AUBERT,B | 06J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<34 90 1 AUBERT 02L BABR Repl. by AUBERT,B 06J

1 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{392}/Γ

Test of lepton family number conservation. Allowed by higher-order electroweak interactions.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|------|----------------------------------|
| $<1.1 \times 10^{-4}$ | 90 | BORNHEIM 04 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.3 × 10⁻⁴ 90 AMMAR 94 CLE2 Repl. by BORNHEIM 04 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{393}/Γ

Test of lepton family number conservation. Allowed by higher-order electroweak interactions.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|------|----------------------------------|
| $<3.8 \times 10^{-5}$ | 90 | BORNHEIM 04 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8.3 × 10⁻⁴ 90 AMMAR 94 CLE2 Repl. by BORNHEIM 04

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$ Γ_{394}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-----------------------|-------------|----------------------------------|
| <22 | 90 | ¹ AUBERT,B | 04J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses the fully reconstructed $B^0 \rightarrow D^{(*)} - \ell + \nu_\ell$ events as a tag.

 $\Gamma(\nu\bar{\nu}\gamma)/\Gamma_{\text{total}}$ Γ_{395}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-----------------------|-------------|----------------------------------|
| <4.7 | 90 | ¹ AUBERT,B | 04J BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses the fully reconstructed $B^0 \rightarrow D^{(*)} - \ell + \nu_\ell$ events as a tag.

A REVIEW GOES HERE – Check our WWW List of Reviews

POLARIZATION IN B^0 DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal (L) or both are transverse and parallel (\parallel) or perpendicular (\perp) to each other with the parameters Γ_L/Γ , Γ_\perp/Γ , and the relative phases ϕ_\parallel and ϕ_\perp . See the definitions in the note on “Polarization in B Decays” review in the B^0 Particle Listings.

 Γ_L/Γ in $B^0 \rightarrow J/\psi(1S) K^*(892)^0$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-----------------------|-------------|----------------------------------|
| 0.566±0.010 OUR AVERAGE | | | | |
| 0.556±0.009±0.010 | | ¹ AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.574±0.012±0.009 | | ITOH | 05 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.59 ± 0.06 ± 0.01 | | ² AFFOLDER | 00N CDF | $p\bar{p}$ at 1.8 TeV |
| 0.52 ± 0.07 ± 0.04 | | ³ JESSOP | 97 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.65 ± 0.10 ± 0.04 | 65 | ABE | 95Z CDF | $p\bar{p}$ at 1.8 TeV |
| 0.97 ± 0.16 ± 0.15 | 13 | ⁴ ALBRECHT | 94G ARG | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.566±0.012±0.005 | | ¹ AUBERT | 05P BABR | Repl. by AUBERT 07AD |
| 0.62 ± 0.02 ± 0.03 | | ⁵ ABE | 02N BELL | Repl. by ITOH 05 |
| 0.597±0.028±0.024 | | ⁶ AUBERT | 01H BABR | Repl. by AUBERT 07AD |
| 0.80 ± 0.08 ± 0.05 | 42 | ⁴ ALAM | 94 CLE2 | Sup. by JESSOP 97 |

¹ Obtained by combining the B^0 and B^+ modes.

² AFFOLDER 00N measurements are based on 190 B^0 candidates obtained from a data sample of 89 pb^{-1} . The P -wave fraction is found to be $0.13^{+0.12}_{-0.09} \pm 0.06$.

³ JESSOP 97 is the average over a mixture of B^0 and B^+ decays. The P -wave fraction is found to be $0.16 \pm 0.08 \pm 0.04$.

⁴ Averaged over an admixture of B^0 and B^+ decays.

⁵ Averaged over an admixture of B^0 and B^+ decays and the P wave fraction is $(19 \pm 2 \pm 3)\%$.

⁶ Averaged over an admixture of B^0 and B^- decays and the P wave fraction is $(16.0 \pm 3.2 \pm 1.4) \times 10^{-2}$.

Γ_{\perp}/Γ in $B^0 \rightarrow J/\psi K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|---------------------|-----------|-------------------------------------|
| 0.219±0.018 OUR AVERAGE | | | Error includes scale factor of 2.1. |
| 0.233±0.010±0.005 | ¹ AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.195±0.012±0.008 | ITOH | 05 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by combining the B^0 and B^+ modes.

 ϕ_{\parallel} in $B^0 \rightarrow J/\psi K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|---------------------|-----------|----------------------------------|
| -2.93±0.08±0.04 | ¹ AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by combining the B^0 and B^+ modes.

 ϕ_{\perp} in $B^0 \rightarrow J/\psi K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|---------------------|-----------|----------------------------------|
| 2.91±0.05±0.03 | ¹ AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by combining the B^0 and B^+ modes.

 Γ_L/Γ in $B^0 \rightarrow \psi(2S) K^*(892)^0$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----------------------|-----------|----------------------------------|
| 0.47±0.05 OUR AVERAGE | | | |
| 0.48±0.05±0.02 | ¹ AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.45±0.11±0.04 | ² RICHICHI | 01 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by combining the B^0 and B^+ modes.

² Averages between charged and neutral B mesons.

 Γ_{\perp}/Γ in $B^0 \rightarrow \psi(2S) K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|---------------------|-----------|----------------------------------|
| 0.30±0.06±0.02 | ¹ AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by combining the B^0 and B^+ modes.

 ϕ_{\parallel} in $B^0 \rightarrow \psi(2S) K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------|---------------------|-----------|----------------------------------|
| -2.8±0.4±0.1 | ¹ AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by combining the B^0 and B^+ modes.

 ϕ_{\perp} in $B^0 \rightarrow \psi(2S) K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------|---------------------|-----------|----------------------------------|
| 2.8±0.3 ±0.1 | ¹ AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by combining the B^0 and B^+ modes.

 Γ_L/Γ in $B^0 \rightarrow \chi_{c1} K^*(892)^0$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|---------------------|-----------|----------------------------------|
| 0.77±0.07±0.04 | ¹ AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by combining the B^0 and B^+ modes.

Γ_{\perp}/Γ in $B^0 \rightarrow \chi_{c1} K^*(892)^0$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|-------------|----------------------------------|
| 0.03±0.04±0.02 | 1 AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by combining the B^0 and B^+ modes.

 $\phi_{||}$ in $B^0 \rightarrow \chi_{c1} K^*(892)^0$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------|--------------------|-------------|----------------------------------|
| 0.0±0.3 ±0.1 | 1 AUBERT | 07AD BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Obtained by combining the B^0 and B^+ modes.

 Γ_L/Γ in $B^0 \rightarrow D_s^{*+} D^{*-}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------|--------------------|-------------|----------------------------------|
| 0.52 ±0.05 OUR AVERAGE | | | |
| 0.519±0.050±0.028 | 1 AUBERT | 03I BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.506±0.139±0.036 | AHMED | 00B CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Measurement performed using partial reconstruction of D^{*-} decay.

 Γ_L/Γ in $B^0 \rightarrow D^{*-} \rho^+$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|----------------------------------|
| 0.885±0.016±0.012 | | CSORNA | 03 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.93 ± 0.05 ± 0.05 | 76 | ALAM | 94 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

 Γ_L/Γ in $B^0 \rightarrow D^{*+} D^{*-}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|-------------|----------------------------------|
| 0.57±0.08±0.02 | MIYAKE | 05 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

 Γ_{\perp}/Γ in $B^0 \rightarrow D^{*+} D^{*-}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------------------------|
| 0.150±0.032 OUR AVERAGE | AUBERT | 07BO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.125±0.044±0.007 | AUBERT,BE | 05A BABR | Repl. by AUBERT 07BO |
| 0.063±0.055±0.009 | AUBERT | 03Q BABR | Repl. by AUBERT,BE 05A |

 Γ_L/Γ in $B^0 \rightarrow D^{*-} \omega \pi^+$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|--------------------|-------------|----------------------------------|
| 0.654±0.042±0.016 | 1 AUBERT | 06L BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Invariant mass of the $[\omega \pi]$ system is restricted in the region 1.1 and 1.9 GeV.

 Γ_L/Γ in $B^0 \rightarrow \phi K^*(892)^0$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|--------------------|-------------|----------------------------------|
| 0.484±0.033 OUR AVERAGE | | | |
| 0.506±0.040±0.015 | AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.45 ± 0.05 ± 0.02 | CHEN | 05A BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------------------------|-----------------------|----------|-----------------------|
| $0.52 \pm 0.05 \pm 0.02$ | ¹ AUBERT,B | 04W BABR | Repl. by AUBERT 07D |
| $0.65 \pm 0.07 \pm 0.02$ | AUBERT | 03V BABR | Repl. by AUBERT,B 04W |
| $0.41 \pm 0.10 \pm 0.04$ | CHEN | 03B BELL | Repl. by CHEN 05A |

¹ AUBERT,B 04W also measures the fraction of parity-odd transverse contribution $f_{\perp} = 0.22 \pm 0.05 \pm 0.02$ and the phases of the parity-even and parity-odd transverse amplitudes relative to the longitudinal amplitude.

Γ_L/Γ in $B^0 \rightarrow K^{*0}\bar{K}^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|------|--------------------------------------|
| $0.80^{+0.10}_{-0.12} \pm 0.06$ | AUBERT | 08I | BABR $e^+e^- \rightarrow \gamma(4S)$ |

Γ_{\perp}/Γ in $B^0 \rightarrow \phi K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|------|--------------------------------------|
| 0.26 ± 0.04 OUR AVERAGE | | | Error includes scale factor of 1.2. |
| $0.227 \pm 0.038 \pm 0.013$ | AUBERT | 07D | BABR $e^+e^- \rightarrow \gamma(4S)$ |
| $0.31^{+0.06}_{-0.05} \pm 0.02$ | ¹ CHEN | 05A | BELL $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------------------------|----------|----------|---------------------|
| $0.22 \pm 0.05 \pm 0.02$ | AUBERT,B | 04W BABR | Repl. by AUBERT 07D |
|--------------------------|----------|----------|---------------------|

¹ This quantity was recalculated by the BELLE authors from numbers in the original paper.

$\phi_{||}$ in $B^0 \rightarrow \phi K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|------|--------------------------------------|
| 2.33 ± 0.14 OUR AVERAGE | | | |
| $2.31 \pm 0.14 \pm 0.08$ | AUBERT | 07D | BABR $e^+e^- \rightarrow \gamma(4S)$ |
| $2.40^{+0.28}_{-0.24} \pm 0.07$ | ¹ CHEN | 05A | BELL $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------------------------------|----------|----------|---------------------|
| $2.34^{+0.23}_{-0.20} \pm 0.05$ | AUBERT,B | 04W BABR | Repl. by AUBERT 07D |
|---------------------------------|----------|----------|---------------------|

¹ This quantity was recalculated by the BELLE authors from numbers in the original paper.

ϕ_{\perp} in $B^0 \rightarrow \phi K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--------------------------------------|
| 2.33 ± 0.14 OUR AVERAGE | | | |
| $2.24 \pm 0.15 \pm 0.09$ | AUBERT | 07D | BABR $e^+e^- \rightarrow \gamma(4S)$ |

| | | | |
|--------------------------|-------------------|-----|--------------------------------------|
| $2.51 \pm 0.25 \pm 0.06$ | ¹ CHEN | 05A | BELL $e^+e^- \rightarrow \gamma(4S)$ |
|--------------------------|-------------------|-----|--------------------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------------------------|----------|----------|---------------------|
| $2.47 \pm 0.25 \pm 0.05$ | AUBERT,B | 04W BABR | Repl. by AUBERT 07D |
|--------------------------|----------|----------|---------------------|

¹ This quantity was recalculated by the BELLE authors from numbers in the original paper.

$\delta_0(B^0 \rightarrow \phi K^{*0})$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|--------------------------------------|
| $2.78 \pm 0.17 \pm 0.09$ | AUBERT | 07D | BABR $e^+e^- \rightarrow \gamma(4S)$ |

A_{CP}^0 in $B^0 \rightarrow \phi K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|----------|----------------------------------|
| 0.02±0.07 OUR AVERAGE | Error includes scale factor of 1.1. | | |
| -0.03±0.08±0.02 | AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.13±0.12±0.04 | ¹ CHEN | 05A BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.06±0.10±0.01 | AUBERT,B | 04W BABR | Repl. by AUBERT 07D |

¹ This quantity was recalculated by the BELLE authors from numbers in the original paper.

 A_{CP}^\perp in $B^0 \rightarrow \phi K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|----------|----------------------------------|
| -0.11±0.12 OUR AVERAGE | Error includes scale factor of 1.1. | | |
| -0.03±0.16±0.05 | AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.20±0.18±0.04 | ¹ CHEN | 05A BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.10±0.24±0.05 | AUBERT,B | 04W BABR | Repl. by AUBERT 07D |

¹ This quantity was recalculated by the BELLE authors from numbers in the original paper.

 $\Delta\phi_\parallel$ in $B^0 \rightarrow \phi K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|----------|----------------------------------|
| 0.10±0.24 OUR AVERAGE | Error includes scale factor of 1.7. | | |
| 0.24±0.14±0.08 | AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.32±0.27±0.07 | ¹ CHEN | 05A BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.27 ^{+0.20} _{-0.23} ±0.05 | AUBERT,B | 04W BABR | Repl. by AUBERT 07D |

¹ This quantity was recalculated by the BELLE authors from numbers in the original paper.

 $\Delta\phi_\perp$ in $B^0 \rightarrow \phi K^{*0}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|----------|----------------------------------|
| 0.04±0.23 OUR AVERAGE | Error includes scale factor of 1.6. | | |
| 0.19±0.15±0.08 | AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.30±0.25±0.06 | ¹ CHEN | 05A BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.36±0.25±0.05 | AUBERT,B | 04W BABR | Repl. by AUBERT 07D |

¹ This quantity was recalculated by the BELLE authors from numbers in the original paper.

 $\Delta\delta_0(B^0 \rightarrow \phi K^{*0})$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------------------------------|----------------------------------|---------|
| 0.21±0.17±0.08 | Error includes scale factor of 1.6. | | |
| AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 Γ_L/Γ in $B^0 \rightarrow \phi K_2^*(1430)^0$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|----------------------------------|---------|
| 0.853^{+0.061}_{-0.069}±0.036 | Error includes scale factor of 1.6. | | |
| AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 Γ_\perp/Γ in $B^0 \rightarrow \phi K_2^*(1430)^0$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|----------------------------------|---------|
| 0.045^{+0.049}_{-0.040}±0.013 | Error includes scale factor of 1.6. | | |
| AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

ϕ_{\parallel} in $B^0 \rightarrow \phi K_2^*(1430)^0$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|------|---------------------------------------|
| 2.90±0.39±0.06 | AUBERT | 07D | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

 ϕ_{\perp} in $B^0 \rightarrow \phi K_2^*(1430)^0$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|---------------------------------------|
| 5.72^{+0.55}_{-0.87}±0.11 | AUBERT | 07D | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

 $\delta_0(B^0 \rightarrow \phi K_2^*(1430)^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|---------------------------------------|
| 3.54^{+0.12}_{-0.14}±0.06 | AUBERT | 07D | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

 Γ_L/Γ in $B^0 \rightarrow K^*(892)^0 \rho^0$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|------|---------------------------------------|
| 0.57±0.09±0.08 | AUBERT,B | 06G | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

 Γ_L/Γ in $B^0 \rightarrow \rho^+ \rho^-$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|---------|
| 0.977^{+0.028}_{-0.024} OUR AVERAGE | | | |

 $0.992 \pm 0.024^{+0.026}_{-0.013}$ AUBERT 07BF BABR $e^+ e^- \rightarrow \gamma(4S)$ $0.941^{+0.034}_{-0.040} \pm 0.030$ SOMOV 06 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.978 \pm 0.014^{+0.021}_{-0.029}$

AUBERT,B 05C BABR Repl. by AUBERT 07BF

 $0.98^{+0.02}_{-0.08} \pm 0.03$

AUBERT 04G BABR Repl. by AUBERT,B 04R

 $0.99 \pm 0.03^{+0.04}_{-0.03}$

AUBERT,B 04R BABR Repl. by AUBERT,B 05C

 Γ_L/Γ in $B^0 \rightarrow \rho^0 \rho^0$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|------|---------------------------------------|
| 0.87±0.13±0.04 | AUBERT | 07G | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

A REVIEW GOES HERE – Check our WWW List of Reviews

 $B^0-\bar{B}^0$ MIXING PARAMETERSFor a discussion of $B^0-\bar{B}^0$ mixing see the note on “ $B^0-\bar{B}^0$ Mixing” in the B^0 Particle Listings above. χ_d is a measure of the time-integrated $B^0-\bar{B}^0$ mixing probability that a produced $B^0(\bar{B}^0)$ decays as a $\bar{B}^0(B^0)$. Mixing violates $\Delta B \neq 2$ rule.

$$\chi_d = \frac{x_d^2}{2(1+x_d^2)}$$

$$\chi_d = \frac{\Delta m_{B^0}}{\Gamma_{B^0}} = (m_{B_H^0} - m_{B_L^0}) \tau_{B^0},$$

where H, L stand for heavy and light states of two B^0 CP eigenstates and

$$\tau_{B^0} = \frac{1}{0.5(\Gamma_{B_H^0} + \Gamma_{B_L^0})}.$$

χ_d

This B^0 - \bar{B}^0 mixing parameter is the probability (integrated over time) that a produced B^0 (or \bar{B}^0) decays as a \bar{B}^0 (or B^0), e.g. for inclusive lepton decays

$$\begin{aligned}\chi_d &= \Gamma(B^0 \rightarrow \ell^- X \text{ (via } \bar{B}^0)) / \Gamma(B^0 \rightarrow \ell^\pm X) \\ &= \Gamma(\bar{B}^0 \rightarrow \ell^+ X \text{ (via } B^0)) / \Gamma(\bar{B}^0 \rightarrow \ell^\pm X)\end{aligned}$$

Where experiments have measured the parameter $r = \chi/(1-\chi)$, we have converted to χ . Mixing violates the $\Delta B \neq 2$ rule.

Note that the measurement of χ at energies higher than the $\Upsilon(4S)$ have not separated χ_d from χ_s where the subscripts indicate $B^0(\bar{b}d)$ or $B_s^0(\bar{b}s)$. They are listed in the $B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE section.

The experiments at $\Upsilon(4S)$ make an assumption about the $B^0\bar{B}^0$ fraction and about the ratio of the B^\pm and B^0 semileptonic branching ratios (usually that it equals one).

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements, includes χ_d calculated from Δm_{B^0} and τ_{B^0} .

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|--------|-----------------------|----------|-----------------------------------|
| 0.1878±0.0024 OUR EVALUATION | | | | |
| 0.182 ±0.015 OUR AVERAGE | | | | |
| 0.198 ±0.013 | ±0.014 | ¹ BEHRENS | 00B CLE2 | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.16 ±0.04 | ±0.04 | ² ALBRECHT | 94 ARG | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.149 ±0.023 | ±0.022 | ³ BARTEL | 93 CLE2 | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.171 ±0.048 | | ⁴ ALBRECHT | 92L ARG | $e^+e^- \rightarrow \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.20 ±0.13 | ±0.12 | ⁵ ALBRECHT | 96D ARG | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.19 ±0.07 | ±0.09 | ⁶ ALBRECHT | 96D ARG | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.24 ±0.12 | | ⁷ ELSEN | 90 JADE | e^+e^- 35–44 GeV |
| 0.158 ^{+0.052} _{-0.059} | | ARTUSO | 89 CLEO | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.17 ±0.05 | | ⁸ ALBRECHT | 87I ARG | $e^+e^- \rightarrow \Upsilon(4S)$ |
| <0.19 | 90 | ⁹ BEAN | 87B CLEO | $e^+e^- \rightarrow \Upsilon(4S)$ |
| <0.27 | 90 | ¹⁰ AVERY | 84 CLEO | $e^+e^- \rightarrow \Upsilon(4S)$ |

¹ BEHRENS 00B uses high-momentum lepton tags and partially reconstructed $\bar{B}^0 \rightarrow D^{*+}\pi^-$, ρ^- decays to determine the flavor of the B meson.

² ALBRECHT 94 reports $r=0.194 \pm 0.062 \pm 0.054$. We convert to χ for comparison. Uses tagged events (lepton + pion from D^*).

³ BARTEL 93 analysis performed using tagged events (lepton+pion from D^*). Using dilepton events they obtain 0.157 ± 0.016 ^{+0.033}_{-0.028}.

⁴ ALBRECHT 92L is a combined measurement employing several lepton-based techniques. It uses all previous ARGUS data in addition to new data and therefore supersedes ALBRECHT 87I. A value of $r = 20.6 \pm 7.0\%$ is directly measured. The value can be used to measure $x = \Delta M/\Gamma = 0.72 \pm 0.15$ for the B_d meson. Assumes $f_{+-}/f_0 = 1.0 \pm 0.05$ and uses $\tau_{B^\pm}/\tau_{B^0} = (0.95 \pm 0.14) (f_{+-}/f_0)$.

⁵ Uses $D^{*+} K^\pm$ correlations.

⁶ Uses $(D^{*+} \ell^-) K^\pm$ correlations.

⁷ These experiments see a combination of B_s and B_d mesons.

⁸ ALBRECHT 87I is inclusive measurement with like-sign dileptons, with tagged B decays plus leptons, and one fully reconstructed event. Measures $r = 0.21 \pm 0.08$. We convert to χ for comparison. Superseded by ALBRECHT 92L.

⁹ BEAN 87B measured $r < 0.24$; we converted to χ .

¹⁰ Same-sign dilepton events. Limit assumes semileptonic BR for B^+ and B^0 equal. If B^0/B^\pm ratio < 0.58 , no limit exists. The limit was corrected in BEAN 87B from $r < 0.30$ to $r < 0.37$. We converted this limit to χ .

$$\Delta m_{B^0} = m_{B_H^0} - m_{B_L^0}$$

Δm_{B^0} is a measure of 2π times the B^0 - \bar{B}^0 oscillation frequency in time-dependent mixing experiments.

The second “OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

The first “OUR EVALUATION”, also provided by the HFAG, includes Δm_d calculated from χ_d measured at $\Upsilon(4S)$.

| VALUE ($10^{12} \text{ } \text{Hz} \text{ s}^{-1}$) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------|----------|------------------------------------|
| 0.507±0.005 OUR EVALUATION | First | | |
| 0.507±0.005 OUR EVALUATION | Second | | |
| 0.506±0.020±0.016 | ¹ ABAZOV | 06W D0 | $p\bar{p}$ at 1.96 TeV |
| 0.511±0.007 ^{+0.007} _{-0.006} | ² AUBERT | 06G BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 0.511±0.005±0.006 | ³ ABE | 05B BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 0.531±0.025±0.007 | ⁴ ABDALLAH | 03B DLPH | $e^+ e^- \rightarrow Z$ |
| 0.503±0.008±0.010 | ⁵ HASTINGS | 03 BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 0.509±0.017±0.020 | ⁶ ZHENG | 03 BELL | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 0.516±0.016±0.010 | ⁷ AUBERT | 02I BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 0.493±0.012±0.009 | ⁸ AUBERT | 02J BABR | $e^+ e^- \rightarrow \Upsilon(4S)$ |
| 0.497±0.024±0.025 | ⁹ ABBIENDI,G | 00B OPAL | $e^+ e^- \rightarrow Z$ |
| 0.503±0.064±0.071 | ¹⁰ ABE | 99K CDF | $p\bar{p}$ at 1.8 TeV |
| 0.500±0.052±0.043 | ¹¹ ABE | 99Q CDF | $p\bar{p}$ at 1.8 TeV |
| 0.516±0.099 ^{+0.029} _{-0.035} | ¹² AFFOLDER | 99C CDF | $p\bar{p}$ at 1.8 TeV |
| 0.471 ^{+0.078} _{-0.068} ^{+0.033} _{-0.034} | ¹³ ABE | 98C CDF | $p\bar{p}$ at 1.8 TeV |
| 0.458±0.046±0.032 | ¹⁴ ACCIARRI | 98D L3 | $e^+ e^- \rightarrow Z$ |
| 0.437±0.043±0.044 | ¹⁵ ACCIARRI | 98D L3 | $e^+ e^- \rightarrow Z$ |
| 0.472±0.049±0.053 | ¹⁶ ACCIARRI | 98D L3 | $e^+ e^- \rightarrow Z$ |

| | | | | | |
|---|----|------------|-----|------|-------------------------|
| 0.523±0.072±0.043 | 17 | ABREU | 97N | DLPH | $e^+ e^- \rightarrow Z$ |
| 0.493±0.042±0.027 | 15 | ABREU | 97N | DLPH | $e^+ e^- \rightarrow Z$ |
| 0.499±0.053±0.015 | 18 | ABREU | 97N | DLPH | $e^+ e^- \rightarrow Z$ |
| 0.480±0.040±0.051 | 14 | ABREU | 97N | DLPH | $e^+ e^- \rightarrow Z$ |
| 0.444±0.029 ^{+0.020} _{-0.017} | 15 | ACKERSTAFF | 97U | OPAL | $e^+ e^- \rightarrow Z$ |
| 0.430±0.043 ^{+0.028} _{-0.030} | 14 | ACKERSTAFF | 97V | OPAL | $e^+ e^- \rightarrow Z$ |
| 0.482±0.044±0.024 | 19 | BUSKULIC | 97D | ALEP | $e^+ e^- \rightarrow Z$ |
| 0.404±0.045±0.027 | 15 | BUSKULIC | 97D | ALEP | $e^+ e^- \rightarrow Z$ |
| 0.452±0.039±0.044 | 14 | BUSKULIC | 97D | ALEP | $e^+ e^- \rightarrow Z$ |
| 0.539±0.060±0.024 | 20 | ALEXANDER | 96V | OPAL | $e^+ e^- \rightarrow Z$ |
| 0.567±0.089 ^{+0.029} _{-0.023} | 21 | ALEXANDER | 96V | OPAL | $e^+ e^- \rightarrow Z$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---|----|------------|-----|------|----------------------------------|
| 0.492±0.018±0.013 | 22 | AUBERT | 03C | BABR | Repl. by AUBERT 06G |
| 0.516±0.016±0.010 | 23 | AUBERT | 02N | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.494±0.012±0.015 | 24 | HARA | 02 | BELL | Repl. by ABE 05B |
| 0.528±0.017±0.011 | 25 | TOMURA | 02 | BELL | Repl. by ABE 05B |
| 0.463±0.008±0.016 | 8 | ABE | 01D | BELL | Repl. by HASTINGS 03 |
| 0.444±0.028±0.028 | 26 | ACCIARRI | 98D | L3 | $e^+ e^- \rightarrow Z$ |
| 0.497±0.035 | 27 | ABREU | 97N | DLPH | $e^+ e^- \rightarrow Z$ |
| 0.467±0.022 ^{+0.017} _{-0.015} | 28 | ACKERSTAFF | 97V | OPAL | $e^+ e^- \rightarrow Z$ |
| 0.446±0.032 | 29 | BUSKULIC | 97D | ALEP | $e^+ e^- \rightarrow Z$ |
| 0.531 ^{+0.050} _{-0.046} ±0.078 | 30 | ABREU | 96Q | DLPH | Sup. by ABREU 97N |
| 0.496 ^{+0.055} _{-0.051} ±0.043 | 14 | ACCIARRI | 96E | L3 | Repl. by ACCIARRI 98D |
| 0.548±0.050 ^{+0.023} _{-0.019} | 31 | ALEXANDER | 96V | OPAL | $e^+ e^- \rightarrow Z$ |
| 0.496±0.046 | 32 | AKERS | 95J | OPAL | Repl. by ACKERSTAFF 97V |
| 0.462 ^{+0.040} _{-0.053} ^{+0.052} _{-0.035} | 14 | AKERS | 95J | OPAL | Repl. by ACKERSTAFF 97V |
| 0.50 ±0.12 ±0.06 | 17 | ABREU | 94M | DLPH | Sup. by ABREU 97N |
| 0.508±0.075±0.025 | 20 | AKERS | 94C | OPAL | Repl. by ALEXANDER 96V |
| 0.57 ±0.11 ±0.02 | 21 | AKERS | 94H | OPAL | Repl. by ALEXANDER 96V |
| 0.50 ^{+0.07} _{-0.06} ^{+0.11} _{-0.10} | 14 | BUSKULIC | 94B | ALEP | Sup. by BUSKULIC 97D |
| 0.52 ^{+0.10} _{-0.11} ^{+0.04} _{-0.03} | 21 | BUSKULIC | 93K | ALEP | Sup. by BUSKULIC 97D |

¹ Uses opposite-side flavor-tagging with $B \rightarrow D^{(*)} \mu \nu_\mu X$ events.

² Measured using a simultaneous fit of the B^0 lifetime and $\bar{B}^0 B^0$ oscillation frequency Δm_d in the partially reconstructed $B^0 \rightarrow D^{*-} \ell \nu$ decays.

³ Measurement performed using a combined fit of CP -violation, mixing and lifetimes.

⁴ Events with a high transverse momentum lepton were removed and an inclusively reconstructed vertex was required.

⁵ HASTINGS 03 measurement based on the time evolution of dilepton events. It also reports $f_+/f_0 = 1.01 \pm 0.03 \pm 0.09$ and CPT violation parameters in B^0 - \bar{B}^0 mixing.

⁶ ZHENG 03 data analyzed using partially reconstructed $\bar{B}^0 \rightarrow D^{*-} \pi^+$ decay and a flavor tag based on the charge of the lepton from the accompanying B decay.

⁷ Uses a tagged sample of fully-reconstructed neutral B decays at $\gamma(4S)$.

⁸ Measured based on the time evolution of dilepton events in $\gamma(4S)$ decays.

- ⁹ Data analyzed using partially reconstructed $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}$ decay and a combination of flavor tags from the rest of the event.
- ¹⁰ Uses di-muon events.
- ¹¹ Uses jet-charge and lepton-flavor tagging.
- ¹² Uses $\ell^- D^{*+} - \ell$ events.
- ¹³ Uses π - B in the same side.
- ¹⁴ Uses ℓ - ℓ .
- ¹⁵ Uses ℓ - Q_{hem} .
- ¹⁶ Uses ℓ - ℓ with impact parameters.
- ¹⁷ Uses $D^{*\pm}$ - Q_{hem} .
- ¹⁸ Uses π_s^\pm ℓ - Q_{hem} .
- ¹⁹ Uses $D^{*\pm}$ - ℓ / Q_{hem} .
- ²⁰ Uses $D^{*\pm}$ ℓ - Q_{hem} .
- ²¹ Uses $D^{*\pm}$ - ℓ .
- ²² AUBERT 03C uses a sample of approximately 14,000 exclusively reconstructed $B^0 \rightarrow D^*(2010)^- \ell \bar{\nu}$ and simultaneously measures the lifetime and oscillation frequency.
- ²³ AUBERT 02N result based on the same analysis and data sample reported in AUBERT 02I.
- ²⁴ Uses a tagged sample of B^0 decays reconstructed in the mode $B^0 \rightarrow D^* \ell \bar{\nu}$.
- ²⁵ Uses a tagged sample of fully-reconstructed hadronic B^0 decays at $\Upsilon(4S)$.
- ²⁶ ACCIARRI 98D combines results from ℓ - ℓ , ℓ - Q_{hem} , and ℓ - ℓ with impact parameters.
- ²⁷ ABREU 97N combines results from $D^{*\pm}$ - Q_{hem} , ℓ - Q_{hem} , π_s^\pm ℓ - Q_{hem} , and ℓ - ℓ .
- ²⁸ ACKERSTAFF 97V combines results from ℓ - ℓ , ℓ - Q_{hem} , $D^{*\pm}$ - ℓ , and $D^{*\pm}$ - Q_{hem} .
- ²⁹ BUSKULIC 97D combines results from $D^{*\pm}$ - ℓ / Q_{hem} , ℓ - Q_{hem} , and ℓ - ℓ .
- ³⁰ ABREU 96Q analysis performed using lepton, kaon, and jet-charge tags.
- ³¹ ALEXANDER 96v combines results from $D^{*\pm}$ - ℓ and $D^{*\pm}$ ℓ - Q_{hem} .
- ³² AKERS 95J combines results from charge measurement, $D^{*\pm}$ ℓ - Q_{hem} and ℓ - ℓ .

$$\chi_d = \Delta m_{B^0}/\Gamma_{B^0}$$

The second "OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

The first "OUR EVALUATION", also provided by the HFAG, includes χ_d measured at $\Upsilon(4S)$.

| VALUE | DOCUMENT ID |
|-----------------------------------|-------------|
| 0.776±0.008 OUR EVALUATION | First |
| 0.776±0.008 OUR EVALUATION | Second |

$$\text{Re}(\lambda_{CP} / |\lambda_{CP}|) \text{ Re}(z)$$

The λ_{CP} characterizes B^0 and \bar{B}^0 decays to states of charmonium plus K^0_L . Parameter z is used to describe CPT violation in mixing, see the review on "CP Violation" in the reviews section.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|--------------------------------|------|------------------------------------|
| 0.014±0.035±0.034 | ¹ AUBERT,B 04C BABR | | $e^+ e^- \rightarrow \Upsilon(4S)$ |

¹ Corresponds to 90% confidence range [-0.072, 0.101].

$\Delta\Gamma \text{Re}(z)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---------------------------------------|
| $-0.0071 \pm 0.0039 \pm 0.0020$ | AUBERT | 06T | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

 $\text{Re}(z)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|---------------------------------------|
| $0.00 \pm 0.12 \pm 0.01$ | 1 HASTINGS | 03 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Measured using inclusive dilepton events from B^0 decay.

 $\text{Im}(z)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|---------------------------------------|
| -0.015 ± 0.008 OUR AVERAGE | | | |
| $-0.0139 \pm 0.0073 \pm 0.0032$ | ¹ AUBERT | 06T | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| $-0.03 \pm 0.01 \pm 0.03$ | ² HASTINGS | 03 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $0.038 \pm 0.029 \pm 0.025$ | ³ AUBERT,B | 04C | BABR Repl. by AUBERT 06T |
| ¹ Assuming $\Delta\Gamma = 0$, the result becomes $\text{Im}(z) = -0.0037 \pm 0.0046$. | | | |
| ² Measured using inclusive dilepton events from B^0 decay. | | | |
| ³ Corresponds to 90% confidence range $[-0.028, 0.104]$. | | | |

CP VIOLATION PARAMETERS **$\text{Re}(\epsilon_{B^0})/(1+|\epsilon_{B^0}|^2)$**

CP impurity in B_d^0 system. It is obtained from either $a_{\ell\ell}$, the charge asymmetry in like-sign dilepton events or a_{CP} , the time-dependent asymmetry of inclusive B^0 and \bar{B}^0 decays.

The second “OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements. It assumes there is no CP violation in B_s mixing.

The first “OUR EVALUATION”, also provided by the HFAG, uses the measurements from B -factories only.

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------|------|---------------------------------------|
| -0.1 ± 1.4 OUR EVALUATION | First | | |
| -1.2 ± 1.0 OUR EVALUATION | Second | | |
| -0.9 ± 0.9 OUR AVERAGE | | | |
| $-2.3 \pm 1.1 \pm 0.8$ | ¹ ABAZOV | 06S | D0 $p\bar{p}$ at 1.96 TeV |
| $0.4 \pm 1.3 \pm 0.9$ | ² AUBERT | 06T | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| $-0.3 \pm 2.0 \pm 2.1$ | ³ NAKANO | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.2 \pm 2.9 \pm 3.6$ | ⁴ AUBERT | 02K | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| -3.2 ± 6.5 | ⁵ BARATE | 01D | ALEP $e^+ e^- \rightarrow Z$ |
| $3.5 \pm 10.3 \pm 1.5$ | ⁶ JAFFE | 01 | CLE2 $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.2 \pm 13.8 \pm 3.2$ | ⁷ ABBIENDI | 99J | OPAL $e^+ e^- \rightarrow Z$ |
| $2 \pm 7 \pm 3$ | ⁸ ACKERSTAFF | 97U | OPAL $e^+ e^- \rightarrow Z$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------------------|-----------------------|----------|----------------------------------|
| $-14.7 \pm 6.7 \pm 5.7$ | ⁹ AUBERT,B | 04C BABR | Repl. by AUBERT 06T |
| $4 \pm 18 \pm 3$ | ¹⁰ BEHRENS | 00B CLE2 | Repl. by JAFFE 01 |
| < 45 | ¹¹ BARTEL | 93 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses the dimuon charge asymmetry.

² AUBERT 06T reports $|q/p| - 1 = (-0.8 \pm 2.7 \pm 1.9) \times 10^{-3}$. We convert to $(1 - |q/p|^2)/4$.

³ Uses the charge asymmetry in like-sign dilepton events and reports $|q/p| = 1.0005 \pm 0.0040 \pm 0.0043$.

⁴ AUBERT 02K uses the charge asymmetry in like-sign dilepton events.

⁵ BARATE 01D measured by investigating time-dependent asymmetries in semileptonic and fully inclusive B_d^0 decays.

⁶ JAFFE 01 finds $a_{\ell\ell} = 0.013 \pm 0.050 \pm 0.005$ and combines with the previous BEHRENS 00B independent measurement.

⁷ Data analyzed using the time-dependent asymmetry of inclusive B^0 decay. The production flavor of B^0 mesons is determined using both the jet charge and the charge of secondary vertex in the opposite hemisphere.

⁸ ACKERSTAFF 97U assumes *CPT* and is based on measuring the charge asymmetry in a sample of B^0 decays defined by lepton and Q_{hem} tags. If *CPT* is not invoked, $\text{Re}(\epsilon_B) = -0.006 \pm 0.010 \pm 0.006$ is found. The indirect *CPT* violation parameter is determined to $\text{Im}(\delta B) = -0.020 \pm 0.016 \pm 0.006$.

⁹ AUBERT 04C reports $|q/p| = 1.029 \pm 0.013 \pm 0.011$ and we converted it to $(1 - |q/p|^2)/4$.

¹⁰ BEHRENS 00B uses high-momentum lepton tags and partially reconstructed $\bar{B}^0 \rightarrow D^{*+} \pi^-$, ρ^- decays to determine the flavor of the B meson.

¹¹ BARTEL 93 finds $a_{\ell\ell} = 0.031 \pm 0.096 \pm 0.032$ which corresponds to $|a_{\ell\ell}| < 0.18$, which yields the above $|\text{Re}(\epsilon_{B^0})/(1 + |\epsilon_{B^0}|^2)|$.

$A_{T/CP}$

$A_{T/CP}$ is defined as

$$\frac{P(\bar{B}^0 \rightarrow B^0) - P(B^0 \rightarrow \bar{B}^0)}{P(\bar{B}^0 \rightarrow B^0) + P(B^0 \rightarrow \bar{B}^0)},$$

the *CPT* invariant asymmetry between the oscillation probabilities $P(\bar{B}^0 \rightarrow B^0)$ and $P(B^0 \rightarrow \bar{B}^0)$.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|----------|----------------------------------|
| $0.005 \pm 0.012 \pm 0.014$ | ¹ AUBERT | 02K BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ AUBERT 02K uses the charge asymmetry in like-sign dilepton events.

$A_{CP}(B^0 \rightarrow D^*(2010)^+ D^-)$

A_{CP} is defined as

$$\frac{B(\bar{B}^0 \rightarrow \bar{f}) - B(B^0 \rightarrow f)}{B(\bar{B}^0 \rightarrow \bar{f}) + B(B^0 \rightarrow f)},$$

the *CP*-violation charge asymmetry of exclusive B^0 and \bar{B}^0 decay.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|------|---------|
| -0.06 ± 0.09 OUR AVERAGE | Error includes scale factor of 1.7. | | |

| | | | |
|---------------------------|---------------------|-----------|----------------------------------|
| $-0.12 \pm 0.06 \pm 0.02$ | AUBERT | 07AI BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.07 \pm 0.08 \pm 0.04$ | ¹ AUSHEV | 04 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------------------------|----------|----------|-----------------------|
| $-0.03 \pm 0.10 \pm 0.02$ | AUBERT,B | 06A BABR | Repl. by AUBERT 07AI |
| $-0.03 \pm 0.11 \pm 0.05$ | AUBERT | 03J BABR | Repl. by AUBERT,B 06B |

¹ Combines results from fully and partially reconstructed $B^0 \rightarrow D^{*\pm} D^{\mp}$ decays.

$A_{CP}(B^0 \rightarrow K^+ \pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|-----------|----------------------------------|
| -0.101±0.015 OUR AVERAGE | | | |
| -0.107±0.018 ^{+0.007} _{-0.004} | AUBERT | 07AF BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.013±0.078±0.012 | ABULENCIA,A | 06D CDF | $p\bar{p}$ at 1.96 TeV |
| -0.101±0.025±0.005 | ¹ CHAO | 04B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.04 ± 0.16 | ² CHEN | 00 CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.088±0.035±0.013 | ³ CHAO | 05A BELL | Repl. by CHAO 04B |
| -0.133±0.030±0.009 | ⁴ AUBERT,B | 04K BABR | Repl. by AUBERT 07AF |
| -0.07 ± 0.08 ± 0.02 | ⁵ AUBERT | 02D BABR | Repl. by AUBERT 02Q |
| -0.102±0.050±0.016 | ⁶ AUBERT | 02Q BABR | Repl. by AUBERT,B 04K |
| -0.06 ± 0.09 ^{+0.01} _{-0.02} | ⁷ CASEY | 02 BELL | Repl. by CHAO 04B |
| 0.044 ^{+0.186} _{-0.167} ^{+0.018} _{-0.021} | ⁸ ABE | 01K BELL | Repl. by CASEY 02 |
| -0.19 ± 0.10 ± 0.03 | ⁹ AUBERT | 01E BABR | Repl. by AUBERT 02Q |

¹ CHAO 04B reports significance of 3.9 standard deviation for deviation of A_{CP} from zero.² Corresponds to 90% confidence range $-0.30 < A_{CP} < 0.22$.³ Corresponds to a 90% CL interval of $-0.15 < A_{CP} < -0.03$.⁴ Based on a total signal yield of $N(K^-\pi^+) + N(K^+\pi^-) = 1606 \pm 51$ events.⁵ Corresponds to 90% confidence range $-0.21 < A_{CP} < 0.07$.⁶ Corresponds to 90% confidence range $-0.188 < A_{CP} < -0.016$.⁷ Corresponds to 90% confidence range $-0.21 < A_{CP} < +0.09$.⁸ Corresponds to 90% confidence range $-0.25 < A_{CP} < 0.37$.⁹ Corresponds to 90% confidence range $-0.35 < A_{CP} < -0.03$. **$A_{CP}(B^0 \rightarrow \eta' K^*(892)^0)$**

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|----------|----------------------------------|
| -0.08±0.25±0.02 | AUBERT | 07E BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

 $A_{CP}(B^0 \rightarrow \eta K^*(892)^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|----------|----------------------------------|
| 0.19±0.05 OUR AVERAGE | | | |
| 0.17±0.08±0.01 | WANG | 07B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.21±0.06±0.02 | AUBERT,B | 06H BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.02±0.11±0.02 | AUBERT,B | 04D BABR | Repl. by AUBERT,B 06H |

 $A_{CP}(B^0 \rightarrow K^0 \bar{K}^0)$

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|---------|----------------------------------|
| -0.58^{+0.73}_{-0.66}±0.04 | LIN | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

 $A_{CP}(B^0 \rightarrow \eta K_0^*(1430)^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|----------|----------------------------------|
| 0.06±0.13±0.02 | AUBERT,B | 06H BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

$A_{CP}(B^0 \rightarrow \eta K_2^*(1430)^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|------|---------------------------------------|
| -0.07±0.19±0.02 | AUBERT,B | 06H | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

 $A_{CP}(B^0 \rightarrow \rho^+ K^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-------------|------|-------------------------------------|
| -0.08±0.24 OUR AVERAGE | | | Error includes scale factor of 1.7. |

$0.22^{+0.22+0.06}_{-0.23-0.02}$
 $-0.28\pm0.17\pm0.08$

| | | | |
|----------|-----|------|----------------------------------|
| 1 CHANG | 04 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 2 AUBERT | 03T | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Corresponds to 90% confidence range $-0.18 < A_{CP} < 0.64$.

² The result reported corresponds to $-A_{CP}$.

 $A_{CP}(B^0 \rightarrow K^+ \pi^- \pi^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|------|---------------------------------------|
| 0.07±0.11±0.01 | 1 CHANG | 04 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Corresponds to 90% confidence range $-0.12 < A_{CP} < 0.26$.

 $A_{CP}(B^0 \rightarrow K^*(892)^+ \pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-------------|------|---------|
| -0.05±0.14 OUR AVERAGE | | | |

$-0.11\pm0.14\pm0.05$
 $0.26^{+0.33+0.10}_{-0.34-0.08}$

| | | | |
|--------------|-----|------|----------------------------------|
| AUBERT | 06I | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 1 EISENSTEIN | 03 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.23\pm0.18^{+0.09}_{-0.06}$

AUBERT,B 040 BABR Repl. by AUBERT 06I

¹ Corresponds to 90% confidence range $-0.31 < A_{CP} < 0.78$.

 $A_{CP}(B^0 \rightarrow K^*(892)^0 \rho^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|------|---------------------------------------|
| 0.09±0.19±0.02 | AUBERT,B | 06G | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

 $A_{CP}(B^0 \rightarrow a_1^- K^+)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|------|---------------------------------------|
| -0.16±0.12±0.01 | AUBERT | 08F | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

 $A_{CP}(B^0 \rightarrow K^*(892)^0 \pi^+ \pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|------|---------------------------------------|
| +0.07±0.04±0.03 | AUBERT | 07AS | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

 $A_{CP}(B^0 \rightarrow K^*(892)^0 K^+ K^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|------|---------------------------------------|
| +0.01±0.05±0.02 | AUBERT | 07AS | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

$A_{CP}(B^0 \rightarrow K^*(892)^0 \phi)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|----------------------------------|
| -0.01±0.06 OUR AVERAGE | | | |
| -0.03±0.07±0.03 | AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.02±0.09±0.02 | ¹ CHEN | 05A BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.01±0.09±0.02 | AUBERT,B | 04W BABR | Repl. by AUBERT 07D |
| 0.04±0.12±0.02 | AUBERT | 03V BABR | Repl. by AUBERT 04W |
| 0.07±0.15 ^{+0.05} _{-0.03} | ² CHEN | 03B BELL | Repl. by CHEN 05A |
| 0.00±0.27±0.03 | ³ AUBERT | 02E BABR | Repl. by AUBERT 03V |

¹ Corresponds to 90% confidence range $-0.14 < A_{CP} < 0.17$.² Corresponds to 90% confidence range $-0.18 < A_{CP} < 0.33$.³ Corresponds to 90% confidence range $-0.44 < A_{CP} < 0.44$. $A_{CP}(B^0 \rightarrow K^*(892)^0 K^- \pi^+)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|--------------------|----------------------------------|----------------|
| +0.22±0.33±0.20 | | | |
| AUBERT | 07AS BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $A_{CP}(B^0 \rightarrow \phi(K\pi)_0^{*0})$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|----------------------------------|----------------|
| 0.17±0.15±0.03 | | | |
| AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $A_{CP}(B^0 \rightarrow \phi K_2^*(1430)^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|--------------------|----------------------------------|----------------|
| -0.12±0.14±0.04 | | | |
| AUBERT | 07D BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $A_{CP}(B^0 \rightarrow \rho^+ \pi^-)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|----------------------------------|
| 0.08±0.12 OUR AVERAGE | | | |
| -0.03±0.07±0.04 | ¹ AUBERT | 07AA BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.21±0.08±0.04 | KUSAKA | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.02±0.16 ^{+0.05} _{-0.02} | WANG | 05 BELL | Repl. by KUSAKA 07 |
| 0.18±0.08±0.03 | ¹ AUBERT | 03T BABR | Repl. by AUBERT 07AA |

¹ The result reported corresponds to $-A_{CP}$. $A_{CP}(B^0 \rightarrow \rho^- \pi^+)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------------------------|
| -0.16±0.23 OUR AVERAGE | | | |
| Error includes scale factor of 1.7. | | | |
| -0.37±0.16 ^{+0.09} _{-0.10} | AUBERT | 07AA BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.08±0.16±0.11 | KUSAKA | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.53±0.29 ^{+0.09} _{-0.04} | WANG | 05 BELL | Repl. by KUSAKA 07 |

$A_{CP}(B^0 \rightarrow \rho^0 \pi^0)$ VALUE**-0.49±0.36±0.28**

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $-0.53^{+0.67+0.10}_{-0.84-0.15}$ DOCUMENT IDKUSAKA 07 BELL $e^+ e^- \rightarrow \gamma(4S)$

DRAGIC 06 BELL Repl. by KUSAKA 07

 $A_{CP}(B^0 \rightarrow a_1(1260)^{\pm} \pi^{\mp})$ VALUE**-0.07±0.07±0.02** $A_{CP}(B^0 \rightarrow b_1 \pi^+)$ VALUE**-0.05±0.10±0.02**DOCUMENT IDAUBERT 070 BABR $e^+ e^- \rightarrow \gamma(4S)$ DOCUMENT IDAUBERT 07BI BABR $e^+ e^- \rightarrow \gamma(4S)$ $A_{CP}(B^0 \rightarrow K^*(1430)\gamma)$ VALUE**-0.08±0.15±0.01**DOCUMENT IDAUBERT,B 04U BABR $e^+ e^- \rightarrow \gamma(4S)$ $A_{CP}(B^0 \rightarrow p\bar{p}K^*(892)^0)$ VALUE**+0.11±0.13±0.06**DOCUMENT IDAUBERT 07AV BABR $e^+ e^- \rightarrow \gamma(4S)$ $A_{CP}(B^0 \rightarrow p\bar{\Lambda}\pi^-)$ VALUE**-0.02±0.10±0.03**DOCUMENT IDWANG 07C BELL $e^+ e^- \rightarrow \gamma(4S)$ $A_{CP}(B^0 \rightarrow b_1 K^+)$ VALUE**-0.07±0.12±0.02**DOCUMENT IDAUBERT 07BI BABR $e^+ e^- \rightarrow \gamma(4S)$ $C_{D^*(2010)^- D^+} (B^0 \rightarrow D^*(2010)^- D^+)$ VALUE**0.23±0.13 OUR AVERAGE** $0.23 \pm 0.15 \pm 0.04$ $0.23 \pm 0.25 \pm 0.06$ DOCUMENT IDAUBERT 07AI BABR $e^+ e^- \rightarrow \gamma(4S)$ 1 AUSHEV 04 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.17 \pm 0.24 \pm 0.04$ $-0.22 \pm 0.37 \pm 0.10$

AUBERT,B 05Z BABR Repl. by AUBERT 07AI

AUBERT 03J BABR Repl. by

AUBERT,B 05Z

1 Combines results from fully and partially reconstructed $B^0 \rightarrow D^* \pm D^\mp$ decays. $S_{D^*(2010)^- D^+} (B^0 \rightarrow D^*(2010)^- D^+)$ VALUE**-0.55±0.21 OUR AVERAGE** $-0.44 \pm 0.22 \pm 0.06$ DOCUMENT IDAUBERT 07AI BABR $e^+ e^- \rightarrow \gamma(4S)$ $-0.96 \pm 0.43 \pm 0.12$ 1 AUSHEV 04 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $-0.29 \pm 0.33 \pm 0.07$

AUBERT,B 05Z BABR Repl. by AUBERT 07AI

 $-0.24 \pm 0.69 \pm 0.12$

AUBERT 03J BABR Repl. by AUBERT,B 05Z

1 Combines results from fully and partially reconstructed $B^0 \rightarrow D^* \pm D^\mp$ decays.

$C_{D^*(2010)^+ D^-} (B^0 \rightarrow D^*(2010)^+ D^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|-------------------------------------|
| 0.01±0.26 OUR AVERAGE | | | Error includes scale factor of 2.0. |
| 0.18±0.15±0.04 | AUBERT | 07AI BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.37±0.22±0.06 | 1 AUSHEV | 04 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.09±0.25±0.06 | AUBERT,B | 05Z BABR | Repl. by AUBERT 07AI |
| -0.47±0.40±0.12 | AUBERT | 03J BABR | Repl. by AUBERT,B 05Z |
| 1 Combines results from fully and partially reconstructed $B^0 \rightarrow D^* \pm D^\mp$ decays. | | | |

 $S_{D^*(2010)^+ D^-} (B^0 \rightarrow D^*(2010)^+ D^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|----------------------------------|
| -0.74±0.19 OUR AVERAGE | | | |
| -0.79±0.21±0.06 | AUBERT | 07AI BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.55±0.39±0.12 | 1 AUSHEV | 04 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.54±0.35±0.07 | AUBERT,B | 05Z BABR | Repl. by AUBERT 07AI |
| -0.82±0.75±0.14 | AUBERT | 03J BABR | Repl. by AUBERT,B 05Z |
| 1 Combines results from fully and partially reconstructed $B^0 \rightarrow D^* \pm D^\mp$ decays. | | | |

 $C_{D^{*+} D^{*-}} (B^0 \rightarrow D^{*+} D^{*-})$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|----------------------------------|
| 0.02±0.10 OUR AVERAGE | | | |
| -0.02±0.11±0.02 | 1 AUBERT | 07BO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.26±0.26±0.06 | 2 MIYAKE | 05 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.28±0.23±0.02 | 3 AUBERT | 03Q BABR | Repl. by AUBERT 07BO |
| 1 Assumes both CP -even and CP -odd states having the CP asymmetry. | | | |
| 2 Belle Collab. quotes $A_{D^{*+} D^{*-}}$ which is equal to $-C_{D^{*+} D^{*-}}$. | | | |
| 3 AUBERT 03Q reports $ \lambda =0.75 \pm 0.19 \pm 0.02$ and $\text{Im}(\lambda)=0.05 \pm 0.29 \pm 0.10$. We convert them to S and C parameters taking into account correlations. | | | |

 $S_{D^{*+} D^{*-}} (B^0 \rightarrow D^{*+} D^{*-})$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|----------------------------------|
| -0.67±0.18 OUR AVERAGE | | | |
| -0.66±0.19±0.04 | 1 AUBERT | 07BO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.75±0.56±0.12 | MIYAKE | 05 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.06±0.37±0.13 | 2 AUBERT | 03Q BABR | Repl. by AUBERT 07BO |
| 1 Assumes both CP -even and CP -odd states having the CP asymmetry. | | | |
| 2 AUBERT 03Q reports $ \lambda =0.75 \pm 0.19 \pm 0.02$ and $\text{Im}(\lambda)=0.05 \pm 0.29 \pm 0.10$. We convert them to S and C parameters taking into account correlations. | | | |

 $C_+ (B^0 \rightarrow D^{*+} D^{*-})$ See the note in the $C_{\pi\pi}$ datablock, but for CP even final state.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|-----------|----------------------------------|
| -0.05±0.14±0.02 | AUBERT | 07BO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.06±0.17±0.03 | 1 AUBERT,BE | 05A BABR | Repl. by AUBERT 07BO |
| 1 AUBERT,BE 05A reports a CP -odd fraction $R_\perp = 0.125 \pm 0.044 \pm 0.007$. | | | |

S_+ ($B^0 \rightarrow D^{*+} D^{*-}$)See the note in the $S_{\pi\pi}$ datablock, but for CP even final state.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|-----------|----------------------------------|
| -0.72±0.19±0.05 | AUBERT | 07BO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.75±0.25±0.03 | ¹ AUBERT,BE | 05A BABR | Repl. by AUBERT 07BO |
| ¹ AUBERT,BE 05A reports a CP -odd fraction $R_\perp = 0.125 \pm 0.044 \pm 0.007$. | | | |

 C_- ($B^0 \rightarrow D^{*+} D^{*-}$)See the note in the $C_{\pi\pi}$ datablock, but for CP odd final state.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|-----------|----------------------------------|
| +0.23±0.67±0.10 | AUBERT | 07BO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.20±0.96±0.11 | ¹ AUBERT,BE | 05A BABR | Repl. by AUBERT 07BO |
| ¹ AUBERT,BE 05A reports a CP -odd fraction $R_\perp = 0.125 \pm 0.044 \pm 0.007$. | | | |

 S_- ($B^0 \rightarrow D^{*+} D^{*-}$)See the note in the $S_{\pi\pi}$ datablock, but for CP odd final state.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|-----------|----------------------------------|
| -1.83±1.04±0.23 | AUBERT | 07BO BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -1.75±1.78±0.22 | ¹ AUBERT,BE | 05A BABR | Repl. by AUBERT 07BO |
| ¹ AUBERT,BE 05A reports a CP -odd fraction $R_\perp = 0.125 \pm 0.044 \pm 0.007$. | | | |

 C ($B^0 \rightarrow D^*(2010)^+ D^*(2010)^- K_S^0$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|----------------------|------|---------------------------------------|
| 0.01±0.28±0.09 | ¹ DALSENO | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Reports value of A which is equal to $-C$. **S ($B^0 \rightarrow D^*(2010)^+ D^*(2010)^- K_S^0$)**

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|----------------------|------|---------------------------------------|
| 0.06^{+0.45}_{-0.44}±0.06 | ¹ DALSENO | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

¹ This value includes an unknown CP dilution factor D due to possible contributions from intermediate resonances and different partial waves. **$C_{D^+ D^-}$ ($B^0 \rightarrow D^+ D^-$)**

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|----------|---------------------------------------|
| -0.4 ±0.5 OUR AVERAGE | | | Error includes scale factor of 3.1. |
| 0.11±0.22±0.07 | | | |
| -0.91±0.23±0.06 | ¹ FRATINA | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.11±0.35±0.06 | AUBERT,B | 05Z BABR | Repl. by AUBERT 07AI |

¹ The paper reports A , which is equal to $-C$.

$S_{D^+ D^-} (B^0 \rightarrow D^+ D^-)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------------------|-------------|---------------------------------------|
| -0.81±0.29 OUR AVERAGE | Error includes scale factor of 1.1. | | |
| -0.54±0.34±0.06 | AUBERT | 07AI | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| -1.13±0.37±0.09 | FRATINA | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.29±0.63±0.06 | AUBERT,B | 05Z | BABR Repl. by AUBERT 07AI |

 $C_{J/\psi(1S)\pi^0} (B^0 \rightarrow J/\psi(1S)\pi^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|----------------------|-------------|---------------------------------------|
| -0.11±0.20 OUR AVERAGE | | | |
| -0.21±0.26±0.06 | AUBERT,B | 06B | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.01±0.29±0.03 | ¹ KATAOKA | 04 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.38±0.41±0.09 | AUBERT | 03N | BABR Repl. by AUBERT,B 06B |
| ¹ BELLE Collab. quotes $A_{J/\psi\pi^0}$ which is equal to $-C_{J/\psi\pi^0}$. | | | |

 $S_{J/\psi(1S)\pi^0} (B^0 \rightarrow J/\psi(1S)\pi^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|---------------------------------------|
| -0.69±0.25 OUR AVERAGE | | | |
| -0.68±0.30±0.04 | AUBERT,B | 06B | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.72±0.42±0.09 | KATAOKA | 04 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.05±0.49±0.16 | AUBERT | 03N | BABR Repl. by AUBERT,B 06B |

 $C_{D_{CP}^{(*)} h^0} (B^0 \rightarrow D_{CP}^{(*)} h^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|--------------------|-------------|----------------|
| -0.23±0.16±0.04 | | | |

 $S_{D_{CP}^{(*)} h^0} (B^0 \rightarrow D_{CP}^{(*)} h^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|--------------------|-------------|----------------|
| -0.56±0.23±0.05 | | | |

 $S_{K_S^0 K_S^0} (B^0 \rightarrow K_S^0 K_S^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|----------------|
| -1.28^{+0.80+0.11}_{-0.73-0.16} | | | |

 $C_{K_S^0 K_S^0} (B^0 \rightarrow K_S^0 K_S^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|--------------------|-------------|----------------|
| -0.40±0.41±0.06 | | | |

$C_{\eta'(958)K} (B^0 \rightarrow \eta'(958)K_S^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|----------|----------------------------------|
| -0.04±0.20 OUR AVERAGE | Error includes scale factor of 2.5. | | |
| -0.21±0.10±0.02 | AUBERT | 05M BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.19±0.11±0.05 | ¹ CHEN | 05B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.26±0.22±0.03 | ¹ ABE | 03C BELL | Repl. by ABE 03H |
| 0.01±0.16±0.04 | ¹ ABE | 03H BELL | Repl. by CHEN 05B |
| 0.10±0.22±0.04 | AUBERT | 03W BABR | Repl. by AUBERT 05M |
| -0.13±0.32 ^{+0.06} _{-0.09} | ¹ CHEN | 02B BELL | Repl. by ABE 03C |

¹ BELLE Collab. quotes $A_{\eta'(958)K_S^0}$ which is equal to $-C_{\eta'(958)K_S^0}$.

 $S_{\eta'(958)K} (B^0 \rightarrow \eta'(958)K_S^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|----------|----------------------------------|
| 0.43±0.17 OUR AVERAGE | Error includes scale factor of 1.5. | | |
| 0.30±0.14±0.02 | AUBERT | 05M BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| +0.65±0.18±0.04 | CHEN | 05B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.71±0.37 ^{+0.05} _{-0.06} | ABE | 03C BELL | Repl. by ABE 03H |
| 0.43±0.27±0.05 | ABE | 03H BELL | Repl. by CHEN 05B |
| 0.02±0.34±0.03 | AUBERT | 03W BABR | Repl. by AUBERT 05M |
| 0.28±0.55 ^{+0.07} _{-0.08} | CHEN | 02B BELL | Repl. by ABE 03C |

 $C_{\eta' K^0} (B^0 \rightarrow \eta' K^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-------------------------------------|----------|----------------------------------|
| -0.09±0.08 OUR AVERAGE | Error includes scale factor of 1.5. | | |
| -0.16±0.07±0.03 | ¹ AUBERT | 07A BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.01±0.07±0.05 | ^{1,2} CHEN | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ The mixing-induced CP violation is reported with a significance of more than 5 standard deviations in this $b \rightarrow s$ penguin dominated mode.

² The paper reports A , which is equal to $-C$.

 $S_{\eta' K^0} (B^0 \rightarrow \eta' K^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|---------------------|----------|----------------------------------|
| 0.61±0.07 OUR AVERAGE | | | |
| 0.58±0.10±0.03 | ¹ AUBERT | 07A BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.64±0.10±0.04 | ¹ CHEN | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ The mixing-induced CP violation is reported with a significance of more than 5 standard deviations in this $b \rightarrow s$ penguin dominated mode.

$C_{\omega K_S^0} (B^0 \rightarrow \omega K_S^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|------|---------------------------------------|
| -0.25±0.31 OUR AVERAGE | Error includes scale factor of 1.6. | | |
| +0.09±0.29±0.06 | ¹ CHAO | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.55 ^{+0.28} _{-0.26} ± 0.03 | AUBERT,B | 06E | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.27±0.48±0.15 | ¹ CHEN | 05B | BELL Repl by CHAO 07 |
| ¹ Belle Collab. quotes $A_{\omega K_S^0}$ which is equal to $-C_{\omega K_S^0}$. | | | |

 $S_{\omega K_S^0} (B^0 \rightarrow \omega K_S^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---------------------------------------|
| 0.35±0.29 OUR AVERAGE | | | |
| +0.11±0.46±0.07 | CHAO | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| +0.51 ^{+0.35} _{-0.39} ± 0.02 | AUBERT,B | 06E | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| +0.76±0.65 ^{+0.13} _{-0.16} | CHEN | 05B | BELL Repl. by CHAO 07 |

 $C_{f_0(980) K_S^0} (B^0 \rightarrow f_0(980) K_S^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|------|---------------------------------------|
| -0.03±0.26 OUR AVERAGE | Error includes scale factor of 1.9. | | |
| -0.41±0.23±0.07 | ¹ AUBERT | 07AX | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| +0.15±0.15±0.07 | ¹ CHAO | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| +0.39±0.27±0.09 | ¹ CHEN | 05B | BELL Repl by CHAO 07 |
| ¹ Quotes $A_{f_0(980) K_S^0}$ which is equal to $-C_{f_0(980) K_S^0}$. | | | |

 $S_{f_0(980) K_S^0} (B^0 \rightarrow f_0(980) K_S^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|------|---------------------------------------|
| -0.02±0.21 OUR AVERAGE | Error includes scale factor of 1.1. | | |
| -0.25±0.26±0.10 | ¹ AUBERT | 07AX | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| +0.18±0.23±0.11 | CHAO | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| +0.47±0.41±0.08 | CHEN | 05B | BELL Repl. by CHAO 07 |
| ¹ Reports β_{eff} . We quote S obtained from epaps: E-PRLTAO-99-076741. | | | |

 $C_{K_S K_S K_S} (B^0 \rightarrow K_S K_S K_S)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|------|---------------------------------------|
| -0.15±0.16 OUR AVERAGE | Error includes scale factor of 1.1. | | |
| +0.02±0.21±0.05 | AUBERT | 07AT | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.31±0.20±0.07 | ¹ CHEN | 07 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.34 ^{+0.28} _{-0.25} ± 0.05 | AUBERT,B | 05 | BABR Repl. by AUBERT 07AT |
| -0.54±0.34±0.09 | ¹ SUMISAWA | 05 | BELL Repl. by CHEN 07 |
| ¹ Belle Collab. quotes $A_{K_S K_S K_S}$ which is equal to $-C_{K_S K_S K_S}$. | | | |

$S_{K_S K_S K_S} (B^0 \rightarrow K_S K_S K_S)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------------------|-------------|----------------------------------|
| -0.4 ± 0.5 OUR AVERAGE | Error includes scale factor of 2.5. | | |
| -0.71 ± 0.24 ± 0.04 | AUBERT | 07AT BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.30 ± 0.32 ± 0.08 | CHEN | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.71 $^{+0.38}_{-0.32}$ ± 0.04 | AUBERT,B | 05 BABR | Repl. by AUBERT 07AT |
| 1.26 ± 0.68 ± 0.20 | SUMISAWA | 05 BELL | Repl. by CHEN 07. |

 $C_{K^+ K^- K_S^0} (B^0 \rightarrow K^+ K^- K_S^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------------------------|
| 0.07 ± 0.08 OUR AVERAGE | | | |
| 0.054 ± 0.102 ± 0.060 | 1,2 AUBERT | 07AX BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.09 ± 0.10 ± 0.05 | 1,2 CHAO | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.10 ± 0.14 ± 0.04 | 2 AUBERT | 05T BABR | Repl. by AUBERT 07AX |
| 0.09 ± 0.12 ± 0.07 | 1 CHEN | 05B BELL | Repl. by CHAO 07 |
| -0.10 ± 0.19 ± 0.10 | 2 AUBERT,B | 04V BABR | Repl. by AUBERT 05T |
| 0.40 ± 0.33 $^{+0.28}_{-0.10}$ | 1 ABE | 03C BELL | Repl. by ABE 03H |
| 0.17 ± 0.16 ± 0.04 | 1,2 ABE | 03H BELL | Repl. by CHEN 05B |

¹ Quotes $A_{K^+ K^- K_S^0}$ which is equal to $-C_{K^+ K^- K_S^0}$.

² Excludes the events from $B^0 \rightarrow \phi K_S^0$ decay. The results are derived from a combined sample of $K^+ K^- K_S^0$ and $K^+ K^- K_L^0$ decays.

 $S_{K^+ K^- K_S^0} (B^0 \rightarrow K^+ K^- K_S^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------------------------|
| -0.74 $^{+0.12}_{-0.10}$ OUR AVERAGE | | | |
| -0.764 ± 0.111 $^{+0.071}_{-0.040}$ | 1,2 AUBERT | 07AX BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.68 ± 0.15 $^{+0.21}_{-0.13}$ | 1 CHAO | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.42 ± 0.17 ± 0.03 | 1,3 AUBERT | 05T BABR | Repl. by AUBERT 07AX |
| -0.49 ± 0.18 ± 0.04 | CHEN | 05B BELL | Repl. by CHAO 07 |
| -0.56 ± 0.25 ± 0.04 | 1,4 AUBERT,B | 04V BABR | Repl. by AUBERT 05T |
| -0.49 ± 0.43 ± 0.11 | ABE | 03C BELL | Repl. by ABE 03H |
| -0.51 ± 0.26 ± 0.05 | 1,5 ABE | 03H BELL | Repl. by CHEN 05B |

¹ Excludes events from $B^0 \rightarrow \phi K_S^0$ decay. The results are derived from a combined sample of $K^+ K^- K_S^0$ and $K^+ K^- K_L^0$ decays.

² Reports β_{eff} . We quote S obtained from epaps: E-PRLTAO-99-076741.

³ The measured CP -even final states fraction is $0.89 \pm 0.08 \pm 0.06$.

⁴ The measured CP -even final states fraction is $0.98 \pm 0.15 \pm 0.04$.

⁵ The measured CP -even final states fraction is $1.03 \pm 0.15 \pm 0.05$.

$C_{K^+ K^- K_S^0}$ ($B^0 \rightarrow K^+ K^- K_S^0$ inclusive)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------|-----------|----------------------------------|
| 0.015 ± 0.077 ± 0.053 | 1,2 AUBERT | 07AX BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

1 Measured using full Dalitz plot fit including ϕ component.2 The results are derived from a combined sample of $K^+ K^- K_S^0$ and $K^+ K^- K_L^0$ decays. $S_{K^+ K^- K_S^0}$ ($B^0 \rightarrow K^+ K^- K_S^0$ inclusive)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-------------|-----------|----------------------------------|
| -0.647 ± 0.116 ± 0.040 | 1 AUBERT | 07AX BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

1 Measured using full Dalitz plot fit including ϕ component. $C_{\phi K_S^0}$ ($B^0 \rightarrow \phi K_S^0$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|------|---------|
| -0.01 ± 0.12 OUR AVERAGE | | | |

| | | | |
|---|------------|-----------|----------------------------------|
| 0.08 ± 0.18 ± 0.04 | 1,2 AUBERT | 07AX BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.07 ± 0.15 ± 0.05 | 1,2 CHEN | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.00 ± 0.23 ± 0.05 | 2 AUBERT | 05T BABR | Repl. by AUBERT 07AX |
| -0.08 ± 0.22 ± 0.09 | 1,2 CHEN | 05B BELL | Repl. by CHEN 07 |
| 0.01 ± 0.33 ± 0.10 | 2 AUBERT,B | 04G BABR | Repl. by AUBERT 05T |
| 0.56 ± 0.41 ± 0.16 | 1 ABE | 03C BELL | Repl. by ABE 03H |
| 0.15 ± 0.29 ± 0.07 | 1 ABE | 03H BELL | Repl. by CHEN 05B |

1 Quotes $A_{\phi K_S^0}$ which is equal to $-C_{\phi K_S^0}$.2 Result combines B -meson final states ϕK_S^0 and ϕK_L^0 by assuming $S_{\phi K_S^0} = -S_{\phi K_L^0}$ $S_{\phi K_S^0}$ ($B^0 \rightarrow \phi K_S^0$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------|------|---------|
| 0.39 ± 0.17 OUR AVERAGE | | | |

| | | | |
|---|------------|-----------|----------------------------------|
| 0.21 ± 0.26 ± 0.11 | 1,2 AUBERT | 07AX BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.50 ± 0.21 ± 0.06 | 1 CHEN | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.50 ± 0.25 ± 0.07 -0.04 | 1 AUBERT | 05T BABR | Repl. by AUBERT 07AX |
| 0.08 ± 0.33 ± 0.09 | 1 CHEN | 05B BELL | Repl. by CHEN 07 |
| 0.47 ± 0.34 ± 0.08 -0.06 | 1 AUBERT,B | 04G BABR | Repl. by AUBERT 05T |
| -0.73 ± 0.64 ± 0.22 | ABE | 03C BELL | Repl. by ABE 03H |
| -0.96 ± 0.50 ± 0.09 -0.11 | ABE | 03H BELL | Repl. by CHEN 05B |

1 Result combines B -meson final states ϕK_S^0 and ϕK_L^0 by assuming $S_{\phi K_S^0} = -S_{\phi K_L^0}$ 2 Reports β_{eff} . We quote S obtained from epaps: E-PRLTAO-99-076741.

$C_{K_S^0 \pi^0} (B^0 \rightarrow K_S^0 \pi^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|----------------------------------|
| 0.14±0.11 OUR AVERAGE | | | |
| +0.24±0.15±0.03 | AUBERT | 08E BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| +0.05±0.14±0.05 | ¹ CHAO | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| +0.06±0.18±0.03 | AUBERT | 05Y BABR | Repl. by AUBERT 08E |
| -0.16±0.29±0.05 | ^{1,2} CHAO | 05A BELL | Repl. by CHEN 05B |
| +0.11±0.20±0.09 | ¹ CHEN | 05B BELL | Repl by CHAO 07 |
| -0.03±0.36±0.11 | ¹ AUBERT | 04M BABR | Repl. by AUBERT,B 04M |
| +0.40 ^{+0.27} _{-0.28} ±0.09 | ³ AUBERT,B | 04M BABR | Repl. by AUBERT 05Y |

¹ Reports A which is equal to $-C$.² Corresponds to a 90% CL interval of $-0.33 < A_{CP} < 0.64$.³ Based on a total signal yield of 122 ± 16 events. $S_{K_S^0 \pi^0} (B^0 \rightarrow K_S^0 \pi^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|----------------------------------|
| 0.38±0.19 OUR AVERAGE | | | |
| +0.40±0.23±0.03 | AUBERT | 08E BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| +0.33±0.35±0.08 | CHAO | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| +0.35 ^{+0.30} _{-0.33} ±0.04 | AUBERT | 05Y BABR | Repl. by AUBERT 08E |
| +0.32±0.61±0.13 | CHEN | 05B BELL | Repl. by CHAO 07 |
| +0.48 ^{+0.38} _{-0.47} ±0.06 | ¹ AUBERT,B | 04M BABR | Repl. by AUBERT 05Y |

¹ Based on a total signal yield of 122 ± 16 events. $C (B^0 \rightarrow K_S^0 \pi^0 \pi^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|----------------------------------|----------------|
| 0.23±0.52±0.13 | | | |
| AUBERT | 07AQ BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $S (B^0 \rightarrow K_S^0 \pi^0 \pi^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|----------------------------------|----------------|
| 0.72±0.71±0.08 | | | |
| AUBERT | 07AQ BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $C_{K_S^0 \pi^0 \gamma} (B^0 \rightarrow K_S^0 \pi^0 \gamma)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------|-------------|----------------------------------|
| 0.0 ±0.4 OUR AVERAGE | | | |
| +0.20±0.20±0.06 | ^{1,2} USHIRODA | 06 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| -1.0 ±0.5 ±0.2 | ³ AUBERT,B | 05P BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |

² USHIRODA 05 BELL Repl. by USHIRODA 06¹ Requires $M_{K_S^0 \pi^0} < 1.8 \text{ GeV}/c^2$.² Reports $A_{K_S^0 \pi^0 \gamma}$, which is $-C_{K_S^0 \pi^0 \gamma}$.³ Requires $1.1 < M_{K_S^0 \pi^0} < 1.8 \text{ GeV}/c^2$.

$S_{K_S^0 \pi^0 \gamma} (B^0 \rightarrow K_S^0 \pi^0 \gamma)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|---------------------------|------|----------------------------------|
| -0.01±0.30 OUR AVERAGE | | | |
| -0.10±0.31±0.07 | ¹ USHIRODA 06 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.9 ±1.0 ±0.2 | ² AUBERT,B 05P | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.58 ^{+0.46} _{-0.38} ±0.11 | USHIRODA 05 | BELL | Repl. by USHIRODA 06 |
| ¹ Requires $M_{K_S^0 \pi^0} < 1.8 \text{ GeV}/c^2$. | | | |
| ² Requires $1.1 < M_{K_S^0 \pi^0} < 1.8 \text{ GeV}/c^2$. | | | |

 $C_{K^*(892)^0 \gamma} (B^0 \rightarrow K^*(892)^0 \gamma)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|------|----------------------------------|
| -0.12±0.30 OUR AVERAGE | | | |
| | Error includes scale factor of 1.8. | | |
| +0.20±0.24±0.05 | ^{1,2} USHIRODA 06 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.40±0.23±0.03 | AUBERT,B 05P | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.57±0.32±0.09 | ³ AUBERT,B 04Z | BABR | Repl. by AUBERT,B 05P |
| ¹ Reports value of A which is equal to $-C$. | | | |
| ² Requires $0.8 < M_{K_S^0 \pi^0} < 1.0 \text{ GeV}/c^2$. | | | |
| ³ Based on a total signal of 105 ± 14 events with $K^*(892)^0 \rightarrow K_S^0 \pi^0$ only. | | | |

 $S_{K^*(892)^0 \gamma} (B^0 \rightarrow K^*(892)^0 \gamma)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|---------------------------|------|----------------------------------|
| -0.27±0.26 OUR AVERAGE | | | |
| -0.32 ^{+0.36} _{-0.33} ±0.05 | ¹ USHIRODA 06 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.21±0.40±0.05 | AUBERT,B 05P | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.79 ^{+0.63} _{-0.50} ±0.10 | ² USHIRODA 05 | BELL | Repl. by USHIRODA 06 |
| 0.25±0.63±0.14 | ³ AUBERT,B 04Z | BABR | Repl. by AUBERT,B 05P |
| ¹ Requires $0.8 < M_{K_S^0 \pi^0} < 1.0 \text{ GeV}/c^2$. | | | |
| ² Assumes $C(B^0 \rightarrow K^*(892)^0 \gamma) = 0$. | | | |
| ³ Based on a total signal of 105 ± 14 events with $K^*(892)^0 \rightarrow K_S^0 \pi^0$ only. | | | |

 $C(B^0 \rightarrow \rho^0 \gamma)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|--------------------------|------|----------------------------------|
| +0.44±0.49±0.14 | | | |
| | ¹ USHIRODA 08 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| ¹ Reports value of A which is equal to $-C$. | | | |

 $S(B^0 \rightarrow \rho^0 \gamma)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|------|----------------------------------|
| -0.83±0.65±0.18 | | | |
| | USHIRODA 08 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

$C_{\pi\pi}$ ($B^0 \rightarrow \pi^+\pi^-$)

$C_{\pi\pi}$ is defined as $(1 - |\lambda|^2)/(1 + |\lambda|^2)$, where the quantity $\lambda = q/p \bar{A}_f/A_f$ is a phase convention independent observable quantity for the final state f . For details, see the review on "CP Violation" in the Reviews section.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|-----------|--------------------------------------|
| -0.38±0.17 OUR AVERAGE | Error includes scale factor of 2.6. | | |
| -0.21±0.09±0.02 | AUBERT | 07AF BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| -0.55±0.08±0.05 | 1 ISHINO | 07 | BELL $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.56±0.12±0.06 | 1 ABE | 05D | BELL Repl. by ISHINO 07 |
| -0.09±0.15±0.04 | AUBERT,BE | 05 | BABR Repl. by AUBERT 07AF |
| -0.58±0.15±0.07 | 1 ABE | 04E | BELL Repl. by ABE 05D |
| -0.77±0.27±0.08 | 1 ABE | 03G | BELL Repl. by ABE 04E. |
| -0.94 ^{+0.31} _{-0.25} ±0.09 | 1 ABE | 02M | BELL Repl. by ABE 03G |
| -0.25 ^{+0.45} _{-0.47} ±0.14 | 2 AUBERT | 02D | BABR Repl. by AUBERT 02Q |
| -0.30±0.25±0.04 | 3 AUBERT | 02Q | BABR Repl. by AUBERT,BE 05 |

¹ Paper reports $A_{\pi\pi}$ which equals to $-C_{\pi\pi}$.

² Corresponds to 90% confidence range $-1.0 < C_{\pi\pi} < 0.47$.

³ Corresponds to 90% confidence range $-0.72 < C_{\pi\pi} < 0.12$.

 $S_{\pi\pi}$ ($B^0 \rightarrow \pi^+\pi^-$)

$S_{\pi\pi} = 2\text{Im}\lambda/(1+|\lambda|^2)$, see the note in the $C_{\pi\pi}$ datablock above.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|--------------------------------------|
| -0.61±0.08 OUR AVERAGE | | | |
| -0.60±0.11±0.03 | AUBERT | 07AF BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| -0.61±0.10±0.04 | ISHINO | 07 | BELL $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.67±0.16±0.06 | 1 ABE | 05D | BELL Repl. by ISHINO 07 |
| -0.30±0.17±0.03 | AUBERT,BE | 05 | BABR Repl. by AUBERT 07AF |
| -1.00±0.21±0.07 | 2 ABE | 04E | BELL Repl. by ABE 05D |
| -1.23±0.41 ^{+0.08} _{-0.07} | ABE | 03G | BELL Repl. by ABE 04E. |
| -1.21 ^{+0.38} _{-0.27} ^{+0.16} _{-0.13} | ABE | 02M | BELL Repl. by ABE 03G |
| 0.03 ^{+0.52} _{-0.56} ±0.11 | 3 AUBERT | 02D | BABR Repl. by AUBERT 02Q |
| 0.02±0.34±0.05 | 4 AUBERT | 02Q | BABR Repl. by AUBERT,BE 05 |

¹ Rule out the CP-conserving case, $C_{\pi\pi} = S_{\pi\pi} = 0$, at the 5.4 sigma level.

² Rule out the CP-conserving case, $C_{\pi\pi} = S_{\pi\pi} = 0$, at the 5.2 sigma level.

³ Corresponds to 90% confidence range $-0.89 < S_{\pi\pi} < 0.85$.

⁴ Corresponds to 90% confidence range $-0.54 < S_{\pi\pi} < 0.58$.

$C_{\pi^0\pi^0}(B^0 \rightarrow \pi^0\pi^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|-----------|---------------------------------|
| -0.48±0.30 OUR AVERAGE | | | |
| -0.49±0.35±0.05 | AUBERT | 07BC BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| -0.44 ^{+0.52} _{-0.53} ±0.17 | ¹ CHAO | 05 BELL | $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.12±0.56±0.06 | ² AUBERT | 05L BABR | Repl. by AUBERT 07BC |
| ¹ BELLE Collab. quotes $A_{\pi^0\pi^0}$ which is equal to $-C_{\pi^0\pi^0}$. | | | |
| ² Corresponds to a 90% CL interval of $-0.88 < A_{CP} < 0.64$. | | | |

 $C_{\rho\pi}(B^0 \rightarrow \rho^+\pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|---------------------------------|
| 0.01±0.14 OUR AVERAGE | | | |
| 0.15±0.09±0.05 | AUBERT | 07AA BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| -0.13±0.09±0.05 | KUSAKA | 07 BELL | $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.25±0.17 ^{+0.02} _{-0.06} | WANG | 05 BELL | Repl. by KUSAKA 07 |
| 0.36±0.18±0.04 | AUBERT | 03T BABR | Repl. by AUBERT 07AA |

 $S_{\rho\pi}(B^0 \rightarrow \rho^+\pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|---------------------------------|
| 0.01±0.09 OUR AVERAGE | | | |
| -0.03±0.11±0.04 | AUBERT | 07AA BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| 0.06±0.13±0.05 | KUSAKA | 07 BELL | $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.28±0.23 ^{+0.10} _{-0.08} | WANG | 05 BELL | Repl. by KUSAKA 07 |
| 0.19±0.24±0.03 | AUBERT | 03T BABR | Repl. by AUBERT 07AA |

 $\Delta C_{\rho\pi}(B^0 \rightarrow \rho^+\pi^-)$

$\Delta C_{\rho\pi}$ describes the asymmetry between the rates $\Gamma(B^0 \rightarrow \rho^+\pi^-) + \Gamma(\bar{B}^0 \rightarrow \rho^-\pi^+)$ and $\Gamma(B^0 \rightarrow \rho^-\pi^+) + \Gamma(\bar{B}^0 \rightarrow \rho^+\pi^-)$.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|---------------------------------|
| 0.37±0.08 OUR AVERAGE | | | |
| 0.39±0.09±0.09 | AUBERT | 07AA BABR | $e^+e^- \rightarrow \gamma(4S)$ |
| 0.36±0.10±0.05 | KUSAKA | 07 BELL | $e^+e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.38±0.18 ^{+0.02} _{-0.04} | WANG | 05 BELL | Repl. by KUSAKA 07 |
| 0.28 ^{+0.18} _{-0.19} ±0.04 | AUBERT | 03T BABR | Repl. by AUBERT 07AA |

$\Delta S_{\rho\pi} (B^0 \rightarrow \rho^+ \pi^-)$

$\Delta S_{\rho\pi}$ is related to the strong phase difference between the amplitudes contributing to $B^0 \rightarrow \rho^+ \pi^-$.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|----------------------------------|
| -0.05±0.10 OUR AVERAGE | | | |
| -0.01±0.14±0.06 | AUBERT | 07AA BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.08±0.13±0.05 | KUSAKA | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.30±0.24±0.09 | WANG | 05 BELL | Repl. by KUSAKA 07 |
| 0.15±0.25±0.03 | AUBERT | 03T BABR | Repl. by AUBERT 07AA |

 $C_{\rho^0\pi^0} (B^0 \rightarrow \rho^0 \pi^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|----------------------------------|---------|
| -0.10±0.40±0.53 | | | |
| AUBERT | 07AA BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $S_{\rho^0\pi^0} (B^0 \rightarrow \rho^0 \pi^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|-----------|----------------------------------|
| 0.1 ±0.4 OUR AVERAGE | | | |
| 0.04±0.44±0.18 | AUBERT | 07AA BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.17±0.57±0.35 | KUSAKA | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

 $C_{a_1\pi} (B^0 \rightarrow a_1(1260)^+ \pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|----------------------------------|---------|
| -0.10±0.15±0.09 | | | |
| AUBERT | 07O BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $S_{a_1\pi} (B^0 \rightarrow a_1(1260)^+ \pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|----------------------------------|---------|
| 0.37±0.21±0.07 | | | |
| AUBERT | 07O BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $\Delta C_{a_1\pi} (B^0 \rightarrow a_1(1260)^+ \pi^-)$

$\Delta C_{a_1\pi}$ describes the asymmetry between the rates $\Gamma(B^0 \rightarrow a_1^+ \pi^-) + \Gamma(\bar{B}^0 \rightarrow a_1^- \pi^+)$ and $\Gamma(B^0 \rightarrow a_1^- \pi^+) + \Gamma(\bar{B}^0 \rightarrow a_1^+ \pi^-)$.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|----------------------------------|---------|
| 0.26±0.15±0.07 | | | |
| AUBERT | 07O BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $\Delta S_{a_1\pi} (B^0 \rightarrow a_1(1260)^+ \pi^-)$

$\Delta S_{a_1\pi}$ is related to the strong phase difference between the amplitudes contributing to $B^0 \rightarrow a_1 \pi$ decays.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|----------------------------------|---------|
| -0.14±0.21±0.06 | | | |
| AUBERT | 07O BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $C (B^0 \rightarrow b_1^- K^+)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|----------------------------------|---------|
| -0.22±0.23±0.05 | | | |
| AUBERT | 07BI BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

 $\Delta C (B^0 \rightarrow b_1^- \pi^+)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|----------------------------------|---------|
| -1.04±0.23±0.08 | | | |
| AUBERT | 07BI BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

$C_{\rho^0 K_S^0} (B^0 \rightarrow \rho^0 K_S^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|-------------|----------------------------------|
| 0.64±0.41±0.20 | AUBERT | 07F BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

 $S_{\rho^0 K_S^0} (B^0 \rightarrow \rho^0 K_S^0)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|-------------|----------------------------------|
| 0.20±0.52±0.24 | AUBERT | 07F BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

 $C_{\rho\rho} (B^0 \rightarrow \rho^+ \rho^-)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------|--------------------|-------------|----------------|
| -0.05±0.13 OUR AVERAGE | | | |

| | | | |
|--|--------------------|-----------|----------------------------------|
| 0.01±0.15±0.06 | AUBERT | 07BF BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| -0.16±0.21±0.08 | ¹ SOMOV | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.00±0.30±0.09 | ¹ SOMOV | 06 BELL | Repl. by SOMOV 07 |
| -0.03±0.18±0.09 | AUBERT,B | 05C BABR | Repl. by AUBERT 07BF |
| -0.17±0.27±0.14 | AUBERT,B | 04R BABR | Repl. by AUBERT,B 05C |

¹ BELLE Collab. quotes A_{CP} which is equal to $-C$.

 $S_{\rho\rho} (B^0 \rightarrow \rho^+ \rho^-)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------|--------------------|-------------|----------------|
| -0.06±0.17 OUR AVERAGE | | | |

| | | | |
|--|----------|-----------|----------------------------------|
| -0.17±0.20 ^{+0.05} _{-0.06} | AUBERT | 07BF BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.19±0.30±0.08 | SOMOV | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.08±0.41±0.09 | SOMOV | 06 BELL | Repl. by SOMOV 07 |
| -0.33±0.24 ^{+0.08} _{-0.14} | AUBERT,B | 05C BABR | Repl. by AUBERT 07BF |
| -0.42±0.42±0.14 | AUBERT,B | 04R BABR | Repl. by AUBERT,B 05C |

 $|\lambda| (B^0 \rightarrow c\bar{c}K^0)$

The same λ quantity, defined in the $C_{\pi\pi}$ datablock above.

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------------|--------------------|-------------|----------------|
| 0.988±0.020 OUR EVALUATION | | | |
| 0.964±0.025 OUR AVERAGE | | | |

| | | | |
|--|---------------------|-----------|----------------------------------|
| 0.952±0.022±0.017 | ¹ AUBERT | 07AY BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 1.007±0.041±0.033 | ² ABE | 05B BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.950±0.031±0.013 | ³ AUBERT | 05F BABR | Repl. by AUBERT 07AY |
| 0.950±0.049±0.025 | ⁴ ABE | 02Z BELL | Repl. by ABE 05B |
| 0.948±0.051±0.030 | ⁵ AUBERT | 02P BABR | Repl. by AUBERT 05F |

¹ Measurement based on $B^0 \rightarrow c\bar{c}K^0$ decays.

² Measurement based on 152×10^6 $B\bar{B}$ pairs.

³ Measurement based on 227×10^6 $B\bar{B}$ pairs.

⁴ Measured with both $\eta_f = \pm 1$ samples.

⁵ Measured with the high purity of $\eta_f = -1$ samples.

$|\lambda| (B^0 \rightarrow J/\psi K^*(892)^0)$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-----------------------|----------|----------------------------------|
| <0.25 | 95 | ¹ AUBERT,B | 04H BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses the measured cosine coefficients C and \bar{C} and assumes $|q/p| = 1$.

$\cos 2\beta (B^0 \rightarrow J/\psi K^*(892)^0)$

$\beta (\phi_1)$ is one of the angles of CMK unitarity triangle, see the review on "CP" Violation in the Reviews section.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

1.7 $^{+0.7}_{-0.9}$ OUR AVERAGE Error includes scale factor of 1.6.

| | | | |
|---------------------------------|---------------------|----------|----------------------------------|
| $2.72^{+0.50}_{-0.79} \pm 0.27$ | ¹ AUBERT | 05P BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $0.87 \pm 0.74 \pm 0.12$ | ² ITOH | 05 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ The measurement is obtained when $\sin 2\beta$ is fixed to 0.726 and the sign of $\cos 2\beta$ is positive with 86% confidence level.

² The measurement is obtained with $\sin 2\beta$ fixed to 0.731.

$\cos 2\beta (B^0 \rightarrow [K_S^0 \pi^+ \pi^-]_{D^{(*)}} h^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

1.0 $^{+0.6}_{-0.7}$ OUR AVERAGE Error includes scale factor of 1.8.

| | | | |
|---|-----------------------|-----------|----------------------------------|
| $0.42 \pm 0.49 \pm 0.16$ | ¹ AUBERT | 07BH BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $1.87^{+0.40}_{-0.53} {}^{+0.22}_{-0.32}$ | ² KROKOVNY | 06 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ AUBERT 07BH evaluates the likelihoods for the positive and negative solutions assuming $\sin(2\beta_{eff}) = 0.678$. It quotes $L_+ / (L_+ + L_-) = 0.86$ corresponding to a likelihood ratio of $L_+/L_- = 6.14$ in favor of the positive solution.

² KROKOVNY 06 evaluates the likelihoods for the positive and negative solutions assuming $\sin(2\beta_{eff}) = 0.689$. It quotes $L_+ / (L_+ + L_-) = 0.983$ corresponding to a likelihood ratio of $L_+/L_- = 57.8$ in favor of the positive solution.

$(S_+ + S_-)/2 (B^0 \rightarrow D^{*-} \pi^+)$

$S_{\pm} = -\frac{2Im(\lambda_{\pm})}{1+|\lambda_{\pm}|^2}$ where λ_+ and λ_- are defined in the $C_{\pi\pi}$ datablock above for $B^0 \rightarrow D^{*-} \pi^+$ and $\bar{B}^0 \rightarrow D^{*+} \pi^-$.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

-0.037 ± 0.012 OUR AVERAGE

| | | | |
|------------------------------|---------------------|----------|----------------------------------|
| $-0.040 \pm 0.023 \pm 0.010$ | ¹ AUBERT | 06Y BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| $-0.039 \pm 0.020 \pm 0.013$ | ² RONGA | 06 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| $-0.034 \pm 0.014 \pm 0.009$ | ³ AUBERT | 05Z BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------------|----------------------|-----|------|---------------------|
| $-0.030 \pm 0.028 \pm 0.018$ | ³ GERSHON | 05 | BELL | Repl. by RONGA 06 |
| $-0.068 \pm 0.038 \pm 0.020$ | ¹ AUBERT | 04V | BABR | Repl. by AUBERT 06Y |
| $-0.063 \pm 0.024 \pm 0.014$ | ³ AUBERT | 04W | BABR | Repl. by AUBERT 05Z |
| $0.060 \pm 0.040 \pm 0.019$ | ¹ SARANGI | 04 | BELL | Repl. by RONGA 06 |

¹ Uses fully reconstructed $B^0 \rightarrow D^* \pm \pi^\mp$ decays.

² Combines the results from fully reconstructed and partially reconstructed $D^* \pi$ events by taking weighted averages. Assumes that systematic errors from physics parameters and fit biases in the two measurements are 100% correlated.

³ Uses partially reconstructed $B^0 \rightarrow D^* \pm \pi^\mp$ decays.

$(S_- - S_+)/2 (B^0 \rightarrow D^{*-} \pi^+)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|------|---------------------------------------|
| -0.006 ± 0.016 OUR AVERAGE | | | |
| $0.049 \pm 0.042 \pm 0.015$ | ¹ AUBERT | 06Y | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| $-0.011 \pm 0.020 \pm 0.013$ | ² RONGA | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |
| $-0.019 \pm 0.022 \pm 0.013$ | ³ AUBERT | 05Z | BABR $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------------|----------------------|-----|------|---------------------|
| $-0.005 \pm 0.028 \pm 0.018$ | ³ GERSHON | 05 | BELL | Repl. by RONGA 06 |
| $0.031 \pm 0.070 \pm 0.033$ | ¹ AUBERT | 04V | BABR | Repl. by AUBERT 06Y |
| $-0.004 \pm 0.037 \pm 0.014$ | ³ AUBERT | 04W | BABR | Repl. by AUBERT 05Z |
| $0.049 \pm 0.040 \pm 0.019$ | ¹ SARANGI | 04 | BELL | Repl. by RONGA 06 |

¹ Uses fully reconstructed $B^0 \rightarrow D^* \pm \pi^\mp$ decays.

² Combines the results from fully reconstructed and partially reconstructed $D^* \pi$ events by taking weighted averages. Assumes that systematic errors from physics parameters and fit biases in the two measurements are 100% correlated.

³ Uses partially reconstructed $B^0 \rightarrow D^* \pm \pi^\mp$ decays.

$(S_+ + S_-)/2 (B^0 \rightarrow D^- \pi^+)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|------|---------------------------------------|
| -0.046 ± 0.023 OUR AVERAGE | | | |
| $-0.010 \pm 0.023 \pm 0.07$ | ¹ AUBERT | 06Y | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| $-0.050 \pm 0.021 \pm 0.012$ | ² RONGA | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------------|----------------------|-----|------|---------------------|
| $-0.022 \pm 0.038 \pm 0.020$ | ¹ AUBERT | 04V | BABR | Repl. by AUBERT 06Y |
| $-0.062 \pm 0.037 \pm 0.018$ | ¹ SARANGI | 04 | BELL | Repl. by RONGA 06 |

¹ Uses fully reconstructed $B^0 \rightarrow D^\pm \pi^\mp$ decays.

² Combines the results from fully reconstructed and partially reconstructed $D\pi$ events by taking weighted averages. Assumes that systematic errors from physics parameters and fit biases in the two measurements are 100% correlated.

$(S_- - S_+)/2 (B^0 \rightarrow D^- \pi^+)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|------|---------------------------------------|
| -0.022 ± 0.021 OUR AVERAGE | | | |
| $-0.033 \pm 0.042 \pm 0.012$ | ¹ AUBERT | 06Y | BABR $e^+ e^- \rightarrow \gamma(4S)$ |
| $-0.019 \pm 0.021 \pm 0.012$ | ² RONGA | 06 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------------|----------------------|-----|------|---------------------|
| $0.025 \pm 0.068 \pm 0.033$ | ¹ AUBERT | 04V | BABR | Repl. by AUBERT 06Y |
| $-0.025 \pm 0.037 \pm 0.018$ | ¹ SARANGI | 04 | BELL | Repl. by RONGA 06 |

¹ Uses fully reconstructed $B^0 \rightarrow D^\pm \pi^\mp$ decays.

² Combines the results from fully reconstructed and partially reconstructed $D\pi$ events by taking weighted averages. Assumes that systematic errors from physics parameters and fit biases in the two measurements are 100% correlated.

$(S_+ + S_-)/2 (B^0 \rightarrow D^- \rho^+)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|---------------------|----------|----------------------------------|
| -0.024±0.031±0.009 | ¹ AUBERT | 06Y BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses fully reconstructed $B^0 \rightarrow D^- \rho^+$ decays.

$(S_- - S_+)/2 (B^0 \rightarrow D^- \rho^+)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|---------------------|----------|----------------------------------|
| -0.098±0.055±0.018 | ¹ AUBERT | 06Y BABR | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ Uses fully reconstructed $B^0 \rightarrow D^- \rho^+$ decays.

$\sin(2\beta)$

For a discussion of CP violation, see the review on "CP Violation" in the Reviews section. $\sin(2\beta)$ is a measure of the CP -violating amplitude in the $B_d^0 \rightarrow J/\psi(1S) K_S^0$.

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.678±0.025 OUR EVALUATION

0.68 ±0.04 OUR AVERAGE Error includes scale factor of 1.4.

| | | | |
|--|-------------------------|-----------|----------------------------------|
| 0.714±0.032±0.018 | ¹ AUBERT | 07AY BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.642±0.031±0.017 | CHEN | 07 BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 1.56 ±0.42 ±0.21 | ² AUBERT | 04R BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.79 ^{+0.41} _{-0.44} | ³ AFFOLDER | 00C CDF | $p\bar{p}$ at 1.8 TeV |
| 0.84 ^{+0.82} _{-1.04} ±0.16 | ⁴ BARATE | 00Q ALEP | $e^+ e^- \rightarrow Z$ |
| 3.2 ^{+1.8} _{-2.0} ±0.5 | ⁵ ACKERSTAFF | 98Z OPAL | $e^+ e^- \rightarrow Z$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--|----------------------|----------|----------------------------------|
| 0.728±0.056±0.023 | ⁶ ABE | 05B BELL | Repl. by CHEN 07 |
| 0.722±0.040±0.023 | ⁷ AUBERT | 05F BABR | Repl. by AUBERT 07AY |
| 0.99 ±0.14 ±0.06 | ⁸ ABE | 02U BELL | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.719±0.074±0.035 | ⁹ ABE | 02Z BELL | Repl. by ABE 05B |
| 0.59 ±0.14 ±0.05 | ¹⁰ AUBERT | 02N BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.741±0.067±0.034 | ¹¹ AUBERT | 02P BABR | Repl. by AUBERT 05F |
| 0.58 ^{+0.32} _{-0.34} ^{+0.09} _{-0.10} | ABASHIAN | 01 BELL | Repl. by ABE 01G |
| 0.99 ±0.14 ±0.06 | ¹² ABE | 01G BELL | Repl. by ABE 02Z |
| 0.34 ±0.20 ±0.05 | AUBERT | 01 BABR | Repl. by AUBERT 01B |
| 0.59 ±0.14 ±0.05 | ¹² AUBERT | 01B BABR | Repl. by AUBERT 02P |
| 1.8 ±1.1 ±0.3 | ¹³ ABE | 98U CDF | Repl. by AFFOLDER 00C |

- ¹ Measurement based on $B^0 \rightarrow c\bar{c}K^0$ decays.
- ² Measurement in which the J/ψ decays to hadrons or to muons that do not satisfy the standard identification criteria.
- ³ AFFOLDER 00C uses about 400 $B^0 \rightarrow J/\psi(1S)K_S^0$ events. The production flavor of B^0 was determined using three tagging algorithms: a same-side tag, a jet-charge tag, and a soft-lepton tag.
- ⁴ BARATE 00Q uses 23 candidates for $B^0 \rightarrow J/\psi(1S)K_S^0$ decays. A combination of jet-charge, vertex-charge, and same-side tagging techniques were used to determine the B^0 production flavor.
- ⁵ ACKERSTAFF 98Z uses 24 candidates for $B_d^0 \rightarrow J/\psi(1S)K_S^0$ decay. A combination of jet-charge and vertex-charge techniques were used to tag the B_d^0 production flavor.
- ⁶ Measurement based on $152 \times 10^6 B\bar{B}$ pairs.
- ⁷ Measurement based on $227 \times 10^6 B\bar{B}$ pairs.
- ⁸ ABE 02U result is based on the same analysis and data sample reported in ABE 01G.
- ⁹ ABE 02Z result is based on $85 \times 10^6 B\bar{B}$ pairs.
- ¹⁰ AUBERT 02N result based on the same analysis and data sample reported in AUBERT 01B.
- ¹¹ AUBERT 02P result is based on $88 \times 10^6 B\bar{B}$ pairs.
- ¹² First observation of CP violation in B^0 meson system.
- ¹³ ABE 98U uses $198 \pm 17 B_d^0 \rightarrow J/\psi(1S)K^0$ events. The production flavor of B^0 was determined using the same side tagging technique.

$C_{J/\psi K^0} (B^0 \rightarrow J/\psi K^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|----------------------|------|----------------------------------|
| $-0.018 \pm 0.021 \pm 0.014$ | ¹ CHEN 07 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ The paper reports A , which is equal to $-C$.

$S_{J/\psi K^0} (B^0 \rightarrow J/\psi K^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|----------------------------------|
| $0.642 \pm 0.031 \pm 0.017$ | CHEN 07 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

$\sin(2\beta_{\text{eff}})(B^0 \rightarrow \phi K^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|------------------|----------------------------------|---------|
| $0.22 \pm 0.27 \pm 0.12$ | AUBERT 07AX BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | |
|---------------------------------|------------------------------|----------------------|
| $0.50 \pm 0.25^{+0.07}_{-0.04}$ | ¹ AUBERT 05T BABR | Repl. by AUBERT 07AX |
|---------------------------------|------------------------------|----------------------|

¹ Obtained by constraining $C = 0$.

$\sin(2\beta_{\text{eff}})(B^0 \rightarrow K^+ K^- K_S^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|------------------|----------------------------------|---------|
| $0.77 \pm 0.11^{+0.07}_{-0.04}$ | AUBERT 07AX BABR | $e^+ e^- \rightarrow \gamma(4S)$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | |
|--------------------------|------------------------------|----------------------|
| $0.55 \pm 0.22 \pm 0.12$ | ¹ AUBERT 05T BABR | Repl. by AUBERT 07AX |
|--------------------------|------------------------------|----------------------|

¹ Obtained by constraining $C = 0$.

$\sin(2\beta_{\text{eff}})(B^0 \rightarrow [K_S^0 \pi^+ \pi^-]_{D(*)} h^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------------------|---------------------------------------|---------|
| 0.45±0.28 OUR AVERAGE | | | |
| 0.29±0.34±0.06 | AUBERT 07BH BABR | e ⁺ e ⁻ → γ(4S) | |
| 0.78±0.44±0.22 | KROKOVNY 06 BELL | e ⁺ e ⁻ → γ(4S) | |

 $|\lambda| (B^0 \rightarrow [K_S^0 \pi^+ \pi^-]_{D(*)} h^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|---------------------------------------|---------|
| 1.01±0.08±0.02 | | | |
| AUBERT 07BH BABR | | e ⁺ e ⁻ → γ(4S) | |

 $|\sin(2\beta + \gamma)|$

β (ϕ_1) and γ (ϕ_3) are angles of CKM unitarity triangle, see the review on "CP Violation" in the Reviews section.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|--------------|------|---------------------------------------|
| >0.40 | 90 | 1 AUBERT 06Y | BABR | e ⁺ e ⁻ → γ(4S) |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| >0.13 | 95 | 2 RONGA 06 | BELL | e ⁺ e ⁻ → γ(4S) |
| >0.07 | 95 | 2 RONGA 06 | BELL | e ⁺ e ⁻ → γ(4S) |
| >0.35 | 90 | 3 AUBERT 05Z | BABR | e ⁺ e ⁻ → γ(4S) |
| >0.69 | 68 | 4 AUBERT 04V | BABR | e ⁺ e ⁻ → γ(4S) |
| >0.58 | 95 | 5 AUBERT 04W | BABR | Repl. by AUBERT 05Z |

¹ Uses fully reconstructed $B^0 \rightarrow D^{(*)}\pm \pi^\mp$ and $D^\pm \rho^\mp$ decays and some theoretical assumptions.

² Combines the results from fully reconstructed and partially reconstructed $D^{(*)}\pi$ events by taking weighted averages. Assumes that systematic errors from physics parameters and fit biases in the two measurements are 100% correlated.

³ Uses partially reconstructed $B^0 \rightarrow D^{*\pm} \pi^\mp$ decays and some theoretical assumptions.

⁴ Uses fully reconstructed $B^0 \rightarrow D^{(*)}\pm \pi^\mp$ decays and some theoretical assumptions, such as the SU(3) symmetry relation.

⁵ Combining this measurement with the results from AUBERT 04V for fully reconstructed $B^0 \rightarrow D^{(*)}\pm \pi^\mp$ and some theoretical assumptions, such as the SU(3) symmetry relation.

 α

For angle $\alpha(\phi_2)$ of the CKM unitarity triangle, see the review on "CP violation" in the reviews section.

| VALUE (°) | DOCUMENT ID | TECN | COMMENT |
|---|----------------|------|---------------------------------------|
| 96 ± 10 OUR AVERAGE | | | |
| 88 ± 17 | 1 SOMOV 06 | BELL | e ⁺ e ⁻ → γ(4S) |
| 100 ± 13 | 2 AUBERT,B 05C | BABR | e ⁺ e ⁻ → γ(4S) |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 78.6 ± 7.3 | 3 AUBERT 070 | BABR | e ⁺ e ⁻ → γ(4S) |
| 102 $^{+16}_{-12}$ ± 14 | 4 AUBERT,B 04R | BABR | Repl. by AUBERT,B 05C |

¹ Obtained using isospin relation and selecting a solution closest to the CKM best fit average; the 90% CL allowed interval is $59^\circ < \phi_2 (\equiv \alpha) < 115^\circ$.

² Obtained using isospin relation and selecting a solution closest to the CKM best fit average; 90% CL allowed interval is $79^\circ < \alpha < 123^\circ$.

³ The angle α_{eff} is obtained using the measured CP parameters of $B^0 \rightarrow a_1(1260)\pm \pi^\mp$ and choosing one of the four solutions that is compatible with the result of SM-based fits.

⁴ Obtained from the measured CP parameters of the longitudinal polarization by selecting the solution closest to the CKM best fit central value of $\alpha = 95^\circ - 98^\circ$.

$B^0 \rightarrow D^{*-} \ell^+ \nu_\ell$ FORM FACTORS

R_1 (form factor ratio $\sim V/A_1$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|--------------|------|----------------------------------|
| 1.42 ±0.07 OUR AVERAGE | | | |
| 1.429±0.061±0.044 | AUBERT 08R | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 1.18 ±0.30 ±0.12 | DUBOSQ 96 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1.396±0.060±0.044 | AUBERT,B 06Z | BABR | Repl. by AUBERT 08R |

R_2 (form factor ratio $\sim A_2/A_1$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|--------------|------|----------------------------------|
| 0.82 ±0.04 OUR AVERAGE | | | |
| 0.827±0.038±0.022 | AUBERT 08R | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.71 ±0.22 ±0.07 | DUBOSQ 96 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.885±0.040±0.026 | AUBERT,B 06Z | BABR | Repl. by AUBERT 08R |

$\rho_{A_1}^2$ (form factor slope)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|--------------|------|----------------------------------|
| 1.16 ±0.09 OUR AVERAGE | | | |
| Error includes scale factor of 1.6. | | | |
| 1.191±0.048±0.028 | AUBERT 08R | BABR | $e^+ e^- \rightarrow \gamma(4S)$ |
| 0.91 ±0.15 ±0.06 | DUBOSQ 96 | CLE2 | $e^+ e^- \rightarrow \gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1.145±0.059±0.046 | AUBERT,B 06Z | BABR | Repl. by AUBERT 08R |

B^0 REFERENCES

| | | | |
|---------------|----------------|----------------------------|--------------------|
| AALTONEN 08I | PRL 100 101802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AUBERT 08B | PR D77 011102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 08C | PR D77 011104R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 08E | PR D77 012003 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 08F | PRL 100 051803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 08I | PRL 100 081801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 08N | PRL 100 021801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 08P | PR D77 032007 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 08Q | PRL 100 151802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 08R | PR D77 032002 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| LIVENTSEV 08 | PR D77 091503R | D. Liventsev <i>et al.</i> | (BELLE Collab.) |
| UCHIDA 08 | PR D77 051101R | Y. Uchida <i>et al.</i> | (BELLE Collab.) |
| USHIRODA 08 | PRL 100 021602 | Y. Ushiroda <i>et al.</i> | (BELLE Collab.) |
| ABAZOV 07S | PRL 99 142001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABULENCIA 07A | PRL 98 122001 | A. Abulencia <i>et al.</i> | (FNAL CDF Collab.) |
| ADAM 07 | PRL 99 041802 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| Also | PR D76 012007 | D.M. Asner <i>et al.</i> | (CLEO Collab.) |
| AUBERT 07A | PRL 98 031801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AA | PR D76 012004 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AC | PR D76 031101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AD | PR D76 031102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AE | PR D76 031103R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AF | PRL 99 021603 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AG | PRL 99 051801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AI | PRL 99 071801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AJ | PRL 99 081801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AN | PR D76 051101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AO | PR D76 051103R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AQ | PR D76 071101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AS | PR D76 071104R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT 07AT | PR D76 091101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |

| | | | | |
|-------------|------|----------------|----------------------------|-----------------|
| AUBERT | 07AV | PR D76 092004 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07AX | PRL 99 161802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07AY | PRL 99 171803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07B | PR D75 012008 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07BC | PR D76 091102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07BF | PR D76 052007 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07BH | PRL 99 231802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07BI | PRL 99 241803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07BO | PR D76 111102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07D | PRL 98 051801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07E | PRL 98 051802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07F | PRL 98 051803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07G | PRL 98 111801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07H | PR D75 031101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07J | PRL 98 091801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07K | PRL 98 081801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07L | PRL 98 151802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07N | PR D75 072002 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07O | PRL 98 181803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07Q | PR D75 051102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07R | PRL 98 211804 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07Y | PR D75 111102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| CHANG | 07A | PRL 98 131803 | M.-C. Chang <i>et al.</i> | (BELLE Collab.) |
| CHANG | 07B | PR D75 071104R | P. Chang <i>et al.</i> | (BELLE Collab.) |
| CHAO | 07 | PR D76 091103R | Y. Chao <i>et al.</i> | (BELLE Collab.) |
| CHEN | 07 | PRL 98 031802 | K.-F. Chen <i>et al.</i> | (BELLE Collab.) |
| CHEN | 07D | PRL 99 221802 | K.-F. Chen <i>et al.</i> | (BELLE Collab.) |
| DALSENO | 07 | PR D76 072004 | J. Dalseno <i>et al.</i> | (BELLE Collab.) |
| FRATINA | 07 | PRL 98 221802 | S. Fratina <i>et al.</i> | (BELLE Collab.) |
| GARMASH | 07 | PR D75 012006 | A. Garmash <i>et al.</i> | (BELLE Collab.) |
| HOKUUE | 07 | PL B648 139 | T. Hokuee <i>et al.</i> | (BELLE Collab.) |
| ISHINO | 07 | PRL 98 211801 | H. Ishino <i>et al.</i> | (BELLE Collab.) |
| KUSAKA | 07 | PRL 98 221602 | A. Kusaka <i>et al.</i> | (BELLE Collab.) |
| KUZMIN | 07 | PR D76 012006 | A. Kuzmin <i>et al.</i> | (BELLE Collab.) |
| LIN | 07 | PRL 98 181804 | S.-W. Lin <i>et al.</i> | (BELLE Collab.) |
| LIN | 07A | PRL 99 121601 | S.-W. Lin <i>et al.</i> | (BELLE Collab.) |
| MATYJA | 07 | PRL 99 191807 | A. Matyja <i>et al.</i> | (BELLE Collab.) |
| MEDVEDEVA | 07 | PR D76 051102R | T. Medvedeva <i>et al.</i> | (BELLE Collab.) |
| PARK | 07 | PR D75 011101R | K.S. Park <i>et al.</i> | (BELLE Collab.) |
| SCHUEMANN | 07 | PR D75 092002 | J. Schuemann <i>et al.</i> | (BELLE Collab.) |
| SOMOV | 07 | PR D76 011104R | A. Somov <i>et al.</i> | (BELLE Collab.) |
| TSAI | 07 | PR D75 111101R | Y.-T. Tsai <i>et al.</i> | (BELLE Collab.) |
| URQUIJO | 07 | PR D75 032001 | P. Urquijo <i>et al.</i> | (BELLE Collab.) |
| WANG | 07B | PR D75 092005 | C.H. Wang <i>et al.</i> | (BELLE Collab.) |
| WANG | 07C | PR D76 052004 | M.-Z. Wang <i>et al.</i> | (BELLE Collab.) |
| XIE | 07 | PR D75 017101 | Q.L. Xie <i>et al.</i> | (BELLE Collab.) |
| ZUPANC | 07 | PR D75 091102R | A. Zupanc <i>et al.</i> | (BELLE Collab.) |
| ABAZOV | 06S | PR D74 092001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 06W | PR D74 112002 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABULENCIA,A | 06D | PRL 97 211802 | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| ACOSTA | 06 | PRL 96 202001 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| AUBERT | 06 | PR D73 011101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06A | PRL 96 011803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06E | PRL 96 052002 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06G | PR D73 012004 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06I | PR D73 031101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06L | PR D74 012001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06N | PR D74 031103R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06S | PRL 96 241802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06T | PRL 96 251802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06V | PRL 97 051802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06W | PR D73 071102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06X | PR D73 071103R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06Y | PR D73 111101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06A | PR D73 112004 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06B | PR D74 011101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06C | PR D74 011102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06E | PR D74 011106R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06G | PRL 97 201801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06H | PRL 97 201802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06J | PR D73 092001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |

| | | | | |
|-----------|-----|-------------------------|----------------------------|-----------------|
| AUBERT,B | 06K | PRL 97 211801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06L | PR D74 031101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06M | PR D74 031102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06O | PR D74 031104R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06P | PR D74 031105R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06Q | PR D74 091101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06R | PR D74 032005 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06S | PR D74 051101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06T | PR D74 051102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06V | PR D74 051106R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06Y | PR D74 091105R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 06Z | PR D74 092004 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 06C | PRL 97 171805 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 06H | PRL 97 261803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 06J | PR D74 111102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 06N | PR D74 072008 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| BLYTH | 06 | PR D74 092002 | S. Blyth <i>et al.</i> | (BELLE Collab.) |
| CHISTOV | 06A | PR D74 111105R | R. Chistov <i>et al.</i> | (BELLE Collab.) |
| DRAGIC | 06 | PR D73 111105R | J. Dragic <i>et al.</i> | (BELLE Collab.) |
| GABYSHEV | 06 | PRL 97 202003 | N. Gabyshev <i>et al.</i> | (BELLE Collab.) |
| GOKHROO | 06 | PRL 97 162002 | G. Gokhroo <i>et al.</i> | (BELLE Collab.) |
| JEN | 06 | PR D74 111101R | C.-M. Jen <i>et al.</i> | (BELLE Collab.) |
| KROKOVNY | 06 | PRL 97 081801 | P. Krokovsky <i>et al.</i> | (BELLE Collab.) |
| MOHAPATRA | 06 | PRL 96 221601 | D. Mohapatra <i>et al.</i> | (BELLE Collab.) |
| NAKANO | 06 | PR D73 112002 | E. Nakano <i>et al.</i> | (BELLE Collab.) |
| RONGA | 06 | PR D73 092003 | F.J. Ronga <i>et al.</i> | (BELLE Collab.) |
| SCHUEMANN | 06 | PRL 97 061802 | J. Schuemann <i>et al.</i> | (BELLE Collab.) |
| SOMOV | 06 | PRL 96 171801 | A. Somov <i>et al.</i> | (BELLE Collab.) |
| SONI | 06 | PL B634 155 | N. Soni <i>et al.</i> | (BELLE Collab.) |
| USHIRODA | 06 | PR D74 111104R | Y. Ushiroda <i>et al.</i> | (BELLE Collab.) |
| VILLA | 06 | PR D73 051107R | S. Villa <i>et al.</i> | (BELLE Collab.) |
| ABAZOV | 05B | PRL 94 042001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 05C | PRL 94 102001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 05D | PRL 94 182001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 05W | PRL 95 171801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABE | 05A | PRL 94 221805 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 05B | PR D71 072003 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| Also | | PR D71 079903 (errat.) | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 05D | PRL 95 101801 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 05G | PRL 95 231802 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABULENCIA | 05 | PRL 95 221805 | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| Also | | PRL 95 249905 (erratum) | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| ACOSTA | 05 | PRL 94 101803 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| AUBERT | 05 | PRL 94 011801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05B | PR D71 031501R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05E | PR D71 051502R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05F | PRL 94 161803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05I | PRL 94 131801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05J | PRL 94 141801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05K | PRL 94 171801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05L | PRL 94 181802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05M | PRL 94 191802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05O | PR D71 031103R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05P | PR D71 032005 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05T | PR D71 091102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05U | PR D71 091103R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05V | PR D71 091104R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05W | PRL 94 221803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05Y | PR D71 111102 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05Z | PR D71 112003 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 05 | PRL 95 011801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 05C | PRL 95 041805 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 05K | PRL 95 131803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 05O | PR D72 051102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 05P | PR D72 051103R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 05Q | PR D72 051106R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 05Z | PRL 95 131802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 05 | PRL 95 151803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 05A | PRL 95 151804 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 05B | PRL 95 171802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 05C | PR D72 091103R | B. Aubert <i>et al.</i> | (BABAR Collab.) |

| | | | | |
|-----------|-----|----------------|----------------------------|------------------|
| AUBERT,BE | 05E | PRL 95 221801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 05F | PR D72 111101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| CHANG | 05 | PR D71 072007 | M.-C. Chang <i>et al.</i> | (BELLE Collab.) |
| CHANG | 05A | PR D71 091106R | P. Chang <i>et al.</i> | (BELLE Collab.) |
| CHAO | 05 | PRL 94 181803 | Y. Chao <i>et al.</i> | (BELLE Collab.) |
| CHAO | 05A | PR D71 031502R | Y. Chao <i>et al.</i> | (BELLE Collab.) |
| CHEN | 05A | PRL 94 221804 | K.-F. Chen <i>et al.</i> | (BELLE Collab.) |
| CHEN | 05B | PR D72 012004 | K.-F. Chen <i>et al.</i> | (BELLE Collab.) |
| DRUTSKOY | 05 | PRL 94 061802 | A. Drutskoy <i>et al.</i> | (BELLE Collab.) |
| GERSHON | 05 | PL B624 11 | T. Gershon <i>et al.</i> | (BELLE Collab.) |
| ITOH | 05 | PRL 95 091601 | R. Itoh <i>et al.</i> | (BELLE Collab.) |
| LIVENTSEV | 05 | PR D72 051109R | D. Liventsev <i>et al.</i> | (BELLE Collab.) |
| MAJUMDER | 05 | PRL 95 041803 | G. Majumder <i>et al.</i> | (BELLE Collab.) |
| MIYAKE | 05 | PL B618 34 | H. Miyake <i>et al.</i> | (BELLE Collab.) |
| MOHAPATRA | 05 | PR D72 011101R | D. Mohapatra <i>et al.</i> | (BELLE Collab.) |
| NISHIDA | 05 | PL B610 23 | S. Nishida <i>et al.</i> | (BELLE Collab.) |
| OKABE | 05 | PL B614 27 | T. Okabe <i>et al.</i> | (BELLE Collab.) |
| SCHUMANN | 05 | PR D72 011103R | J. Schumann <i>et al.</i> | (BELLE Collab.) |
| SUMISAWA | 05 | PRL 95 061801 | K. Sumisawa <i>et al.</i> | (BELLE Collab.) |
| USHIRODA | 05 | PRL 94 231601 | Y. Ushiroda <i>et al.</i> | (BELLE Collab.) |
| WANG | 05 | PRL 94 121801 | C.C. Wang <i>et al.</i> | (BELLE Collab.) |
| WANG | 05A | PL B617 141 | M.-Z. Wang <i>et al.</i> | (BELLE Collab.) |
| XIE | 05 | PR D72 051105R | Q.L. Xie <i>et al.</i> | (BELLE Collab.) |
| YANG | 05 | PRL 94 111802 | H. Yang <i>et al.</i> | (BELLE Collab.) |
| ZHANG | 05B | PR D71 091107R | L.M. Zhang <i>et al.</i> | (BELLE Collab.) |
| ABDALLAH | 04D | EPJ C33 213 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ABDALLAH | 04E | EPJ C33 307 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ABE | 04E | PRL 93 021601 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ACOSTA | 04D | PRL 93 032001 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| AUBERT | 04A | PR D69 011102 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 04B | PR D69 032004 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 04C | PRL 92 111801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 04G | PR D69 031102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 04H | PRL 92 061801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 04M | PRL 92 201802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 04R | PR D69 052001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 04U | PR D69 091503R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 04V | PRL 92 251801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 04W | PRL 92 251802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 04Y | PRL 93 041801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 04Z | PRL 93 051802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04B | PR D70 011101R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04C | PR D70 012007 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| Also | | PRL 92 181801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04D | PR D70 032006 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04G | PRL 93 071801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04H | PRL 93 081801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04J | PRL 93 091802 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04K | PRL 93 131801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04M | PRL 93 131805 | B. Aubert | (BABAR Collab.) |
| AUBERT,B | 04O | PR D70 091103R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04R | PRL 93 231801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04S | PRL 93 181801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04T | PR D70 091104R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04U | PR D70 091105R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04V | PRL 93 181805 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04W | PRL 93 231804 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04X | PRL 93 181806 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04Z | PRL 93 201801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 04 | PR D70 111102R | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 04A | PR D70 112006 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 04B | PR D70 091106 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUSHEV | 04 | PRL 93 201802 | T. Aushev <i>et al.</i> | (BELLE Collab.) |
| BORNHEIM | 04 | PRL 93 241802 | A. Bornheim <i>et al.</i> | (CLEO Collab.) |
| CHANG | 04 | PL B599 148 | P. Chang <i>et al.</i> | (BELLE Collab.) |
| CHAO | 04 | PR D69 111102R | Y. Chao <i>et al.</i> | (BELLE Collab.) |
| CHAO | 04B | PRL 93 191802 | Y. Chao <i>et al.</i> | (BELLE Collab.) |
| DRAGIC | 04 | PRL 93 131802 | J. Dragic | (BELLE Collab.) |
| DRUTSKOY | 04 | PRL 92 051801 | A. Drutskoy <i>et al.</i> | (BELLE Collab.) |
| GARMASH | 04 | PR D69 012001 | A. Garmash <i>et al.</i> | (BELLE Collab.) |
| KATAOKA | 04 | PRL 93 261801 | S.U. Kataoka <i>et al.</i> | (BELLE Collab.) |

| | | | | |
|------------|-----|-------------------------|-------------------------------|------------------|
| MAJUMDER | 04 | PR D70 111103R | G. Majumder <i>et al.</i> | (BELLE Collab.) |
| NAKAO | 04 | PR D69 112001 | M. Nakao <i>et al.</i> | (BELLE Collab.) |
| SARANGI | 04 | PRL 93 031802 | T.R. Sarangi <i>et al.</i> | (BELLE Collab.) |
| WANG | 04 | PRL 92 131801 | M.Z. Wang <i>et al.</i> | (BELLE Collab.) |
| WANG | 04A | PR D70 012001 | C.H. Wang <i>et al.</i> | (BELLE Collab.) |
| ABDALLAH | 03B | EPJ C28 155 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ABE | 03B | PR D67 032003 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 03C | PR D67 031102R | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 03G | PR D68 012001 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 03H | PRL 91 261602 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ADAM | 03 | PR D67 032001 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| ATHAR | 03 | PR D68 072003 | S.B. Athar <i>et al.</i> | (CLEO Collab.) |
| AUBERT | 03B | PRL 90 091801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03C | PR D67 072002 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03D | PRL 90 181803 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03E | PRL 90 181801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03H | PR D67 091101R | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03I | PR D67 092003 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03J | PRL 90 221801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03K | PRL 90 231801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03L | PRL 91 021801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03N | PRL 91 061802 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03O | PRL 91 071801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03Q | PRL 91 131801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03S | PRL 91 241801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03T | PRL 91 201802 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03U | PRL 91 221802 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03V | PRL 91 171802 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03W | PR D68 161801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 03X | PR D68 092001 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| BORNHEIM | 03 | PR D68 052002 | A. Bornheim <i>et al.</i> | (CLEO Collab.) |
| CHANG | 03 | PR D68 111101R | M.-C. Chang <i>et al.</i> | (BELLE Collab.) |
| CHEN | 03B | PRL 91 201801 | K.-F. Chen <i>et al.</i> | (BELLE Collab.) |
| CSORNA | 03 | PR D67 112002 | S.E. Csorna <i>et al.</i> | (CLEO Collab.) |
| EISENSTEIN | 03 | PR D68 017101 | B.I. Eisenstein <i>et al.</i> | (CLEO Collab.) |
| FANG | 03 | PRL 90 071801 | F. Fang <i>et al.</i> | (BELLE Collab.) |
| GABYSHEV | 03 | PRL 90 121802 | N. Gabyshev <i>et al.</i> | (BELLE Collab.) |
| HASTINGS | 03 | PR D67 052004 | N.C. Hastings <i>et al.</i> | (BELLE Collab.) |
| ISHIKAWA | 03 | PRL 91 261601 | A. Ishikawa <i>et al.</i> | (BELLE Collab.) |
| KROKOVNY | 03 | PRL 90 141802 | P. Krokovny <i>et al.</i> | (BELLE Collab.) |
| KROKOVNY | 03B | PRL 91 262002 | P. Krokovny <i>et al.</i> | (BELLE Collab.) |
| LEE | 03 | PRL 91 261801 | S.H. Lee <i>et al.</i> | (BELLE Collab.) |
| SATPATHY | 03 | PL B553 159 | A. Satpathy <i>et al.</i> | (BELLE Collab.) |
| WANG | 03 | PRL 90 201802 | M.-Z. Wang <i>et al.</i> | (BELLE Collab.) |
| ZHENG | 03 | PR D67 092004 | Y. Zheng <i>et al.</i> | (BELLE Collab.) |
| ABE | 02 | PRL 88 021801 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02E | PL B526 258 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02F | PL B526 247 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02H | PRL 88 171801 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02J | PRL 88 052002 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02K | PRL 88 181803 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02M | PRL 89 071801 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02N | PL B538 11 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02O | PR D65 091103R | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02Q | PRL 89 122001 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02U | PR D66 032007 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02W | PRL 89 151802 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 02Z | PR D66 071102R | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ACOSTA | 02C | PR D65 092009 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ACOSTA | 02G | PR D66 112002 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| AFFOLDER | 02B | PRL 88 071801 | T. Affolder <i>et al.</i> | (CDF Collab.) |
| AHMED | 02B | PR D66 031101R | S. Ahmed <i>et al.</i> | (CLEO Collab.) |
| ASNER | 02 | PR D65 031103R | D.M. Asner <i>et al.</i> | (CLEO Collab.) |
| AUBERT | 02 | PR D65 032001 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02C | PRL 88 101805 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02D | PR D65 051502R | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02E | PR D65 051101R | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02H | PRL 89 011802 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| Also | | PRL 89 169903 (erratum) | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02I | PRL 88 221802 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02J | PRL 88 221803 | B. Aubert <i>et al.</i> | (BaBar Collab.) |

| | | | | |
|-----------------|-----|-------------------------|----------------------------------|------------------|
| AUBERT | 02K | PRL 88 231801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02L | PRL 88 241801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02M | PRL 89 061801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02N | PR D66 032003 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02P | PRL 89 201802 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 02Q | PRL 89 281802 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| BRIERE | 02 | PRL 89 081803 | R. Briere <i>et al.</i> | (CLEO Collab.) |
| CASEY | 02 | PR D66 092002 | B.C.K. Casey <i>et al.</i> | (BELLE Collab.) |
| CHEN | 02B | PL B546 196 | K.-F. Chen <i>et al.</i> | (BELLE Collab.) |
| COAN | 02 | PRL 88 062001 | T.E. Coan <i>et al.</i> | (CLEO Collab.) |
| Also | | PRL 88 069902 (erratum) | T.E. Coan <i>et al.</i> | (CLEO Collab.) |
| DRUTSKOY | 02 | PL B542 171 | A. Drutskoy <i>et al.</i> | (BELLE Collab.) |
| DYTMAN | 02 | PR D66 091101R | S.A. Dytman <i>et al.</i> | (CLEO Collab.) |
| ECKHART | 02 | PRL 89 251801 | E. Eckhart <i>et al.</i> | (CLEO Collab.) |
| EDWARDS | 02 | PR D65 012002 | K.W. Edwards <i>et al.</i> | (CLEO Collab.) |
| GABYSHEV | 02 | PR D66 091102R | N. Gabyshev <i>et al.</i> | (BELLE Collab.) |
| GODANG | 02 | PRL 88 021802 | R. Godang <i>et al.</i> | (CLEO Collab.) |
| GORDON | 02 | PL B542 183 | A. Gordon <i>et al.</i> | (BELLE Collab.) |
| HARA | 02 | PRL 89 251803 | K. Hara <i>et al.</i> | (BELLE Collab.) |
| KROKOVNY | 02 | PRL 89 231804 | P. Korkovny <i>et al.</i> | (BELLE Collab.) |
| MAHAPATRA | 02 | PRL 88 101803 | R. Mahapatra <i>et al.</i> | (CLEO Collab.) |
| NISHIDA | 02 | PRL 89 231801 | S. Nishida <i>et al.</i> | (BELLE Collab.) |
| TOMURA | 02 | PL B542 207 | T. Tomura <i>et al.</i> | (BELLE Collab.) |
| ABASHIAN | 01 | PRL 86 2509 | A. Abashian <i>et al.</i> | (BELLE Collab.) |
| ABE | 01D | PRL 86 3228 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 01G | PRL 87 091802 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 01H | PRL 87 101801 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 01I | PRL 87 111801 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 01K | PR D64 071101 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 01L | PRL 87 161601 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABE | 01M | PL B517 309 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ABREU | 01H | PL B510 55 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ALEXANDER | 01B | PR D64 092001 | J.P. Alexander <i>et al.</i> | (CLEO Collab.) |
| AMMAR | 01B | PRL 87 271801 | R. Ammar <i>et al.</i> | (CLEO Collab.) |
| ANDERSON | 01 | PRL 86 2732 | S. Anderson <i>et al.</i> | (CLEO Collab.) |
| ANDERSON | 01B | PRL 87 181803 | S. Anderson <i>et al.</i> | (CLEO Collab.) |
| AUBERT | 01 | PRL 86 2515 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 01B | PRL 87 091801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 01D | PRL 87 151801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 01E | PRL 87 151802 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 01F | PRL 87 201803 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 01G | PRL 87 221802 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 01H | PRL 87 241801 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| AUBERT | 01I | PRL 87 241803 | B. Aubert <i>et al.</i> | (BaBar Collab.) |
| BARATE | 01D | EPJ C20 431 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BRIERE | 01 | PRL 86 3718 | R.A. Biere <i>et al.</i> | (CLEO Collab.) |
| EDWARDS | 01 | PRL 86 30 | K.W. Edwards <i>et al.</i> | (CLEO Collab.) |
| JAFFE | 01 | PRL 86 5000 | D. Jaffe <i>et al.</i> | (CLEO Collab.) |
| RICHICHI | 01 | PR D63 031103R | S.J. Richichi <i>et al.</i> | (CLEO Collab.) |
| ABBIENDI | 00Q | PL B482 15 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI,G | 00B | PL B493 266 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABE | 00C | PR D62 071101R | K. Abe <i>et al.</i> | (SLD Collab.) |
| AFFOLDER | 00C | PR D61 072005 | T. Affolder <i>et al.</i> | (CDF Collab.) |
| AFFOLDER | 00N | PRL 85 4668 | T. Affolder <i>et al.</i> | (CDF Collab.) |
| AHMED | 00B | PR D62 112003 | S. Ahmed <i>et al.</i> | (CLEO Collab.) |
| ANASTASSOV | 00 | PRL 84 1393 | A. Anastassov <i>et al.</i> | (CLEO Collab.) |
| ARTUSO | 00 | PRL 84 4292 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| AVERY | 00 | PR D62 051101 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| BARATE | 00Q | PL B492 259 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BARATE | 00R | PL B492 275 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BEHRENS | 00 | PR D61 052001 | B.H. Behrens <i>et al.</i> | (CLEO Collab.) |
| BEHRENS | 00B | PL B490 36 | B.H. Behrens <i>et al.</i> | (CLEO Collab.) |
| BERGFELD | 00B | PR D62 091102R | T. Bergfeld <i>et al.</i> | (CLEO Collab.) |
| CHEN | 00 | PRL 85 525 | S. Chen <i>et al.</i> | (CLEO Collab.) |
| COAN | 00 | PRL 84 5283 | T.E. Coan <i>et al.</i> | (CLEO Collab.) |
| CRONIN-HEN...00 | | PRL 85 515 | D. Cronin-Hennessy <i>et al.</i> | (CLEO Collab.) |
| CSORNA | 00 | PR D61 111101 | S.E. Csorna <i>et al.</i> | (CLEO Collab.) |
| JESSOP | 00 | PRL 85 2881 | C.P. Jessop <i>et al.</i> | (CLEO Collab.) |
| LIPELES | 00 | PR D62 032005 | E. Lipeles <i>et al.</i> | (CLEO Collab.) |
| RICHICHI | 00 | PRL 85 520 | S.J. Richichi <i>et al.</i> | (CLEO Collab.) |
| ABBIENDI | 99J | EPJ C12 609 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |

| | | | | |
|-------------|-----|------------------------|------------------------------|------------------|
| ABE | 99K | PR D60 051101 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 99Q | PR D60 072003 | F. Abe <i>et al.</i> | (CDF Collab.) |
| AFFOLDER | 99B | PRL 83 3378 | T. Affolder <i>et al.</i> | (CDF Collab.) |
| AFFOLDER | 99C | PR D60 112004 | T. Affolder <i>et al.</i> | (CDF Collab.) |
| ARTUSO | 99 | PRL 82 3020 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| BARTEL | 99 | PRL 82 3746 | J. Bartelt <i>et al.</i> | (CLEO Collab.) |
| COAN | 99 | PR D59 111101 | T.E. Coan <i>et al.</i> | (CLEO Collab.) |
| ABBOTT | 98B | PL B423 419 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABE | 98 | PR D57 R3811 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 98B | PR D57 5382 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 98C | PRL 80 2057 | F. Abe <i>et al.</i> | (CDF Collab.) |
| Also | | PR D59 032001 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 98O | PR D58 072001 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 98Q | PR D58 092002 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 98U | PRL 81 5513 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 98V | PRL 81 5742 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ACCIARRI | 98D | EPJ C5 195 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 98S | PL B438 417 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACKERSTAFF | 98Z | EPJ C5 379 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| BARATE | 98Q | EPJ C4 387 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BEHRENS | 98 | PRL 80 3710 | B.H. Behrens <i>et al.</i> | (CLEO Collab.) |
| BERGFELD | 98 | PRL 81 272 | T. Bergfeld <i>et al.</i> | (CLEO Collab.) |
| BRANDENB... | 98 | PRL 80 2762 | G. Brandenbrug <i>et al.</i> | (CLEO Collab.) |
| GODANG | 98 | PRL 80 3456 | R. Godang <i>et al.</i> | (CLEO Collab.) |
| NEMATI | 98 | PR D57 5363 | B. Nematici <i>et al.</i> | (CLEO Collab.) |
| ABE | 97J | PRL 79 590 | K. Abe <i>et al.</i> | (SLD Collab.) |
| ABREU | 97F | ZPHY C74 19 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| Also | | ZPHY C75 579 (erratum) | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 97N | ZPHY C76 579 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 97B | PL B391 474 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 97C | PL B391 481 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACKERSTAFF | 97G | PL B395 128 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 97U | ZPHY C76 401 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 97V | ZPHY C76 417 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ARTUSO | 97 | PL B399 321 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| ASNER | 97 | PRL 79 799 | D. Asner <i>et al.</i> | (CLEO Collab.) |
| ATHANAS | 97 | PRL 79 2208 | M. Athanas <i>et al.</i> | (CLEO Collab.) |
| BUSKULIC | 97 | PL B395 373 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 97D | ZPHY C75 397 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| FU | 97 | PRL 79 3125 | X. Fu <i>et al.</i> | (CLEO Collab.) |
| JESSOP | 97 | PRL 79 4533 | C.P. Jessop <i>et al.</i> | (CLEO Collab.) |
| ABE | 96B | PR D53 3496 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 96C | PRL 76 4462 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 96H | PRL 76 2015 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 96L | PRL 76 4675 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 96Q | PR D54 6596 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 96P | ZPHY C71 539 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 96Q | ZPHY C72 17 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 96E | PL B383 487 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ADAM | 96D | ZPHY C72 207 | W. Adam <i>et al.</i> | (DELPHI Collab.) |
| ALBRECHT | 96D | PL B374 256 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALEXANDER | 96T | PRL 77 5000 | J.P. Alexander <i>et al.</i> | (CLEO Collab.) |
| ALEXANDER | 96V | ZPHY C72 377 | G. Alexander <i>et al.</i> | (OPAL Collab.) |
| ASNER | 96 | PR D53 1039 | D.M. Asner <i>et al.</i> | (CLEO Collab.) |
| BARISH | 96B | PRL 76 1570 | B.C. Barish <i>et al.</i> | (CLEO Collab.) |
| BISHAI | 96 | PL B369 186 | M. Bishai <i>et al.</i> | (CLEO Collab.) |
| BUSKULIC | 96J | ZPHY C71 31 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 96V | PL B384 471 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| DUBOSQ | 96 | PRL 76 3898 | J.E. Duboscq <i>et al.</i> | (CLEO Collab.) |
| GIBAUT | 96 | PR D53 4734 | D. Gibaut <i>et al.</i> | (CLEO Collab.) |
| PDG | 96 | PR D54 1 | R. M. Barnett <i>et al.</i> | |
| ABE | 95Z | PRL 75 3068 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 95N | PL B357 255 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 95Q | ZPHY C68 13 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 95H | PL B363 127 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 95I | PL B363 137 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ADAM | 95 | ZPHY C68 363 | W. Adam <i>et al.</i> | (DELPHI Collab.) |
| AKERS | 95J | ZPHY C66 555 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| AKERS | 95T | ZPHY C67 379 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| ALEXANDER | 95 | PL B341 435 | J. Alexander <i>et al.</i> | (CLEO Collab.) |
| Also | | PL B347 469 (erratum) | J. Alexander <i>et al.</i> | (CLEO Collab.) |

| | | | | |
|---|-----|-----------------------|-----------------------------|------------------------|
| BARISH | 95 | PR D51 1014 | B.C. Barish <i>et al.</i> | (CLEO Collab.) |
| BUSKULIC | 95N | PL B359 236 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| ABE | 94D | PRL 72 3456 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 94M | PL B338 409 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| AKERS | 94C | PL B327 411 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| AKERS | 94H | PL B336 585 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| AKERS | 94J | PL B337 196 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| AKERS | 94L | PL B337 393 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| ALAM | 94 | PR D50 43 | M.S. Alam <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 94 | PL B324 249 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 94G | PL B340 217 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| AMMAR | 94 | PR D49 5701 | R. Ammar <i>et al.</i> | (CLEO Collab.) |
| ATHANAS | 94 | PRL 73 3503 | M. Athanas <i>et al.</i> | (CLEO Collab.) |
| Also | | PRL 74 3090 (erratum) | M. Athanas <i>et al.</i> | (CLEO Collab.) |
| BUSKULIC | 94B | PL B322 441 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| PDG | 94 | PR D50 1173 | L. Montanet <i>et al.</i> | (CERN, LBL, BOST+) |
| PROCARIO | 94 | PRL 73 1472 | M. Procario <i>et al.</i> | (CLEO Collab.) |
| STONE | 94 | HEPSY 93-11 | S. Stone | |
| Published in <i>B Decays</i> , 2nd Edition, World Scientific, Singapore | | | | |
| ABREU | 93D | ZPHY C57 181 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 93G | PL B312 253 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACTON | 93C | PL B307 247 | P.D. Acton <i>et al.</i> | (OPAL Collab.) |
| ALBRECHT | 93 | ZPHY C57 533 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 93E | ZPHY C60 11 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALEXANDER | 93B | PL B319 365 | J. Alexander <i>et al.</i> | (CLEO Collab.) |
| AMMAR | 93 | PRL 71 674 | R. Ammar <i>et al.</i> | (CLEO Collab.) |
| BARTELT | 93 | PRL 71 1680 | J.E. Bartelt <i>et al.</i> | (CLEO Collab.) |
| BATTLE | 93 | PRL 71 3922 | M. Battle <i>et al.</i> | (CLEO Collab.) |
| BEAN | 93B | PRL 70 2681 | A. Bean <i>et al.</i> | (CLEO Collab.) |
| BUSKULIC | 93D | PL B307 194 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| Also | | PL B325 537 (erratum) | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 93K | PL B313 498 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| SANGHERA | 93 | PR D47 791 | S. Sanghera <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 92C | PL B275 195 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 92G | ZPHY C54 1 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 92L | ZPHY C55 357 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BORTOLETTO | 92 | PR D45 21 | D. Bortoletto <i>et al.</i> | (CLEO Collab.) |
| HENDERSON | 92 | PR D45 2212 | S. Henderson <i>et al.</i> | (CLEO Collab.) |
| KRAMER | 92 | PL B279 181 | G. Kramer, W.F. Palmer | (HAMB, OSU) |
| ALBAJAR | 91C | PL B262 163 | C. Albajar <i>et al.</i> | (UA1 Collab.) |
| ALBAJAR | 91E | PL B273 540 | C. Albajar <i>et al.</i> | (UA1 Collab.) |
| ALBRECHT | 91B | PL B254 288 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 91C | PL B255 297 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 91E | PL B262 148 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BERKELMAN | 91 | ARNPS 41 1 | K. Berkelman, S. Stone | (CORN, SYRA) |
| "Decays of <i>B</i> Mesons" | | | | |
| FULTON | 91 | PR D43 651 | R. Fulton <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 90B | PL B241 278 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 90J | ZPHY C48 543 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ANTREASYAN | 90B | ZPHY C48 553 | D. Antreasyan <i>et al.</i> | (Crystal Ball Collab.) |
| BORTOLETTO | 90 | PRL 64 2117 | D. Bortoletto <i>et al.</i> | (CLEO Collab.) |
| ELSEN | 90 | ZPHY C46 349 | E. Elsen <i>et al.</i> | (JADE Collab.) |
| ROSNER | 90 | PR D42 3732 | J.L. Rosner | |
| WAGNER | 90 | PRL 64 1095 | S.R. Wagner <i>et al.</i> | (Mark II Collab.) |
| ALBRECHT | 89C | PL B219 121 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 89G | PL B229 304 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 89J | PL B229 175 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 89L | PL B232 554 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ARTUSO | 89 | PRL 62 2233 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| AVERRILL | 89 | PR D39 123 | D.A. Averill <i>et al.</i> | (HRS Collab.) |
| VERY | 89B | PL B223 470 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| BEBEK | 89 | PRL 62 8 | C. Bebek <i>et al.</i> | (CLEO Collab.) |
| BORTOLETTO | 89 | PRL 62 2436 | D. Bortoletto <i>et al.</i> | (CLEO Collab.) |
| BORTOLETTO | 89B | PL 63 1667 | D. Bortoletto <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 88F | PL B209 119 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 88K | PL B215 424 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 87C | PL B185 218 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 87D | PL B199 451 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 87I | PL B192 245 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 87J | PL B197 452 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| AVERY | 87 | PL B183 429 | P. Avery <i>et al.</i> | (CLEO Collab.) |

| | | | | |
|----------|-----|-------------|----------------------------------|-----------------|
| BEAN | 87B | PRL 58 183 | A. Bean <i>et al.</i> | (CLEO Collab.) |
| BEBEK | 87 | PR D36 1289 | C. Bebek <i>et al.</i> | (CLEO Collab.) |
| ALAM | 86 | PR D34 3279 | M.S. Alam <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 86F | PL B182 95 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| PDG | 86 | PL 170B 1 | M. Aguilar-Benitez <i>et al.</i> | (CERN, CIT+) |
| CHEN | 85 | PR D31 2386 | A. Chen <i>et al.</i> | (CLEO Collab.) |
| HAAS | 85 | PRL 55 1248 | J. Haas <i>et al.</i> | (CLEO Collab.) |
| AVERY | 84 | PRL 53 1309 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| GILES | 84 | PR D30 2279 | R. Giles <i>et al.</i> | (CLEO Collab.) |
| BEHRENDS | 83 | PRL 50 881 | S. Behrends <i>et al.</i> | (CLEO Collab.) |
