

$\Xi(1690)$ $I(J^P) = \frac{1}{2}(?)$ Status: ***

AUBERT 08AK, in a study of $\Lambda_c^+ \rightarrow \Xi^- \pi^+ K^+$, finds some evidence that the $\Xi(1690)$ has $J^P = 1/2^-$.

IONISI 78 sees a threshold enhancement in both the neutral and negatively charged $\Sigma\bar{K}$ mass spectra in $K^- p \rightarrow (\Sigma\bar{K})K\pi$ at 4.2 GeV/c. The data from the $\Sigma\bar{K}$ channels alone cannot distinguish between a resonance and a large scattering length. Weaker evidence at the same mass is seen in the corresponding $\Lambda\bar{K}$ channels, and a coupled-channel analysis yields results consistent with a new Ξ .

BIAGI 81 sees an enhancement at 1700 MeV in the diffractively produced ΛK^- system. A peak is also observed in the $\Lambda\bar{K}^0$ mass spectrum at 1660 MeV that is consistent with a 1720 MeV resonance decaying to $\Sigma^0\bar{K}^0$, with the γ from the Σ^0 decay not detected.

BIAGI 87 provides further confirmation of this state in diffractive dissociation of Ξ^- into ΛK^- . The significance claimed is 6.7 standard deviations.

ADAMOVICH 98 sees a peak of 1400 ± 300 events in the $\Xi^- \pi^+$ spectrum produced by 345 GeV/c Σ^- -nucleus interactions.

 $\Xi(1690)$ MASSES**MIXED CHARGES**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
1690±10 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.	

 $\Xi(1690)^0$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1686±4	1400	ADAMOVICH 98	WA89	Σ^- nucleus, 345 GeV/c
1699±5	175	¹ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c
1684±5	183	² DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

 $\Xi(1690)^-$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1691.1± 1.9±2.0	104	BIAGI 87	SPEC	Ξ^- Be 116 GeV
1700 ± 10	150	³ BIAGI 81	SPEC	Ξ^- H 100, 135 GeV
1694 ± 6	45	⁴ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

 $\Xi(1690)$ WIDTHS**MIXED CHARGES**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
<30 OUR ESTIMATE	

$\Xi(1690)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10 ± 6	1400	ADAMOVICH 98	WA89	Σ^- nucleus, 345 GeV/c
44 ± 23	175	¹ DIONISI	78	HBC $K^- p$ 4.2 GeV/c
20 ± 4	183	² DIONISI	78	HBC $K^- p$ 4.2 GeV/c

 $\Xi(1690)^-$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 8	90	104	BIAGI	87	SPEC Ξ^- Be 116 GeV
47 ± 14		150	³ BIAGI	81	SPEC Ξ^- H 100, 135 GeV
26 ± 6		45	⁴ DIONISI	78	HBC $K^- p$ 4.2 GeV/c

 $\Xi(1690)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \Lambda \bar{K}$	seen
$\Gamma_2 \Sigma \bar{K}$	seen
$\Gamma_3 \Xi \pi$	seen
$\Gamma_4 \Xi^- \pi^+ \pi^0$	
$\Gamma_5 \Xi^- \pi^+ \pi^-$	possibly seen
$\Gamma_6 \Xi(1530) \pi$	

 $\Xi(1690)$ BRANCHING RATIOS **$\Gamma(\Lambda \bar{K})/\Gamma_{\text{total}}$**

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
seen	104	BIAGI	87	SPEC	Ξ^- Be 116 GeV

 Γ_1/Γ **$\Gamma(\Sigma \bar{K})/\Gamma(\Lambda \bar{K})$**

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.75 ± 0.39	75	ABE	02C	BELL	$e^+ e^- \approx \gamma(4S)$
2.7 ± 0.9		DIONISI	78	HBC	$0 K^- p$ 4.2 GeV/c
3.1 ± 1.4		DIONISI	78	HBC	$- K^- p$ 4.2 GeV/c

 Γ_2/Γ_1 **$\Gamma(\Xi \pi)/\Gamma(\Sigma \bar{K})$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.09	DIONISI	78	HBC	$0 K^- p$ 4.2 GeV/c

 Γ_3/Γ_2 **$\Gamma(\Xi \pi)/\Gamma_{\text{total}}$**

VALUE	DOCUMENT ID	TECN	COMMENT
seen	ADAMOVICH 98	WA89	Σ^- nucleus, 345 GeV/c

 Γ_3/Γ **$\Gamma(\Xi^- \pi^+ \pi^0)/\Gamma(\Sigma \bar{K})$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.04	DIONISI	78	HBC	$0 K^- p$ 4.2 GeV/c

 Γ_4/Γ_2

$\Gamma(\Xi^-\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_5/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
possibly seen	4	BIAGI	87	SPEC	Ξ^- Be 116 GeV
$\Gamma(\Xi^-\pi^+\pi^-)/\Gamma(\Sigma\bar{K})$			Γ_5/Γ_2		
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<0.03		DIONISI	78	HBC	$K^- p$ 4.2 GeV/c
$\Gamma(\Xi(1530)\pi)/\Gamma(\Sigma\bar{K})$			Γ_6/Γ_2		
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<0.06		DIONISI	78	HBC	$K^- p$ 4.2 GeV/c

$\Xi(1690)$ FOOTNOTES

¹ From a fit to the $\Sigma^+ K^-$ spectrum.

² From a coupled-channel analysis of the $\Sigma^+ K^-$ and $\Lambda\bar{K}^0$ spectra.

³ A fit to the inclusive spectrum from $\Xi^- N \rightarrow \Lambda K^- X$.

⁴ From a coupled-channel analysis of the $\Sigma^0 K^-$ and ΛK^- spectra.

$\Xi(1690)$ REFERENCES

AUBERT	08AK	PR D78 034008	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	02C	PL B524 33	K. Abe <i>et al.</i>	(KEK BELLE Collab.)
ADAMOVICH	98	EPJ C5 621	M.I. Adamovich <i>et al.</i>	(CERN WA89 Collab.)
BIAGI	87	ZPHY C34 15	S.F. Biagi <i>et al.</i>	(BRIS, CERN, GEVA+) I
BIAGI	81	ZPHY C9 305	S.F. Biagi <i>et al.</i>	(BRIS, CAVE, GEVA+) II
DIONISI	78	PL 80B 145	C. Dionisi <i>et al.</i>	(CERN, AMST, NIJM+) I