

**$\rho_3(1690)$**  $I^G(J^{PC}) = 1^+(3^{--})$  **$\rho_3(1690)$  MASS**VALUE (MeV)DOCUMENT ID**1688.8±2.1 OUR AVERAGE**

Includes data from the 5 datablocks that follow this one.

**2 $\pi$  MODE**VALUE (MeV)EVTSDOCUMENT IDTECNCHGCOMMENT

The data in this block is included in the average printed for a previous datablock.

**1686± 4 OUR AVERAGE**

1677±14		EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow 2\pi p$
1679±11	476	BALTAY	78B	HBC	0	15 $\pi^+ p \rightarrow \pi^+ \pi^- n$
1678±12	175	<sup>1</sup> ANTIPOV	77	CIBS	0	25 $\pi^- p \rightarrow p3\pi$
1690± 7	600	<sup>1</sup> ENGLER	74	DBC	0	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$
1693± 8		<sup>2</sup> GRAYER	74	ASPK	0	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1678±12		MATTHEWS	71C	DBC	0	7 $\pi^+ N$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1734±10		<sup>3</sup> CORDEN	79	OMEG		12–15 $\pi^- p \rightarrow n2\pi$
1692±12		<sup>2,4</sup> ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1737±23		ARMENISE	70	DBC	0	9 $\pi^+ N$
1650±35	122	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\pi$
1687±21		STUNTEBECK	70	HDBC	0	8 $\pi^- p$ , 5.4 $\pi^+ d$
1683±13		ARMENISE	68	DBC	0	5.1 $\pi^+ d$
1670±30		GOLDBERG	65	HBC	0	6 $\pi^+ d$ , 8 $\pi^- p$

<sup>1</sup> Mass errors enlarged by us to  $\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.<sup>2</sup> Uses same data as HYAMS 75.<sup>3</sup> From a phase shift solution containing a  $f_2'(1525)$  width two times larger than the  $K\bar{K}$  result.<sup>4</sup> From phase-shift analysis. Error takes account of spread of different phase-shift solutions. **$K\bar{K}$  AND  $K\bar{K}\pi$  MODES**VALUE (MeV)EVTSDOCUMENT IDTECNCHGCOMMENT

The data in this block is included in the average printed for a previous datablock.

**1696± 4 OUR AVERAGE**

1699± 5		ALPER	80	CNTR	0	62 $\pi^- p \rightarrow K^+ K^- n$
1698±12	6k	<sup>5,6</sup> MARTIN	78D	SPEC		10 $\pi p \rightarrow K_S^0 K^- p$
1692± 6		BLUM	75	ASPK	0	18.4 $\pi^- p \rightarrow nK^+ K^-$
1690±16		ADERHOLZ	69	HBC	+	8 $\pi^+ p \rightarrow K\bar{K}\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1694± 8		<sup>7</sup> COSTA...	80	OMEG		10 $\pi^- p \rightarrow K^+ K^- n$

<sup>5</sup> From a fit to  $J^P = 3^-$  partial wave.<sup>6</sup> Systematic error on mass scale subtracted.<sup>7</sup> They cannot distinguish between  $\rho_3(1690)$  and  $\omega_3(1670)$ .**( $4\pi$ ) $\pm$  MODE**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
The data in this block is included in the average printed for a previous datablock.					

**1686 $\pm$  5 OUR AVERAGE** Error includes scale factor of 1.1.

1694 $\pm$ 6		<sup>8</sup> EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p 4\pi$
1665 $\pm$ 15	177	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p 4\pi$
1670 $\pm$ 10		THOMPSON	74	HBC	+	13 $\pi^+ p$
1687 $\pm$ 20		CASON	73	HBC	—	8,18.5 $\pi^- p$
1685 $\pm$ 14		<sup>9</sup> CASON	73	HBC	—	8,18.5 $\pi^- p$
1680 $\pm$ 40	144	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N 4\pi$
1689 $\pm$ 20	102	<sup>9</sup> BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N 2\rho$
1705 $\pm$ 21		CASO	70	HBC	—	11.2 $\pi^- p \rightarrow n \rho 2\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1718 $\pm$ 10		<sup>10</sup> EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p 4\pi$
1673 $\pm$ 9		<sup>11</sup> EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p 4\pi$
1733 $\pm$ 9	66	<sup>9</sup> KLIGER	74	HBC	—	4.5 $\pi^- p \rightarrow p 4\pi$
1630 $\pm$ 15		HOLMES	72	HBC	+	10–12 $K^+ p$
1720 $\pm$ 15		BALTAY	68	HBC	+	7, 8.5 $\pi^+ p$

<sup>8</sup> From  $\rho^- \rho^0$  mode, not independent of the other two EVANGELISTA 81 entries.<sup>9</sup> From  $\rho^\pm \rho^0$  mode.<sup>10</sup> From  $a_2(1320)^- \pi^0$  mode, not independent of the other two EVANGELISTA 81 entries.<sup>11</sup> From  $a_2(1320)^0 \pi^-$  mode, not independent of the other two EVANGELISTA 81 entries. **$\omega\pi$  MODE**

<u>VALUE (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
The data in this block is included in the average printed for a previous datablock.					

**1681 $\pm$  7 OUR AVERAGE**

1670 $\pm$ 25		<sup>12</sup> ALDE	95	GAM2		38 $\pi^- p \rightarrow \omega \pi^0 n$
1690 $\pm$ 15		EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow \omega \pi p$
1666 $\pm$ 14		GESSAROLI	77	HBC		11 $\pi^- p \rightarrow \omega \pi p$
1686 $\pm$ 9		THOMPSON	74	HBC	+	13 $\pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1654 $\pm$ 24		BARNHAM	70	HBC	+	10 $K^+ p \rightarrow \omega \pi X$

<sup>12</sup> Supersedes ALDE 92C.

## $\eta\pi^+\pi^-$ MODE

(For difficulties with MMS experiments, see the  $a_2(1320)$  mini-review in the 1973 edition.)

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.				

### **$1682 \pm 12$ OUR AVERAGE**

$1685 \pm 10 \pm 20$	AMELIN	00	VES	$37 \pi^- p \rightarrow \eta\pi^+\pi^- n$
$1680 \pm 15$	FUKUI	88	SPEC 0	$8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1700 \pm 47$	<sup>13</sup> ANDERSON	69	MMS	$16 \pi^- p$ backward
$1632 \pm 15$	<sup>13,14</sup> FOCACCI	66	MMS	$7-12 \pi^- p \rightarrow p\text{MM}$
$1700 \pm 15$	<sup>13,14</sup> FOCACCI	66	MMS	$7-12 \pi^- p \rightarrow p\text{MM}$
$1748 \pm 15$	<sup>13,14</sup> FOCACCI	66	MMS	$7-12 \pi^- p \rightarrow p\text{MM}$


<sup>13</sup> Seen in 2.5–3 GeV/c  $\bar{p}p$ .  $2\pi^+ 2\pi^-$ , with 0, 1, 2  $\pi^+\pi^-$  pairs in  $\rho$  band not seen by OREN 74 (2.3 GeV/c  $\bar{p}p$ ) with more statistics. (Jan. 1976)

<sup>14</sup> Not seen by BOWEN 72.

## $\rho_3(1690)$ WIDTH

### **$2\pi$ , $K\bar{K}$ , AND $K\bar{K}\pi$ MODES**

VALUE (MeV)	DOCUMENT ID
<b><math>161 \pm 10</math> OUR AVERAGE</b>	Includes data from the 5 datablocks that follow this one. Error includes scale factor of 1.5. See the ideogram below.



$\rho_3(1690)$  width,  $2\pi$ ,  $K\bar{K}$ , and  $K\bar{K}\pi$  modes (MeV)

## 2π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					

**186±14 OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

220±29		DENNEY	83	LASS	10 $\pi^+ N$
246±37		EVANGELIS...	81	OMEG	— 12 $\pi^- p \rightarrow 2\pi p$
116±30	476	BALTAY	78B	HBC	0 15 $\pi^+ p \rightarrow$ $\pi^+ \pi^- n$
162±50	175	15 ANTIPOV	77	CIBS	0 25 $\pi^- p \rightarrow p 3\pi$
167±40	600	ENGLER	74	DBC	0 6 $\pi^+ n \rightarrow$ $\pi^+ \pi^- p$
200±18		16 GRAYER	74	ASPK	0 17 $\pi^- p \rightarrow$ $\pi^+ \pi^- n$
156±36		MATTHEWS	71C	DBC	0 7 $\pi^+ N$
171±65		ARMENISE	70	DBC	0 9 $\pi^+ d$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
322±35		17 CORDEN	79	OMEG	12–15 $\pi^- p \rightarrow$ $n 2\pi$
240±30	16,18 ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow$ $\pi^+ \pi^- n$
180±30	122	BARTSCH	70B	HBC	+ 8 $\pi^+ p \rightarrow N 2\pi$


$267^{+72}_{-46}$	STUNTEBECK 70	HDBC 0	$8 \pi^- p, 5.4 \pi^+ d$
$188 \pm 49$	ARMENISE 68	DBC 0	$5.1 \pi^+ d$
$180 \pm 40$	GOLDBERG 65	HBC 0	$6 \pi^+ d, 8 \pi^- p$

<sup>15</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>16</sup> Uses same data as HYAMS 75 and BECKER 79.

<sup>17</sup> From a phase shift solution containing a  $f_2'(1525)$  width two times larger than the  $K\bar{K}$  result.

<sup>18</sup> From phase-shift analysis. Error takes account of spread of different phase-shift solutions.



## $K\bar{K}$ AND $K\bar{K}\pi$ MODES

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

### 204±18 OUR AVERAGE

$199 \pm 40$	6000	19 MARTIN	78D SPEC	$10 \pi^- p \rightarrow K_S^0 K^- p$
$205 \pm 20$		BLUM	75 ASPK 0	$18.4 \pi^- p \rightarrow n K^+ K^-$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
$219 \pm 4$		ALPER	80 CNTR 0	$62 \pi^- p \rightarrow K^+ K^- n$
$186 \pm 11$		20 COSTA...	80 OMEG	$10 \pi^- p \rightarrow K^+ K^- n$
$112 \pm 60$		ADERHOLZ	69 HBC +	$8 \pi^+ p \rightarrow K\bar{K}\pi$

<sup>19</sup> From a fit to  $J^P = 3^-$  partial wave.

<sup>20</sup> They cannot distinguish between  $\rho_3(1690)$  and  $\omega_3(1670)$ .

**( $4\pi$ ) $\pm$  MODE**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
The data in this block is included in the average printed for a previous datablock.					

**129 $\pm$ 10 OUR AVERAGE**

123 $\pm$ 13		<sup>21</sup> EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p 4\pi$
105 $\pm$ 30	177	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p 4\pi$
169 $^{+70}_{-48}$		CASON	73	HBC	—	8,18.5 $\pi^- p$
135 $\pm$ 30	144	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N 4\pi$
160 $\pm$ 30	102	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N 2\rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
230 $\pm$ 28		<sup>22</sup> EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p 4\pi$
184 $\pm$ 33		<sup>23</sup> EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p 4\pi$
150	66	<sup>24</sup> KLIGER	74	HBC	—	4.5 $\pi^- p \rightarrow p 4\pi$
106 $\pm$ 25		THOMPSON	74	HBC	+	13 $\pi^+ p$
125 $^{+83}_{-35}$		<sup>24</sup> CASON	73	HBC	—	8,18.5 $\pi^- p$
130 $\pm$ 30		HOLMES	72	HBC	+	10–12 $K^+ p$
180 $\pm$ 30	90	<sup>24</sup> BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N a_2 \pi$
100 $\pm$ 35		BALTAY	68	HBC	+	7, 8.5 $\pi^+ p$

<sup>21</sup> From  $\rho^- \rho^0$  mode, not independent of the other two EVANGELISTA 81 entries.<sup>22</sup> From  $a_2(1320)^- \pi^0$  mode, not independent of the other two EVANGELISTA 81 entries.<sup>23</sup> From  $a_2(1320)^0 \pi^-$  mode, not independent of the other two EVANGELISTA 81 entries.<sup>24</sup> From  $\rho^\pm \rho^0$  mode. **$\omega\pi$  MODE**

<u>VALUE (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
The data in this block is included in the average printed for a previous datablock.					

**190 $\pm$ 40 OUR AVERAGE**

230 $\pm$ 65		<sup>25</sup> ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega \pi^0 n$
190 $\pm$ 65		EVANGELIS...	81	OMEG	—
160 $\pm$ 56		GESSAROLI	77	HBC	12 $\pi^- p \rightarrow \omega \pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
89 $\pm$ 25		THOMPSON	74	HBC	+
130 $^{+73}_{-43}$		BARNHAM	70	HBC	+
25 Supersedes ALDE 92C.					

 **$\eta\pi^+\pi^-$  MODE**(For difficulties with MMS experiments, see the  $a_2(1320)$  mini-review in the 1973 edition.)

<u>VALUE (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
The data in this block is included in the average printed for a previous datablock.					

**126 $\pm$ 40 OUR AVERAGE** Error includes scale factor of 1.8.

220 $\pm$ 30 $\pm$ 50		AMELIN	00	VES	37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
106 $\pm$ 27		FUKUI	88	SPEC 0	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

195	<sup>26</sup> ANDERSON	69	MMS	—	16 $\pi^- p$ backward
< 21	<sup>26,27</sup> FOCACCI	66	MMS	—	7–12 $\pi^- p \rightarrow p\text{MM}$
< 30	<sup>26,27</sup> FOCACCI	66	MMS	—	7–12 $\pi^- p \rightarrow p\text{MM}$
< 38	<sup>26,27</sup> FOCACCI	66	MMS	—	7–12 $\pi^- p \rightarrow p\text{MM}$

<sup>26</sup> Seen in 2.5–3 GeV/c  $\bar{p}p$ .  $2\pi^+ 2\pi^-$ , with 0, 1, 2  $\pi^+ \pi^-$  pairs in  $\rho^0$  band not seen by OREN 74 (2.3 GeV/c  $\bar{p}p$ ) with more statistics. (Jan. 1979)

<sup>27</sup> Not seen by BOWEN 72.

## $\rho_3(1690)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor
$\Gamma_1$ $4\pi$	(71.1 $\pm$ 1.9) %	
$\Gamma_2$ $\pi^\pm \pi^+ \pi^- \pi^0$	(67 $\pm$ 22) %	
$\Gamma_3$ $\omega \pi$	(16 $\pm$ 6) %	
$\Gamma_4$ $\pi \pi$	(23.6 $\pm$ 1.3) %	
$\Gamma_5$ $K\bar{K}\pi$	( 3.8 $\pm$ 1.2) %	
$\Gamma_6$ $K\bar{K}$	( 1.58 $\pm$ 0.26) %	1.2
$\Gamma_7$ $\eta \pi^+ \pi^-$	seen	
$\Gamma_8$ $\rho(770)\eta$	seen	
$\Gamma_9$ $\pi \pi \rho$	seen	
Excluding $2\rho$ and $a_2(1320)\pi$ .		
$\Gamma_{10}$ $a_2(1320)\pi$	seen	
$\Gamma_{11}$ $\rho \rho$	seen	
$\Gamma_{12}$ $\phi \pi$		
$\Gamma_{13}$ $\eta \pi$		
$\Gamma_{14}$ $\pi^\pm 2\pi^+ 2\pi^- \pi^0$		

## CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 10 measurements and one constraint to determine 4 parameters. The overall fit has a  $\chi^2 = 14.7$  for 7 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$$\begin{array}{c|ccc} & & & \\ x_4 & -77 & & \\ x_5 & -74 & 17 & \\ x_6 & -15 & 2 & 0 \\ \hline & x_1 & x_4 & x_5 \end{array}$$

**$\rho_3(1690)$  BRANCHING RATIOS** **$\Gamma(\pi\pi)/\Gamma_{\text{total}}$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	<b><math>\Gamma_4/\Gamma</math></b>
<b><math>0.236 \pm 0.013</math> OUR FIT</b>					
<b><math>0.243 \pm 0.013</math> OUR AVERAGE</b>					
$0.259^{+0.018}_{-0.019}$	BECKER	79	ASPK	0	$17 \pi^- p$ polarized
$0.23 \pm 0.02$	CORDEN	79	OMEG		$12-15 \pi^- p \rightarrow n 2\pi$
$0.22 \pm 0.04$	28 MATTHEWS	71C	HDBC	0	$7 \pi^+ n \rightarrow \pi^- p$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$0.245 \pm 0.006$	29 ESTABROOKS	75	RVUE		$17 \pi^- p \rightarrow \pi^+ \pi^- n$

28 One-pion-exchange model used in this estimation.

29 From phase-shift analysis of HYAMS 75 data.

 **$\Gamma(\pi\pi)/\Gamma(\pi^\pm \pi^+ \pi^- \pi^0)$** 


<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	<b><math>\Gamma_4/\Gamma_2</math></b>
<b><math>0.35 \pm 0.11</math></b>	CASON	73	HBC	—	$8, 18.5 \pi^- p$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.2	HOLMES	72	HBC	+	$10-12 K^+ p$
<0.12	BALLAM	71B	HBC	—	$16 \pi^- p$

 **$\Gamma(\pi\pi)/\Gamma(4\pi)$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	<b><math>\Gamma_4/\Gamma_1</math></b>
<b><math>0.332 \pm 0.026</math> OUR FIT</b>	Error includes scale factor of 1.1.				
<b><math>0.30 \pm 0.10</math></b>	BALTAY	78B	HBC	0	$15 \pi^+ p \rightarrow p 4\pi$

 **$\Gamma(K\bar{K})/\Gamma(\pi\pi)$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	<b><math>\Gamma_6/\Gamma_4</math></b>
<b><math>0.067 \pm 0.011</math> OUR FIT</b>	Error includes scale factor of 1.2.				
<b><math>0.118^{+0.040}_{-0.032}</math> OUR AVERAGE</b>	Error includes scale factor of 1.7. See the ideogram below.				
$0.191^{+0.040}_{-0.037}$	GORLICH	80	ASPK	0	$17, 18 \pi^- p$ polarized
$0.08 \pm 0.03$	BARTSCH	70B	HBC	+	$8 \pi^+ p$
$0.08^{+0.08}_{-0.03}$	CRENNELL	68B	HBC		$6.0 \pi^- p$



### $\Gamma(K\bar{K}\pi)/\Gamma(\pi\pi)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.16±0.05 OUR FIT</b>				
<b>0.16±0.05</b>	30 BARTSCH	70B HBC	+	$8\pi^+ p$
30 Increased by us to correspond to $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$ .				

### $[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$      $(\Gamma_9 + \Gamma_{10} + \Gamma_{11})/\Gamma_2$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.94±0.09 OUR AVERAGE</b>				
0.96±0.21	BALTAY	78B HBC	+	$15\pi^+ p \rightarrow p4\pi$
0.88±0.15	BALLAM	71B HBC	-	$16\pi^- p$
1 ± 0.15	BARTSCH	70B HBC	+	$8\pi^+ p$
consistent with 1	CASO	68 HBC	-	$11\pi^- p$

### $\Gamma(\rho\rho)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

### $\Gamma_{11}/\Gamma_2$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
0.12±0.11		BALTAY	78B HBC	+	$15\pi^+ p \rightarrow p4\pi$
0.56	66	KLIGER	74 HBC	-	$4.5\pi^- p \rightarrow p4\pi$
0.13±0.09		31 THOMPSON	74 HBC	+	$13\pi^+ p$
0.7 ± 0.15		BARTSCH	70B HBC	+	$8\pi^+ p$

<sup>31</sup>  $\rho\rho$  and  $a_2(1320)\pi$  modes are indistinguishable.

$\Gamma(\rho\rho)/[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]$   $\Gamma_{11}/(\Gamma_9+\Gamma_{10}+\Gamma_{11})$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.48±0.16 CASO 68 HBC – 11  $\pi^- p$

 $\Gamma(a_2(1320)\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$   $\Gamma_{10}/\Gamma_2$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.66±0.08	BALTAJ	78B	HBC	+	15 $\pi^+ p \rightarrow p 4\pi$
0.36±0.14	<sup>32</sup> THOMPSON	74	HBC	+	13 $\pi^+ p$
not seen	CASON	73	HBC	–	8,18.5 $\pi^- p$
0.6 ±0.15	BARTSCH	70B	HBC	+	8 $\pi^+ p$
0.6	BALTAJ	68	HBC	+	7,8.5 $\pi^+ p$

<sup>32</sup>  $\rho\rho$  and  $a_2(1320)\pi$  modes are indistinguishable.

 $\Gamma(\omega\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$   $\Gamma_3/\Gamma_2$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	------------	----------------

**0.23±0.05 OUR AVERAGE** Error includes scale factor of 1.2.

0.33±0.07	THOMPSON	74	HBC	+	13 $\pi^+ p$
0.12±0.07	BALLAM	71B	HBC	–	16 $\pi^- p$
0.25±0.10	BALTAJ	68	HBC	+	7,8.5 $\pi^+ p$
0.25±0.10	JOHNSTON	68	HBC	–	7.0 $\pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.11	95	BALTAJ	78B	HBC	+	15 $\pi^+ p \rightarrow p 4\pi$
<0.09		KLIGER	74	HBC	–	4.5 $\pi^- p \rightarrow p 4\pi$

 $\Gamma(\phi\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$   $\Gamma_{12}/\Gamma_2$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.11 BALTAJ 68 HBC + 7,8.5  $\pi^+ p$

 $\Gamma(\pi^\pm 2\pi^+ 2\pi^- \pi^0)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$   $\Gamma_{14}/\Gamma_2$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.15 BALTAJ 68 HBC + 7,8.5  $\pi^+ p$

 $\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$   $\Gamma_{13}/\Gamma_2$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.02 THOMPSON 74 HBC + 13  $\pi^+ p$

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<b>0.0158±0.0026 OUR FIT</b>	Error includes scale factor of 1.2.			
<b>0.0130±0.0024 OUR AVERAGE</b>				

0.013 ± 0.003	COSTA...	80	OMEG 0	10 $\pi^- p \rightarrow K^+ K^- n$
0.013 ± 0.004	33 MARTIN	78B	SPEC	10 $\pi^- p \rightarrow K_S^0 K^- p$

<sup>33</sup> From  $(\Gamma_4 \Gamma_6)^{1/2} = 0.056 \pm 0.034$  assuming  $B(\rho_3(1690) \rightarrow \pi \pi) = 0.24$ .

 $\Gamma(\omega\pi)/[\Gamma(\omega\pi) + \Gamma(\rho\rho)]$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				

0.22±0.08 CASON 73 HBC — 8,18.5  $\pi^- p$

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	FUKUI	88	SPEC 8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

 $\Gamma(a_2(1320)\pi)/\Gamma(\rho(770)\eta)$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.5±2.0</b>	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

 $\Gamma_3/(\Gamma_3 + \Gamma_{11})$  $\Gamma_7/\Gamma$  $\Gamma_{10}/\Gamma_8$  **$\rho_3(1690)$  REFERENCES**

AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
COSTA...	80	NP B175 402	G. Costa de Beauregard <i>et al.</i>	(BARI, BONN+)
GORLICH	80	NP B174 16	L. Gorlich <i>et al.</i>	(CRAC, MPIM, CERN+)
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)
MARTIN	78B	NP B140 158	A.D. Martin <i>et al.</i>	(DURH, GEVA)
MARTIN	78D	PL 74B 417	A.D. Martin <i>et al.</i>	(DURH, GEVA)
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+)
BLUM	75	PL 57B 403	W. Blum <i>et al.</i>	(CERN, MPIM) JP
ESTABROOKS	75	NP B95 322	P.G. Estabrooks, A.D. Martin	(DURH)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)
KLIGER	74	SJNP 19 428	G.K. Kliger <i>et al.</i>	(ITEP)
		Translated from YAF 19 839.		
OREN	74	NP B71 189	Y. Oren <i>et al.</i>	(ANL, OXF)
THOMPSON	74	NP B69 220	G. Thompson <i>et al.</i>	(PURD)
CASON	73	PR D7 1971	N.M. Cason <i>et al.</i>	(NDAM)
BOWEN	72	PRL 29 890	D.R. Bowen <i>et al.</i>	(NEAS, STON)
HOLMES	72	PR D6 3336	R. Holmes <i>et al.</i>	(ROCH)
BALLAM	71B	PR D3 2606	J. Ballam <i>et al.</i>	(SLAC)
MATTHEWS	71C	NP B33 1	J.A.J. Matthews <i>et al.</i>	(TNTO, WISC) JP

ARMENISE	70	LNC 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)
BARNHAM	70	PRL 24 1083	K.W.J. Barnham <i>et al.</i>	(BIRM)
BARTSCH	70B	NP B22 109	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN)
CASO	70	LNC 3 707	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)
ANDERSON	69	PRL 22 1390	E.W. Anderson <i>et al.</i>	(BNL, CMU)
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+) I
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE) I
CASO	68	NC 54A 983	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)
CRENNELL	68B	PL 28B 136	D.J. Crennell <i>et al.</i>	(BNL)
JOHNSTON	68	PRL 20 1414	T.F. Johnston <i>et al.</i>	(TNTO, WISC) IJP
FOCACCI	66	PRL 17 890	M.N. Focacci <i>et al.</i>	(CERN)
GOLDBERG	65	PL 17 354	M. Goldberg <i>et al.</i>	(CERN, EPOL, ORSAY+)