

$\Sigma(1670)$ Bumps $I(J^P) = 1(?)$

OMITTED FROM SUMMARY TABLE

Formation experiments are listed separately in the preceding entry.

Probably there are two states at the same mass with the same quantum numbers, one decaying to $\Sigma\pi$ and $\Lambda\pi$, the other to $\Lambda(1405)\pi$.
See the note in front of the preceding entry.

 **$\Sigma(1670)$ MASS
(PRODUCTION EXPERIMENTS)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
≈ 1670 OUR ESTIMATE					
1670 \pm 4		1 CARROLL	76	DPWA	Isospin-1 total σ
1675 \pm 10		2 HEPP	76	DBC	$K^- N$ 1.6–1.75 GeV/c
1665 \pm 1		APSELL	74	HBC	$K^- p$ 2.87 GeV/c
1688 \pm 2 or 1683 \pm 5	1.2k	BERTHON	74	HBC	Quasi-2-body σ
1670 \pm 6		AGUILAR-...	70B	HBC	$K^- p \rightarrow \Sigma\pi\pi$ 4 GeV
1668 \pm 10		AGUILAR-...	70B	HBC	$K^- p \rightarrow \Sigma 3\pi$ 4 GeV
1660 \pm 10		ALVAREZ	63	HBC	$K^- p$ 1.51 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1668 \pm 10	150	3 FERRERSORIA81	OMEG	–	$\pi^- p$ 9,12 GeV/c
1655 to 1677		TIMMERMANS76	HBC	+	$K^- p$ 4.2 GeV/c
1665 \pm 5		BUGG	68	CNTR	$K^- p, d$ total σ
1661 \pm 9	70	PRIMER	68	HBC	See BARNES 69E
1685		ALEXANDER	62C	HBC	$\pi^- p$ 2–2.2 GeV/c

 **$\Sigma(1670)$ WIDTH
(PRODUCTION EXPERIMENTS)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
67.0 \pm 2.4					
		APSELL	74	HBC	$K^- p$ 2.87 GeV/c
110 \pm 12		AGUILAR-...	70B	HBC	$K^- p \rightarrow \Sigma\pi\pi$ 4 GeV
135 \pm 40 -30		AGUILAR-...	70B	HBC	$K^- p \rightarrow \Sigma 3\pi$ 4 GeV
40 \pm 10		ALVAREZ	63	HBC	+
• • • We do not use the following data for averages, fits, limits, etc. • • •					
90 \pm 20	150	3 FERRERSORIA81	OMEG	–	$\pi^- p$ 9,12 GeV/c
52		1 CARROLL	76	DPWA	Isospin-1 total σ
48 to 63		TIMMERMANS76	HBC	+	$K^- p$ 4.2 GeV/c
30 \pm 15		BUGG	68	CNTR	
60 \pm 20	70	PRIMER	68	HBC	+
45		ALEXANDER	62C	HBC	$\pi^- p$ 2–2.2 GeV/c

$\Sigma(1670)$ DECAY MODES (PRODUCTION EXPERIMENTS)

Mode	
Γ_1	$N\bar{K}$
Γ_2	$\Lambda\pi$
Γ_3	$\Sigma\pi$
Γ_4	$\Lambda\pi\pi$
Γ_5	$\Sigma\pi\pi$
Γ_6	$\Sigma(1385)\pi$
Γ_7	$\Lambda(1405)\pi$

$\Sigma(1670)$ BRANCHING RATIOS (PRODUCTION EXPERIMENTS)

$\Gamma(N\bar{K})/\Gamma(\Sigma\pi)$		Γ_1/Γ_3			
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<0.03		TIMMERMANS76	HBC	+	$K^- p$ 4.2 GeV/c
<0.10		BERTHON 74	HBC	0	Quasi-2-body σ
<0.2		AGUILAR-...	70B	HBC	
<0.26		BARNES	69E	HBC	+
0.025		BUGG	68	CNTR	0 Assuming $J = 3/2$
<0.24	0	PRIMER	68	HBC	+
<0.6		LONDON	66	HBC	+
<0.19	0	ALVAREZ	63	HBC	+
$\geq 0.5 \pm 0.25$		SMITH	63	HBC	-0
$\Gamma(\Lambda\pi)/\Gamma(\Sigma\pi)$		Γ_2/Γ_3			
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.76 ± 0.09		ESTES	74	HBC	0 $K^- p$ 2.1,2.6 GeV/c
0.45 ± 0.15		BARNES	69E	HBC	+
0.15 ± 0.07		HUWE	69	HBC	+
0.11 ± 0.06	33	BUTTON-...	68	HBC	+
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$\leq 0.45 \pm 0.07$		TIMMERMANS76	HBC	+	$K^- p$ 4.2 GeV/c
0.55 ± 0.11		BERTHON 74	HBC	0	Quasi-2-body σ
0	0	PRIMER	68	HBC	+
<0.6		LONDON	66	HBC	+
1.2	130	ALVAREZ	63	HBC	+
1.2		SMITH	63	HBC	-0

$\Gamma(\Lambda\pi\pi)/\Gamma(\Sigma\pi)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_4/Γ_3
<0.6		LONDON	66	HBC	+	$K^- p$ 2.25 GeV/c
0.56	90	ALVAREZ	63	HBC	+	$K^- p$ 1.15 GeV/c
0.17		SMITH	63	HBC	-0	

 $\Gamma(\Sigma\pi\pi)/\Gamma(\Sigma\pi)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_5/Γ_3
largest at small angles		ESTES	74	HBC	0	$K^- p$ 2.1,2.6 GeV/c

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.2	² HEPP	76	DBC	-	$K^- N$ 1.6–1.75 GeV/c	
0.56	180	ALVAREZ	63	HBC	+	$K^- p$ 1.15 GeV/c

 $\Gamma(\Lambda(1405)\pi)/\Gamma(\Sigma\pi)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_7/Γ_3
1.8 ± 0.3 to 0.02 ± 0.07		3,4 TIMMERMANS76	HBC	+	$K^- p$ 4.2 GeV/c	
largest at small angles		ESTES	74	HBC	±	$K^- p$ 2.1,2.6 GeV/c
3.0 ± 1.6	50	LONDON	66	HBC	+	$K^- p$ 2.25 GeV/c

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.58 ± 0.20	17	PRIMER	68	HBC	+	See BARNES 69E
-------------	----	--------	----	-----	---	----------------

 $\Gamma(\Sigma\pi)/\Gamma(\Sigma\pi\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_3/Γ_5
varies with prod. angle					
5 APSELL	74	HBC	+	$K^- p$ 2.87 GeV/c	
1.39 ± 0.16	BERTHON	74	HBC	0	Quasi-2-body σ
2.5 to 0.24	4 EBERHARD	69	HBC		$K^- p$ 2.6 GeV/c
<0.4	BIRMINGHAM	66	HBC	+	$K^- p$ 3.5 GeV/c
0.30 ± 0.15	LONDON	66	HBC	+	$K^- p$ 2.25 GeV/c

 $\Gamma(\Lambda(1405)\pi)/\Gamma(\Sigma\pi\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_7/Γ_5
0.97 ± 0.08	TIMMERMANS76	HBC		$K^- p$ 4.2 GeV/c	
1.00 ± 0.02	APSELL	74	HBC		$K^- p$ 2.87 GeV/c
0.90 ^{+0.10} _{-0.16}	EBERHARD	65	HBC	+	$K^- p$ 2.45 GeV/c

 $\Gamma(\Lambda(1405)\pi)/\Gamma(\Sigma(1385)\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_7/Γ_6
<0.8	EBERHARD	65	HBC	+	$K^- p$ 2.45 GeV/c

$\Gamma(\Lambda\pi\pi)/\Gamma(\Sigma\pi\pi)$

<u>VALUE</u>
0.35 ± 0.2

$\Gamma(\Lambda\pi)/\Gamma(\Sigma\pi\pi)$

<u>VALUE</u>
<0.2

$\Gamma(\Lambda\pi)/[\Gamma(\Lambda\pi) + \Gamma(\Sigma\pi)]$

<u>VALUE</u>
<0.6

$\Gamma(\Sigma(1385)\pi)/\Gamma(\Sigma\pi)$

<u>VALUE</u>
$\leq 0.21 \pm 0.05$

Γ_4/Γ_5

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
BIRMINGHAM 66	HBC	+	$K^- p$ 3.5 GeV/c

Γ_2/Γ_5

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
BIRMINGHAM 66	HBC	+	$K^- p$ 3.5 GeV/c

$\Gamma_2/(\Gamma_2+\Gamma_3)$

<u>DOCUMENT ID</u>	<u>TECN</u>
AGUILAR-...	70B

Γ_6/Γ_3

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
TIMMERMANS76	HBC	$K^- p$ 4.2 GeV/c

$\Sigma(1670)$ QUANTUM NUMBERS (PRODUCTION EXPERIMENTS)

<u>VALUE</u>	<u>EVTS</u>
$J^P = 3/2^-$	400
$J^P = 3/2^-$	
$J^P = 3/2^+$	

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
BUTTON-...	68	\pm	$\Sigma^0\pi$
EBERHARD	67	+	$\Lambda(1405)\pi$
LEVEQUE	65		$\Lambda(1405)\pi$

$\Sigma(1670)$ FOOTNOTES

¹ Total cross-section bump with $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}} = 0.23$.

² Enhancements in $\Sigma\pi$ and $\Sigma\pi\pi$ cross sections.

³ Backward production in the $\Lambda\pi^-K^+$ final state.

⁴ Depending on production angle.

⁵ APSELL 74, ESTES 74, and TIMMERMANS 76 find strong branching ratio dependence on production angle, as in earlier production experiments.

$\Sigma(1670)$ REFERENCES (PRODUCTION EXPERIMENTS)

FERRERSORIA 81	NP B178 373	A. Ferrer Soria <i>et al.</i>	(CERN, CDEF, EPOL+)
CARROLL 76	PRL 37 806	A.S. Carroll <i>et al.</i>	(BNL) I
HEPP 76	NP B115 82	V. Hepp <i>et al.</i>	(CERN, HEID, MPIM) I
TIMMERMANS 76	NP B112 77	J.J.M. Timmermans <i>et al.</i>	(NIJM, CERN+) JP
APSELL 74	PR D10 1419	S.P. Apsell <i>et al.</i>	(BRAN, UMD, SYRA+) I
BERTHON 74	NC 21A 146	A. Berthon <i>et al.</i>	(CDEF, RHEL, SACL+)
ESTES 74	Thesis LBL-3827	R.D. Estes	(LBL)
AGUILAR-...	PRL 25 58	M. Aguilar-Benitez <i>et al.</i>	(BNL, SYRA)
BARNES 69E	BNL 13823	V.E. Barnes <i>et al.</i>	(BNL, SYRA)
EBERHARD 69	PRL 22 200	P.H. Eberhard <i>et al.</i>	(LRL)
HUWE 69	PR 181 1824	D.O. Huwe	(LRL)
BUGG 68	PR 168 1466	D.V. Bugg <i>et al.</i>	(RHEL, BIRM, CAVE) I
BUTTON-...	PRL 21 1123	J. Button-Shafer	(MASA, LRL) JP
PRIMER 68	PRL 20 610	M. Primer <i>et al.</i>	(SYRA, BNL)
EBERHARD 67	PR 163 1446	P. Eberhard <i>et al.</i>	(LRL, ILL) IJP

BIRMINGHAM	66	PR 152 1148	M. Haque <i>et al.</i>	(BIRM, GLAS, LOIC, OXF+)
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IJ
EBERHARD	65	PRL 14 466	P.H. Eberhard <i>et al.</i>	(LRL, ILL) I
LEVEQUE	65	PL 18 69	A. Leveque <i>et al.</i>	(SACL, EPOL, GLAS+) JP
ALVAREZ	63	PRL 10 184	L.W. Alvarez <i>et al.</i>	(LRL) I
SMITH	63	Athens Conf. 67	G.A. Smith	(LRL)
ALEXANDER	62C	CERN Conf. 320	G. Alexander <i>et al.</i>	(LRL) I
