

## Extra Dimensions

For explanation of terms used and discussion of significant model dependence of following limits, see the “Extra Dimensions” review. Footnotes describe originally quoted limit.  $n$  indicates the number of extra dimensions.

Limits not encoded here are summarized in the “Extra Dimensions” review.

A REVIEW GOES HERE – Check our WWW List of Reviews

---

### Limits on $R$ from Deviations in Gravitational Force Law

This section includes limits on the size of extra dimensions from deviations in the Newtonian ( $1/r^2$ ) gravitational force law at short distances. Deviations are parametrized by a gravitational potential of the form  $V = -(G m m'/r) [1 + \alpha \exp(-r/R)]$ . For  $\delta$  toroidal extra dimensions of equal size,  $\alpha = 8\delta/3$ . Quoted bounds are for  $\delta = 2$  unless otherwise noted.

<i>VALUE</i> ( $\mu\text{m}$ )	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>COMMENT</i>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
95	1	MASUDA 09	Torsion pendulum
	2	GERACI 08	Microcantilever
	3	TRENKEL 08	Newton's constant
	4	DECCA 07A	Torsion oscillator
< 30	95	5 KAPNER 07	Torsion pendulum
< 47	95	6 TU 07	Torsion pendulum
		7 SMULLIN 05	Microcantilever
< 130	95	8 HOYLE 04	Torsion pendulum
		9 CHIAVERINI 03	Microcantilever
≤ 200	95	10 LONG 03	Microcantilever
< 190	95	11 HOYLE 01	Torsion pendulum
		12 HOSKINS 85	Torsion pendulum

<sup>1</sup> MASUDA 09 obtain improved constraints on non-Newtonian forces with strengths  $10^9 \lesssim |\alpha| \lesssim 10^{11}$  and length scales  $R = 1.0\text{--}2.9 \mu\text{m}$ . See their Fig. 3. This bound does not place limits on the size of extra flat dimensions.

<sup>2</sup> GERACI 08 obtain improved constraints on non-Newtonian forces with strengths  $|\alpha| > 14,000$  and length scales  $R = 5\text{--}15 \mu\text{m}$ . See their Fig. 9. This bound does not place limits on the size of extra flat dimensions.

<sup>3</sup> TRENKEL 08 uses two independent measurements of Newton's constant  $G$  to constrain new forces with strength  $|\alpha| \simeq 10^{-4}$  and length scales  $R = 0.02\text{--}1 \text{ m}$ . See their Fig. 1. This bound does not place limits on the size of extra flat dimensions.

<sup>4</sup> DECCA 07A search for new forces and obtain bounds in the region with strengths  $|\alpha| \simeq 10^{13}\text{--}10^{18}$  and length scales  $R = 20\text{--}86 \text{ nm}$ . See their Fig. 6. This bound does not place limits on the size of extra flat dimensions.

<sup>5</sup> KAPNER 07 search for new forces, probing a range of  $\alpha \simeq 10^{-3}\text{--}10^5$  and length scales  $R \simeq 10\text{--}1000 \mu\text{m}$ . For  $\delta = 1$  the bound on  $R$  is  $44 \mu\text{m}$ . For  $\delta = 2$ , the bound is expressed in terms of  $M_*$ , here translated to a bound on the radius. See their Fig. 6 for details on the bound.

<sup>6</sup> TU 07 search for new forces probing a range of  $|\alpha| \simeq 10^{-1}\text{--}10^5$  and length scales  $R \simeq 20\text{--}1000 \mu\text{m}$ . For  $\delta = 1$  the bound on  $R$  is  $53 \mu\text{m}$ . See their Fig. 3 for details on the bound.

- <sup>7</sup> SMULLIN 05 search for new forces, and obtain bounds in the region with strengths  $\alpha \simeq 10^3\text{--}10^8$  and length scales  $R = 6\text{--}20\ \mu\text{m}$ . See their Figs. 1 and 16 for details on the bound. This work does not place limits on the size of extra flat dimensions.
- <sup>8</sup> HOYLE 04 search for new forces, probing  $\alpha$  down to  $10^{-2}$  and distances down to  $10\ \mu\text{m}$ . Quoted bound on  $R$  is for  $\delta = 2$ . For  $\delta = 1$ , bound goes to  $160\ \mu\text{m}$ . See their Fig. 34 for details on the bound.
- <sup>9</sup> CHIAVERINI 03 search for new forces, probing  $\alpha$  above  $10^4$  and  $\lambda$  down to  $3\ \mu\text{m}$ , finding no signal. See their Fig. 4 for details on the bound. This bound does not place limits on the size of extra flat dimensions.
- <sup>10</sup> LONG 03 search for new forces, probing  $\alpha$  down to 3, and distances down to about  $10\ \mu\text{m}$ . See their Fig. 4 for details on the bound.
- <sup>11</sup> HOYLE 01 search for new forces, probing  $\alpha$  down to  $10^{-2}$  and distances down to  $20\ \mu\text{m}$ . See their Fig. 4 for details on the bound. The quoted bound is for  $\alpha \geq 3$ .
- <sup>12</sup> HOSKINS 85 search for new forces, probing distances down to 4 mm. See their Fig. 13 for details on the bound. This bound does not place limits on the size of extra flat dimensions.

## Limits on $R$ from On-Shell Production of Gravitons: $\delta = 2$

This section includes limits on on-shell production of gravitons in collider and astrophysical processes. Bounds quoted are on  $R$ , the assumed common radius of the flat extra dimensions, for  $\delta = 2$  extra dimensions. Studies often quote bounds in terms of derived parameter; experiments are actually sensitive to the masses of the KK gravitons:  $m_{\vec{n}} = |\vec{n}|/R$ . See the Review on “Extra Dimensions” for details. Bounds are given in  $\mu\text{m}$  for  $\delta=2$ .

VALUE ( $\mu\text{m}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
< 245	95	13 AALTONEN	08AC CDF	$p\bar{p} \rightarrow \gamma G, jG$
< 615	95	14 ABAZOV	08S D0	$p\bar{p} \rightarrow \gamma G$
< 0.916	95	15 DAS	08	Supernova cooling
< 350	95	16 ABULENCIA,A	06 CDF	$p\bar{p} \rightarrow jG$
< 270	95	17 ABDALLAH	05B DLPFH	$e^+ e^- \rightarrow \gamma G$
< 210	95	18 ACHARD	04E L3	$e^+ e^- \rightarrow \gamma G$
< 480	95	19 ACOSTA	04C CDF	$\bar{p}p \rightarrow jG$
< 0.00038	95	20 CASSE	04	Neutron star $\gamma$ sources
< 610	95	21 ABAZOV	03 D0	$\bar{p}p \rightarrow jG$
< 0.96	95	22 HANNESTAD	03	Supernova cooling
< 0.096	95	23 HANNESTAD	03	Diffuse $\gamma$ background
< 0.051	95	24 HANNESTAD	03	Neutron star $\gamma$ sources
< 0.00016	95	25 HANNESTAD	03	Neutron star heating
< 300	95	26 HEISTER	03C ALEP	$e^+ e^- \rightarrow \gamma G$
		27 FAIRBAIRN	01	Cosmology
< 0.66	95	28 HANHART	01	Supernova cooling
		29 CASSISI	00	Red giants
<1300	95	30 ACCIARRI	99S L3	$e^+ e^- \rightarrow ZG$

## Limits on $R$ from On-Shell Production of Gravitons: $\delta \geq 3$

This section includes limits similar to those in the previous section, but for  $\delta = 3$  extra dimensions. Bounds are given in nm for  $\delta = 3$ . Entries are also shown for papers examining models with  $\delta > 3$ .

<i>VALUE</i> (nm)	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
< 2.8	95	13 AALTONEN	08AC CDF	$p\bar{p} \rightarrow \gamma G, jG$
< 4.56	95	14 ABAZOV	08S D0	$p\bar{p} \rightarrow \gamma G$
< 2.09	95	15 DAS	08	Supernova cooling
< 3.6	95	16 ABULENCIA,A	06 CDF	$p\bar{p} \rightarrow jG$
< 3.5	95	17 ABDALLAH	05B DLPH	$e^+ e^- \rightarrow \gamma G$
< 2.9	95	18 ACHARD	04E L3	$e^+ e^- \rightarrow \gamma G$
	95	19 ACOSTA	04C CDF	$\bar{p}p \rightarrow jG$
< 0.0042	95	20 CASSE	04	Neutron star $\gamma$ sources
< 6.1	95	21 ABAZOV	03 D0	$\bar{p}p \rightarrow jG$
< 1.14	95	22 HANNESTAD	03	Supernova cooling
< 0.025	95	23 HANNESTAD	03	Diffuse $\gamma$ background
< 0.11	95	24 HANNESTAD	03	Neutron star $\gamma$ sources
< 0.0026	95	25 HANNESTAD	03	Neutron star heating
< 3.9	95	26 HEISTER	03C ALEP	$e^+ e^- \rightarrow \gamma G$
	27 FAIRBAIRN	01		Cosmology
< 0.8	95	28 HANHART	01	Supernova cooling
	29 CASSISI	00		Red giants
<18	95	30 ACCIARRI	99S L3	$e^+ e^- \rightarrow ZG$

<sup>13</sup> AALTONEN 08AC search for  $p\bar{p} \rightarrow \gamma G$  and  $p\bar{p} \rightarrow jG$  at  $\sqrt{s} = 1.96$  TeV with  $2.0 \text{ fb}^{-1}$  and  $1.1 \text{ fb}^{-1}$  respectively, in order to place bounds on the fundamental scale and size of the extra dimensions. See their Table III for limits on all  $\delta \leq 6$ .

<sup>14</sup> ABAZOV 08S search for  $p\bar{p} \rightarrow \gamma G$ , using  $1 \text{ fb}^{-1}$  of data at  $\sqrt{s} = 1.96$  TeV to place bounds on  $M_D$  for two to eight extra dimensions, from which these bounds on  $R$  are derived. See their paper for intermediate values of  $\delta$ .

<sup>15</sup> DAS 08 obtain a limit on  $R$  from Kaluza-Klein graviton cooling of SN1987A due to plasmon-plasmon annihilation.

<sup>16</sup> ABULENCIA,A 06 search for  $p\bar{p} \rightarrow jG$  using  $368 \text{ pb}^{-1}$  of data at  $\sqrt{s} = 1.96$  TeV. See their Table II for bounds for all  $\delta \leq 6$ .

<sup>17</sup> ABDALLAH 05B search for  $e^+ e^- \rightarrow \gamma G$  at  $\sqrt{s} = 180\text{--}209$  GeV to place bounds on the size of extra dimensions and the fundamental scale. Limits for all  $\delta \leq 6$  are given in their Table 6. These limits supersede those in ABREU 00Z.

<sup>18</sup> ACHARD 04E search for  $e^+ e^- \rightarrow \gamma G$  at  $\sqrt{s} = 189\text{--}209$  GeV to place bounds on the size of extra dimensions and the fundamental scale. See their Table 8 for limits with  $\delta \leq 8$ . These limits supersede those in ACCIARRI 99R.

<sup>19</sup> ACOSTA 04C search for  $\bar{p}p \rightarrow jG$  at  $\sqrt{s} = 1.8$  TeV to place bounds on the size of extra dimensions and the fundamental scale. See their paper for bounds on  $\delta = 4, 6$ .

<sup>20</sup> CASSE 04 obtain a limit on  $R$  from the gamma-ray emission of point  $\gamma$  sources that arises from the photon decay of gravitons around newly born neutron stars, applying the technique of HANNESTAD 03 to neutron stars in the galactic bulge. Limits for all  $\delta \leq 7$  are given in their Table I.

<sup>21</sup> ABAZOV 03 search for  $p\bar{p} \rightarrow jG$  at  $\sqrt{s}=1.8$  TeV to place bounds on  $M_D$  for 2 to 7 extra dimensions, from which these bounds on  $R$  are derived. See their paper for bounds on intermediate values of  $\delta$ . We quote results without the approximate NLO scaling introduced in the paper.

<sup>22</sup> HANNESTAD 03 obtain a limit on  $R$  from graviton cooling of supernova SN1987a. Limits for all  $\delta \leq 7$  are given in their Tables V and VI.

- 23 HANNESTAD 03 obtain a limit on  $R$  from gravitons emitted in supernovae and which subsequently decay, contaminating the diffuse cosmic  $\gamma$  background. Limits for all  $\delta \leq 7$  are given in their Tables V and VI. These limits supersede those in HANNESTAD 02.
- 24 HANNESTAD 03 obtain a limit on  $R$  from gravitons emitted in two recent supernovae and which subsequently decay, creating point  $\gamma$  sources. Limits for all  $\delta \leq 7$  are given in their Tables V and VI. These limits are corrected in the published erratum.
- 25 HANNESTAD 03 obtain a limit on  $R$  from the heating of old neutron stars by the surrounding cloud of trapped KK gravitons. Limits for all  $\delta \leq 7$  are given in their Tables V and VI. These limits supersede those in HANNESTAD 02.
- 26 HEISTER 03C use the process  $e^+ e^- \rightarrow \gamma G$  at  $\sqrt{s} = 189\text{--}209$  GeV to place bounds on the size of extra dimensions and the scale of gravity. See their Table 4 for limits with  $\delta \leq 6$  for derived limits on  $M_D$ .
- 27 FAIRBAIRN 01 obtains bounds on  $R$  from over production of KK gravitons in the early universe. Bounds are quoted in paper in terms of fundamental scale of gravity. Bounds depend strongly on temperature of QCD phase transition and range from  $R < 0.13 \mu\text{m}$  to  $0.001 \mu\text{m}$  for  $\delta=2$ ; bounds for  $\delta=3,4$  can be derived from Table 1 in the paper.
- 28 HANHART 01 obtain bounds on  $R$  from limits on graviton cooling of supernova SN 1987a using numerical simulations of proto-neutron star neutrino emission.
- 29 CASSISI 00 obtain rough bounds on  $M_D$  (and thus  $R$ ) from red giant cooling for  $\delta=2,3$ . See their paper for details.
- 30 ACCIARRI 99S search for  $e^+ e^- \rightarrow Z G$  at  $\sqrt{s}=189$  GeV. Limits on the gravity scale are found in their Table 2, for  $\delta \leq 4$ .

## Mass Limits on $M_{TT}$

This section includes limits on the cut-off mass scale,  $M_{TT}$ , of dimension-8 operators from KK graviton exchange in models of large extra dimensions. Ambiguities in the UV-divergent summation are absorbed into the parameter  $\lambda$ , which is taken to be  $\lambda = \pm 1$  in the following analyses. Bounds for  $\lambda = -1$  are shown in parenthesis after the bound for  $\lambda = +1$ , if appropriate. Different papers use slightly different definitions of the mass scale. The definition used here is related to another popular convention by  $M_{TT}^4 = (2/\pi) \Lambda_T^4$ , as discussed in the above Review on "Extra Dimensions."

<u>VALUE (TeV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
> 1.48	95	31 ABAZOV	09AE D0	$p\bar{p} \rightarrow$ dijet, angular distrib.
> 1.45	95	32 ABAZOV	09D D0	$p\bar{p} \rightarrow e^+ e^-, \gamma\gamma$
> 1.1 ( $> 1.0$ )	95	33 SCHAEL	07A ALEP	$e^+ e^- \rightarrow e^+ e^-$
> 0.898 ( $> 0.998$ )	95	34 ABDALLAH	06C DLPH	$e^+ e^- \rightarrow \ell^+ \ell^-$
> 0.853 ( $> 0.939$ )	95	35 GERDES	06	$p\bar{p} \rightarrow e^+ e^-, \gamma\gamma$
> 0.96 ( $> 0.93$ )	95	36 ABAZOV	05V D0	$p\bar{p} \rightarrow \mu^+ \mu^-$
> 0.78 ( $> 0.79$ )	95	37 CHEKANOV	04B ZEUS	$e^\pm p \rightarrow e^\pm X$
> 0.805 ( $> 0.956$ )	95	38 ABBIENDI	03D OPAL	$e^+ e^- \rightarrow \gamma\gamma$
> 0.7 ( $> 0.7$ )	95	39 ACHARD	03D L3	$e^+ e^- \rightarrow ZZ$
> 0.82 ( $> 0.78$ )	95	40 ADLOFF	03 H1	$e^\pm p \rightarrow e^\pm X$
> 1.28 ( $> 1.25$ )	95	41 GIUDICE	03 RVUE	
> 20.6 ( $> 15.7$ )	95	42 GIUDICE	03 RVUE	Dim-6 operators
> 0.80 ( $> 0.85$ )	95	43 HEISTER	03C ALEP	$e^+ e^- \rightarrow \gamma\gamma$
> 0.84 ( $> 0.99$ )	95	44 ACHARD	02D L3	$e^+ e^- \rightarrow \gamma\gamma$
> 1.2 ( $> 1.1$ )	95	45 ABBOTT	01 D0	$p\bar{p} \rightarrow e^+ e^-, \gamma\gamma$
> 0.60 ( $> 0.63$ )	95	46 ABBIENDI	00R OPAL	$e^+ e^- \rightarrow \mu^+ \mu^-$

> 0.63	(> 0.50)	95	46	ABBIENDI	00R	OPAL	$e^+ e^- \rightarrow \tau^+ \tau^-$
> 0.68	(> 0.61)	95	46	ABBIENDI	00R	OPAL	$e^+ e^- \rightarrow \mu^+ \mu^-, \tau^+ \tau^-$
			47	ABREU	00A	DLPH	$e^+ e^- \rightarrow \gamma\gamma$
> 0.680	(> 0.542)	95	48	ABREU	00S	DLPH	$e^+ e^- \rightarrow \mu^+ \mu^-, \tau^+ \tau^-$
> 15–28		99.7	49	CHANG	00B	RVUE	Electroweak
> 0.98		95	50	CHEUNG	00	RVUE	$e^+ e^- \rightarrow \gamma\gamma$
> 0.29–0.38		95	51	GRAESSER	00	RVUE	$(g-2)_\mu$
> 0.50–1.1		95	52	HAN	00	RVUE	Electroweak
> 2.0	(> 2.0)	95	53	MATHEWS	00	RVUE	$\bar{p}p \rightarrow jj$
> 1.0	(> 1.1)	95	54	MELE	00	RVUE	$e^+ e^- \rightarrow VV$
			55	ABBIENDI	99P	OPAL	
			56	ACCIARRI	99M	L3	
			57	ACCIARRI	99S	L3	
> 1.412	(> 1.077)	95	58	BOURILKOV	99		$e^+ e^- \rightarrow e^+ e^-$

31 ABAZOV 09AE use dijet angular distributions in  $0.7 \text{ fb}^{-1}$  of data from  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  to place lower bounds on  $\Lambda_T$  (equivalent to their  $M_S$ ), here converted to  $M_{TT}$ .

32 ABAZOV 09D use  $1.05 \text{ fb}^{-1}$  of data from  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  to place lower bounds on  $\Lambda_T$  (equivalent to their  $M_S$ ), here converted to  $M_{TT}$ .

33 SCHAEL 07A use  $e^+ e^-$  collisions at  $\sqrt{s} = 189\text{--}209 \text{ GeV}$  to place lower limits on  $\Lambda_T$ , here converted to limits on  $M_{TT}$ .

34 ABDALLAH 06C use  $e^+ e^-$  collisions at  $\sqrt{s} \sim 130\text{--}207 \text{ GeV}$  to place lower limits on  $M_{TT}$ , which is equivalent to their definition of  $M_S$ . Bound shown includes all possible final state leptons,  $\ell = e, \mu, \tau$ . Bounds on individual leptonic final states can be found in their Table 31.

35 GERDES 06 use 100 to 110  $\text{pb}^{-1}$  of data from  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8 \text{ TeV}$ , as recorded by the CDF Collaboration during Run I of the Tevatron. Bound shown includes a  $K$ -factor of 1.3. Bounds on individual  $e^+ e^-$  and  $\gamma\gamma$  final states are found in their Table I.

36 ABAZOV 05V use  $246 \text{ pb}^{-1}$  of data from  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  to search for deviations in the differential cross section to  $\mu^+ \mu^-$  from graviton exchange.

37 CHEKANOV 04B search for deviations in the differential cross section of  $e^\pm p \rightarrow e^\pm X$  with  $130 \text{ pb}^{-1}$  of combined data and  $Q^2$  values up to  $40,000 \text{ GeV}^2$  to place a bound on  $M_{TT}$ .

38 ABBIENDI 03D use  $e^+ e^-$  collisions at  $\sqrt{s}=181\text{--}209 \text{ GeV}$  to place bounds on the ultra-violet scale  $M_{TT}$ , which is equivalent to their definition of  $M_S$ .

39 ACHARD 03D look for deviations in the cross section for  $e^+ e^- \rightarrow ZZ$  from  $\sqrt{s} = 200\text{--}209 \text{ GeV}$  to place a bound on  $M_{TT}$ .

40 ADLOFF 03 search for deviations in the differential cross section of  $e^\pm p \rightarrow e^\pm X$  at  $\sqrt{s}=301$  and  $319 \text{ GeV}$  to place bounds on  $M_{TT}$ .

41 GIUDICE 03 review existing experimental bounds on  $M_{TT}$  and derive a combined limit.

42 GIUDICE 03 place bounds on  $\Lambda_6$ , the coefficient of the gravitationally-induced dimension-6 operator  $(2\pi\lambda/\Lambda_6^2)(\sum \bar{f}\gamma_\mu\gamma^5 f)(\sum \bar{f}\gamma^\mu\gamma^5 f)$ , using data from a variety of experiments. Results are quoted for  $\lambda=\pm 1$  and are independent of  $\delta$ .

43 HEISTER 03C use  $e^+ e^-$  collisions at  $\sqrt{s}=189\text{--}209 \text{ GeV}$  to place bounds on the scale of dim-8 gravitational interactions. Their  $M_S^\pm$  is equivalent to our  $M_{TT}$  with  $\lambda=\pm 1$ .

44 ACHARD 02 search for s-channel graviton exchange effects in  $e^+ e^- \rightarrow \gamma\gamma$  at  $E_{\text{cm}} = 192\text{--}209 \text{ GeV}$ .

45 ABBOTT 01 search for variations in differential cross sections to  $e^+ e^-$  and  $\gamma\gamma$  final states at the Tevatron.

46 ABBIENDI 00R uses  $e^+ e^-$  collisions at  $\sqrt{s} = 189 \text{ GeV}$ .

- 47 ABREU 00A search for *s*-channel graviton exchange effects in  $e^+e^- \rightarrow \gamma\gamma$  at  $E_{cm}=189\text{--}202$  GeV.
- 48 ABREU 00S uses  $e^+e^-$  collisions at  $\sqrt{s}=183$  and 189 GeV. Bounds on  $\mu$  and  $\tau$  individual final states given in paper.
- 49 CHANG 00B derive  $3\sigma$  limit on  $M_{TT}$  of (28,19,15) TeV for  $\delta=(2,4,6)$  respectively assuming the presence of a torsional coupling in the gravitational action. Highly model dependent.
- 50 CHEUNG 00 obtains limits from anomalous diphoton production at OPAL due to graviton exchange. Original limit for  $\delta=4$ . However, unknown *UV* theory renders  $\delta$  dependence unreliable. Original paper works in HLZ convention.
- 51 GRAESSER 00 obtains a bound from graviton contributions to  $g-2$  of the muon through loops of 0.29 TeV for  $\delta=2$  and 0.38 TeV for  $\delta=4,6$ . Limits scale as  $\lambda^{1/2}$ . However calculational scheme not well-defined without specification of high-scale theory. See the "Extra Dimensions Review."
- 52 HAN 00 calculates corrections to gauge boson self-energies from KK graviton loops and constrain them using  $S$  and  $T$ . Bounds on  $M_{TT}$  range from 0.5 TeV ( $\delta=6$ ) to 1.1 TeV ( $\delta=2$ ); see text. Limits have strong dependence,  $\lambda^{\delta+2}$ , on unknown  $\lambda$  coefficient.
- 53 MATHEWS 00 search for evidence of graviton exchange in CDF and D $\emptyset$  dijet production data. See their Table 2 for slightly stronger  $\delta$ -dependent bounds. Limits expressed in terms of  $\tilde{M}_S^4 = M_{TT}^4/8$ .
- 54 MELE 00 obtains bound from KK graviton contributions to  $e^+e^- \rightarrow VV$  ( $V=\gamma,W,Z$ ) at LEP. Authors use Hewett conventions.
- 55 ABBIENDI 99P search for *s*-channel graviton exchange effects in  $e^+e^- \rightarrow \gamma\gamma$  at  $E_{cm}=189$  GeV. The limits  $G_+ > 660$  GeV and  $G_- > 634$  GeV are obtained from combined  $E_{cm}=183$  and 189 GeV data, where  $G_\pm$  is a scale related to the fundamental gravity scale.
- 56 ACCIARRI 99M search for the reaction  $e^+e^- \rightarrow \gamma G$  and *s*-channel graviton exchange effects in  $e^+e^- \rightarrow \gamma\gamma, W^+W^-, ZZ, e^+e^-, \mu^+\mu^-, \tau^+\tau^-, q\bar{q}$  at  $E_{cm}=183$  GeV. Limits on the gravity scale are listed in their Tables 1 and 2.
- 57 ACCIARRI 99S search for the reaction  $e^+e^- \rightarrow ZG$  and *s*-channel graviton exchange effects in  $e^+e^- \rightarrow \gamma\gamma, W^+W^-, ZZ, e^+e^-, \mu^+\mu^-, \tau^+\tau^-, q\bar{q}$  at  $E_{cm}=189$  GeV. Limits on the gravity scale are listed in their Tables 1 and 2.
- 58 BOURILKOV 99 performs global analysis of LEP data on  $e^+e^-$  collisions at  $\sqrt{s}=183$  and 189 GeV. Bound is on  $\Lambda_T$ .
- 

## Direct Limits on Gravitational or String Mass Scale

This section includes limits on the fundamental gravitational scale and/or the string scale from processes which depend directly on one or the other of these scales.

VALUE (TeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$\gtrsim 1\text{--}2$	59 ANCHORDOQ.02B	RVUE	Cosmic Rays
$>0.49$	60 ACCIARRI 00P	L3	$e^+e^- \rightarrow e^+e^-$
59 ANCHORDOQUI 02B derive bound on $M_D$ from non-observation of black hole production in high-energy cosmic rays. Bound is stronger for larger $\delta$ , but depends sensitively on threshold for black hole production.			
60 ACCIARRI 00P uses $e^+e^-$ collisions at $\sqrt{s}=183$ and 189 GeV. Bound on string scale $M_s$ from massive string modes. $M_s$ is defined in hep-ph/0001166 by $M_s(1/\pi)^{1/8}\alpha^{-1/4} = M$ where $(4\pi G)^{-1} = M^{n+2}R^n$ .			

---

## Limits on $1/R = M_c$

This section includes limits on  $1/R = M_c$ , the compactification scale in models with TeV extra dimensions, due to exchange of Standard Model KK excitations. Bounds assume fermions are not in the bulk, unless stated otherwise. See the "Extra Dimensions" review for discussion of model dependence.

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
>1.59	95	61 ABAZOV	09AE D0	$p\bar{p} \rightarrow$ dijet, angular dist.
>0.6	95	62 HAISCH	07 RVUE	$\bar{B} \rightarrow X_s \gamma$
>0.6	90	63 GOGOLADZE	06 RVUE	Electroweak
>3.3	95	64 CORNET	00 RVUE	Electroweak
> 3.3–3.8	95	65 RIZZO	00 RVUE	Electroweak
61 ABAZOV 09AE use dijet angular distributions in $0.7 \text{ fb}^{-1}$ of data from $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ to place a lower bound on the compactification scale.				
62 HAISCH 07 use inclusive $\bar{B}$ -meson decays to place a Higgs mass independent bound on the compactification scale $1/R$ in the minimal universal extra dimension model.				
63 GOGOLADZE 06 use electroweak precision observables to place a lower bound on the compactification scale in models with universal extra dimensions. Bound assumes a 115 GeV Higgs mass. See their Fig. 3 for the bound as a function of the Higgs mass.				
64 CORNET 00 translates a bound on the coefficient of the 4-fermion operator $(\bar{\ell}\gamma_\mu\tau^a\ell)(\bar{\ell}\gamma^\mu\tau^a\ell)$ derived by Hagiwara and Matsumoto into a limit on the mass scale of KK $W$ bosons.				
65 RIZZO 00 obtains limits from global electroweak fits in models with a Higgs in the bulk (3.8 TeV) or on the standard brane (3.3 TeV).				

## Limits on Kaluza-Klein Gravitons in Warped Extra Dimensions

This sections places limits on the mass of the first Kaluza-Klein (KK) excitation of the graviton in the warped extra dimension model of Randall and Sundrum. Experimental bounds depend strongly on the warp parameter,  $k$ . See the "Extra Dimensions" review for a full discussion.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
66 AALTONEN	08S CDF	$p\bar{p} \rightarrow G \rightarrow ZZ$	
67 ABAZOV	08J D0	$p\bar{p} \rightarrow G \rightarrow e^+e^-, \gamma\gamma$	
68 AALTONEN	07G CDF	$p\bar{p} \rightarrow G \rightarrow \gamma\gamma$	
69 AALTONEN	07H CDF	$p\bar{p} \rightarrow G \rightarrow e\bar{e}$	
70 ABAZOV	05N D0	$p\bar{p} \rightarrow G \rightarrow \ell\ell, \gamma\gamma$	
71 ABULENCIA	05A CDF	$p\bar{p} \rightarrow G \rightarrow \ell\bar{\ell}$	
66 AALTONEN 08S use $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ to search for KK gravitons in warped extra dimensions. They search for graviton resonances decaying to four electrons via two $Z$ bosons using $1.1 \text{ fb}^{-1}$ of data. See their Fig. 8 for limits on $\sigma \cdot B(G \rightarrow ZZ)$ versus the graviton mass.			
67 ABAZOV 08J use $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ to search for KK gravitons in warped extra dimensions. They search for graviton resonances decaying to electrons and photons using $1 \text{ fb}^{-1}$ of data. For warp parameter values of $k/\bar{M}_P$ between 0.01 and 0.1 the lower limit on the mass of the lightest excitation is between 300 and 900 GeV. See their Fig. 4 for more details.			
68 AALTONEN 07G use $p\bar{p}$ collisions at 1.96 TeV to search for KK gravitons in warped extra dimensions. They search for graviton resonances decaying to photons using $1.2 \text{ fb}^{-1}$ of data. For warp parameter values of $k/\bar{M}_P = 0.1, 0.05$ , and 0.01 the bounds			

on the graviton mass are 850, 694, and 230 GeV, respectively. See their Fig. 3 for more details.

- 69 AALTONEN 07H use  $p\bar{p}$  collisions at 1.96 TeV to search for KK gravitons in warped extra dimensions. They search for graviton resonances decaying to electrons using  $1.3 \text{ fb}^{-1}$  of data. For a warp parameter value of  $k/\bar{M}_P = 0.1$  the bound on the graviton mass is 807 GeV. See their Fig. 4 for more details. A combined analysis with the diphoton data of AALTONEN 07G yields for  $k/\bar{M}_P = 0.1$  a graviton mass lower bound of 889 GeV.
- 70 ABAZOV 05N use  $p\bar{p}$  collisions at 1.96 TeV to search for KK gravitons in warped extra dimensions. They search for graviton resonances decaying to muons, electrons or photons, using  $260 \text{ pb}^{-1}$  of data. For warp parameter values of  $k/\bar{M}_P = 0.1, 0.05, \text{ and } 0.01$ , the bounds on the graviton mass are 785, 650 and 250 GeV respectively. See their Fig. 3 for more details.
- 71 ABULENCIA 05A use  $p\bar{p}$  collisions at 1.96 TeV to search for KK gravitons in warped extra dimensions. They search for graviton resonances decaying to muons or electrons, using  $200 \text{ pb}^{-1}$  of data. For warp parameter values of  $k/\bar{M}_P = 0.1, 0.05, \text{ and } 0.01$ , the bounds on the graviton mass are 710, 510 and 170 GeV respectively.

## Limits on Mass of Radion

This section includes limits on mass of radion, usually in context of Randall-Sundrum models. See the “Extra Dimension Review” for discussion of model dependence.

<u>VALUE (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
$\gtrsim 35$	72 ABBIENDI 05	OPAL	$e^+ e^- \rightarrow Z$ radion
$> 120$	73 MAHANTA 00		$Z \rightarrow \text{radion } \ell\bar{\ell}$
	74 MAHANTA 00B		$p\bar{p} \rightarrow \text{radion} \rightarrow \gamma\gamma$
72	ABBIENDI 05 use $e^+ e^-$ collisions at $\sqrt{s} = 91$ GeV and $\sqrt{s} = 189\text{--}209$ GeV to place bounds on the radion mass in the RS model. See their Fig. 5 for bounds that depend on the radion-Higgs mixing parameter $\xi$ and on $\Lambda_W = \Lambda_\phi/\sqrt{6}$ . No parameter-independent bound is obtained.		
73	MAHANTA 00 obtain bound on radion mass in the RS model. Bound is from Higgs boson search at LEP I.		
74	MAHANTA 00B uses $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV; production via gluon-gluon fusion. Authors assume a radion vacuum expectation value of 1 TeV.		

## REFERENCES FOR Extra Dimensions

ABAZOV	09AE	PRL 103 191803	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	09D	PRL 102 051601	V.M. Abazov <i>et al.</i>	(D0 Collab.)
MASUDA	09	PRL 102 171101	M. Masuda, M. Sasaki	(ICRR)
AALTONEN	08AC	PRL 101 181602	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	08S	PR D78 012008	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	08J	PRL 100 091802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	08S	PRL 101 011601	V.M. Abazov <i>et al.</i>	(D0 Collab.)
DAS	08	PR D78 063011	P.K. Das, V.H.S. Kumar, P.K. Suresh	
GERACI	08	PR D78 022002	A.A. Geraci <i>et al.</i>	(STAN)
TRENKEL	08	PR D77 122001	C. Trenkel	
AALTONEN	07G	PRL 99 171801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	07H	PRL 99 171802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
DECCA	07A	EPJ C51 963	R.S. Decca <i>et al.</i>	
HAISCH	07	PR D76 034014	U. Haisch, A. Weiler	
KAPNER	07	PRL 98 021101	D.J. Kapner <i>et al.</i>	
SCHAEL	07A	EPJ C49 411	S. Schael <i>et al.</i>	(ALEPH Collab.)
TU	07	PRL 98 201101	L.-C. Tu <i>et al.</i>	
ABDALLAH	06C	EPJ C45 589	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABULENCIA,A	06	PRL 97 171802	A. Abulencia <i>et al.</i>	(CDF Collab.)
GERDES	06	PR D73 112008	D. Gerdes <i>et al.</i>	
GOGOLADZE	06	PR D74 093012	I. Gogoladze, C. Macesanu	

ABAZOV	05N	PRL 95 091801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	05V	PRL 95 161602	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBIENDI	05	PL B609 20	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABDALLAH	05B	EPJ C38 395	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABULENCIA	05A	PRL 95 252001	A. Abulencia <i>et al.</i>	(CDF Collab.)
SMULLIN	05	PR D72 122001	S.J. Smullin <i>et al.</i>	
ACHARD	04E	PL B587 16	P. Achard <i>et al.</i>	(L3 Collab.)
ACOSTA	04C	PRL 92 121802	D. Acosta <i>et al.</i>	(CDF Collab.)
CASSE	04	PRL 92 111102	M. Casse <i>et al.</i>	
CHEKANOV	04B	PL B591 23	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
HOYLE	04	PR D70 042004	C.D. Hoyle <i>et al.</i>	(WASH)
ABAZOV	03	PRL 90 251802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBIENDI	03D	EPJ C26 331	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ACHARD	03D	PL B572 133	P. Achard <i>et al.</i>	(L3 Collab.)
ADLOFF	03	PL B568 35	C. Adloff <i>et al.</i>	(H1 Collab.)
CHIAPERINI	03	PRL 90 151101	J. Chiaverini <i>et al.</i>	
GIUDICE	03	NP B663 377	G.F. Giudice, A. Strumia	
HANNESTAD	03	PR D67 125008	S. Hannestad, G.G. Raffelt	
Also		PR D69 029901(erratum)	S. Hannestad, G.G. Raffelt	
HEISTER	03C	EPJ C28 1	A. Heister <i>et al.</i>	(ALEPH Collab.)
LONG	03	Nature 421 922	J.C. Long <i>et al.</i>	
ACHARD	02	PL B524 65	P. Achard <i>et al.</i>	(L3 Collab.)
ACHARD	02D	PL B531 28	P. Achard <i>et al.</i>	(L3 Collab.)
ANCHORDOQ...02B		PR D66 103002	L. Anchordoqui <i>et al.</i>	
HANNESTAD	02	PRL 88 071301	S. Hannestad, G. Raffelt	
ABBOTT	01	PRL 86 1156	B. Abbott <i>et al.</i>	(D0 Collab.)
FAIRBAIRN	01	PL B508 335	M. Fairbairn	
HANHART	01	PL B509 1	C. Hanhart <i>et al.</i>	
HOYLE	01	PRL 86 1418	C.D. Hoyle <i>et al.</i>	
ABBIENDI	00R	EPJ C13 553	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	00A	PL B491 67	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00S	PL B485 45	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00Z	EPJ C17 53	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	00P	PL B489 81	M. Acciari <i>et al.</i>	(L3 Collab.)
CASSISI	00	PL B481 323	S. Cassisi <i>et al.</i>	
CHANG	00B	PRL 85 3765	L.N. Chang <i>et al.</i>	
CHEUNG	00	PR D61 015005	K. Cheung	
CORNET	00	PR D61 037701	F. Cornet, M. Relano, J. Rico	
GRAESSER	00	PR D61 074019	M.L. Graesser	
HAN	00	PR D62 125018	T. Han, D. Marfatia, R.-J. Zhang	
MAHANTA	00	PL B480 176	U. Mahanta, S. Rakshit	
MAHANTA	00B	PL B483 196	U. Mahanta, A. Datta	
MATHEWS	00	JHEP 0007 008	P. Mathews, S. Raychaudhuri, K. Sridhar	
MELE	00	PR D61 117901	S. Mele, E. Sanchez	
RIZZO	00	PR D61 016007	T.G. Rizzo, J.D. Wells	
ABBIENDI	99P	PL B465 303	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ACCIARRI	99M	PL B464 135	M. Acciari <i>et al.</i>	(L3 Collab.)
ACCIARRI	99R	PL B470 268	M. Acciari <i>et al.</i>	(L3 Collab.)
ACCIARRI	99S	PL B470 281	M. Acciari <i>et al.</i>	(L3 Collab.)
BOURILKOV	99	JHEP 9908 006	D. Bourilkov	
HOSKINS	85	PR D32 3084	J.K. Hoskins <i>et al.</i>	