

$\chi_{b2}(1P)$
 $I^G(JPC) = 0^+(2^{++})$
 J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$. $J = 2$ from SKWARNICKI 87.

 $\chi_{b2}(1P)$ MASSVALUE (MeV)DOCUMENT ID**9912.21±0.26±0.31 OUR EVALUATION**

From average γ energy below, using $\Upsilon(2S)$
mass = 10023.26 ± 0.31 MeV

 γ ENERGY IN $\Upsilon(2S)$ DECAYVALUE (MeV)DOCUMENT IDTECNCOMMENT**110.44±0.29 OUR AVERAGE**

Error includes scale factor of 1.1.

110.58 $\pm 0.08 \pm 0.30$	ARTUSO	05	CLEO	$\Upsilon(2S) \rightarrow \gamma X$
110.8 $\pm 0.3 \pm 0.6$	EDWARDS	99	CLE2	$\Upsilon(2S) \rightarrow \gamma \chi(1P)$
107.0 $\pm 1.1 \pm 1.3$	WALK	86	CBAL	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
110.6 $\pm 0.3 \pm 0.9$	ALBRECHT	85E	ARG	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
110.4 $\pm 0.8 \pm 2.2$	NERNST	85	CBAL	$\Upsilon(2S) \rightarrow \gamma X$
109.5 $\pm 0.7 \pm 1.0$	HAAS	84	CLEO	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
108.2 $\pm 0.3 \pm 2.0$	KLOPFEN...	83	CUSB	$\Upsilon(2S) \rightarrow \gamma X$
108.8 ± 4.0	PAUSS	83	CUSB	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

 $\chi_{b2}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \gamma \Upsilon(1S)$	(22 ± 4) %	
$\Gamma_2 D^0 X$	< 7.9 %	90%
$\Gamma_3 \pi^+ \pi^- K^+ K^- \pi^0$	(8 ± 5) $\times 10^{-5}$	
$\Gamma_4 2\pi^+ \pi^- K^- K_S^0$	< 1.0 $\times 10^{-4}$	90%
$\Gamma_5 2\pi^+ \pi^- K^- K_S^0 2\pi^0$	(5.3 ± 2.4) $\times 10^{-4}$	
$\Gamma_6 2\pi^+ 2\pi^- 2\pi^0$	(3.5 ± 1.4) $\times 10^{-4}$	
$\Gamma_7 2\pi^+ 2\pi^- K^+ K^-$	(1.1 ± 0.4) $\times 10^{-4}$	
$\Gamma_8 2\pi^+ 2\pi^- K^+ K^- \pi^0$	(2.1 ± 0.9) $\times 10^{-4}$	
$\Gamma_9 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(3.9 ± 1.8) $\times 10^{-4}$	
$\Gamma_{10} 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 5 $\times 10^{-4}$	90%
$\Gamma_{11} 3\pi^+ 3\pi^-$	(7.0 ± 3.1) $\times 10^{-5}$	
$\Gamma_{12} 3\pi^+ 3\pi^- 2\pi^0$	(1.0 ± 0.4) $\times 10^{-3}$	
$\Gamma_{13} 3\pi^+ 3\pi^- K^+ K^-$	< 8 $\times 10^{-5}$	90%
$\Gamma_{14} 3\pi^+ 3\pi^- K^+ K^- \pi^0$	(3.6 ± 1.5) $\times 10^{-4}$	
$\Gamma_{15} 4\pi^+ 4\pi^-$	(8 ± 4) $\times 10^{-5}$	
$\Gamma_{16} 4\pi^+ 4\pi^- 2\pi^0$	(1.8 ± 0.7) $\times 10^{-3}$	

$\chi_{b2}(1P)$ BRANCHING RATIOS **$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$**

VALUE	DOCUMENT ID	TECN	COMMENT
0.22±0.04 OUR AVERAGE			
0.27±0.06±0.06	WALK 86	CBAL	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
0.20±0.05	KLOPFEN... 83	CUSB	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

 Γ_1/Γ **$\Gamma(D^0 X)/\Gamma_{\text{total}}$**

VALUE	CL%
<7.9 × 10⁻²	90

DOCUMENT ID	TECN	COMMENT
1,2 BRIERE	CLEO	$\Upsilon(2S) \rightarrow \gamma D^0 X$

¹ For $p_{D^0} > 2.5$ GeV/c.² The authors also present their result as $(5.4 \pm 1.9 \pm 0.5) \times 10^{-2}$. **$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$** **$\Gamma_3/\Gamma$**

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
0.84±0.50±0.04	8	3 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$

³ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] = (6 \pm 3 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$ **Γ_4/Γ**

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	4 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$

⁴ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] < 7 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

 $\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$ **Γ_5/Γ**

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
5.3±2.4±0.3	11	5 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$

⁵ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] = (38 \pm 14 \pm 10) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ **Γ_6/Γ**

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
3.5±1.4±0.2	19	6 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

⁶ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] = (25 \pm 8 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.4±0.1	14	7 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$

⁷ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (8 \pm 2 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1±0.9±0.1	13	8 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$

⁸ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (15 \pm 5 \pm 4) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.9±1.8±0.2	11	9 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$

⁹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (28 \pm 11 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	10 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$

¹⁰ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] < 36 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.70±0.31±0.03	9	11 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$

¹¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (5 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.2±3.6±0.5	34	12 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$

¹² ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (73 \pm 16 \pm 20) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<0.8	90	13 ASNER	08A CLEO	$\gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
13 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.				

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

<i>VALUE</i> (units 10^{-4})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
3.6 ± 1.5 ± 0.2	14	14 ASNER	08A CLEO	$\gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
14 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (26 \pm 8 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

<i>VALUE</i> (units 10^{-4})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
0.84 ± 0.40 ± 0.04	7	15 ASNER	08A CLEO	$\gamma(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$
15 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (6 \pm 2 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

<i>VALUE</i> (units 10^{-4})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
18 ± 7 ± 1	29	16 ASNER	08A CLEO	$\gamma(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
16 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P))] = (132 \pm 31 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\chi_{b2}(1P)$ REFERENCES

ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
SKWARNICKI	87	PRL 58 972	T. Skwarnicki <i>et al.</i>	(Crystal Ball Collab.) J
WALK	86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFENSTEIN	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)