

$K_2(1580)$

$I(J^P) = \frac{1}{2}(2^-)$

OMMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K^- \pi^+ \pi^-$ system. Needs confirmation.

$K_2(1580)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 1580	OTTER	79	— 10,14,16 $K^- p$

$K_2(1580)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 110	OTTER	79	— 10,14,16 $K^- p$

$K_2(1580)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	seen
Γ_2 $K_2^*(1430)\pi$	possibly seen

$K_2(1580)$ BRANCHING RATIOS

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	
seen	
OTTER	79 HBC — 10,14,16 $K^- p$
$\Gamma(K_2^*(1430)\pi)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	
possibly seen	
OTTER	79 HBC — 10,14,16 $K^- p$

$K_2(1580)$ REFERENCES

OTTER	79	NP B147 1	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP
-------	----	-----------	------------------------	-------------------------------