

THE $\rho(770)$

Updated May 2010 by S. Eidelman (Novosibirsk) and G. Venzanoni (Frascati).

The determination of the parameters of the $\rho(770)$ is beset with many difficulties because of its large width. In physical region fits for hadroproduced ρ mesons, the line shape does not correspond to a relativistic Breit-Wigner function with a P -wave width, but requires some additional shape parameter. This dependence on parameterization was demonstrated long ago [1]. Bose-Einstein correlations are another source of shifts in the $\rho(770)$ line shape, particularly in multiparticle final state systems [2].

The same model dependence afflicts any other source of resonance parameters, such as the energy-dependence of the phase shift δ_1^1 , or the pole position. It is, therefore, not surprising that a study of $\rho(770)$ dominance in the decays of the η and η' reveals the need for specific dynamical effects, in addition to the $\rho(770)$ pole [3,4].

The cleanest determination of the $\rho(770)$ mass and width comes from e^+e^- annihilation and τ -lepton decays. Barate *et al.* [5] shows that the charged $\rho(770)$ parameters measured in τ -lepton decays are consistent with those of the neutral one determined from e^+e^- data [6]. This conclusion is qualitatively supported by the high-statistics study of Anderson *et al.* [7]. However, model-independent comparison of the two-pion mass spectrum in τ decays, and the $e^+e^- \rightarrow \pi^+\pi^-$ cross section, gives indications of discrepancies between the overall normalization: τ data are about 3% higher than e^+e^- data [7,8]. A detailed analysis using such two-pion mass spectra from τ decays measured by OPAL [9], CLEO [7], and ALEPH [10], as well as recent pion form factor measurements in e^+e^- annihilation by CMD-2 [11,12], show that the discrepancy can be as high as 10% above the ρ meson [13,14]. This discrepancy remains after recent measurements of the two-pion cross section in e^+e^- annihilation at KLOE [15,16] and SND [17,18]. The effect is not accounted for by isospin breaking [19–21], but the accuracy of its calculation may be overestimated [22,23]. Ghozzi [24] suggests that this effect can be explained if the

charged ρ mass were higher than that of the neutral one by a few MeV. Existing theoretical models of the possible mass difference predict either a much smaller value [25], or a heavier neutral ρ meson [26]. Experimental accuracy is not yet sufficient for unambiguous conclusions. The size of the effect is also sensitive to the possible width difference [27,28]. The discrepancy between e^+e^- and τ becomes smaller after a new measurement of the pion form factor using radiative return at BaBar [29], a high-statistics study of τ decays into two pions at Belle [30] and reanalysis of isospin breaking effects [31]. Benayoun *et al.* [32,33] performs a detailed analysis of the whole set of the ρ , ω , and ϕ decays, consistently taking into account mixing effects in the hidden local symmetry model, and claims that in this approach τ decays to two pions can be naturally accounted for.

References

1. J. Pisut and M. Roos, Nucl. Phys. **B6**, 325 (1968).
2. G.D. Lafferty, Z. Phys. **C60**, 659 (1993).
3. A. Abele *et al.*, Phys. Lett. **B402**, 195 (1997).
4. M. Benayoun *et al.*, Eur. Phys. J. **C31**, 525 (2003).
5. R. Barate *et al.*, Z. Phys. **C76**, 15 (1997).
6. L.M. Barkov *et al.*, Nucl. Phys. **B256**, 365 (1985).
7. S. Anderson *et al.*, Phys. Rev. **D61**, 112002 (2000).
8. S. Eidelman and V. Ivanchenko, Nucl. Phys. (Proc. Supp.) **B76**, 319 (1999).
9. K. Ackerstaff *et al.*, Eur. Phys. J. **C7**, 571 (1999).
10. S. Schael *et al.*, Phys. Reports **421**, 191 (2005).
11. R.R. Akhmetshin *et al.*, Phys. Lett. **B527**, 161 (2002).
12. R.R. Akhmetshin *et al.*, Phys. Lett. **B578**, 285 (2004).
13. M. Davier *et al.*, Eur. Phys. J. **C27**, 497 (2003).
14. M. Davier *et al.*, Eur. Phys. J. **C31**, 503 (2003).
15. A. Aloisio *et al.*, Phys. Lett. **B606**, 12 (2005).
16. F. Ambrosino *et al.*, Phys. Lett. **B670**, 285 (2009).
17. M.N. Achasov *et al.*, Sov. Phys. JETP **101**, 1053 (2005).
18. M.N. Achasov *et al.*, Sov. Phys. JETP **103**, 380 (2006).
19. R. Alemany *et al.*, Eur. Phys. J. **C2**, 123 (1998).
20. V. Cirigliano *et al.*, Phys. Lett. **B513**, 361 (2001).
21. V. Cirigliano *et al.*, Eur. Phys. J. **C23**, 121 (2002).

22. K. Maltman and C.E. Wolfe, Phys. Rev. **D73**, 013004 (2006).
23. C.E. Wolfe and K. Maltman, Phys. Rev. **D80**, 114024 (2009).
24. S. Ghozzi and F. Jegerlehner, Phys. Lett. **B583**, 222 (2004).
25. J. Bijnens *et al.*, Phys. Lett. **B374**, 210 (1996).
26. M.N. Achasov *et al.*, Sov. Phys. JETP Lett. **69**, 7 (1999).
27. G. Toledo Sanchez, J.L. Garcia-Luna, and V. Gonzalez-Enciso, Phys. Rev. **D76**, 033001 (2007).
28. F.V. Florez-Baez *et al.*, Phys. Rev. **D76**, 096010 (2007).
29. B. Aubert *et al.*, Phys. Rev. Lett. **103**, 231801 (2009).
30. M. Fujikawa *et al.*, Phys. Rev. **D78**, 072006 (2008).
31. M. Davier *et al.*, Eur. Phys. J. **C66**, 127 (2010).
32. M. Benayoun *et al.*, Eur. Phys. J. **C55**, 199 (2008).
33. M. Benayoun *et al.*, Eur. Phys. J. **C65**, 211 (2010).