

$\Lambda(2100)$ G_{07} $I(J^P) = 0(\frac{7}{2}^-)$ Status: ***

Discovered by COOL 66 and by WOHL 66. Most of the results published before 1973 are now obsolete and have been omitted. They may be found in our 1982 edition Physics Letters **111B** 1 (1982).

This entry only includes results from partial-wave analyses. Parameters of peaks seen in cross sections and in invariant-mass distributions around 2100 MeV used to be listed in a separate entry immediately following. It may be found in our 1986 edition Physics Letters **170B** 1 (1986).

 $\Lambda(2100)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2090 to 2110 (≈ 2100) OUR ESTIMATE			
2104 \pm 10	GOPAL 80	DPWA $\bar{K}N \rightarrow \bar{K}N$	
2106 \pm 30	DEBELLEFON 78	DPWA $\bar{K}N \rightarrow \bar{K}N$	
2110 \pm 10	GOPAL 77	DPWA $\bar{K}N$ multichannel	
2105 \pm 10	HEMINGWAY 75	DPWA $K^- p \rightarrow \bar{K}N$	
2115 \pm 10	KANE 74	DPWA $K^- p \rightarrow \Sigma\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2094	BACCARI 77	DPWA $K^- p \rightarrow \Lambda\omega$	
2094	DECLAIS 77	DPWA $\bar{K}N \rightarrow \bar{K}N$	
2110 or 2089	¹ NAKKASYAN 75	DPWA $K^- p \rightarrow \Lambda\omega$	

 $\Lambda(2100)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
100 to 250 (≈ 200) OUR ESTIMATE			
157 \pm 40	DEBELLEFON 78	DPWA $\bar{K}N \rightarrow \bar{K}N$	
250 \pm 30	GOPAL 77	DPWA $\bar{K}N$ multichannel	
241 \pm 30	HEMINGWAY 75	DPWA $K^- p \rightarrow \bar{K}N$	
152 \pm 15	KANE 74	DPWA $K^- p \rightarrow \Sigma\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
98	BACCARI 77	DPWA $K^- p \rightarrow \Lambda\omega$	
250	DECLAIS 77	DPWA $\bar{K}N \rightarrow \bar{K}N$	
244 or 302	¹ NAKKASYAN 75	DPWA $K^- p \rightarrow \Lambda\omega$	

 $\Lambda(2100)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	25–35 %
Γ_2 $\Sigma\pi$	~ 5 %
Γ_3 $\Lambda\eta$	<3 %
Γ_4 ΞK	<3 %

Γ_5	$\Lambda\omega$	<8 %
Γ_6	$N\bar{K}^*(892)$	10–20 %
Γ_7	$N\bar{K}^*(892)$, $S=1/2$, G -wave	
Γ_8	$N\bar{K}^*(892)$, $S=3/2$, D -wave	

The above branching fractions are our estimates, not fits or averages.

$\Lambda(2100)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.25 to 0.35 OUR ESTIMATE				
0.34±0.03	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.24±0.06	DEBELLEFON 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.31±0.03	HEMINGWAY 75	DPWA	$K^- p \rightarrow \bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.29	DECLAIS 77	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.30±0.03	GOPAL 77	DPWA	See GOPAL 80	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Sigma\pi$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
+0.12±0.04	GOPAL 77	DPWA	$\bar{K}N$ multichannel	
+0.11±0.01	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Lambda\eta$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
-0.050±0.020	RADER 73	MPWA	$K^- p \rightarrow \Lambda\eta$	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Xi K$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
0.035±0.018	LITCHFIELD 71	DPWA	$K^- p \rightarrow \Xi K$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.003	MULLER 69B	DPWA	$K^- p \rightarrow \Xi K$	
0.05	TRIPP 67	RVUE	$K^- p \rightarrow \Xi K$	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Lambda\omega$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
-0.070	² BACCARI 77	DPWA	GD_{37} wave	
+0.011	² BACCARI 77	DPWA	GG_{17} wave	
+0.008	² BACCARI 77	DPWA	GG_{37} wave	
0.122 or 0.154	¹ NAKKASYAN 75	DPWA	$K^- p \rightarrow \Lambda\omega$	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow N\bar{K}^*(892)$, $S=3/2$, D -wave $(\Gamma_1\Gamma_8)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.21±0.04	CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$

$(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow N\bar{K}^*(892)$, $S=1/2$, G -wave $(\Gamma_1\Gamma_7)^{1/2}/\Gamma$			
VALUE	DOCUMENT ID	TECN	COMMENT
-0.04 ± 0.03	³ CAMERON	78B DPWA	$K^- p \rightarrow N\bar{K}^*$

$\Lambda(2100)$ FOOTNOTES

- ¹ The NAKKASYAN 75 values are from the two best solutions found. Each has the $\Lambda(2100)$ and one additional resonance (P_3 or F_5).
- ² Note that the three for BACCARI 77 entries are for three different waves.
- ³ The published sign has been changed to be in accord with the baryon-first convention. The upper limit on the G_3 wave is 0.03.

$\Lambda(2100)$ REFERENCES

PDG	86	PL 170B 1	M. Aguilar-Benitez <i>et al.</i>	(CERN, CIT+)
PDG	82	PL 111B 1	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159	G.P. Gopal	(RHEL) IJP
CAMERON	78B	NP B146 327	W. Cameron <i>et al.</i>	(RHEL, LOIC) IJP
DEBELLEFON	78	NC 42A 403	A. de Bellefon <i>et al.</i>	(CDEF, SACL) IJP
BACCARI	77	NC 41A 96	B. Baccari <i>et al.</i>	(SACL, CDEF) IJP
DECLAIS	77	CERN 77-16	Y. Declais <i>et al.</i>	(CAEN, CERN) IJP
GOPAL	77	NP B119 362	G.P. Gopal <i>et al.</i>	(LOIC, RHEL) IJP
HEMINGWAY	75	NP B91 12	R.J. Hemingway <i>et al.</i>	(CERN, HEIDH, MPIM) IJP
NAKKASYAN	75	NP B93 85	A. Nakkasyan	(CERN) IJP
KANE	74	LBL-2452	D.F. Kane	(LBL) IJP
RADER	73	NC 16A 178	R.K. Rader <i>et al.</i>	(SACL, HEID, CERN+)
LITCHFIELD	71	NP B30 125	P.J. Litchfield <i>et al.</i>	(RHEL, CDEF, SACL) IJP
MULLER	69B	Thesis UCRL 19372	R.A. Muller	(LRL)
TRIPP	67	NP B3 10	R.D. Tripp <i>et al.</i>	(LRL, SLAC, CERN+)
COOL	66	PRL 16 1228	R.L. Cool <i>et al.</i>	(BNL)
WOHL	66	PRL 17 107	C.G. Wohl, F.T. Solmitz, M.L. Stevenson	(LRL) IJP