

$K_4^*(2045)$ $I(J^P) = \frac{1}{2}(4^+)$ **$K_4^*(2045)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2045 ± 9 OUR AVERAGE	Error includes scale factor of 1.1.				
$2062 \pm 14 \pm 13$		1 ASTON	86	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
2039 ± 10	400	2,3 CLELAND	82	SPEC \pm	$50 K^+ p \rightarrow K_S^0 \pi^\pm p$
2070^{+100}_{-40}		4 ASTON	81C	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2079 ± 7	431	TORRES	86	MPSF	$400 \mu A \rightarrow 4KX$
2088 ± 20	650	BAUBILLIER	82	HBC -	$8.25 K^- p \rightarrow K_S^0 \pi^- p$
2115 ± 46	488	CARMONY	77	HBC 0	$9 K^+ d \rightarrow K^+ \pi' s X$

¹ From a fit to all moments.
² From a fit to 8 moments.
³ Number of events evaluated by us.
⁴ From energy-independent partial-wave analysis.

 $K_4^*(2045)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
198 ± 30 OUR AVERAGE					
$221 \pm 48 \pm 27$		5 ASTON	86	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
189 ± 35	400	6,7 CLELAND	82	SPEC \pm	$50 K^+ p \rightarrow K_S^0 \pi^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
61 ± 58	431	TORRES	86	MPSF	$400 \mu A \rightarrow 4KX$
170^{+100}_{-50}	650	BAUBILLIER	82	HBC -	$8.25 K^- p \rightarrow K_S^0 \pi^- p$
240^{+500}_{-100}		8 ASTON	81C	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
300 ± 200		CARMONY	77	HBC 0	$9 K^+ d \rightarrow K^+ \pi' s X$

⁵ From a fit to all moments.
⁶ From a fit to 8 moments.
⁷ Number of events evaluated by us.
⁸ From energy-independent partial-wave analysis.

$K_4^*(2045)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(9.9±1.2) %
Γ_2 $K^*(892)\pi\pi$	(9 ± 5) %
Γ_3 $K^*(892)\pi\pi\pi$	(7 ± 5) %
Γ_4 $\rho K\pi$	(5.7±3.2) %
Γ_5 $\omega K\pi$	(5.0±3.0) %
Γ_6 $\phi K\pi$	(2.8±1.4) %
Γ_7 $\phi K^*(892)$	(1.4±0.7) %

 $K_4^*(2045)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.099±0.012	ASTON 88 LASS 0 11 $K^- p \rightarrow K^- \pi^+ n$
$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$	Γ_2/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.89±0.53	BAUBILLIER 82 HBC – 8.25 $K^- p \rightarrow p K_S^0 3\pi$
$\Gamma(K^*(892)\pi\pi\pi)/\Gamma(K\pi)$	Γ_3/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.75±0.49	BAUBILLIER 82 HBC – 8.25 $K^- p \rightarrow p K_S^0 3\pi$
$\Gamma(\rho K\pi)/\Gamma(K\pi)$	Γ_4/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.58±0.32	BAUBILLIER 82 HBC – 8.25 $K^- p \rightarrow p K_S^0 3\pi$
$\Gamma(\omega K\pi)/\Gamma(K\pi)$	Γ_5/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.50±0.30	BAUBILLIER 82 HBC – 8.25 $K^- p \rightarrow p K_S^0 3\pi$
$\Gamma(\phi K\pi)/\Gamma_{\text{total}}$	Γ_6/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.028±0.014	⁹ TORRES 86 MPSF 400 $pA \rightarrow 4KX$
$\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$	Γ_7/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.014±0.007	⁹ TORRES 86 MPSF 400 $pA \rightarrow 4KX$

⁹ Error determination is model dependent.

$K_4^*(2045)$ REFERENCES

ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
TORRES	86	PR D34 707	S. Torres <i>et al.</i>	(VPI, ARIZ, FNAL, FSU+)
BAUBILLIER	82	PL 118B 447	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)
ASTON	81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
CARMONY	77	PR D16 1251	D.D. Carmony <i>et al.</i>	(PURD, UCD, IUPU)