

$\eta'(958)$ $I^G(J^{PC}) = 0^+(0^-+)$ **$\eta'(958)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
957.78 ±0.06 OUR AVERAGE				
957.793±0.054±0.036	3.9k	LIBBY 08	CLEO	$J/\psi \rightarrow \gamma\eta'$
957.9 ±0.2 ±0.6	4800	WURZINGER 96	SPEC	$1.68 pd \rightarrow {}^3He\eta'$
957.46 ±0.33		DUANE 74	MMS	$\pi^- p \rightarrow n\Lambda\eta'$
958.2 ±0.5	1414	DANBURG 73	HBC	$2.2 K^- p \rightarrow \Lambda\eta'$
958 ±1	400	JACOBS 73	HBC	$2.9 K^- p \rightarrow \Lambda\eta'$
956.1 ±1.1	3415	¹ BASILE 71	CNTR	$1.6 \pi^- p \rightarrow n\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
957.5 ±0.2		BAI 04J	BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
959 ±1	630	² BELADIDZE 92C	VES	$36 \pi^- Be \rightarrow \pi^- \eta' \eta Be$
958 ±1	340	² ARMSTRONG 91B	OMEG	$300 pp \rightarrow pp\eta\pi^+\pi^-$
958.2 ±0.4	622	² AUGUSTIN 90	DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
957.8 ±0.2	2420	² AUGUSTIN 90	DM2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
956.3 ±1.0	143	² GIDAL 87	MRK2	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
957.4 ±1.4	535	³ BASILE 71	CNTR	$1.6 \pi^- p \rightarrow n\eta'$
957 ±1		RITTENBERG 69	HBC	$1.7-2.7 K^- p$

¹ Using all η' decays.² Systematic uncertainty not estimated.³ Using η' decays into neutrals. Not independent of the other listed BASILE 71 η' mass measurement. **$\eta'(958)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.194±0.009 OUR FIT					
0.30 ±0.09 OUR AVERAGE					
0.40 ±0.22	4800	WURZINGER 96	SPEC		$1.68 pd \rightarrow {}^3He\eta'$
0.28 ±0.10	1000	BINNIE 79	MMS 0		$\pi^- p \rightarrow n\Lambda\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.20 ±0.04		BAI 04J	BES2		$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

 $\eta'(958)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \pi^+\pi^-\eta$	(43.2 ±0.7) %	
$\Gamma_2 \rho^0\gamma$ (including non-resonant $\pi^+\pi^-\gamma$)	(29.3 ±0.5) %	
$\Gamma_3 \pi^0\pi^0\eta$	(21.7 ±0.8) %	
$\Gamma_4 \omega\gamma$	(2.75±0.22) %	
$\Gamma_5 \gamma\gamma$	(2.22±0.08) %	

Γ_6	$3\pi^0$	$(1.68 \pm 0.22) \times 10^{-3}$		
Γ_7	$\mu^+ \mu^- \gamma$	$(1.09 \pm 0.27) \times 10^{-4}$		
Γ_8	$\pi^+ \pi^- \mu^+ \mu^-$	$< 2.2 \times 10^{-4}$	90%	
Γ_9	$\pi^+ \pi^- \pi^0$	$(3.6^{+1.1}_{-0.9}) \times 10^{-3}$		
Γ_{10}	$\pi^0 \rho^0$	$< 4 \%$	90%	
Γ_{11}	$2(\pi^+ \pi^-)$	$< 2.4 \times 10^{-4}$	90%	
Γ_{12}	$\pi^+ \pi^- 2\pi^0$	$< 2.5 \times 10^{-3}$	90%	
Γ_{13}	$2(\pi^+ \pi^-)$ neutrals	$< 1 \%$	95%	
Γ_{14}	$2(\pi^+ \pi^-)\pi^0$	$< 1.9 \times 10^{-3}$	90%	
Γ_{15}	$2(\pi^+ \pi^-)2\pi^0$	$< 1 \%$	95%	
Γ_{16}	$3(\pi^+ \pi^-)$	$< 5 \times 10^{-4}$	90%	
Γ_{17}	$\pi^+ \pi^- e^+ e^-$	$(2.4^{+1.3}_{-1.0}) \times 10^{-3}$		
Γ_{18}	$\gamma e^+ e^-$	$< 9 \times 10^{-4}$	90%	
Γ_{19}	$\pi^0 \gamma \gamma$	$< 8 \times 10^{-4}$	90%	
Γ_{20}	$4\pi^0$	$< 5 \times 10^{-4}$	90%	
Γ_{21}	$e^+ e^-$	$< 2.1 \times 10^{-7}$	90%	
Γ_{22}	invisible	$< 9 \times 10^{-4}$	90%	

**Charge conjugation (C), Parity (P),
Lepton family number (LF) violating modes**

Γ_{23}	$\pi^+ \pi^-$	P, CP	$< 2.9 \times 10^{-3}$	90%
Γ_{24}	$\pi^0 \pi^0$	P, CP	$< 1.0 \times 10^{-3}$	90%
Γ_{25}	$\pi^0 e^+ e^-$	C	$[a] < 1.4 \times 10^{-3}$	90%
Γ_{26}	$\eta e^+ e^-$	C	$[a] < 2.4 \times 10^{-3}$	90%
Γ_{27}	3γ	C	$< 1.0 \times 10^{-4}$	90%
Γ_{28}	$\mu^+ \mu^- \pi^0$	C	$[a] < 6.0 \times 10^{-5}$	90%
Γ_{29}	$\mu^+ \mu^- \eta$	C	$[a] < 1.5 \times 10^{-5}$	90%
Γ_{30}	$e \mu$	LF	$< 4.7 \times 10^{-4}$	90%

[a] C parity forbids this to occur as a single-photon process.

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, 2 combinations of partial widths obtained from integrated cross section, and 13 branching ratios uses 39 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 29.6$ for 31 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

	x_2	x_3	x_4	x_5	x_6	x_9	x_{17}	Γ	
	x_1	x_2	x_3	x_4	x_5	x_6	x_9	x_{17}	
x_2	1								
x_3	-76	-57							
x_4	-19	-23	6						
x_5	-33	-25	36	-1					
x_6	-23	-17	28	1	10				
x_9	-1	-5	-7	-2	-4	-2			
x_{17}	-4	-6	-5	-2	-3	-2	0		
Γ	28	6	-23	4	-74	-6	4	4	

	Mode	Rate (MeV)
Γ_1	$\pi^+ \pi^- \eta$	0.084 ± 0.005
Γ_2	$\rho^0 \gamma$ (including non-resonant $\pi^+ \pi^- \gamma$)	0.0568 ± 0.0030
Γ_3	$\pi^0 \pi^0 \eta$	0.0421 ± 0.0023
Γ_4	$\omega \gamma$	0.0053 ± 0.0005
Γ_5	$\gamma \gamma$	0.00429 ± 0.00014
Γ_6	$3\pi^0$	$(3.3 \pm 0.5) \times 10^{-4}$
Γ_9	$\pi^+ \pi^- \pi^0$	$(7.0 \pm 2.1) \times 10^{-4}$
Γ_{17}	$\pi^+ \pi^- e^+ e^-$	$(4.7 \pm 2.5) \times 10^{-4}$

$\eta'(958)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$		Γ_5		
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.29 ± 0.14 OUR FIT				
4.28 ± 0.19 OUR AVERAGE				
$4.17 \pm 0.10 \pm 0.27$	2000	⁴ ACCIARRI	98Q L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \gamma$
$4.53 \pm 0.29 \pm 0.51$	266	KARCH	92 CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$
$3.61 \pm 0.13 \pm 0.48$		⁵ BEHREND	91 CELL	$e^+ e^- \rightarrow e^+ e^- \eta'(958)$
$4.6 \pm 1.1 \pm 0.6$	23	BARU	90 MD1	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \gamma$

$4.57 \pm 0.25 \pm 0.44$		BUTLER	90	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta'(958)$
$5.08 \pm 0.24 \pm 0.71$	547	⁶ ROE	90	ASP	$e^+ e^- \rightarrow e^+ e^- 2\gamma$
$3.8 \pm 0.7 \pm 0.6$	34	AIHARA	88C	TPC	$e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$
$4.9 \pm 0.5 \pm 0.5$	136	⁷ WILLIAMS	88	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$4.7 \pm 0.6 \pm 0.9$	143	⁸ GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$
4.0 ± 0.9		⁹ BARTEL	85E	JADE	$e^+ e^- \rightarrow e^+ e^- 2\gamma$

⁴ No non-resonant $\pi^+\pi^-$ contribution found.⁵ Reevaluated by us using $B(\eta' \rightarrow \rho(770)\gamma) = (30.2 \pm 1.3)\%$.⁶ Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.⁷ Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.⁸ Superseded by BUTLER 90.⁹ Systematic error not evaluated.

$\eta'(958) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $\gamma\gamma$ and with the total width is obtained from the integrated cross section into channel(i) in the $\gamma\gamma$ annihilation.

$$\Gamma(\gamma\gamma) \times \Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}} \quad \Gamma_5\Gamma_2/\Gamma$$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.26±0.04 OUR FIT				
1.26±0.07 OUR AVERAGE				Error includes scale factor of 1.2.
1.09±0.04±0.13		BEHREND	91	CELL $e^+ e^- \rightarrow e^+ e^- \rho(770)^0\gamma$
1.35±0.09±0.21		AIHARA	87	TPC $e^+ e^- \rightarrow e^+ e^- \rho\gamma$
1.13±0.04±0.13	867	ALBRECHT	87B	ARG $e^+ e^- \rightarrow e^+ e^- \rho\gamma$
1.53±0.09±0.21		ALTHOFF	84E	TASS $e^+ e^- \rightarrow e^+ e^- \rho\gamma$
1.14±0.08±0.11	243	BERGER	84B	PLUT $e^+ e^- \rightarrow e^+ e^- \rho\gamma$
1.73±0.34±0.35	95	JENNI	83	MRK2 $e^+ e^- \rightarrow e^+ e^- \rho\gamma$
1.49±0.13±0.027	213	BARTEL	82B	JADE $e^+ e^- \rightarrow e^+ e^- \rho\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.85±0.31±0.24	43	BEHREND	83B	CELL $e^+ e^- \rightarrow e^+ e^- \rho\gamma$

$$\Gamma(\gamma\gamma) \times \Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}} \quad \Gamma_5\Gamma_3/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.93±0.05 OUR FIT			
0.92±0.06±0.11	10 KARCH	92	CBAL $e^+ e^- \rightarrow e^+ e^- \eta\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.95±0.05±0.08	11 KARCH	90	CBAL $e^+ e^- \rightarrow e^+ e^- \eta\pi^0\pi^0$
1.00±0.08±0.10	11,12 ANTREASYAN	87	CBAL $e^+ e^- \rightarrow e^+ e^- \eta\pi^0\pi^0$
10 Reevaluated by us using $B(\eta \rightarrow \gamma\gamma) = (39.21 \pm 0.34)\%$. Supersedes ANTREASYAN 87 and KARCH 90.			
11 Superseded by KARCH 92.			
12 Using $BR(\eta \rightarrow 2\gamma) = (38.9 \pm 0.5)\%$.			

$\eta'(958)$ DECAY PARAMETERS

$$|\text{MATRIX ELEMENT}|^2 = |1 + \alpha y|^2 + cx + dx^2$$

α decay parameter


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.059±0.011 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
-0.072±0.012±0.006	7k	13 AMELIN	05A VES	28 $\pi^- A \rightarrow \eta' \pi^- A^*$
-0.021±0.025	6.7k	14 BRIERE	00 CLEO	10.6 $e^+ e^- \rightarrow \text{hadrons}$
-0.058±0.013		15,16 ALDE	86 GAM2	38 $\pi^- p \rightarrow n \eta 2\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.08 ± 0.03		15,16 KALBFLEISCH 74	RVUE	$\eta' \rightarrow \eta \pi^+ \pi^-$

¹³ This is a real part of α while $\text{Im}(\alpha) = 0.0 \pm 0.1 \pm 0.0$.

¹⁴ Assuming $\text{Im}(\alpha) = 0$, $c = 0$, and $d = 0$.

¹⁵ May not necessarily be the same for $\eta' \rightarrow \eta \pi^+ \pi^-$ and $\eta' \rightarrow \eta \pi^0 \pi^0$.

¹⁶ Assuming $\text{Im}(\alpha) = 0$, $c = 0$.



$\eta'(958) \rightarrow \eta\pi\pi$ DECAY PARAMETERS

$$|\text{MATRIX ELEMENT}|^2 \propto 1 + a Y + b Y^2 + c X + d X^2$$

X and Y are Dalitz variables and a , b , c , and d are real-valued parameters.
May be different for $\eta'(958) \rightarrow \eta\pi^+\pi^-$ and $\eta'(958) \rightarrow \eta\pi^0\pi^0$ decays.

a decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.066 ± 0.016 ± 0.003	15k	17 BLIK	09 GAM4	32.5 $\pi^- p \rightarrow \eta' n$
-0.127 ± 0.016 ± 0.008	20k	18 DOROFEEV	07 VES	27 $\pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

17 From $\eta' \rightarrow \eta\pi^0\pi^0$ decay. Parameters a , b , c , d are strongly correlated.

18 From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

b decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.063 ± 0.028 ± 0.004	15k	19 BLIK	09 GAM4	32.5 $\pi^- p \rightarrow \eta' n$
-0.106 ± 0.028 ± 0.014	20k	20 DOROFEEV	07 VES	27 $\pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

19 From $\eta' \rightarrow \eta\pi^0\pi^0$ decay. Parameters a , b , c , d are strongly correlated.

20 From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

c decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.107 ± 0.096 ± 0.003	15k	21 BLIK	09 GAM4	32.5 $\pi^- p \rightarrow \eta' n$
0.015 ± 0.011 ± 0.014	20k	22 DOROFEEV	07 VES	27 $\pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

21 From $\eta' \rightarrow \eta\pi^0\pi^0$ decay. Parameters a , b , c , d are strongly correlated.

22 From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

d decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.018 ± 0.078 ± 0.006	15k	23 BLIK	09 GAM4	32.5 $\pi^- p \rightarrow \eta' n$
-0.082 ± 0.017 ± 0.008	20k	24 DOROFEEV	07 VES	27 $\pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

23 From $\eta' \rightarrow \eta\pi^0\pi^0$ decay. Parameters a , b , c , d are strongly correlated. If $c \equiv 0$ from Bose-Einstein symmetry, $d = -0.067 \pm 0.020 \pm 0.003$.

24 From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

$\eta'(958)$ β PARAMETER
 $|\text{MATRIX ELEMENT}|^2 = (1 + 2\beta Z)$

See the “Note on η' Decay Parameters” in our 1994 edition Physical Review
D50 1173 (1994), p. 1454.

β decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.46 ± 0.22 OUR AVERAGE				Error includes scale factor of 1.4.
-0.59 ± 0.18	235	BLIK	08	GAMS 32 $\pi^- p \rightarrow \eta' n$
-0.1 ± 0.3		ALDE	87B	GAM2 38 $\pi^- p \rightarrow n 3\pi^0$

$\eta'(958)$ BRANCHING RATIOS

$\Gamma(\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$

$$\Gamma_1/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.432 ± 0.007 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.424 \pm 0.011 \pm 0.004$ 1.2k 25 PEDLAR 09 CLEO $J/\psi \rightarrow \gamma \eta'$

25 Not independent of other η' branching fractions and ratios in PEDLAR 09.

$\Gamma(\pi^+ \pi^- \eta(\text{charged decay}))/\Gamma_{\text{total}}$

$$0.286 \Gamma_1/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1237 ± 0.0020 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.123 ± 0.014 107 RITTENBERG 69 HBC 1.7–2.7 $K^- p$

0.10 ± 0.04 10 LONDON 66 HBC 2.24 $K^- p \rightarrow \Lambda 2\pi^+ 2\pi^- \pi^0$

0.07 ± 0.04 7 BADIER 65B HBC 3 $K^- p$

$\Gamma(\pi^+ \pi^- \eta(\text{neutral decay}))/\Gamma_{\text{total}}$

$$0.714 \Gamma_1/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.309 ± 0.005 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.314 ± 0.026 281 RITTENBERG 69 HBC 1.7–2.7 $K^- p$

$\Gamma(\rho^0 \gamma(\text{including non-resonant } \pi^+ \pi^- \gamma))/\Gamma_{\text{total}}$

$$\Gamma_2/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.293 ± 0.006 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.287 \pm 0.007 \pm 0.004$ 0.2k 26 PEDLAR 09 CLEO $J/\psi \rightarrow \gamma \eta'$

0.329 ± 0.033 298 RITTENBERG 69 HBC 1.7–2.7 $K^- p$

0.2 ± 0.1 20 LONDON 66 HBC 2.24 $K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$

0.34 ± 0.09 35 BADIER 65B HBC 3 $K^- p$

26 Not independent of other η' branching fractions and ratios in PEDLAR 09.

$\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / \Gamma(\pi^+ \pi^- \eta)$ Γ_2 / Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.678 ± 0.017 OUR FIT			
0.683 ± 0.020 OUR AVERAGE			
0.677 ± 0.024 ± 0.011	PEDLAR 09	CLE3	$J/\psi \rightarrow \eta' \gamma$
0.69 ± 0.03	ABLIKIM 06E	BES2	$J/\psi \rightarrow \eta' \gamma$

 $\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / \Gamma(\pi^+ \pi^- \eta (\text{neutral decay}))$ $\Gamma_2 / 0.714 \Gamma_1$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.949 ± 0.024 OUR FIT				
0.97 ± 0.09 OUR AVERAGE				
0.70 ± 0.22	AMSLER 04B	CBAR	$0 \bar{p} p \rightarrow \pi^+ \pi^- \eta$	
1.07 ± 0.17	BELADIDZE 92C	VES	$36 \pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$	
0.92 ± 0.14	DANBURG 73	HBC	$2.2 K^- p \rightarrow \Lambda X^0$	
1.11 ± 0.18	JACOBS 73	HBC	$2.9 K^- p \rightarrow \Lambda X^0$	

 $\Gamma(\pi^0 \pi^0 \eta) / \Gamma_{\text{total}}$ Γ_3 / Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.217 ± 0.008 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.235 ± 0.013 ± 0.004 3.2k ²⁷ PEDLAR 09 CLEO $J/\psi \rightarrow \gamma \eta'$

²⁷ Not independent of other η' branching fractions and ratios in PEDLAR 09.

 $\Gamma(\pi^0 \pi^0 \eta (\text{3}\pi^0 \text{decay})) / \Gamma_{\text{total}}$ $0.321 \Gamma_3 / \Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0697 ± 0.0026 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.11 ± 0.06 4 BENSINGER 70 DBC $2.2 \pi^+ d$

 $\Gamma(\pi^0 \pi^0 \eta) / \Gamma(\pi^+ \pi^- \eta)$ Γ_3 / Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.502 ± 0.026 OUR FIT			
0.555 ± 0.043 ± 0.013	PEDLAR 09	CLE3	$J/\psi \rightarrow \eta' \gamma$

 $\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / \Gamma(\pi \pi \eta)$ $\Gamma_2 / (\Gamma_1 + \Gamma_3)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.451 ± 0.012 OUR FIT			

0.43 ± 0.02 ± 0.02 BARBERIS 98C OMEG 450 $p p \rightarrow p_f \eta' p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ± 0.15 DAVIS 68 HBC $5.5 K^- p$

 $\Gamma(\omega \gamma) / \Gamma_{\text{total}}$ Γ_4 / Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0275 ± 0.0023 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0234 ± 0.0030 ± 0.0004 70 ²⁸ PEDLAR 09 CLEO $J/\psi \rightarrow \gamma \eta'$

²⁸ Not independent of other η' branching fractions and ratios in PEDLAR 09.

$\Gamma(\omega\gamma)/\Gamma(\pi^+\pi^-\eta)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_4/Γ_1
0.064±0.005 OUR FIT					
0.055±0.007±0.001		PEDLAR	09	CLE3 $J/\psi \rightarrow \eta'\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.068±0.013	68	ZANFINO	77	ASPK $8.4\pi^-p$	

 $\Gamma(\omega\gamma)/\Gamma(\pi^0\pi^0\eta)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_4/Γ_3
0.126±0.011 OUR FIT				
0.147±0.016	ALDE	87B	GAM2 $38\pi^-p \rightarrow n4\gamma$	

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/[\Gamma(\pi^+\pi^-\eta)+\Gamma(\pi^0\pi^0\eta)+\Gamma(\omega\gamma)]$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_2/(\Gamma_1+\Gamma_3+\Gamma_4)$
0.433±0.012 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.25 ± 0.14	DAUBER	64	HBC	$1.95K^-p$
-------------	--------	----	-----	------------

 $[\Gamma(\pi^0\pi^0\eta(\text{charged decay}))+\Gamma(\omega(\text{charged decay})\gamma)]/\Gamma_{\text{total}}$

$(0.286\Gamma_3+0.89\Gamma_4)/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0865±0.0032 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.045 ± 0.029	42	RITTENBERG	69	HBC $1.7\text{--}2.7K^-p$
---------------	----	------------	----	---------------------------

 $\Gamma(\pi^+\pi^-\text{ neutrals})/\Gamma_{\text{total}}$

$(0.714\Gamma_1+0.286\Gamma_3+0.89\Gamma_4)/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.395±0.004 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4 ± 0.1	39	LONDON	66	HBC $2.24K^-p \rightarrow \Lambda\pi^+\pi^-\text{ neutrals}$
0.35 ± 0.06	33	BADIER	65B	HBC $3K^-p$

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

Γ_5/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

2.22±0.08 OUR FIT**2.00±0.15 OUR AVERAGE**

$1.98^{+0.31}_{-0.27}\pm 0.07$	114	29 WICHT	08	BELL $B^\pm \rightarrow K^\pm\gamma\gamma$
2.00 ± 0.18		30 STANTON	80	SPEC $8.45\pi^-p \rightarrow n\pi^+\pi^-2\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.25\pm 0.16\pm 0.03$	0.3k	31 PEDLAR	09	CLEO $J/\psi \rightarrow \gamma\eta'$
1.8 ± 0.2	6000	32 APEL	79	NICE $15\text{--}40\pi^-p \rightarrow n2\gamma$
2.5 ± 0.7		DUANE	74	MMS $\pi^-p \rightarrow n\text{MM}$
1.71 ± 0.33	68	DALPIAZ	72	CNTR $1.6\pi^-p \rightarrow nX^0$
$2.0^{+0.8}_{-0.6}$	31	HARVEY	71	OSPK $3.65\pi^-p \rightarrow nX^0$

²⁹ WICHT 08 reports $[\Gamma(\eta'(958) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [\mathcal{B}(B^+ \rightarrow \eta' K^+)] = (1.40^{+0.16+0.15}_{-0.15-0.12}) \times 10^{-6}$ which we divide by our best value $\mathcal{B}(B^+ \rightarrow \eta' K^+) = (7.06 \pm 0.25) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³⁰ Includes APEL 79 result.

³¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.

³² Data is included in STANTON 80 evaluation.

$\Gamma(\gamma\gamma)/\Gamma(\pi^+\pi^-\eta)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_1
0.0512±0.0023 OUR FIT				
0.053 ±0.004 ±0.001	PEDLAR 09	CLE3	$J/\psi \rightarrow \eta'\gamma$	

$\Gamma(\gamma\gamma)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
0.0756±0.0034 OUR FIT				
0.080 ±0.008	ABLIKIM 06E	BES2	$J/\psi \rightarrow \eta'\gamma$	

$\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_3
0.102±0.004 OUR FIT				
0.105±0.010 OUR AVERAGE			Error includes scale factor of 1.9.	
0.091±0.009	AMSLER 93	CBAR	0.0 $\bar{p}p$	
0.112±0.002±0.006	ALDE 87B	GAM2	38 $\pi^- p \rightarrow n2\gamma$	

$\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta(\text{neutral decay}))$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_5/0.714\Gamma_3$
0.143±0.006 OUR FIT					

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.188±0.058 16 APEL 72 OSPK 3.8 $\pi^- p \rightarrow nX^0$

$\Gamma(\text{ neutrals})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$(0.714\Gamma_3+0.09\Gamma_4+\Gamma_5)/\Gamma$
0.180±0.006 OUR FIT					

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.185±0.022 535 BASILE 71 CNTR 1.6 $\pi^- p \rightarrow nX^0$

0.189±0.026 123 RITTENBERG 69 HBC 1.7–2.7 $K^- p$

$\Gamma(3\pi^0)/\Gamma(\pi^0\pi^0\eta)$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_3
78±10 OUR FIT					
78±10 OUR AVERAGE					
86±19	235	BLIK 08	GAMS	32 $\pi^- p \rightarrow \eta' n$	
74±15		ALDE 87B	GAM2	38 $\pi^- p \rightarrow n6\gamma$	
75±18		BINON 84	GAM2	30–40 $\pi^- p \rightarrow n6\gamma$	

$\Gamma(\mu^+\mu^-\gamma)/\Gamma(\gamma\gamma)$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_5
4.9±1.2	33	VIKTOROV 80	CNTR	25,33 $\pi^- p \rightarrow 2\mu\gamma$	

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<2.4	90	33 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
33 Not independent of measured value of Γ_8/Γ_1 from NAIK 09.				

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\pi^+\pi^-\eta)$ Γ_8/Γ_1

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.5	90	34 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
34 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 1.3 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.				

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE</u> (units 10^{-2})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.36 $^{+0.11}_{-0.09}$ OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.37^{+0.11}_{-0.09} \pm 0.04$	35 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<9	95 DANBURG	73 HBC	$2.2 K^- p \rightarrow \Lambda X^0$
<5	90 RITTENBERG 69	HBC	$1.7-2.7 K^- p$

35 Not independent of measured value of Γ_9/Γ_1 from NAIK 09.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_9/Γ_1

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.3 $^{+2.5}_{-2.1}$ OUR FIT				
8.25 $^{+2.49}_{-2.12}$ ± 0.04	20 36 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$	

36 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (21^{+6}_{-5} \pm 2) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0\rho^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.04	90 RITTENBERG 65	HBC	$2.7 K^- p$	

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				

< 2.4	90 37 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<100	90 RITTENBERG 69	HBC	$1.7-2.7 K^- p$

37 Not independent of measured value of Γ_{11}/Γ_1 from NAIK 09.

$\Gamma(2(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ Γ_{11}/Γ_1

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.6	90	38 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
38 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 1.4 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.				

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<27	90	39 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
39 Not independent of measured value of Γ_{12}/Γ_1 from NAIK 09.				

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{12}/Γ_1

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6	90	40 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
40 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 15 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.				

 $\Gamma(2(\pi^+\pi^-)\text{ neutrals})/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.01	95	DANBURG 73	HBC	$2.2 K^- p \rightarrow \Lambda X^0$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.01	90	RITTENBERG 69	HBC	1.7–2.7 $K^- p$

 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.002	90	41 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<0.01	90	RITTENBERG 69	HBC	1.7–2.7 $K^- p$
41 Not independent of measured value of Γ_{14}/Γ_1 from NAIK 09.				

 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{14}/Γ_1

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	42 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
42 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 11 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.				

 $\Gamma(2(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.01	95	KALBFLEISCH 64B	HBC	$K^- p \rightarrow \Lambda 2(\pi^+\pi^-)+\text{MM}$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.01	90	LONDON 66	HBC	Compilation

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.53	90	43 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<5	95	KALBFLEISCH 64B	HBC	$K^- p \rightarrow \Lambda 2(\pi^+\pi^-)$

43 Not independent of measured value of Γ_{16}/Γ_1 from NAIK 09. $\Gamma(3(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ Γ_{16}/Γ_1

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.2	90	44 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
44 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 3(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 3.0 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.				

 $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.4^{+1.3}_{-1.0} OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.5^{+1.2}_{-0.9} \pm 0.5$	45 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<6	90	RITTENBERG 65	HBC $K^- p$

45 Not independent of measured value of Γ_{17}/Γ_1 from NAIK 09. $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\pi^+\pi^-\eta)$ Γ_{17}/Γ_1

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.6^{+3.0}_{-2.2} OUR FIT				

5.50^{+2.99}_{-2.29} ± 0.03	8	46 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
---	---	---------	---------	----------------------------------

46 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-e^+e^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (14^{+7}_{-5} \pm 3) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma(\gamma e^+e^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.9	90	BRIERE	00 CLEO	$10.6 e^+e^-$

 $\Gamma(\pi^0\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_{19}/Γ_3

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<37	90	ALDE	87B GAM2	$38 \pi^- p \rightarrow n4\gamma$

 $\Gamma(4\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_{20}/Γ_3

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<23	90	ALDE	87B GAM2	$38 \pi^- p \rightarrow n8\gamma$

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-7})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.1	90	VOROBIEV	88	$e^+ e^- \rightarrow \pi^+ \pi^- \eta$

 Γ_{21}/Γ $\Gamma(\text{invisible})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9.5 90 47 NAIK 09 CLEO $J/\psi \rightarrow \gamma \eta'$

47 Not independent of measured value of Γ_{22}/Γ_1 from NAIK 09.

 $\Gamma(\text{invisible})/\Gamma(\gamma\gamma)$

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.69 90 ABLIKIM 06Q BES $J/\psi \rightarrow \phi \eta'$

 $\Gamma(\text{invisible})/\Gamma(\pi^+ \pi^- \eta)$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.1	90	48 NAIK 09 CLEO	$J/\psi \rightarrow \gamma \eta'$	

48 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \text{invisible})/\Gamma(\eta'(958) \rightarrow \pi^+ \pi^- \eta)] / [\mathcal{B}(\eta \rightarrow 2\gamma)] < 5.4 \times 10^{-3}$ which we multiply by our best value $\mathcal{B}(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.

 Γ_{22}/Γ_5 $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 29	90	49 MORI 07A BELL	$\gamma\gamma \rightarrow \pi^+ \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 3.3	90	50 MORI 07A BELL	$\gamma\gamma \rightarrow \pi^+ \pi^-$	
<800	95	DANBURG 73 HBC	$2.2 K^- p \rightarrow \Lambda X^0$	
<200	90	RITTENBERG 69 HBC	1.7–2.7 $K^- p$	

49 Taking into account interference with the $\gamma\gamma \rightarrow \pi^+ \pi^-$ continuum.

50 Without interference with the $\gamma\gamma \rightarrow \pi^+ \pi^-$ continuum.

 Γ_{23}/Γ $\Gamma(\pi^0 \pi^0)/\Gamma(\pi^0 \pi^0 \eta)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<45	90	ALDE 87B GAM2	$\pi^- p \rightarrow n 4\gamma$	

 Γ_{24}/Γ_3 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.4	90	BRIERE 00 CLEO	$10.6 e^+ e^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13 90 RITTENBERG 65 HBC 2.7 $K^- p$

 Γ_{25}/Γ $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 2.4	90	BRIERE 00 CLEO	$10.6 e^+ e^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<11 90 RITTENBERG 65 HBC 2.7 $K^- p$

 Γ_{26}/Γ

$\Gamma(3\gamma)/\Gamma(\pi^0\pi^0\eta)$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>
<4.6	90

 Γ_{27}/Γ_3

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ALDE	GAM2	$38 \pi^- p \rightarrow n 3\gamma$

 $\Gamma(\mu^+\mu^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>
<6.0	90

 Γ_{28}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
DZHELYADIN 81	CNTR	$30 \pi^- p \rightarrow \eta' n$

 $\Gamma(\mu^+\mu^-\eta)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>
<1.5	90

 Γ_{29}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
DZHELYADIN 81	CNTR	$30 \pi^- p \rightarrow \eta' n$

 $\Gamma(e\mu)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>
<4.7	90

 Γ_{30}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
BRIERE	CLEO	$10.6 e^+ e^-$

 $\eta'(958)$ C-NONCONSERVING DECAY PARAMETER

See the note on η decay parameters in the Stable Particle Particle Listings for definition of this parameter.

DECAY ASYMMETRY PARAMETER FOR $\pi^+\pi^-\gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.03 ± 0.04 OUR AVERAGE				
-0.019 ± 0.056		AIHARA	87	$2\gamma \rightarrow \pi^+\pi^-\gamma$
-0.069 ± 0.078	295	GRIGORIAN	75	$2.1 \pi^- p$
0.00 ± 0.10	103	KALBFLEISCH	75	$HBC \quad 2.18 K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.07 ± 0.08	152	RITTENBERG	65	$HBC \quad 2.1-2.7 K^- p$

 $\eta'(958)$ REFERENCES

BLIK	09	PAN 72 231 Translated from YAF 72 258.	A.M. Blik <i>et al.</i>	(IHEP (Protvino))
NAIK	09	PRL 102 061801	P. Naik <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
BLIK	08	PAN 71 2124 Translated from YAF 71 2161.	A. Blik <i>et al.</i>	(GAMS-4 π Collab.)
LIBBY	08	PRL 101 182002	J. Libby <i>et al.</i>	(CLEO Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
DOROFEEV	07	PL B651 22	V. Dorofeev <i>et al.</i>	(VES Collab.)
MORI	07A	JPSJ 76 074102	T. Mori <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06Q	PRL 97 202002	M. Ablikim <i>et al.</i>	(BES Collab.)
AMELIN	05A	PAN 68 372 Translated from YAF 68 401.	D.V. Amelin <i>et al.</i>	(VES Collab.)
AMSLER	04B	EPJ C33 23	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
BRIERE	00	PRL 84 26	R. Briere <i>et al.</i>	(CLEO Collab.)
ACCIARRI	98Q	PL B418 399	M. Acciari <i>et al.</i>	(L3 Collab.)
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)
WURZINGER	96	PL B374 283	R. Wurzinger <i>et al.</i>	(BONN, ORSAY, SACL+)
PDG	94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)
AMSLER	93	ZPHY C58 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BELADIDZE	92C	SJNP 55 1535 Translated from YAF 55 2748.	G.M. Beladidze, S.I. Bityukov, G.V. Borisov	(SERP+)

KARCH	92	ZPHY C54 33	K. Karch <i>et al.</i>	(Crystal Ball Collab.)
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BEHREND	91	ZPHY C49 401	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BARU	90	ZPHY C48 581	S.E. Baru <i>et al.</i>	(MD-1 Collab.)
BUTLER	90	PR D42 1368	F. Butler <i>et al.</i>	(Mark II Collab.)
KARCH	90	PL B249 353	K. Karch <i>et al.</i>	(Crystal Ball Collab.)
ROE	90	PR D41 17	N.A. Roe <i>et al.</i>	(ASP Collab.)
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)
VOROBIEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)
		Translated from YAF 48	436.	
WILLIAMS	88	PR D38 1365	D.A. Williams <i>et al.</i>	(Crystal Ball Collab.)
AIHARA	87	PR D35 2650	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.) JP
ALBRECHT	87B	PL B199 457	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALDE	87B	ZPHY C36 603	D.M. Alde <i>et al.</i>	(LANL, BELG, SERP, LAPP)
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)
ALDE	86	PL B177 115	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
BARTEL	85E	PL 160B 421	W. Bartel <i>et al.</i>	(JADE Collab.)
ALTHOFF	84E	PL 147B 487	M. Althoff <i>et al.</i>	(TASSO Collab.)
BERGER	84B	PL 142B 125	C. Berger	(PLUTO Collab.)
BINON	84	PL 140B 264	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP+)
BEHREND	83B	PL 125B 518 (erratum)	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
Also		PL 114B 378	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
JENNI	83	PR D27 1031	P. Jenni <i>et al.</i>	(SLAC, LBL)
BARTEL	82B	PL 113B 190	W. Bartel <i>et al.</i>	(JADE Collab.)
DZHELYADIN	81	PL 105B 239	R.I. Dzhelyadin <i>et al.</i>	(SERP)
STANTON	80	PL B92 353	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+)
VIKTOROV	80	SJNP 32 520	V.A. Viktorov <i>et al.</i>	(SERP)
		Translated from YAF 32	1005.	
APEL	79	PL 83B 131	W.D. Apel, K.H. Augenstein, E. Bertolucci	(KARLK+)
BINNIE	79	PL 83B 141	D.M. Binnie <i>et al.</i>	(LOIC)
ZANFINO	77	PRL 38 930	C. Zanfino <i>et al.</i>	(CARL, MCGI, OHIO+)
GRIGORIAN	75	NP B91 232	A. Grigorian <i>et al.</i>	(+)
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)
DUANE	74	PRL 32 425	A. Duane <i>et al.</i>	(LOIC, SHMP)
KALBFLEISCH	74	PR D10 916	G.R. Kalbfleisch	(BNL)
DANBURG	73	PR D8 3744	J.S. Danburg <i>et al.</i>	(BNL, MICH) JP
JACOBS	73	PR D8 18	S.M. Jacobs <i>et al.</i>	(BRAN, UMD, SYRA+) JP
APEL	72	PL 40B 680	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)
DALPIAZ	72	PL 42B 377	P.F. Dalpiaz <i>et al.</i>	(CERN)
BASILE	71	NC 3A 371	M. Basile <i>et al.</i>	(CERN, BGNA, STRB)
HARVEY	71	PRL 27 885	E.H. Harvey <i>et al.</i>	(MINN, MICH)
BENSINGER	70	PL 33B 505	J.R. Bensinger <i>et al.</i>	(WISC)
RITTENBERG	69	Thesis UCRL 18863	A. Rittenberg	(LRL) I
DAVIS	68	PL 27B 532	R. Davis <i>et al.</i>	(NWES, ANL)
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IJP
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)
RITTENBERG	65	PRL 15 556	A. Rittenberg, G.R. Kalbfleisch	(LRL, BNL)
DAUBER	64	PRL 13 449	P.M. Dauber <i>et al.</i>	(UCLA) JP
KALBFLEISCH	64B	PRL 13 349	G.R. Kalbfleisch, O.I. Dahl, A. Rittenberg	(LRL) JP
