

$\chi_{b2}(2P)$
 $I^G(JPC) = 0^+(2^{++})$
J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

 $\chi_{b2}(2P)$ MASSVALUE (GeV)DOCUMENT ID
 $10.26865 \pm 0.00022 \pm 0.00050$ OUR EVALUATION From γ energy below, using $\Upsilon(3S)$
mass = 10355.2 ± 0.5 MeV

 $m_{\chi_{b2}(2P)} - m_{\chi_{b1}(2P)}$ VALUE (MeV)DOCUMENT IDTECNCOMMENT **$13.5 \pm 0.4 \pm 0.5$**

1 HEINTZ

92

CSB2

 $e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma \gamma$
¹ From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

 γ ENERGY IN $\Upsilon(3S)$ DECAYVALUE (MeV)EVTSDOCUMENT IDTECNCOMMENT **86.19 ± 0.22 OUR EVALUATION**

Treating systematic errors as correlated

 86.40 ± 0.18 OUR AVERAGE $86.04 \pm 0.06 \pm 0.27$

ARTUSO

05

CLEO

 $\Upsilon(3S) \rightarrow \gamma X$ 86 ± 1

101

CRAWFORD

92B

CLE2

 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ 86.7 ± 0.4

10319

2 HEINTZ

92

CSB2

 $e^+ e^- \rightarrow \gamma X$ 86.9 ± 0.4

157

3 HEINTZ

92

CSB2

 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ $86.4 \pm 0.1 \pm 0.4$

30741

MORRISON

91

CLE2

 $e^+ e^- \rightarrow \gamma X$
² A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

³ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

 $\chi_{b2}(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \omega \Upsilon(1S)$	$(1.10^{+0.34}_{-0.30})\%$	
$\Gamma_2 \gamma \Upsilon(2S)$	$(16.2 \pm 2.4)\%$	
$\Gamma_3 \gamma \Upsilon(1S)$	$(7.1 \pm 1.0)\%$	
$\Gamma_4 \pi\pi\chi_{b2}(1P)$	$(6.0 \pm 2.1) \times 10^{-3}$	
$\Gamma_5 D^0 X$	$< 2.4\%$	90%
$\Gamma_6 \pi^+\pi^- K^+ K^- \pi^0$	$< 1.1 \times 10^{-4}$	90%
$\Gamma_7 2\pi^+\pi^- K^- K_S^0$	$< 9 \times 10^{-5}$	90%
$\Gamma_8 2\pi^+\pi^- K^- K_S^0 2\pi^0$	$< 7 \times 10^{-4}$	90%

Γ_9	$2\pi^+ 2\pi^- 2\pi^0$	$(3.9 \pm 1.6) \times 10^{-4}$		
Γ_{10}	$2\pi^+ 2\pi^- K^+ K^-$	$(9 \pm 4) \times 10^{-5}$		
Γ_{11}	$2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(2.4 \pm 1.1) \times 10^{-4}$		
Γ_{12}	$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(4.7 \pm 2.3) \times 10^{-4}$		
Γ_{13}	$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$< 4 \times 10^{-4}$		90%
Γ_{14}	$3\pi^+ 3\pi^-$	$(9 \pm 4) \times 10^{-5}$		
Γ_{15}	$3\pi^+ 3\pi^- 2\pi^0$	$(1.2 \pm 0.4) \times 10^{-3}$		
Γ_{16}	$3\pi^+ 3\pi^- K^+ K^-$	$(1.4 \pm 0.7) \times 10^{-4}$		
Γ_{17}	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(4.2 \pm 1.7) \times 10^{-4}$		
Γ_{18}	$4\pi^+ 4\pi^-$	$(9 \pm 5) \times 10^{-5}$		
Γ_{19}	$4\pi^+ 4\pi^- 2\pi^0$	$(1.3 \pm 0.5) \times 10^{-3}$		

 $\chi_{b2}(2P)$ BRANCHING RATIOS **$\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.10^{+0.32+0.11}_{-0.28-0.10}$	$20.1^{+5.8}_{-5.1}$	⁴ CRONIN-HEN..04	CLE3	$\Upsilon(3S) \rightarrow \gamma \omega \Upsilon(1S)$

⁴ Using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.4 \pm 0.8)\%$ and $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 2 (2.48 \pm 0.06)\%$.

 $\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.162 ± 0.024 OUR AVERAGE			
$0.135 \pm 0.025 \pm 0.035$	⁵ CRAWFORD 92B	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
$0.173 \pm 0.021 \pm 0.019$	⁶ HEINTZ 92	CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

⁵ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (4.98 \pm 0.94 \pm 0.62) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 0.135 \pm 0.003 \pm 0.017$.

⁶ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.1 \pm 0.5 \pm 0.4)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

 $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.071 ± 0.010 OUR AVERAGE			
$0.072 \pm 0.014 \pm 0.013$	⁷ CRAWFORD 92B	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
$0.070 \pm 0.010 \pm 0.006$	⁸ HEINTZ 92	CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

⁷ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(2S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (5.03 \pm 0.94 \pm 0.63) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 0.135 \pm 0.003 \pm 0.017$.

⁸ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.1 \pm 0.5 \pm 0.4)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

 $\Gamma(\pi\pi\chi_{b2}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$6.0 \pm 1.6 \pm 1.4$	⁹ CAWLFIELD 06	CLE3	$\Upsilon(3S) \rightarrow 2(\gamma \pi \ell)$

⁹ CAWLFIELD 06 quote $\Gamma(\chi_b(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV assuming I-spin conservation, no D-wave contribution, $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$ keV, and $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$ keV.

$\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-2}$	90	10,11 BRIERE	08 CLEO	$\gamma(3S) \rightarrow \gamma D^0 X$

10 For $p_{D^0} > 2.5$ GeV/c.11 The authors also present their result as $(0.2 \pm 1.4 \pm 0.1) \times 10^{-2}$. $\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	12 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

12 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 14 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

 $\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	13 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$

13 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 12 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

 $\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	14 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$

14 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 87 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

 $\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.9 \pm 1.6 \pm 0.5$	23	15 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

15 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (51 \pm 16 \pm 13) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.9 \pm 0.4 \pm 0.1$	11	16 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$

16 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±1.0±0.3	16	17 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$

17 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (32 \pm 11 \pm 8) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.7±2.2±0.6	14	18 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$

18 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (62 \pm 23 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	19 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$

19 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 58 \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.9±0.4±0.1	14	20 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$

20 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12±4±1	45	21 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$

21 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (159 \pm 33 \pm 43) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.4±0.7±0.2	12	22 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$

22 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (19 \pm 7 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.2±1.7±0.5	16	23 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$

²³ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (55 \pm 16 \pm 15) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.9±0.4±0.1	9	24 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$

²⁴ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
13±5±2	27	25 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$

²⁵ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (165 \pm 46 \pm 50) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\chi_{b2}(2P)$ REFERENCES

ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
CAWLFIELD	06	PR D73 012003	C. Cawlfieeld <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...04	04	PRL 92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO3 Collab.)
CRAWFORD	92B	PL B294 139	G. Crawford, R. Fulton	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)