

$f_2(1810)$ $I^G(J^{PC}) = 0^+(2^{++})$

OMMITTED FROM SUMMARY TABLE

Needs confirmation.

 $f_2(1810)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1815 ± 12 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.			
1800 ± 30	40	ALDE	88D GAM4	$300 \pi^- p \rightarrow \pi^- p 4\pi^0$
1806 ± 10	1600	ALDE	87 GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$
1870 ± 40	1	ALDE	86D GAM4	$100 \pi^- p \rightarrow \eta\eta n$
1857^{+35}_{-24}	2	COSTA...	80 OMEG	$10 \pi^- p \rightarrow K^+ K^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1858^{+18}_{-71}	³ LONGACRE	86 RVUE	Compilation
1799 ± 15	⁴ CASON	82 STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$


¹ Seen in only one solution.

² Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

³ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

⁴ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.

WEIGHTED AVERAGE
 1815 ± 12 (Error scaled by 1.4)



$f_2(1810)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
197± 22 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.			
160± 30	40	ALDE	88D GAM4	300 $\pi^- p \rightarrow \pi^- p 4\pi^0$
190± 20	1600	ALDE	87 GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$
250± 30	5	ALDE	86D GAM4	100 $\pi^- p \rightarrow \eta\eta n$
185 ⁺¹⁰² ₋₁₃₉	6	COSTA...	80 OMEG	10 $\pi^- p \rightarrow K^+ K^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
388 ⁺¹⁵ ₋₂₁	7	LONGACRE	86 RVUE	Compilation
280 ⁺⁴² ₋₃₅	8	CASON	82 STRC	8 $\pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$


⁵ Seen in only one solution.

⁶ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

⁷ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

⁸ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.

WEIGHTED AVERAGE
197±22 (Error scaled by 1.5)



$f_2(1810)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	
Γ_2 $\eta\eta$	
Γ_3 $4\pi^0$	seen
Γ_4 K^+K^-	

$f_2(1810)$ BRANCHING RATIOS

$$\frac{\Gamma(\pi\pi)}{\Gamma_{\text{total}}} \quad \frac{\Gamma_1}{\Gamma}$$

VALUE	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
not seen	AMSLER 02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
not seen	PROKOSHKIN 97	GAM2	$38 \pi^- p \rightarrow \pi^0\pi^0 n$
$0.21^{+0.02}_{-0.03}$	9 LONGACRE 86	RVUE	Compilation
0.44 ± 0.03	10 CASON 82	STRC	$8 \pi^+ p \rightarrow \Delta^{++}\pi^0\pi^0$

⁹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

¹⁰ Included in LONGACRE 86 global analysis.

$$\frac{\Gamma(\eta\eta)}{\Gamma_{\text{total}}} \quad \frac{\Gamma_2}{\Gamma}$$

VALUE	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.008^{+0.028}_{-0.003}$	11 LONGACRE 86	RVUE	Compilation
¹¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.			

$$\frac{\Gamma(\pi\pi)}{\Gamma(4\pi^0)} \quad \frac{\Gamma_1/\Gamma_3}{\Gamma}$$

VALUE	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
<0.75	ALDE 87	GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$

$$\frac{\Gamma(4\pi^0)}{\Gamma(\eta\eta)} \quad \frac{\Gamma_3/\Gamma_2}{\Gamma}$$

VALUE	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.8 ± 0.3	ALDE 87	GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$

$$\frac{\Gamma(K^+K^-)}{\Gamma_{\text{total}}} \quad \frac{\Gamma_4}{\Gamma}$$

VALUE	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.003^{+0.019}_{-0.002}$	12 LONGACRE 86	RVUE	Compilation
seen	COSTA... 80	OMEG	$10 \pi^- p \rightarrow K^+K^- n$
¹² From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.			

$f_2(1810)$ REFERENCES

AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
PROKOSHIN	97	SPD 42 117	Y.D. Prokoshkin <i>et al.</i>	(SERP)
		Translated from DANS 353 323.		
ALDE	88D	SJNP 47 810	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
		Translated from YAF 47 1273.		
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)
COSTA...	80	NP B175 402	G. Costa de Beauregard <i>et al.</i>	(BARI, BONN+)