



$$J = 1$$

A REVIEW GOES HERE – Check our WWW List of Reviews

W MASS

The W -mass listed here corresponds to the mass parameter in a Breit-Wigner distribution with mass-dependent width. To obtain the world average, common systematic uncertainties between experiments are properly taken into account. The LEP-2 average W mass based on published results is 80.376 ± 0.033 GeV [CERN-PH-EP/2006-042]. The combined Tevatron data yields an average W mass of 80.420 ± 0.031 GeV [FERMILAB-TM-2439-E].

OUR FIT uses these average LEP and Tevatron mass values and combines them assuming no correlations.

| <u>VALUE (GeV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| 80.399 ± 0.023 OUR FIT | | | | |
| 80.401 ± 0.043 | 500k | 1 ABAZOV | 09AB D0 | $E_{cm}^{p\bar{p}} = 1.96$ TeV |
| 80.336 ± 0.055 ± 0.039 | 10.3k | 2 ABDALLAH | 08A DLPH | $E_{cm}^{ee} = 161$ –209 GeV |
| 80.413 ± 0.034 ± 0.034 | 115k | 3 AALTONEN | 07F CDF | $E_{cm}^{p\bar{p}} = 1.96$ TeV |
| 80.415 ± 0.042 ± 0.031 | 11830 | 4 ABBIENDI | 06 OPAL | $E_{cm}^{ee} = 170$ –209 GeV |
| 80.270 ± 0.046 ± 0.031 | 9909 | 5 ACHARD | 06 L3 | $E_{cm}^{ee} = 161$ –209 GeV |
| 80.440 ± 0.043 ± 0.027 | 8692 | 6 SCHAEEL | 06 ALEP | $E_{cm}^{ee} = 161$ –209 GeV |
| 80.483 ± 0.084 | 49247 | 7 ABAZOV | 02D D0 | $E_{cm}^{p\bar{p}} = 1.8$ TeV |
| 80.433 ± 0.079 | 53841 | 8 AFFOLDER | 01E CDF | $E_{cm}^{p\bar{p}} = 1.8$ TeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 82.87 ± 1.82 $\begin{smallmatrix} +0.30 \\ -0.16 \end{smallmatrix}$ | 1500 | 9 AKTAS | 06 H1 | $e^\pm p \rightarrow \bar{\nu}_e(\nu_e)X, \sqrt{s} \approx 300$ GeV |
| 80.3 ± 2.1 ± 1.2 ± 1.0 | 645 | 10 CHEKANOV | 02C ZEUS | $e^- p \rightarrow \nu_e X, \sqrt{s} = 318$ GeV |
| 81.4 $\begin{smallmatrix} +2.7 \\ -2.6 \end{smallmatrix}$ ± 2.0 $\begin{smallmatrix} +3.3 \\ -3.0 \end{smallmatrix}$ | 1086 | 11 BREITWEG | 00D ZEUS | $e^+ p \rightarrow \bar{\nu}_e X, \sqrt{s} \approx 300$ GeV |
| 80.84 ± 0.22 ± 0.83 | 2065 | 12 ALITTI | 92B UA2 | See W/Z ratio below |
| 80.79 ± 0.31 ± 0.84 | | 13 ALITTI | 90B UA2 | $E_{cm}^{p\bar{p}} = 546,630$ GeV |
| 80.0 ± 3.3 ± 2.4 | 22 | 14 ABE | 89I CDF | $E_{cm}^{p\bar{p}} = 1.8$ TeV |
| 82.7 ± 1.0 ± 2.7 | 149 | 15 ALBAJAR | 89 UA1 | $E_{cm}^{p\bar{p}} = 546,630$ GeV |
| 81.8 $\begin{smallmatrix} +6.0 \\ -5.3 \end{smallmatrix}$ ± 2.6 | 46 | 16 ALBAJAR | 89 UA1 | $E_{cm}^{p\bar{p}} = 546,630$ GeV |
| 89 ± 3 ± 6 | 32 | 17 ALBAJAR | 89 UA1 | $E_{cm}^{p\bar{p}} = 546,630$ GeV |
| 81. ± 5. | 6 | ARNISON | 83 UA1 | $E_{cm}^{ee} = 546$ GeV |
| 80. $\begin{smallmatrix} +10. \\ -6. \end{smallmatrix}$ | 4 | BANNER | 83B UA2 | Repl. by ALITTI 90B |

- ¹ ABAZOV 09AB study the transverse mass, transverse electron momentum, and transverse missing energy in a sample of 0.5 million $W \rightarrow e\nu$ decays selected in Run-II data. The quoted result combines all three methods, accounting for correlations.
- ² ABDALLAH 08A use direct reconstruction of the kinematics of $W^+ W^- \rightarrow q\bar{q}\ell\nu$ and $W^+ W^- \rightarrow q\bar{q}q\bar{q}$ events for energies 172 GeV and above. The W mass was also extracted from the dependence of the WW cross section close to the production threshold and combined appropriately to obtain the final result. The systematic error includes ± 0.025 GeV due to final state interactions and ± 0.009 GeV due to LEP energy uncertainty.
- ³ AALTONEN 07F obtain high purity $W \rightarrow e\nu_e$ and $W \rightarrow \mu\nu_\mu$ candidate samples totaling 63,964 and 51,128 events respectively. The W mass value quoted above is derived by simultaneously fitting the transverse mass and the lepton, and neutrino p_T distributions.
- ⁴ ABBIENDI 06 use direct reconstruction of the kinematics of $W^+ W^- \rightarrow q\bar{q}\ell\nu_\ell$ and $W^+ W^- \rightarrow q\bar{q}q\bar{q}$ events. The result quoted here is obtained combining this mass value with the results using $W^+ W^- \rightarrow \ell\nu_\ell\ell'\nu_{\ell'}$ events in the energy range 183–207 GeV (ABBIENDI 03C) and the dependence of the WW production cross-section on m_{WW} at threshold. The systematic error includes ± 0.009 GeV due to the uncertainty on the LEP beam energy.
- ⁵ ACHARD 06 use direct reconstruction of the kinematics of $W^+ W^- \rightarrow q\bar{q}\ell\nu_\ell$ and $W^+ W^- \rightarrow q\bar{q}q\bar{q}$ events in the C.M. energy range 189–209 GeV. The result quoted here is obtained combining this mass value with the results obtained from a direct W mass reconstruction at 172 and 183 GeV and with those from the dependence of the WW production cross-section on m_{WW} at 161 and 172 GeV (ACCIARRI 99).
- ⁶ SCHAEEL 06 use direct reconstruction of the kinematics of $W^+ W^- \rightarrow q\bar{q}\ell\nu_\ell$ and $W^+ W^- \rightarrow q\bar{q}q\bar{q}$ events in the C.M. energy range 183–209 GeV. The result quoted here is obtained combining this mass value with those obtained from the dependence of the W pair production cross-section on m_{WW} at 161 and 172 GeV (BARATE 97 and BARATE 97S respectively). The systematic error includes ± 0.009 GeV due to possible effects of final state interactions in the $q\bar{q}q\bar{q}$ channel and ± 0.009 GeV due to the uncertainty on the LEP beam energy.
- ⁷ ABAZOV 02D improve the measurement of the W -boson mass including $W \rightarrow e\nu_e$ events in which the electron is close to a boundary of a central electromagnetic calorimeter module. Properly combining the results obtained by fitting $m_T(W)$, $p_T(e)$, and $p_T(\nu)$, this sample provides a mass value of 80.574 ± 0.405 GeV. The value reported here is a combination of this measurement with all previous $D\bar{D}$ W -boson mass measurements.
- ⁸ AFFOLDER 01E fit the transverse mass spectrum of 30115 $W \rightarrow e\nu_e$ events ($M_{WW} = 80.473 \pm 0.065 \pm 0.092$ GeV) and of 14740 $W \rightarrow \mu\nu_\mu$ events ($M_{WW} = 80.465 \pm 0.100 \pm 0.103$ GeV) obtained in the run IB (1994-95). Combining the electron and muon results, accounting for correlated uncertainties, yields $M_{WW} = 80.470 \pm 0.089$ GeV. They combine this value with their measurement of ABE 95P reported in run IA (1992-93) to obtain the quoted value.
- ⁹ AKTAS 06 fit the Q^2 dependence ($300 < Q^2 < 30,000$ GeV²) of the charged-current differential cross section with a propagator mass. The first error is experimental and the second corresponds to uncertainties due to input parameters and model assumptions.
- ¹⁰ CHEKANOV 02C fit the Q^2 dependence ($200 < Q^2 < 60000$ GeV²) of the charged-current differential cross sections with a propagator mass fit. The last error is due to the uncertainty on the probability density functions.
- ¹¹ BREITWEG 00D fit the Q^2 dependence ($200 < Q^2 < 22500$ GeV²) of the charged-current differential cross sections with a propagator mass fit. The last error is due to the uncertainty on the probability density functions.
- ¹² ALITTI 92B result has two contributions to the systematic error (± 0.83); one (± 0.81) cancels in m_W/m_Z and one (± 0.17) is noncancelling. These were added in quadrature. We choose the ALITTI 92B value without using the LEP m_Z value, because we perform our own combined fit.

- ¹³ There are two contributions to the systematic error (± 0.84): one (± 0.81) which cancels in m_W/m_Z and one (± 0.21) which is non-cancelling. These were added in quadrature.
¹⁴ ABE 89I systematic error dominated by the uncertainty in the absolute energy scale.
¹⁵ ALBAJAR 89 result is from a total sample of 299 $W \rightarrow e\nu$ events.
¹⁶ ALBAJAR 89 result is from a total sample of 67 $W \rightarrow \mu\nu$ events.
¹⁷ ALBAJAR 89 result is from $W \rightarrow \tau\nu$ events.

W/Z MASS RATIO

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|--------------------|----------------------|---------|-------------------------------|
| 0.8819 ± 0.0012 | OUR AVERAGE | | | |
| 0.8821 ± 0.0011 ± 0.0008 | 28323 | ¹⁸ ABBOTT | 98N D0 | $E_{cm}^{p\bar{p}} = 1.8$ TeV |
| 0.88114 ± 0.00154 ± 0.00252 | 5982 | ¹⁹ ABBOTT | 98P D0 | $E_{cm}^{p\bar{p}} = 1.8$ TeV |
| 0.8813 ± 0.0036 ± 0.0019 | 156 | ²⁰ ALITTI | 92B UA2 | $E_{cm}^{p\bar{p}} = 630$ GeV |

- ¹⁸ ABBOTT 98N obtain this from a study of 28323 $W \rightarrow e\nu_e$ and 3294 $Z \rightarrow e^+e^-$ decays. Of this latter sample, 2179 events are used to calibrate the electron energy scale.
¹⁹ ABBOTT 98P obtain this from a study of 5982 $W \rightarrow e\nu_e$ events. The systematic error includes an uncertainty of ± 0.00175 due to the electron energy scale.
²⁰ Scale error cancels in this ratio.

$m_Z - m_W$

| VALUE (GeV) | DOCUMENT ID | TECN | COMMENT |
|-------------------------|---|------|-----------------------------------|
| 10.4 ± 1.4 ± 0.8 | ALBAJAR 89 | UA1 | $E_{cm}^{p\bar{p}} = 546,630$ GeV |
| • • • | We do not use the following data for averages, fits, limits, etc. • • • | | |
| 11.3 ± 1.3 ± 0.9 | ANSARI 87 | UA2 | $E_{cm}^{p\bar{p}} = 546,630$ GeV |

$m_{W^+} - m_{W^-}$

Test of *CPT* invariance.

| VALUE (GeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------|------|-------------|---------|-------------------------------|
| -0.19 ± 0.58 | 1722 | ABE | 90G CDF | $E_{cm}^{p\bar{p}} = 1.8$ TeV |

W WIDTH

The W width listed here corresponds to the width parameter in a Breit-Wigner distribution with mass-dependent width. To obtain the world average, common systematic uncertainties between experiments are properly taken into account. The LEP-2 average W width based on published results is 2.196 ± 0.083 GeV [CERN-PH-EP/2006-042]. The combined Tevatron data yields an average W width of 2.046 ± 0.049 GeV [FERMILAB-TM-2460-E].

OUR FIT uses these average LEP and Tevatron width values and combines them assuming no correlations.

| <u>VALUE (GeV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--------------------------------|
| 2.085±0.042 OUR FIT | | | | |
| 2.028±0.072 | 5272 | 21 ABAZOV | 09AK D0 | $E_{cm}^{p\bar{p}} = 1.96$ GeV |
| 2.032±0.045±0.057 | 6055 | 22 AALTONEN | 08B CDF | $E_{cm}^{p\bar{p}} = 1.96$ TeV |
| 2.404±0.140±0.101 | 10.3k | 23 ABDALLAH | 08A DLPH | $E_{cm}^{ee} = 183$ –209 GeV |
| 1.996±0.096±0.102 | 10729 | 24 ABBIENDI | 06 OPAL | $E_{cm}^{ee} = 170$ –209 GeV |
| 2.18 ±0.11 ±0.09 | 9795 | 25 ACHARD | 06 L3 | $E_{cm}^{ee} = 172$ –209 GeV |
| 2.14 ±0.09 ±0.06 | 8717 | 26 SCHAEEL | 06 ALEP | $E_{cm}^{ee} = 183$ –209 GeV |
| 2.23 ^{+0.15} _{-0.14} ±0.10 | 294 | 27 ABAZOV | 02E D0 | Direct meas. |
| 2.05 ±0.10 ±0.08 | 662 | 28 AFFOLDER | 00M CDF | Direct meas. |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 2.152±0.066 | 79176 | 29 ABBOTT | 00B D0 | Extracted value |
| 2.064±0.060±0.059 | | 30 ABE | 95W CDF | Extracted value |
| 2.10 ^{+0.14} _{-0.13} ±0.09 | 3559 | 31 ALITTI | 92 UA2 | Extracted value |
| 2.18 ^{+0.26} _{-0.24} ±0.04 | | 32 ALBAJAR | 91 UA1 | Extracted value |

²¹ ABAZOV 09AK obtain this result fitting the high-end tail (100–200 GeV) of the transverse mass spectrum in $W \rightarrow e\nu$ decays.

²² AALTONEN 08B obtain this result fitting the high-end tail (90–200 GeV) of the transverse mass spectrum in semileptonic $W \rightarrow e\nu_e$ and $W \rightarrow \mu\nu_\mu$ decays.

²³ ABDALLAH 08A use direct reconstruction of the kinematics of $W^+ W^- \rightarrow q\bar{q}l\nu$ and $W^+ W^- \rightarrow q\bar{q}q\bar{q}$ events. The systematic error includes ± 0.065 GeV due to final state interactions.

²⁴ ABBIENDI 06 use direct reconstruction of the kinematics of $W^+ W^- \rightarrow q\bar{q}l\nu_\ell$ and $W^+ W^- \rightarrow q\bar{q}q\bar{q}$ events. The systematic error includes ± 0.003 GeV due to the uncertainty on the LEP beam energy.

²⁵ ACHARD 06 use direct reconstruction of the kinematics of $W^+ W^- \rightarrow q\bar{q}l\nu_\ell$ and $W^+ W^- \rightarrow q\bar{q}q\bar{q}$ events in the C.M. energy range 189–209 GeV. The result quoted here is obtained combining this value of the width with the result obtained from a direct W mass reconstruction at 172 and 183 GeV (ACCIARRI 99).

²⁶ SCHAEEL 06 use direct reconstruction of the kinematics of $W^+ W^- \rightarrow q\bar{q}l\nu_\ell$ and $W^+ W^- \rightarrow q\bar{q}q\bar{q}$ events. The systematic error includes ± 0.05 GeV due to possible effects of final state interactions in the $q\bar{q}q\bar{q}$ channel and ± 0.01 GeV due to the uncertainty on the LEP beam energy.

²⁷ ABAZOV 02E obtain this result fitting the high-end tail (90–200 GeV) of the transverse-mass spectrum in semileptonic $W \rightarrow e\nu_e$ decays.

²⁸ AFFOLDER 00M fit the high transverse mass (100–200 GeV) $W \rightarrow e\nu_e$ and $W \rightarrow \mu\nu_\mu$ events to obtain $\Gamma(W) = 2.04 \pm 0.11(\text{stat}) \pm 0.09(\text{syst})$ GeV. This is combined with the earlier CDF measurement (ABE 95C) to obtain the quoted result.

²⁹ ABBOTT 00B measure $R = 10.43 \pm 0.27$ for the $W \rightarrow e\nu_e$ decay channel. They use the SM theoretical predictions for $\sigma(W)/\sigma(Z)$ and $\Gamma(W \rightarrow e\nu_e)$ and the world average for $B(Z \rightarrow ee)$. The value quoted here is obtained combining this result (2.169 ± 0.070 GeV) with that of ABBOTT 99H.

³⁰ ABE 95W measured $R = 10.90 \pm 0.32 \pm 0.29$. They use $m_W = 80.23 \pm 0.18$ GeV, $\sigma(W)/\sigma(Z) = 3.35 \pm 0.03$, $\Gamma(W \rightarrow e\nu) = 225.9 \pm 0.9$ MeV, $\Gamma(Z \rightarrow e^+e^-) = 83.98 \pm 0.18$ MeV, and $\Gamma(Z) = 2.4969 \pm 0.0038$ GeV.

- ³¹ ALITTI 92 measured $R = 10.4_{-0.6}^{+0.7} \pm 0.3$. The values of $\sigma(Z)$ and $\sigma(W)$ come from $O(\alpha_s^2)$ calculations using $m_W = 80.14 \pm 0.27$ GeV, and $m_Z = 91.175 \pm 0.021$ GeV along with the corresponding value of $\sin^2\theta_W = 0.2274$. They use $\sigma(W)/\sigma(Z) = 3.26 \pm 0.07 \pm 0.05$ and $\Gamma(Z) = 2.487 \pm 0.010$ GeV.
- ³² ALBAJAR 91 measured $R = 9.5_{-1.0}^{+1.1}$ (stat. + syst.). $\sigma(W)/\sigma(Z)$ is calculated in QCD at the parton level using $m_W = 80.18 \pm 0.28$ GeV and $m_Z = 91.172 \pm 0.031$ GeV along with $\sin^2\theta_W = 0.2322 \pm 0.0014$. They use $\sigma(W)/\sigma(Z) = 3.23 \pm 0.05$ and $\Gamma(Z) = 2.498 \pm 0.020$ GeV. This measurement is obtained combining both the electron and muon channels.

W⁺ DECAY MODES

W^- modes are charge conjugates of the modes below.

| Mode | Fraction (Γ_i/Γ) | Confidence level |
|---------------------------|--------------------------------|------------------|
| Γ_1 $\ell^+ \nu$ | [a] $(10.80 \pm 0.09) \%$ | |
| Γ_2 $e^+ \nu$ | $(10.75 \pm 0.13) \%$ | |
| Γ_3 $\mu^+ \nu$ | $(10.57 \pm 0.15) \%$ | |
| Γ_4 $\tau^+ \nu$ | $(11.25 \pm 0.20) \%$ | |
| Γ_5 hadrons | $(67.60 \pm 0.27) \%$ | |
| Γ_6 $\pi^+ \gamma$ | $< 8 \times 10^{-5}$ | 95% |
| Γ_7 $D_s^+ \gamma$ | $< 1.3 \times 10^{-3}$ | 95% |
| Γ_8 cX | $(33.4 \pm 2.6) \%$ | |
| Γ_9 $c\bar{s}$ | $(31_{-11}^{+13}) \%$ | |
| Γ_{10} invisible | [b] $(1.4 \pm 2.9) \%$ | |

[a] ℓ indicates each type of lepton (e , μ , and τ), not sum over them.

[b] This represents the width for the decay of the W boson into a charged particle with momentum below detectability, $p < 200$ MeV.

W PARTIAL WIDTHS

$\Gamma(\text{invisible})$

Γ_{10}

This represents the width for the decay of the W boson into a charged particle with momentum below detectability, $p < 200$ MeV.

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-------------------------|----------------------|----------|--|
| $30_{-48}^{+52} \pm 33$ | ³³ BARATE | 99I ALEP | $E_{\text{cm}}^{ee} = 161+172+183$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | |
|----------------------|----------|--|
| ³⁴ BARATE | 99L ALEP | $E_{\text{cm}}^{ee} = 161+172+183$ GeV |
|----------------------|----------|--|

³³ BARATE 99I measure this quantity using the dependence of the total cross section σ_{WW} upon a change in the total width. The fit is performed to the WW measured cross sections at 161, 172, and 183 GeV. This partial width is < 139 MeV at 95%CL.

³⁴ BARATE 99L use W -pair production to search for effectively invisible W decays, tagging with the decay of the other W boson to Standard Model particles. The partial width for effectively invisible decay is < 27 MeV at 95%CL.

W BRANCHING RATIOS

Overall fits are performed to determine the branching ratios of the W . LEP averages on $W \rightarrow e\nu_e$, $W \rightarrow \mu\nu_\mu$, and $W \rightarrow \tau\nu_\tau$, and their correlations are first obtained by combining results from the four experiments taking properly into account the common systematics. The procedure is described in the note LEPEWWG/XSEC/2001-02, 30 March 2001, at <http://lepewwg.web.cern.ch/LEPEWWG/lepww/4f/PDG01>. The LEP average values so obtained, using published data, are given in the note LEPEWWG/XSEC/2005-01 accessible at <http://lepewwg.web.cern.ch/LEPEWWG/lepww/4f/PDG05/>. These results, together with results from the $p\bar{p}$ colliders are then used in fits to obtain the world average W branching ratios. A first fit determines three individual leptonic branching ratios, $B(W \rightarrow e\nu_e)$, $B(W \rightarrow \mu\nu_\mu)$, and $B(W \rightarrow \tau\nu_\tau)$. This fit has a $\chi^2 = 4.7$ for 10 degrees of freedom. A second fit assumes lepton universality and determines the leptonic branching ratio $B(W \rightarrow \ell\nu_\ell)$ and the hadronic branching ratio is derived as $B(W \rightarrow \text{hadrons}) = 1 - 3 B(W \rightarrow \ell\nu)$. This fit has a $\chi^2 = 11.3$ for 12 degrees of freedom.

The LEP $W \rightarrow \ell\nu$ data are obtained by the Collaborations using individual leptonic channels and are, therefore, not included in the overall fits to avoid double counting.

Note: The LEP combination including the new OPAL results, ABBI-ENDI 07A, could not be performed in time for this *Review*. Thus, the OUR FIT values quoted below use the previous OPAL results as in ABBI-ENDI,G 00.

$\Gamma(\ell^+\nu)/\Gamma_{\text{total}}$

ℓ indicates average over e , μ , and τ modes, not sum over modes.

Γ_1/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------|----------------------|------|---|
| 10.80 ± 0.09 OUR FIT | | | | |
| 10.86 ± 0.12 ± 0.08 | 16438 | ABBIENDI | 07A | OPAL $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 10.85 ± 0.14 ± 0.08 | 13600 | ABDALLAH | 04G | DLPH $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 10.83 ± 0.14 ± 0.10 | 11246 | ACHARD | 04J | L3 $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 10.96 ± 0.12 ± 0.05 | 16116 | SCHAEL | 04A | ALEP $E_{\text{cm}}^{ee} = 183\text{--}209$ GeV |
| 11.02 ± 0.52 | 11858 | ³⁵ ABBOTT | 99H | D0 $E_{\text{cm}}^{pp} = 1.8$ TeV |
| 10.4 ± 0.8 | 3642 | ³⁶ ABE | 92I | CDF $E_{\text{cm}}^{pp} = 1.8$ TeV |

³⁵ ABBOTT 99H measure $R \equiv [\sigma_W B(W \rightarrow \ell\nu_\ell)] / [\sigma_Z B(Z \rightarrow \ell\ell)] = 10.90 \pm 0.52$ combining electron and muon channels. They use $M_W = 80.39 \pm 0.06$ GeV and the SM theoretical predictions for $\sigma(W)/\sigma(Z)$ and $B(Z \rightarrow \ell\ell)$.

³⁶ $1216 \pm 38^{+27}_{-31}$ $W \rightarrow \mu\nu$ events from ABE 92I and $2426 W \rightarrow e\nu$ events of ABE 91C. ABE 92I give the inverse quantity as 9.6 ± 0.7 and we have inverted.

$\Gamma(e^+ \nu) / \Gamma_{\text{total}}$ **Γ_2 / Γ**

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

10.75 ± 0.13 OUR FIT

| | | | | | |
|---------------------|------|----------|-----|------|--|
| 10.71 ± 0.25 ± 0.11 | 2374 | ABBIENDI | 07A | OPAL | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 10.55 ± 0.31 ± 0.14 | 1804 | ABDALLAH | 04G | DLPH | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 10.78 ± 0.29 ± 0.13 | 1576 | ACHARD | 04J | L3 | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 10.78 ± 0.27 ± 0.10 | 2142 | SCHAEL | 04A | ALEP | $E_{\text{cm}}^{ee} = 183\text{--}209$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------|---------------|--------|-----|------|--------------------------------------|
| 10.61 ± 0.28 | ³⁷ | ABAZOV | 04D | TEVA | $E_{\text{cm}}^{p\bar{p}} = 1.8$ TeV |
|--------------|---------------|--------|-----|------|--------------------------------------|

³⁷ ABAZOV 04D take into account all correlations to properly combine the CDF (ABE 95W) and DØ (ABBOTT 00B) measurements of the ratio R in the electron channel. The ratio R is defined as $[\sigma_W \cdot B(W \rightarrow e\nu_e)] / [\sigma_Z \cdot B(Z \rightarrow ee)]$. The combination gives $R^{\text{Tevatron}} = 10.59 \pm 0.23$. σ_W / σ_Z is calculated at next-to-next-to-leading order (3.360 ± 0.051). The branching fraction $B(Z \rightarrow ee)$ is taken from this Review as (3.363 ± 0.004)%.

$\Gamma(\mu^+ \nu) / \Gamma_{\text{total}}$ **Γ_3 / Γ**

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

10.57 ± 0.15 OUR FIT

| | | | | | |
|---------------------|------|----------|-----|------|--|
| 10.78 ± 0.24 ± 0.10 | 2397 | ABBIENDI | 07A | OPAL | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 10.65 ± 0.26 ± 0.08 | 1998 | ABDALLAH | 04G | DLPH | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 10.03 ± 0.29 ± 0.12 | 1423 | ACHARD | 04J | L3 | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 10.87 ± 0.25 ± 0.08 | 2216 | SCHAEL | 04A | ALEP | $E_{\text{cm}}^{ee} = 183\text{--}209$ GeV |

$\Gamma(\tau^+ \nu) / \Gamma_{\text{total}}$ **Γ_4 / Γ**

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

11.25 ± 0.20 OUR FIT

| | | | | | |
|---------------------|------|----------|-----|------|--|
| 11.14 ± 0.31 ± 0.17 | 2177 | ABBIENDI | 07A | OPAL | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 11.46 ± 0.39 ± 0.19 | 2034 | ABDALLAH | 04G | DLPH | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 11.89 ± 0.40 ± 0.20 | 1375 | ACHARD | 04J | L3 | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 11.25 ± 0.32 ± 0.20 | 2070 | SCHAEL | 04A | ALEP | $E_{\text{cm}}^{ee} = 183\text{--}209$ GeV |

$\Gamma(\text{hadrons}) / \Gamma_{\text{total}}$ **Γ_5 / Γ**

OUR FIT value is obtained by a fit to the lepton branching ratio data assuming lepton universality.

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

67.60 ± 0.27 OUR FIT

| | | | | | |
|---------------------|-------|----------|-----|------|--|
| 67.41 ± 0.37 ± 0.23 | 16438 | ABBIENDI | 07A | OPAL | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 67.45 ± 0.41 ± 0.24 | 13600 | ABDALLAH | 04G | DLPH | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 67.50 ± 0.42 ± 0.30 | 11246 | ACHARD | 04J | L3 | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| 67.13 ± 0.37 ± 0.15 | 16116 | SCHAEL | 04A | ALEP | $E_{\text{cm}}^{ee} = 183\text{--}209$ GeV |

$\Gamma(\mu^+\nu)/\Gamma(e^+\nu)$ Γ_3/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|-------------|--------------------|-------------|----------------|
|--------------|-------------|--------------------|-------------|----------------|

0.983±0.018 OUR FIT

| | | | | |
|------------------|------|----------------------|---------|---|
| 0.89 ±0.10 | 13k | ³⁸ ABACHI | 95D D0 | $E_{cm}^{p\bar{p}} = 1.8 \text{ TeV}$ |
| 1.02 ±0.08 | 1216 | ³⁹ ABE | 92I CDF | $E_{cm}^{p\bar{p}} = 1.8 \text{ TeV}$ |
| 1.00 ±0.14 ±0.08 | 67 | ALBAJAR | 89 UA1 | $E_{cm}^{p\bar{p}} = 546,630 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------------------------|----|---------|---------|---------------------|
| 1.24 ^{+0.6} _{-0.4} | 14 | ARNISON | 84D UA1 | Repl. by ALBAJAR 89 |
|--------------------------------------|----|---------|---------|---------------------|

³⁸ ABACHI 95D obtain this result from the measured $\sigma_W B(W \rightarrow \mu\nu) = 2.09 \pm 0.23 \pm 0.11 \text{ nb}$ and $\sigma_W B(W \rightarrow e\nu) = 2.36 \pm 0.07 \pm 0.13 \text{ nb}$ in which the first error is the combined statistical and systematic uncertainty, the second reflects the uncertainty in the luminosity.

³⁹ ABE 92I obtain $\sigma_W B(W \rightarrow \mu\nu) = 2.21 \pm 0.07 \pm 0.21$ and combine with ABE 91C $\sigma_W B(W \rightarrow e\nu)$ to give a ratio of the couplings from which we derive this measurement.

$\Gamma(\tau^+\nu)/\Gamma(e^+\nu)$ Γ_4/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|-------------|--------------------|-------------|----------------|
|--------------|-------------|--------------------|-------------|----------------|

1.046±0.023 OUR FIT

| | | | | |
|------------------|-----|----------------------|---------|---|
| 0.961±0.061 | 980 | ⁴⁰ ABBOTT | 00D D0 | $E_{cm}^{p\bar{p}} = 1.8 \text{ TeV}$ |
| 0.94 ±0.14 | 179 | ⁴¹ ABE | 92E CDF | $E_{cm}^{p\bar{p}} = 1.8 \text{ TeV}$ |
| 1.04 ±0.08 ±0.08 | 754 | ⁴² ALITTI | 92F UA2 | $E_{cm}^{p\bar{p}} = 630 \text{ GeV}$ |
| 1.02 ±0.20 ±0.12 | 32 | ALBAJAR | 89 UA1 | $E_{cm}^{p\bar{p}} = 546,630 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------|-----|---------|---------|---------------------|
| 0.995±0.112±0.083 | 198 | ALITTI | 91C UA2 | Repl. by ALITTI 92F |
| 1.02 ±0.20 ±0.10 | 32 | ALBAJAR | 87 UA1 | Repl. by ALBAJAR 89 |

⁴⁰ ABBOTT 00D measure $\sigma_W \times B(W \rightarrow \tau\nu_\tau) = 2.22 \pm 0.09 \pm 0.10 \pm 0.10 \text{ nb}$. Using the ABBOTT 00B result $\sigma_W \times B(W \rightarrow e\nu_e) = 2.31 \pm 0.01 \pm 0.05 \pm 0.10 \text{ nb}$, they quote the ratio of the couplings from which we derive this measurement.

⁴¹ ABE 92E use two procedures for selecting $W \rightarrow \tau\nu_\tau$ events. The missing E_T trigger leads to $132 \pm 14 \pm 8$ events and the τ trigger to $47 \pm 9 \pm 4$ events. Proper statistical and systematic correlations are taken into account to arrive at $\sigma B(W \rightarrow \tau\nu) = 2.05 \pm 0.27 \text{ nb}$. Combined with ABE 91C result on $\sigma B(W \rightarrow e\nu)$, ABE 92E quote a ratio of the couplings from which we derive this measurement.

⁴² This measurement is derived by us from the ratio of the couplings of ALITTI 92F.

$\Gamma(\pi^+\gamma)/\Gamma(e^+\nu)$ Γ_6/Γ_2

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|----------------|
|--------------|------------|--------------------|-------------|----------------|

| | | | | |
|------------------------|----|-----------------------|---------|--|
| $< 7 \times 10^{-4}$ | 95 | ABE | 98H CDF | $E_{cm}^{p\bar{p}} = 1.8 \text{ TeV}$ |
| $< 4.9 \times 10^{-3}$ | 95 | ⁴³ ALITTI | 92D UA2 | $E_{cm}^{p\bar{p}} = 630 \text{ GeV}$ |
| $< 58 \times 10^{-3}$ | 95 | ⁴⁴ ALBAJAR | 90 UA1 | $E_{cm}^{p\bar{p}} = 546, 630 \text{ GeV}$ |

⁴³ ALITTI 92D limit is 3.8×10^{-3} at 90%CL.

⁴⁴ ALBAJAR 90 obtain < 0.048 at 90%CL.

$\Gamma(D_s^+\gamma)/\Gamma(e^+\nu)$ Γ_7/Γ_2

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|----------------|
|--------------|------------|--------------------|-------------|----------------|

| | | | | |
|------------------------|----|-----|---------|---------------------------------------|
| $< 1.2 \times 10^{-2}$ | 95 | ABE | 98P CDF | $E_{cm}^{p\bar{p}} = 1.8 \text{ TeV}$ |
|------------------------|----|-----|---------|---------------------------------------|

$\Gamma(cX)/\Gamma(\text{hadrons})$ **Γ_8/Γ_5**

| <u>VALUE</u> | <u>EVT5</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|------------------------|-------------|--|
| 0.49 ± 0.04 | OUR AVERAGE | | | |
| 0.481 ± 0.042 ± 0.032 | 3005 | ⁴⁵ ABBIENDI | 00V OPAL | $E_{\text{cm}}^{ee} = 183 + 189 \text{ GeV}$ |
| 0.51 ± 0.05 ± 0.03 | 746 | ⁴⁶ BARATE | 99M ALEP | $E_{\text{cm}}^{ee} = 172 + 183 \text{ GeV}$ |

⁴⁵ ABBIENDI 00V tag $W \rightarrow cX$ decays using measured jet properties, lifetime information, and leptons produced in charm decays. From this result, and using the additional measurements of $\Gamma(W)$ and $B(W \rightarrow \text{hadrons})$, $|V_{cs}|$ is determined to be $0.969 \pm 0.045 \pm 0.036$.

⁴⁶ BARATE 99M tag c jets using a neural network algorithm. From this measurement $|V_{cs}|$ is determined to be $1.00 \pm 0.11 \pm 0.07$.

$R_{cs} = \Gamma(c\bar{s})/\Gamma(\text{hadrons})$ **Γ_9/Γ_5**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|---------------------|-------------|--|
| 0.46^{+0.18}_{-0.14} ± 0.07 | ⁴⁷ ABREU | 98N DLPH | $E_{\text{cm}}^{ee} = 161+172 \text{ GeV}$ |

⁴⁷ ABREU 98N tag c and s jets by identifying a charged kaon as the highest momentum particle in a hadronic jet. They also use a lifetime tag to independently identify a c jet, based on the impact parameter distribution of charged particles in a jet. From this measurement $|V_{cs}|$ is determined to be $0.94^{+0.32}_{-0.26} \pm 0.13$.

AVERAGE PARTICLE MULTIPLICITIES IN HADRONIC W DECAY

Summed over particle and antiparticle, when appropriate.

$\langle N_{\pi^\pm} \rangle$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------|-----------------------|-------------|--|
| 15.70 ± 0.35 | ⁴⁸ ABREU,P | 00F DLPH | $E_{\text{cm}}^{ee} = 189 \text{ GeV}$ |

⁴⁸ ABREU,P 00F measure $\langle N_{\pi^\pm} \rangle = 31.65 \pm 0.48 \pm 0.76$ and $15.51 \pm 0.38 \pm 0.40$ in the fully hadronic and semileptonic final states respectively. The value quoted is a weighted average without assuming any correlations.

$\langle N_{K^\pm} \rangle$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|-----------------------|-------------|--|
| 2.20 ± 0.19 | ⁴⁹ ABREU,P | 00F DLPH | $E_{\text{cm}}^{ee} = 189 \text{ GeV}$ |

⁴⁹ ABREU,P 00F measure $\langle N_{K^\pm} \rangle = 4.38 \pm 0.42 \pm 0.12$ and $2.23 \pm 0.32 \pm 0.17$ in the fully hadronic and semileptonic final states respectively. The value quoted is a weighted average without assuming any correlations.

$\langle N_p \rangle$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|-----------------------|-------------|--|
| 0.92 ± 0.14 | ⁵⁰ ABREU,P | 00F DLPH | $E_{\text{cm}}^{ee} = 189 \text{ GeV}$ |

⁵⁰ ABREU,P 00F measure $\langle N_p \rangle = 1.82 \pm 0.29 \pm 0.16$ and $0.94 \pm 0.23 \pm 0.06$ in the fully hadronic and semileptonic final states respectively. The value quoted is a weighted average without assuming any correlations.

$\langle N_{\text{charged}} \rangle$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|--------------------|-------------|--|
| 19.39 ± 0.08 OUR AVERAGE | | | |
| 19.38 ± 0.05 ± 0.08 | 51 ABBIENDI | 06A OPAL | $E_{\text{cm}}^{ee} = 189\text{--}209$ GeV |
| 19.44 ± 0.17 | 52 ABREU,P | 00F DLPH | $E_{\text{cm}}^{ee} = 183\text{+}189$ GeV |
| 19.3 ± 0.3 ± 0.3 | 53 ABBIENDI | 99N OPAL | $E_{\text{cm}}^{ee} = 183$ GeV |
| 19.23 ± 0.74 | 54 ABREU | 98C DLPH | $E_{\text{cm}}^{ee} = 172$ GeV |

⁵¹ ABBIENDI 06A measure $\langle N_{\text{charged}} \rangle = 38.74 \pm 0.12 \pm 0.26$ when both W bosons decay hadronically and $\langle N_{\text{charged}} \rangle = 19.39 \pm 0.11 \pm 0.09$ when one W boson decays semileptonically. The value quoted here is obtained under the assumption that there is no color reconnection between W bosons; the value is a weighted average taking into account correlations in the systematic uncertainties.

⁵² ABREU,P 00F measure $\langle N_{\text{charged}} \rangle = 39.12 \pm 0.33 \pm 0.36$ and $38.11 \pm 0.57 \pm 0.44$ in the fully hadronic final states at 189 and 183 GeV respectively, and $\langle N_{\text{charged}} \rangle = 19.49 \pm 0.31 \pm 0.27$ and $19.78 \pm 0.49 \pm 0.43$ in the semileptonic final states. The value quoted is a weighted average without assuming any correlations.

⁵³ ABBIENDI 99N use the final states $W^+ W^- \rightarrow q\bar{q}\ell\bar{\nu}_\ell$ to derive this value.

⁵⁴ ABREU 98C combine results from both the fully hadronic as well semileptonic $W W$ final states after demonstrating that the W decay charged multiplicity is independent of the topology within errors.

TRIPLE GAUGE COUPLINGS (TGC'S)

A REVIEW GOES HERE – Check our WWW List of Reviews

g_1^Z

OUR FIT below is obtained by combining the measurements taking into account properly the common systematic errors (see LEPEWWG/TGC/2005-01 at <http://lepewwg.web.cern.ch/LEPEWWG/lepww/tgc>).

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| 0.984^{+0.022}_{-0.019} OUR FIT | | | | |
| 1.07 ^{+0.08} _{-0.12} | 1880 | 55 ABDALLAH | 08C DLPH | $E_{\text{cm}}^{ee} = 189\text{--}209$ GeV |
| 1.001 ± 0.027 ± 0.013 | 9310 | 56 SCHAEEL | 05A ALEP | $E_{\text{cm}}^{ee} = 183\text{--}209$ GeV |
| 0.987 ^{+0.034} _{-0.033} | 9800 | 57 ABBIENDI | 04D OPAL | $E_{\text{cm}}^{ee} = 183\text{--}209$ GeV |
| 0.966 ^{+0.034} _{-0.032} ± 0.015 | 8325 | 58 ACHARD | 04D L3 | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 1.04 ± 0.09 | | 59 ABAZOV | 09AD D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |
| | | 60 ABAZOV | 09AJ D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |
| | 13 | 61 ABAZOV | 07Z D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |
| | 2.3 | 62 ABAZOV | 05S D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |
| 0.98 ± 0.07 ± 0.01 | 2114 | 63 ABREU | 01I DLPH | $E_{\text{cm}}^{ee} = 183\text{+}189$ GeV |
| | 331 | 64 ABBOTT | 99I D0 | $E_{\text{cm}}^{p\bar{p}} = 1.8$ TeV |

- 55 ABDALLAH 08C determine this triple gauge coupling from the measurement of the spin density matrix elements in $e^+e^- \rightarrow W^+W^- \rightarrow (qq)(\ell\nu)$, where $\ell = e$ or μ . Values of all other couplings are fixed to their standard model values.
- 56 SCHAEEL 05A study single-photon, single- W , and WW -pair production from 183 to 209 GeV. The result quoted here is derived from the WW -pair production sample. Each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values.
- 57 ABBIENDI 04D combine results from W^+W^- in all decay channels. Only CP -conserving couplings are considered and each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values. The 95% confidence interval is $0.923 < g_1^Z < 1.054$.
- 58 ACHARD 04D study WW -pair production, single- W production and single-photon production with missing energy from 189 to 209 GeV. The result quoted here is obtained from the WW -pair production sample including data from 161 to 183 GeV, ACCIARRI 99Q. Each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values.
- 59 ABAZOV 09AD study the $p\bar{p} \rightarrow \ell\nu 2\text{jet}$ process arising in WW and WZ production. They select 12,473 (14,392) events in the electron (muon) channel with an expected di-boson signal of 436 (527) events. The results on the anomalous couplings are derived from an analysis of the p_T spectrum of the 2-jet system and quoted at 68% C.L. and for a form factor of 2 TeV. This measurement is not used for obtaining the mean as it is for a specific form factor.
- 60 ABAZOV 09AJ study the $p\bar{p} \rightarrow 2\ell 2\nu$ process arising in WW production. They select 100 events with an expected WW signal of 65 events. An analysis of the p_T spectrum of the two charged leptons leads to 95% C.L. limits of $0.86 < g_1^Z < 1.3$, for a form factor $\Lambda = 2$ TeV.
- 61 ABAZOV 07Z set limits on anomalous TGCs using the measured cross section and $p_T(Z)$ distribution in WZ production with both the W and the Z decaying leptonically into electrons and muons. Setting other couplings to their standard model values, the 95% C.L. limits for a form factor scale $\Lambda = 1.5$ TeV are $-0.15 < \Delta g_1^Z < 0.35$, and for $\Lambda = 2$ TeV are $-0.14 < \Delta g_1^Z < 0.34$.
- 62 ABAZOV 05S study $p\bar{p} \rightarrow WZ$ production with a subsequent trilepton decay to $\ell\nu\ell'\bar{\ell}'$ (ℓ and $\ell' = e$ or μ). Three events (estimated background 0.71 ± 0.08 events) with WZ decay characteristics are observed from which they derive limits on the anomalous WWZ couplings. The 95% CL limit for a form factor scale $\Lambda = 1.5$ TeV is $0.51 < g_1^Z < 1.66$, fixing λ_Z and κ_Z to their Standard Model values.
- 63 ABREU 01I combine results from e^+e^- interactions at 189 GeV leading to W^+W^- and $W e\nu_e$ final states with results from ABREU 99L at 183 GeV. The 95% confidence interval is $0.84 < g_1^Z < 1.13$.
- 64 ABBOTT 99I perform a simultaneous fit to the $W\gamma$, $WW \rightarrow$ dilepton, $WW/WZ \rightarrow e\nu jj$, $WW/WZ \rightarrow \mu\nu jj$, and $WZ \rightarrow$ trilepton data samples. For $\Lambda = 2.0$ TeV, the 95%CL limits are $0.63 < g_1^Z < 1.57$, fixing λ_Z and κ_Z to their Standard Model values, and assuming Standard Model values for the $WW\gamma$ couplings.

κ_γ

OUR FIT below is obtained by combining the measurements taking into account properly the common systematic errors (see LEPEWWG/TGC/2005-01 at <http://lepewwg.web.cern.ch/LEPEWWG/lepww/tgc>).

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|-------------|----------|---|
| 0.973^{+0.044}_{-0.045} | | | | OUR FIT |
| 0.68 ^{+0.17} _{-0.15} | 1880 | 65 ABDALLAH | 08C DLPH | $E_{cm}^{ee} = 189\text{--}209$ GeV |
| 0.971 \pm 0.055 \pm 0.030 | 10689 | 66 SCHAEEL | 05A ALEP | $E_{cm}^{ee} = 183\text{--}209$ GeV |
| 0.88 ^{+0.09} _{-0.08} | 9800 | 67 ABBIENDI | 04D OPAL | $E_{cm}^{ee} = 183\text{--}209$ GeV |
| 1.013 ^{+0.067} _{-0.064} \pm 0.026 | 10575 | 68 ACHARD | 04D L3 | $E_{cm}^{ee} = 161\text{--}209$ GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| | 53 | 69 AARON | 09B H1 | $E_{cm}^{ep} = 0.3$ TeV |
| 1.07 ^{+0.26} _{-0.29} | | 70 ABAZOV | 09AD D0 | $E_{cm}^{p\bar{p}} = 1.96$ TeV |
| | | 71 ABAZOV | 09AJ D0 | $E_{cm}^{p\bar{p}} = 1.96$ TeV |
| | | 72 ABAZOV | 08R D0 | $E_{cm}^{p\bar{p}} = 1.96$ TeV |
| | 1617 | 73 AALTONEN | 07L CDF | $E_{cm}^{p\bar{p}} = 1.96$ GeV |
| | 17 | 74 ABAZOV | 06H D0 | $E_{cm}^{p\bar{p}} = 1.96$ TeV |
| | 141 | 75 ABAZOV | 05J D0 | $E_{cm}^{p\bar{p}} = 1.96$ TeV |
| 1.25 ^{+0.21} _{-0.20} \pm 0.06 | 2298 | 76 ABREU | 01I DLPH | $E_{cm}^{ee} = 183\text{+}189$ GeV |
| | | 77 BREITWEG | 00 ZEUS | $e^+ p \rightarrow e^+ W^\pm X$, $\sqrt{s} \approx 300$ GeV |
| 0.92 \pm 0.34 | 331 | 78 ABBOTT | 99I D0 | $E_{cm}^{p\bar{p}} = 1.8$ TeV |

⁶⁵ ABDALLAH 08C determine this triple gauge coupling from the measurement of the spin density matrix elements in $e^+ e^- \rightarrow W^+ W^- \rightarrow (qq)(\ell\nu)$, where $\ell = e$ or μ . Values of all other couplings are fixed to their standard model values.

⁶⁶ SCHAEEL 05A study single-photon, single- W , and WW -pair production from 183 to 209 GeV. Each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values.

⁶⁷ ABBIENDI 04D combine results from $W^+ W^-$ in all decay channels. Only CP -conserving couplings are considered and each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values. The 95% confidence interval is $0.73 < \kappa_\gamma < 1.07$.

⁶⁸ ACHARD 04D study WW -pair production, single- W production and single-photon production with missing energy from 189 to 209 GeV. The result quoted here is obtained including data from 161 to 183 GeV, ACCIARRI 99Q. Each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values.

⁶⁹ AARON 09B study single- W production in ep collisions at 0.3 TeV C.M. energy. They select 53 $W \rightarrow e/\mu$ events with a standard model expectation of 54.1 ± 7.4 events. Fitting the transverse momentum spectrum of the hadronic recoil system they obtain a 95% C.L. limit of $-3.7 < \kappa_\gamma < -1.5$ or $0.3 < \kappa_\gamma < 1.5$, where the ambiguity is due to the quadratic dependence of the cross section to the coupling parameter.

⁷⁰ ABAZOV 09AD study the $p\bar{p} \rightarrow \ell\nu$ 2jet process arising in WW and WZ production. They select 12,473 (14,392) events in the electron (muon) channel with an expected di-boson signal of 436 (527) events. The results on the anomalous couplings are derived from an analysis of the p_T spectrum of the 2-jet system and quoted at 68% C.L. and for a form factor of 2 TeV. This measurement is not used for obtaining the mean as it is for a specific form factor.

- 71 ABAZOV 09AJ study the $p\bar{p} \rightarrow 2\ell 2\nu$ process arising in WW production. They select 100 events with an expected WW signal of 65 events. An analysis of the p_T spectrum of the two charged leptons leads to 95% C.L. limits of $0.46 < \kappa_\gamma < 1.83$, for a form factor $\Lambda = 2$ TeV.
- 72 ABAZOV 08R use 0.7 fb^{-1} $p\bar{p}$ data at 1.96 TeV to select 263 $W\gamma + X$ events, of which 187 constitute signal, with the W decaying into an electron or a muon, which is required to be well separated from a photon with $E_T > 9$ GeV. A likelihood fit to the photon E_T spectrum yields a 95% CL limit $0.49 < \kappa_\gamma < 1.51$ with other couplings fixed to their Standard Model values.
- 73 AALTONEN 07L set limits on anomalous TGCs using the $p_T(W)$ distribution in WW and WZ production with the W decaying to an electron or muon and the Z to 2 jets. Setting other couplings to their standard model value, the 95% C.L. limits are $0.54 < \kappa_\gamma < 1.39$ for a form factor scale $\Lambda = 1.5$ TeV.
- 74 ABAZOV 06H study $p\bar{p} \rightarrow WW$ production with a subsequent decay $WW \rightarrow e^+\nu_e e^-\bar{\nu}_e$, $WW \rightarrow e^\pm\nu_e e^\mp\nu_\mu$ or $WW \rightarrow \mu^+\nu_\mu\mu^-\bar{\nu}_\mu$. The 95% C.L. limit for a form factor scale $\Lambda = 1$ TeV is $-0.05 < \kappa_\gamma < 2.29$, fixing $\lambda_\gamma=0$. With the assumption that the $WW\gamma$ and WWZ couplings are equal the 95% C.L. one-dimensional limit ($\Lambda = 2$ TeV) is $0.68 < \kappa < 1.45$.
- 75 ABAZOV 05J perform a likelihood fit to the photon E_T spectrum of $W\gamma + X$ events, where the W decays to an electron or muon which is required to be well separated from the photon. For $\Lambda = 2.0$ TeV the 95% CL limits are $0.12 < \kappa_\gamma < 1.96$. In the fit λ_γ is kept fixed to its Standard Model value.
- 76 ABREU 01I combine results from e^+e^- interactions at 189 GeV leading to W^+W^- , $W e\nu_e$, and $\nu\bar{\nu}\gamma$ final states with results from ABREU 99L at 183 GeV. The 95% confidence interval is $0.87 < \kappa_\gamma < 1.68$.
- 77 BREITWEG 00 search for W production in events with large hadronic p_T . For $p_T > 20$ GeV, the upper limit on the cross section gives the 95%CL limit $-3.7 < \kappa_\gamma < 2.5$ (for $\lambda_\gamma=0$).
- 78 ABBOTT 99I perform a simultaneous fit to the $W\gamma$, $WW \rightarrow$ dilepton, $WW/WZ \rightarrow e\nu jj$, $WW/WZ \rightarrow \mu\nu jj$, and $WZ \rightarrow$ trilepton data samples. For $\Lambda = 2.0$ TeV, the 95%CL limits are $0.75 < \kappa_\gamma < 1.39$.

λ_γ

OUR FIT below is obtained by combining the measurements taking into account properly the common systematic errors (see LEPEWWG/TGC/2005-01 at <http://lepewwg.web.cern.ch/LEPEWWG/lepww/tgc>).

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| $-0.028^{+0.020}_{-0.021}$ OUR FIT | | | | |
| $0.16^{+0.12}_{-0.13}$ | 1880 | 79 ABDALLAH | 08C DLPH | $E_{\text{cm}}^{ee} = 189\text{--}209$ GeV |
| $-0.012 \pm 0.027 \pm 0.011$ | 10689 | 80 SCHAEEL | 05A ALEP | $E_{\text{cm}}^{ee} = 183\text{--}209$ GeV |
| $-0.060^{+0.034}_{-0.033}$ | 9800 | 81 ABBIENDI | 04D OPAL | $E_{\text{cm}}^{ee} = 183\text{--}209$ GeV |
| $-0.021^{+0.035}_{-0.034} \pm 0.017$ | 10575 | 82 ACHARD | 04D L3 | $E_{\text{cm}}^{ee} = 161\text{--}209$ GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| | 53 | 83 AARON | 09B H1 | $E_{\text{cm}}^{ep} = 0.3$ TeV |
| 0.00 ± 0.06 | | 84 ABAZOV | 09AD D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |
| | | 85 ABAZOV | 09AJ D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |

| | | | | | | |
|------|------------|------------|----------|-----|----------|---|
| | | 86 | ABAZOV | 08R | D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96 \text{ TeV}$ |
| | 1617 | 87 | AALTONEN | 07L | CDF | $E_{\text{cm}}^{p\bar{p}} = 1.96 \text{ GeV}$ |
| | 17 | 88 | ABAZOV | 06H | D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96 \text{ TeV}$ |
| | 141 | 89 | ABAZOV | 05J | D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96 \text{ TeV}$ |
| 0.05 | ± 0.09 | ± 0.01 | 2298 | 90 | ABREU | 01I DLPH $E_{\text{cm}}^{ee} = 183+189 \text{ GeV}$ |
| | | | | 91 | BREITWEG | 00 ZEUS $e^+ p \rightarrow e^+ W^\pm X$, $\sqrt{s} \approx 300 \text{ GeV}$ |
| 0.00 | $+0.10$ | | 331 | 92 | ABBOTT | 99I D0 $E_{\text{cm}}^{p\bar{p}} = 1.8 \text{ TeV}$ |
| | -0.09 | | | | | |

- 79 ABDALLAH 08C determine this triple gauge coupling from the measurement of the spin density matrix elements in $e^+e^- \rightarrow W^+W^- \rightarrow (qq)(\ell\nu)$, where $\ell = e$ or μ . Values of all other couplings are fixed to their standard model values.
- 80 SCHAEEL 05A study single-photon, single- W , and WW -pair production from 183 to 209 GeV. Each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values.
- 81 ABBIENDI 04D combine results from W^+W^- in all decay channels. Only CP -conserving couplings are considered and each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values. The 95% confidence interval is $-0.13 < \lambda_\gamma < 0.01$.
- 82 ACHARD 04D study WW -pair production, single- W production and single-photon production with missing energy from 189 to 209 GeV. The result quoted here is obtained including data from 161 to 183 GeV, ACCIARRI 99Q. Each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values.
- 83 AARON 09B study single- W production in ep collisions at 0.3 TeV C.M. energy. They select 53 $W \rightarrow e/\mu$ events with a standard model expectation of 54.1 ± 7.4 events. Fitting the transverse momentum spectrum of the hadronic recoil system they obtain a 95% C.L. limit of $-2.5 < \lambda_\gamma < 2.5$.
- 84 ABAZOV 09AD study the $p\bar{p} \rightarrow \ell\nu 2\text{jet}$ process arising in WW and WZ production. They select 12,473 (14,392) events in the electron (muon) channel with an expected di-boson signal of 436 (527) events. The results on the anomalous couplings are derived from an analysis of the p_T spectrum of the 2-jet system and quoted at 68% C.L. and for a form factor of 2 TeV. This measurement is not used for obtaining the mean as it is for a specific form factor.
- 85 ABAZOV 09AJ study the $p\bar{p} \rightarrow 2\ell 2\nu$ process arising in WW production. They select 100 events with an expected WW signal of 65 events. An analysis of the p_T spectrum of the two charged leptons leads to 95% C.L. limits of $-0.14 < \lambda_\gamma < 0.18$, for a form factor $\Lambda = 2 \text{ TeV}$.
- 86 ABAZOV 08R use $0.7 \text{ fb}^{-1} p\bar{p}$ data at 1.96 TeV to select 263 $W\gamma + X$ events, of which 187 constitute signal, with the W decaying into an electron or a muon, which is required to be well separated from a photon with $E_T > 9 \text{ GeV}$. A likelihood fit to the photon E_T spectrum yields a 95% CL limit $-0.12 < \lambda_\gamma < 0.13$ with other couplings fixed to their Standard Model values.
- 87 AALTONEN 07L set limits on anomalous TGCs using the $p_T(W)$ distribution in WW and WZ production with the W decaying to an electron or muon and the Z to 2 jets. Setting other couplings to their standard model value, the 95% C.L. limits are $-0.18 < \lambda_\gamma < 0.17$ for a form factor scale $\Lambda = 1.5 \text{ TeV}$.
- 88 ABAZOV 06H study $p\bar{p} \rightarrow WW$ production with a subsequent decay $WW \rightarrow e^+\nu_e e^-\bar{\nu}_e$, $WW \rightarrow e^\pm\nu_e\mu^\mp\nu_\mu$ or $WW \rightarrow \mu^+\nu_\mu\mu^-\bar{\nu}_\mu$. The 95% C.L. limit for a form factor scale $\Lambda = 1 \text{ TeV}$ is $-0.97 < \lambda_\gamma < 1.04$, fixing $\kappa_\gamma=1$. With the assumption that the $WW\gamma$ and WWZ couplings are equal the 95% C.L. one-dimensional limit ($\Lambda = 2 \text{ TeV}$) is $-0.29 < \lambda < 0.30$.

- 89 ABAZOV 05J perform a likelihood fit to the photon E_T spectrum of $W\gamma + X$ events, where the W decays to an electron or muon which is required to be well separated from the photon. For $\Lambda = 2.0$ TeV the 95% CL limits are $-0.20 < \lambda_\gamma < 0.20$. In the fit κ_γ is kept fixed to its Standard Model value.
- 90 ABREU 01I combine results from e^+e^- interactions at 189 GeV leading to W^+W^- , $W e \nu_e$, and $\nu \bar{\nu} \gamma$ final states with results from ABREU 99L at 183 GeV. The 95% confidence interval is $-0.11 < \lambda_\gamma < 0.23$.
- 91 BREITWEG 00 search for W production in events with large hadronic p_T . For $p_T > 20$ GeV, the upper limit on the cross section gives the 95%CL limit $-3.2 < \lambda_\gamma < 3.2$ for κ_γ fixed to its Standard Model value.
- 92 ABBOTT 99I perform a simultaneous fit to the $W\gamma$, $WW \rightarrow$ dilepton, $WW/WZ \rightarrow e\nu jj$, $WW/WZ \rightarrow \mu\nu jj$, and $WZ \rightarrow$ trilepton data samples. For $\Lambda = 2.0$ TeV, the 95%CL limits are $-0.18 < \lambda_\gamma < 0.19$.

κ_Z

This coupling is CP -conserving (C - and P - separately conserving).

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|--------|---|
| $0.924^{+0.059}_{-0.056} \pm 0.024$ | 7171 | 93 ACHARD | 04D L3 | $E_{\text{cm}}^{ee} = 189\text{--}209$ GeV |
| • • • | | | | We do not use the following data for averages, fits, limits, etc. • • • |
| | 17 | 94 ABAZOV | 06H D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |
| | 2.3 | 95 ABAZOV | 05S D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |

- 93 ACHARD 04D study WW -pair production, single- W production and single-photon production with missing energy from 189 to 209 GeV. The result quoted here is obtained using the WW -pair production sample. Each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values.
- 94 ABAZOV 06H study $\bar{p}p \rightarrow WW$ production with a subsequent decay $WW \rightarrow e^+ \nu_e e^- \bar{\nu}_e$, $WW \rightarrow e^\pm \nu_e \mu^\mp \nu_\mu$ or $WW \rightarrow \mu^+ \nu_\mu \mu^- \bar{\nu}_\mu$. The 95% C.L. limit for a form factor scale $\Lambda = 2$ TeV is $0.55 < \kappa_Z < 1.55$, fixing $\lambda_Z = 0$. With the assumption that the $WW\gamma$ and WWZ couplings are equal the 95% C.L. one-dimensional limit ($\Lambda = 2$ TeV) is $0.68 < \kappa < 1.45$.
- 95 ABAZOV 05S study $\bar{p}p \rightarrow WZ$ production with a subsequent trilepton decay to $\ell \nu \ell' \bar{\ell}'$ (ℓ and $\ell' = e$ or μ). Three events (estimated background 0.71 ± 0.08 events) with WZ decay characteristics are observed from which they derive limits on the anomalous WWZ couplings. The 95% CL limit for a form factor scale $\Lambda = 1$ TeV is $-1.0 < \kappa_Z < 3.4$, fixing λ_Z and g_1^Z to their Standard Model values.

λ_Z

This coupling is CP -conserving (C - and P - separately conserving).

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|--------|---|
| $-0.088^{+0.060}_{-0.057} \pm 0.023$ | 7171 | 96 ACHARD | 04D L3 | $E_{\text{cm}}^{ee} = 189\text{--}209$ GeV |
| • • • | | | | We do not use the following data for averages, fits, limits, etc. • • • |
| | 13 | 97 ABAZOV | 07Z D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |
| | 17 | 98 ABAZOV | 06H D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |
| | 2.3 | 99 ABAZOV | 05S D0 | $E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV |

- ⁹⁶ ACHARD 04D study WW -pair production, single- W production and single-photon production with missing energy from 189 to 209 GeV. The result quoted here is obtained using the WW -pair production sample. Each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values.
- ⁹⁷ ABAZOV 07Z set limits on anomalous TGCs using the measured cross section and $p_T(Z)$ distribution in WZ production with both the W and the Z decaying leptonically into electrons and muons. Setting other couplings to their standard model values, the 95% C.L. limits for a form factor scale $\Lambda = 1.5$ TeV are $-0.18 < \lambda_Z < 0.22$, and for $\Lambda = 2$ TeV are $-0.17 < \lambda_Z < 0.21$.
- ⁹⁸ ABAZOV 06H study $\bar{p}p \rightarrow WW$ production with a subsequent decay $WW \rightarrow e^+ \nu_e e^- \bar{\nu}_e$, $WW \rightarrow e^\pm \nu_e \mu^\mp \nu_\mu$ or $WW \rightarrow \mu^+ \nu_\mu \mu^- \bar{\nu}_\mu$. The 95% C.L. limit for a form factor scale $\Lambda = 2$ TeV is $-0.39 < \lambda_Z < 0.39$, fixing $\kappa_Z=1$. With the assumption that the $WW\gamma$ and WWZ couplings are equal the 95% C.L. one-dimensional limit ($\Lambda = 2$ TeV) is $-0.29 < \lambda < 0.30$.
- ⁹⁹ ABAZOV 05S study $\bar{p}p \rightarrow WZ$ production with a subsequent trilepton decay to $\ell\nu\ell'\bar{\ell}'$ (ℓ and $\ell' = e$ or μ). Three events (estimated background 0.71 ± 0.08 events) with WZ decay characteristics are observed from which they derive limits on the anomalous WWZ couplings. The 95% CL limit for a form factor scale $\Lambda = 1.5$ TeV is $-0.48 < \lambda_Z < 0.48$, fixing g_1^Z and κ_Z to their Standard Model values.

g_5^Z

This coupling is CP -conserving but C - and P -violating.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------------------|----------|----------------------------------|
| 0.93 ± 0.09 OUR AVERAGE | | Error includes scale factor of 1.1. | | |
| $0.96^{+0.13}_{-0.12}$ | 9800 | 100 ABBIENDI | 04D OPAL | $E_{cm}^{ee} = 183-209$ GeV |
| $1.00 \pm 0.13 \pm 0.05$ | 7171 | 101 ACHARD | 04D L3 | $E_{cm}^{ee} = 189-209$ GeV |
| $0.56^{+0.23}_{-0.22} \pm 0.12$ | 1154 | 102 ACCIARRI | 99Q L3 | $E_{cm}^{ee} = 161+172+ 183$ GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.84 ± 0.23 | | 103 EBOLI | 00 THEO | LEP1, SLC+ Tevatron |

- ¹⁰⁰ ABBIENDI 04D combine results from $W^+ W^-$ in all decay channels. Only CP -conserving couplings are considered and each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values. The 95% confidence interval is $0.72 < g_5^Z < 1.21$.
- ¹⁰¹ ACHARD 04D study WW -pair production, single- W production and single-photon production with missing energy from 189 to 209 GeV. The result quoted here is obtained using the WW -pair production sample. Each parameter is determined from a single-parameter fit in which the other parameters assume their Standard Model values.
- ¹⁰² ACCIARRI 99Q study W -pair, single- W , and single photon events.
- ¹⁰³ EBOLI 00 extract this indirect value of the coupling studying the non-universal one-loop contributions to the experimental value of the $Z \rightarrow b\bar{b}$ width ($\Lambda=1$ TeV is assumed).

g_4^Z

This coupling is CP -violating (C -violating and P -conserving).

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------|----------|-----------------------------|
| -0.30 ± 0.17 OUR AVERAGE | | | | |
| $-0.39^{+0.19}_{-0.20}$ | 1880 | 104 ABDALLAH | 08C DLPH | $E_{cm}^{ee} = 189-209$ GeV |
| $-0.02^{+0.32}_{-0.33}$ | 1065 | 105 ABBIENDI | 01H OPAL | $E_{cm}^{ee} = 189$ GeV |

- 104 ABDALLAH 08C determine this triple gauge coupling from the measurement of the spin density matrix elements in $e^+e^- \rightarrow W^+W^- \rightarrow (qq)(\ell\nu)$, where $\ell = e$ or μ . Values of all other couplings are fixed to their standard model values.
- 105 ABBIENDI 01H study W -pair events, with one leptonically and one hadronically decaying W . The coupling is extracted using information from the W production angle together with decay angles from the leptonically decaying W .

$\tilde{\kappa}_Z$

This coupling is CP -violating (C -conserving and P -violating).

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|------------------|------|--|
| $-0.12^{+0.06}_{-0.04}$ OUR AVERAGE | | | | |
| $-0.09^{+0.08}_{-0.05}$ | 1880 | 106 ABDALLAH 08C | DLPH | $E_{\text{cm}}^{ee} = 189\text{--}209$ GeV |
| $-0.20^{+0.10}_{-0.07}$ | 1065 | 107 ABBIENDI 01H | OPAL | $E_{\text{cm}}^{ee} = 189$ GeV |

- 106 ABDALLAH 08C determine this triple gauge coupling from the measurement of the spin density matrix elements in $e^+e^- \rightarrow W^+W^- \rightarrow (qq)(\ell\nu)$, where $\ell = e$ or μ . Values of all other couplings are fixed to their standard model values.
- 107 ABBIENDI 01H study W -pair events, with one leptonically and one hadronically decaying W . The coupling is extracted using information from the W production angle together with decay angles from the leptonically decaying W .

$\tilde{\lambda}_Z$

This coupling is CP -violating (C -conserving and P -violating).

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|------------------|------|--|
| -0.09 ± 0.07 OUR AVERAGE | | | | |
| -0.08 ± 0.07 | 1880 | 108 ABDALLAH 08C | DLPH | $E_{\text{cm}}^{ee} = 189\text{--}209$ GeV |
| $-0.18^{+0.24}_{-0.16}$ | 1065 | 109 ABBIENDI 01H | OPAL | $E_{\text{cm}}^{ee} = 189$ GeV |

- 108 ABDALLAH 08C determine this triple gauge coupling from the measurement of the spin density matrix elements in $e^+e^- \rightarrow W^+W^- \rightarrow (qq)(\ell\nu)$, where $\ell = e$ or μ . Values of all other couplings are fixed to their standard model values.
- 109 ABBIENDI 01H study W -pair events, with one leptonically and one hadronically decaying W . The coupling is extracted using information from the W production angle together with decay angles from the leptonically decaying W .

W ANOMALOUS MAGNETIC MOMENT

The full magnetic moment is given by $\mu_W = e(1 + \kappa + \lambda)/2m_W$. In the Standard Model, at tree level, $\kappa = 1$ and $\lambda = 0$. Some papers have defined $\Delta\kappa = 1 - \kappa$ and assume that $\lambda = 0$. Note that the electric quadrupole moment is given by $-e(\kappa - \lambda)/m_W^2$. A description of the parameterization of these moments and additional references can be found in HAGIWARA 87 and BAUR 88. The parameter Λ appearing in the theoretical limits below is a regularization cutoff which roughly corresponds to the energy scale where the structure of the W boson becomes manifest.

| VALUE ($e/2m_W$) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|---------------|------|--|
| $2.22^{+0.20}_{-0.19}$ | 2298 | 110 ABREU 01I | DLPH | $E_{\text{cm}}^{ee} = 183\text{--}189$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----|-----------|-----|------|
| 111 | ABE | 95G | CDF |
| 112 | ALITTI | 92C | UA2 |
| 113 | SAMUEL | 92 | THEO |
| 114 | SAMUEL | 91 | THEO |
| 115 | GRIFOLS | 88 | THEO |
| 116 | GROTCH | 87 | THEO |
| 117 | VANDERBIJ | 87 | THEO |
| 118 | GRAU | 85 | THEO |
| 119 | SUZUKI | 85 | THEO |
| 120 | HERZOG | 84 | THEO |

- 110 ABREU 01l combine results from e^+e^- interactions at 189 GeV leading to W^+W^- , $W e \nu_e$, and $\nu \bar{\nu} \gamma$ final states with results from ABREU 99L at 183 GeV to determine Δg_1^Z , $\Delta \kappa_\gamma$, and λ_γ . $\Delta \kappa_\gamma$ and λ_γ are simultaneously floated in the fit to determine μ_W .
- 111 ABE 95G report $-1.3 < \kappa < 3.2$ for $\lambda=0$ and $-0.7 < \lambda < 0.7$ for $\kappa=1$ in $p\bar{p} \rightarrow e\nu_e\gamma X$ and $\mu\nu_\mu\gamma X$ at $\sqrt{s} = 1.8$ TeV.
- 112 ALITTI 92C measure $\kappa = 1_{-2.2}^{+2.6}$ and $\lambda = 0_{-1.8}^{+1.7}$ in $p\bar{p} \rightarrow e\nu\gamma + X$ at $\sqrt{s} = 630$ GeV. At 95%CL they report $-3.5 < \kappa < 5.9$ and $-3.6 < \lambda < 3.5$.
- 113 SAMUEL 92 use preliminary CDF and UA2 data and find $-2.4 < \kappa < 3.7$ at 96%CL and $-3.1 < \kappa < 4.2$ at 95%CL respectively. They use data for $W\gamma$ production and radiative W decay.
- 114 SAMUEL 91 use preliminary CDF data for $p\bar{p} \rightarrow W\gamma X$ to obtain $-11.3 \leq \Delta\kappa \leq 10.9$. Note that their $\kappa = 1 - \Delta\kappa$.
- 115 GRIFOLS 88 uses deviation from ρ parameter to set limit $\Delta\kappa \lesssim 65 (M_W^2/\Lambda^2)$.
- 116 GROTCH 87 finds the limit $-37 < \Delta\kappa < 73.5$ (90% CL) from the experimental limits on $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ assuming three neutrino generations and $-19.5 < \Delta\kappa < 56$ for four generations. Note their $\Delta\kappa$ has the opposite sign as our definition.
- 117 VANDERBIJ 87 uses existing limits to the photon structure to obtain $|\Delta\kappa| < 33 (m_W/\Lambda)$. In addition VANDERBIJ 87 discusses problems with using the ρ parameter of the Standard Model to determine $\Delta\kappa$.
- 118 GRAU 85 uses the muon anomaly to derive a coupled limit on the anomalous magnetic dipole and electric quadrupole (λ) moments $1.05 > \Delta\kappa \ln(\Lambda/m_W) + \lambda/2 > -2.77$. In the Standard Model $\lambda = 0$.
- 119 SUZUKI 85 uses partial-wave unitarity at high energies to obtain $|\Delta\kappa| \lesssim 190 (m_W/\Lambda)^2$. From the anomalous magnetic moment of the muon, SUZUKI 85 obtains $|\Delta\kappa| \lesssim 2.2/\ln(\Lambda/m_W)$. Finally SUZUKI 85 uses deviations from the ρ parameter and obtains a very qualitative, order-of-magnitude limit $|\Delta\kappa| \lesssim 150 (m_W/\Lambda)^4$ if $|\Delta\kappa| \ll 1$.
- 120 HERZOG 84 consider the contribution of W -boson to muon magnetic moment including anomalous coupling of $WW\gamma$. Obtain a limit $-1 < \Delta\kappa < 3$ for $\Lambda \gtrsim 1$ TeV.

ANOMALOUS W/Z QUARTIC COUPLINGS

A REVIEW GOES HERE – Check our WWW List of Reviews

$a_0/\Lambda^2, a_c/\Lambda^2, a_n/\Lambda^2$

Using the $WW\gamma$ final state, the LEP combined 95% CL limits on the anomalous contributions to the $WW\gamma\gamma$ and $WWZ\gamma$ vertices (as of summer 2003) are given below:

(See P. Wells, "Experimental Tests of the Standard Model," Int. Europhysics Conference on High-Energy Physics, Aachen, Germany, 17–23 July 2003)

$$\begin{aligned}
 -0.02 < a_0^W/\Lambda^2 < 0.02 \text{ GeV}^{-2}, \\
 -0.05 < a_c^W/\Lambda^2 < 0.03 \text{ GeV}^{-2}, \\
 -0.15 < a_n/\Lambda^2 < 0.15 \text{ GeV}^{-2}.
 \end{aligned}$$

| VALUE | DOCUMENT ID | TECN |
|-------|-------------|------|
|-------|-------------|------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----|----------|-----|------|
| 121 | ABBIENDI | 04B | OPAL |
| 122 | ABBIENDI | 04L | OPAL |
| 123 | HEISTER | 04A | ALEP |
| 124 | ABDALLAH | 03I | DLPH |
| 125 | ACHARD | 02F | L3 |

- 121 ABBIENDI 04B select 187 $e^+e^- \rightarrow W^+W^-\gamma$ events in the C.M. energy range 180–209 GeV, where $E_\gamma > 2.5$ GeV, the photon has a polar angle $|\cos\theta_\gamma| < 0.975$ and is well isolated from the nearest jet and charged lepton, and the effective masses of both fermion-antifermion systems agree with the W mass within $3\Gamma_W$. The measured differential cross section as a function of the photon energy and photon polar angle is used to extract the 95% CL limits: $-0.020 \text{ GeV}^{-2} < a_0/\Lambda^2 < 0.020 \text{ GeV}^{-2}$, $-0.053 \text{ GeV}^{-2} < a_c/\Lambda^2 < 0.037 \text{ GeV}^{-2}$ and $-0.16 \text{ GeV}^{-2} < a_n/\Lambda^2 < 0.15 \text{ GeV}^{-2}$.
- 122 ABBIENDI 04L select 20 $e^+e^- \rightarrow \nu\bar{\nu}\gamma\gamma$ acoplanar events in the energy range 180–209 GeV and 176 $e^+e^- \rightarrow q\bar{q}\gamma\gamma$ events in the energy range 130–209 GeV. These samples are used to constrain possible anomalous $W^+W^-\gamma\gamma$ and $ZZ\gamma\gamma$ quartic couplings. Further combining with the $W^+W^-\gamma$ sample of ABBIENDI 04B the following one-parameter 95% CL limits are obtained: $-0.007 < a_0^Z/\Lambda^2 < 0.023 \text{ GeV}^{-2}$, $-0.029 < a_c^Z/\Lambda^2 < 0.029 \text{ GeV}^{-2}$, $-0.020 < a_0^W/\Lambda^2 < 0.020 \text{ GeV}^{-2}$, $-0.052 < a_c^W/\Lambda^2 < 0.037 \text{ GeV}^{-2}$.
- 123 In the CM energy range 183 to 209 GeV HEISTER 04A select 30 $e^+e^- \rightarrow \nu\bar{\nu}\gamma\gamma$ events with two acoplanar, high energy and high transverse momentum photons. The photon-photon acoplanarity is required to be $> 5^\circ$, $E_\gamma/\sqrt{s} > 0.025$ (the more energetic photon having energy $> 0.2\sqrt{s}$), $p_{T_\gamma}/E_{\text{beam}} > 0.05$ and $|\cos\theta_\gamma| < 0.94$. A likelihood fit to the photon energy and recoil missing mass yields the following one-parameter 95% CL limits: $-0.012 < a_0^Z/\Lambda^2 < 0.019 \text{ GeV}^{-2}$, $-0.041 < a_c^Z/\Lambda^2 < 0.044 \text{ GeV}^{-2}$, $-0.060 < a_0^W/\Lambda^2 < 0.055 \text{ GeV}^{-2}$, $-0.099 < a_c^W/\Lambda^2 < 0.093 \text{ GeV}^{-2}$.
- 124 ABDALLAH 03I select 122 $e^+e^- \rightarrow W^+W^-\gamma$ events in the C.M. energy range 189–209 GeV, where $E_\gamma > 5$ GeV, the photon has a polar angle $|\cos\theta_\gamma| < 0.95$ and is well isolated from the nearest charged fermion. A fit to the photon energy spectra yields $a_c/\Lambda^2 = 0.000^{+0.019}_{-0.040} \text{ GeV}^{-2}$, $a_0/\Lambda^2 = -0.004^{+0.018}_{-0.010} \text{ GeV}^{-2}$, $\tilde{a}_0/\Lambda^2 = -0.007^{+0.019}_{-0.008} \text{ GeV}^{-2}$, $a_n/\Lambda^2 = -0.09^{+0.16}_{-0.05} \text{ GeV}^{-2}$, and $\tilde{a}_n/\Lambda^2 = +0.05^{+0.07}_{-0.15} \text{ GeV}^{-2}$, keeping the other parameters fixed to their Standard Model values (0). The 95% CL limits are: $-0.063 \text{ GeV}^{-2} < a_c/\Lambda^2 < +0.032 \text{ GeV}^{-2}$, $-0.020 \text{ GeV}^{-2} < a_0/\Lambda^2 < +0.020 \text{ GeV}^{-2}$, $-0.020 \text{ GeV}^{-2} < \tilde{a}_0/\Lambda^2 < +0.020 \text{ GeV}^{-2}$, $-0.18 \text{ GeV}^{-2} < a_n/\Lambda^2 < +0.14 \text{ GeV}^{-2}$, $-0.16 \text{ GeV}^{-2} < \tilde{a}_n/\Lambda^2 < +0.17 \text{ GeV}^{-2}$.
- 125 ACHARD 02F select 86 $e^+e^- \rightarrow W^+W^-\gamma$ events at 192–207 GeV, where $E_\gamma > 5$ GeV and the photon is well isolated. They also select 43 acoplanar $e^+e^- \rightarrow \nu\bar{\nu}\gamma\gamma$ events in this energy range, where the photon energies are > 5 GeV and > 1 GeV and the photon polar angles are between 14° and 166° . All these 43 events are in the recoil mass region corresponding to the Z (75–110 GeV). Using the shape and normalization of the photon spectra in the $W^+W^-\gamma$ events, and combining with the 42 event sample from

189 GeV data (ACCIARRI 00T), they obtain: $a_0/\Lambda^2 = 0.000 \pm 0.010 \text{ GeV}^{-2}$, $a_c/\Lambda^2 = -0.013 \pm 0.023 \text{ GeV}^{-2}$, and $a_n/\Lambda^2 = -0.002 \pm 0.076 \text{ GeV}^{-2}$. Further combining the analyses of $W^+ W^- \gamma$ events with the low recoil mass region of $\nu \bar{\nu} \gamma \gamma$ events (including samples collected at 183 + 189 GeV), they obtain the following one-parameter 95% CL limits: $-0.015 \text{ GeV}^{-2} < a_0/\Lambda^2 < 0.015 \text{ GeV}^{-2}$, $-0.048 \text{ GeV}^{-2} < a_c/\Lambda^2 < 0.026 \text{ GeV}^{-2}$, and $-0.14 \text{ GeV}^{-2} < a_n/\Lambda^2 < 0.13 \text{ GeV}^{-2}$.

W REFERENCES

| | | | | |
|------------|------|-------------------------|---|-------------------|
| AARON | 09B | EPJ C64 251 | F.D. Aaron <i>et al.</i> | (H1 Collab.) |
| ABAZOV | 09AB | PRL 103 141801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 09AD | PR D80 053012 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 09AJ | PRL 103 191801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 09AK | PRL 103 231802 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| AALTONEN | 08B | PRL 100 071801 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 08R | PRL 100 241805 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABDALLAH | 08A | EPJ C55 1 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ABDALLAH | 08C | EPJ C54 345 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| AALTONEN | 07F | PRL 99 151801 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| Also | | PR D77 112001 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 07L | PR D76 111103R | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 07Z | PR D76 111104R | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABBIENDI | 07A | EPJ C52 767 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABAZOV | 06H | PR D74 057101 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| Also | | PR D74 059904 (erratum) | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABBIENDI | 06 | EPJ C45 307 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI | 06A | EPJ C45 291 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ACHARD | 06 | EPJ C45 569 | P. Achard <i>et al.</i> | (L3 Collab.) |
| AKTAS | 06 | PL B632 35 | A. Aktas <i>et al.</i> | (H1 Collab.) |
| SCHAEEL | 06 | EPJ C47 309 | S. Schael <i>et al.</i> | (ALEPH Collab.) |
| ABAZOV | 05J | PR D71 091108R | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 05S | PRL 95 141802 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| SCHAEEL | 05A | PL B614 7 | S. Schael <i>et al.</i> | (ALEPH Collab.) |
| ABAZOV | 04D | PR D70 092008 | V.M. Abazov <i>et al.</i> | (CDF, D0 Collab.) |
| ABBIENDI | 04B | PL B580 17 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI | 04D | EPJ C33 463 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI | 04L | PR D70 032005 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABDALLAH | 04G | EPJ C34 127 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ACHARD | 04D | PL B586 151 | P. Achard <i>et al.</i> | (L3 Collab.) |
| ACHARD | 04J | PL B600 22 | P. Achard <i>et al.</i> | (L3 Collab.) |
| HEISTER | 04A | PL B602 31 | A. Heister <i>et al.</i> | (ALEPH Collab.) |
| SCHAEEL | 04A | EPJ C38 147 | S. Schael <i>et al.</i> | (ALEPH Collab.) |
| ABBIENDI | 03C | EPJ C26 321 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABDALLAH | 03I | EPJ C31 139 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ABAZOV | 02D | PR D66 012001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 02E | PR D66 032008 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ACHARD | 02F | PL B527 29 | P. Achard <i>et al.</i> | (L3 Collab.) |
| CHEKANOV | 02C | PL B539 197 | S. Chekanov <i>et al.</i> | (ZEUS Collab.) |
| ABBIENDI | 01H | EPJ C19 229 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABREU | 01I | PL B502 9 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| AFFOLDER | 01E | PR D64 052001 | T. Affolder <i>et al.</i> | (CDF Collab.) |
| ABBIENDI | 00V | PL B490 71 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI,G | 00 | PL B493 249 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBOTT | 00B | PR D61 072001 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABBOTT | 00D | PRL 84 5710 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABREU,P | 00F | EPJ C18 203 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| Also | | EPJ C25 493 (erratum) | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 00T | PL B490 187 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| AFFOLDER | 00M | PRL 85 3347 | T. Affolder <i>et al.</i> | (CDF Collab.) |
| BREITWEG | 00 | PL B471 411 | J. Breitweg <i>et al.</i> | (ZEUS Collab.) |
| BREITWEG | 00D | EPJ C12 411 | J. Breitweg <i>et al.</i> | (ZEUS Collab.) |
| EBOLI | 00 | MPL A15 1 | O. Eboli, M. Gonzalez-Garcia, S. Novaes | |
| ABBIENDI | 99N | PL B453 153 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBOTT | 99H | PR D60 052003 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABBOTT | 99I | PR D60 072002 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABREU | 99L | PL B459 382 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 99 | PL B454 386 | M. Acciarri <i>et al.</i> | (L3 Collab.) |

| | | | | |
|-----------|-----|-----------------------|---------------------------------|------------------|
| ACCIARRI | 99Q | PL B467 171 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| BARATE | 99I | PL B453 107 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BARATE | 99L | PL B462 389 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BARATE | 99M | PL B465 349 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| ABBOTT | 98N | PR D58 092003 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABBOTT | 98P | PR D58 012002 | B. Abbott <i>et al.</i> | (D0 Collab.) |
| ABE | 98H | PR D58 031101 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 98P | PR D58 091101 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 98C | PL B416 233 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 98N | PL B439 209 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| BARATE | 97 | PL B401 347 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BARATE | 97S | PL B415 435 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| ABACHI | 95D | PRL 75 1456 | S. Abachi <i>et al.</i> | (D0 Collab.) |
| ABE | 95C | PRL 74 341 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 95G | PRL 74 1936 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 95P | PRL 75 11 | F. Abe <i>et al.</i> | (CDF Collab.) |
| Also | | PR D52 4784 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 95W | PR D52 2624 | F. Abe <i>et al.</i> | (CDF Collab.) |
| Also | | PRL 73 220 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 92E | PRL 68 3398 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 92I | PRL 69 28 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ALITTI | 92 | PL B276 365 | J. Alitti <i>et al.</i> | (UA2 Collab.) |
| ALITTI | 92B | PL B276 354 | J. Alitti <i>et al.</i> | (UA2 Collab.) |
| ALITTI | 92C | PL B277 194 | J. Alitti <i>et al.</i> | (UA2 Collab.) |
| ALITTI | 92D | PL B277 203 | J. Alitti <i>et al.</i> | (UA2 Collab.) |
| ALITTI | 92F | PL B280 137 | J. Alitti <i>et al.</i> | (UA2 Collab.) |
| SAMUEL | 92 | PL B280 124 | M.A. Samuel <i>et al.</i> | (OKSU, CARL) |
| ABE | 91C | PR D44 29 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ALBAJAR | 91 | PL B253 503 | C. Albajar <i>et al.</i> | (UA1 Collab.) |
| ALITTI | 91C | ZPHY C52 209 | J. Alitti <i>et al.</i> | (UA2 Collab.) |
| SAMUEL | 91 | PRL 67 9 | M.A. Samuel <i>et al.</i> | (OKSU, CARL) |
| Also | | PRL 67 2920 (erratum) | M.A. Samuel <i>et al.</i> | |
| ABE | 90G | PRL 65 2243 | F. Abe <i>et al.</i> | (CDF Collab.) |
| Also | | PR D43 2070 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ALBAJAR | 90 | PL B241 283 | C. Albajar <i>et al.</i> | (UA1 Collab.) |
| ALITTI | 90B | PL B241 150 | J. Alitti <i>et al.</i> | (UA2 Collab.) |
| ABE | 89I | PRL 62 1005 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ALBAJAR | 89 | ZPHY C44 15 | C. Albajar <i>et al.</i> | (UA1 Collab.) |
| BAUR | 88 | NP B308 127 | U. Baur, D. Zeppenfeld | (FSU, WISC) |
| GRIFOLS | 88 | IJMP A3 225 | J.A. Grifols, S. Peris, J. Sola | (BARC, DESY) |
| Also | | PL B197 437 | J.A. Grifols, S. Peris, J. Sola | (BARC, DESY) |
| ALBAJAR | 87 | PL B185 233 | C. Albajar <i>et al.</i> | (UA1 Collab.) |
| ANSARI | 87 | PL B186 440 | R. Ansari <i>et al.</i> | (UA2 Collab.) |
| GROTCH | 87 | PR D36 2153 | H. Grotch, R.W. Robinett | (PSU) |
| HAGIWARA | 87 | NP B282 253 | K. Hagiwara <i>et al.</i> | (KEK, UCLA, FSU) |
| VANDERBIJ | 87 | PR D35 1088 | J.J. van der Bij | (FNAL) |
| GRAU | 85 | PL 154B 283 | A. Grau, J.A. Grifols | (BARC) |
| SUZUKI | 85 | PL 153B 289 | M. Suzuki | (LBL) |
| ARNISON | 84D | PL 134B 469 | G.T.J. Arnison <i>et al.</i> | (UA1 Collab.) |
| HERZOG | 84 | PL 148B 355 | F. Herzog | (WISC) |
| Also | | PL 155B 468 (erratum) | F. Herzog | (WISC) |
| ARNISON | 83 | PL 122B 103 | G.T.J. Arnison <i>et al.</i> | (UA1 Collab.) |
| BANNER | 83B | PL 122B 476 | M. Banner <i>et al.</i> | (UA2 Collab.) |