

$B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE

$B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE MEAN LIFE

Each measurement of the B mean life is an average over an admixture of various bottom mesons and baryons which decay weakly. Different techniques emphasize different admixtures of produced particles, which could result in a different B mean life.

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements and asymmetric lifetime errors, but ignores the small differences due to different techniques.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.568±0.009 OUR EVALUATION				
1.570±0.005±0.008		¹ ABDALLAH 04E	DLPH	$e^+e^- \rightarrow Z$
1.533±0.015 ^{+0.035} _{-0.031}		² ABE 98B	CDF	$p\bar{p}$ at 1.8 TeV
1.549±0.009±0.015		³ ACCIARRI 98	L3	$e^+e^- \rightarrow Z$
1.611±0.010±0.027		⁴ ACKERSTAFF 97F	OPAL	$e^+e^- \rightarrow Z$
1.582±0.011±0.027		⁴ ABREU 96E	DLPH	$e^+e^- \rightarrow Z$
1.533±0.013±0.022	19.8k	⁵ BUSKULIC 96F	ALEP	$e^+e^- \rightarrow Z$
1.564±0.030±0.036		⁶ ABE,K 95B	SLD	$e^+e^- \rightarrow Z$
1.542±0.021±0.045		⁷ ABREU 94L	DLPH	$e^+e^- \rightarrow Z$
1.523±0.034±0.038	5372	⁸ ACTON 93L	OPAL	$e^+e^- \rightarrow Z$
1.511±0.022±0.078		⁹ BUSKULIC 93O	ALEP	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.575±0.010±0.026		¹⁰ ABREU 96E	DLPH	$e^+e^- \rightarrow Z$
1.50 ^{+0.24} _{-0.21} ± 0.03		¹¹ ABREU 94P	DLPH	$e^+e^- \rightarrow Z$
1.46 ± 0.06 ± 0.06	5344	¹² ABE 93J	CDF	Repl. by ABE 98B
1.23 ^{+0.14} _{-0.13} ± 0.15	188	¹³ ABREU 93D	DLPH	Sup. by ABREU 94L
1.49 ± 0.11 ± 0.12	253	¹⁴ ABREU 93G	DLPH	Sup. by ABREU 94L
1.51 ^{+0.16} _{-0.14} ± 0.11	130	¹⁵ ACTON 93C	OPAL	$e^+e^- \rightarrow Z$
1.535±0.035±0.028	7357	⁸ ADRIANI 93K	L3	Repl. by ACCIARRI 98
1.28 ± 0.10		¹⁶ ABREU 92	DLPH	Sup. by ABREU 94L
1.37 ± 0.07 ± 0.06	1354	¹⁷ ACTON 92	OPAL	Sup. by ACTON 93L
1.49 ± 0.03 ± 0.06		¹⁸ BUSKULIC 92F	ALEP	Sup. by BUSKULIC 96F
1.35 ^{+0.19} _{-0.17} ± 0.05		¹⁹ BUSKULIC 92G	ALEP	$e^+e^- \rightarrow Z$
1.32 ± 0.08 ± 0.09	1386	²⁰ ADEVA 91H	L3	Sup. by ADRIANI 93K
1.32 ^{+0.31} _{-0.25} ± 0.15	37	²¹ ALEXANDER 91G	OPAL	$e^+e^- \rightarrow Z$

1.29	± 0.06	± 0.10	2973	²² DECAMP	91C	ALEP	Sup. by BUSKULIC 92F
1.36	$+0.25$			²³ HAGEMANN	90	JADE	$E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$
1.13	± 0.15			²⁴ LYONS	90	RVUE	
1.35	± 0.10	± 0.24		BRAUNSCH...	89B	TASS	$E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$
0.98	± 0.12	± 0.13		ONG	89	MRK2	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
1.17	$+0.27$	$+0.17$		KLEM	88	DLCO	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
1.29	± 0.20	± 0.21		²⁵ ASH	87	MAC	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
1.02	$+0.42$		301	²⁶ BROM	87	HRS	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

¹ Measurement performed using an inclusive reconstruction and B flavor identification technique.

² Measured using inclusive $J/\psi(1S) \rightarrow \mu^+ \mu^-$ vertex.

³ ACCIARRI 98 uses inclusively reconstructed secondary vertex and lepton impact parameter.

⁴ ACKERSTAFF 97F uses inclusively reconstructed secondary vertices.

⁵ BUSKULIC 96F analyzed using 3D impact parameter.

⁶ ABE,K 95B uses an inclusive topological technique.

⁷ ABREU 94L uses charged particle impact parameters. Their result from inclusively reconstructed secondary vertices is superseded by ABREU 96E.

⁸ ACTON 93L and ADRIANI 93K analyzed using lepton (e and μ) impact parameter at Z .

⁹ BUSKULIC 93O analyzed using dipole method.

¹⁰ Combines ABREU 96E secondary vertex result with ABREU 94L impact parameter result.

¹¹ From proper time distribution of $b \rightarrow J/\psi(1S)\text{anything}$.

¹² ABE 93J analyzed using $J/\psi(1S) \rightarrow \mu\mu$ vertices.

¹³ ABREU 93D data analyzed using $D/D^*\ell\text{anything}$ event vertices.

¹⁴ ABREU 93G data analyzed using charged and neutral vertices.

¹⁵ ACTON 93C analysed using $D/D^*\ell\text{anything}$ event vertices.

¹⁶ ABREU 92 is combined result of muon and hadron impact parameter analyses. Hadron tracks gave $(12.7 \pm 0.4 \pm 1.2) \times 10^{-13} \text{ s}$ for an admixture of B species weighted by production fraction and mean charge multiplicity, while muon tracks gave $(13.0 \pm 1.0 \pm 0.8) \times 10^{-13} \text{ s}$ for an admixture weighted by production fraction and semileptonic branching fraction.

¹⁷ ACTON 92 is combined result of muon and electron impact parameter analyses.

¹⁸ BUSKULIC 92F uses the lepton impact parameter distribution for data from the 1991 run.

¹⁹ BUSKULIC 92G use $J/\psi(1S)$ tags to measure the average b lifetime. This is comparable to other methods only if the $J/\psi(1S)$ branching fractions of the different b -flavored hadrons are in the same ratio.

²⁰ Using $Z \rightarrow e^+ X$ or $\mu^+ X$, ADEVA 91H determined the average lifetime for an admixture of B hadrons from the impact parameter distribution of the lepton.

²¹ Using $Z \rightarrow J/\psi(1S)X$, $J/\psi(1S) \rightarrow \ell^+ \ell^-$, ALEXANDER 91G determined the average lifetime for an admixture of B hadrons from the decay point of the $J/\psi(1S)$.

²² Using $Z \rightarrow eX$ or μX , DECAMP 91C determines the average lifetime for an admixture of B hadrons from the signed impact parameter distribution of the lepton.

²³ HAGEMANN 90 uses electrons and muons in an impact parameter analysis.

²⁴ LYONS 90 combine the results of the B lifetime measurements of ONG 89, BRAUN-SCHWEIG 89B, KLEM 88, and ASH 87, and JADE data by private communication. They use statistical techniques which include variation of the error with the mean life, and possible correlations between the systematic errors. This result is not independent of the measured results used in our average.

²⁵ We have combined an overall scale error of 15% in quadrature with the systematic error of ± 0.7 to obtain ± 2.1 systematic error.

²⁶ Statistical and systematic errors were combined by BROM 87.

CHARGED b -HADRON ADMIXTURE MEAN LIFE

VALUE (10^{-12} s)	DOCUMENT ID	TECN	COMMENT
1.72±0.08±0.06	27 ADAM	95	DLPH $e^+ e^- \rightarrow Z$

²⁷ ADAM 95 data analyzed using vertex-charge technique to tag b -hadron charge.

NEUTRAL b -HADRON ADMIXTURE MEAN LIFE

VALUE (10^{-12} s)	DOCUMENT ID	TECN	COMMENT
1.58±0.11±0.09	28 ADAM	95	DLPH $e^+ e^- \rightarrow Z$

²⁸ ADAM 95 data analyzed using vertex-charge technique to tag b -hadron charge.

MEAN LIFE RATIO $\tau_{\text{charged } b\text{-hadron}}/\tau_{\text{neutral } b\text{-hadron}}$

VALUE	DOCUMENT ID	TECN	COMMENT
1.09^{+0.11}_{-0.10}±0.08	29 ADAM	95	DLPH $e^+ e^- \rightarrow Z$

²⁹ ADAM 95 data analyzed using vertex-charge technique to tag b -hadron charge.

$$|\Delta\tau_b|/\tau_{b,\bar{b}}$$

$\tau_{b,\bar{b}}$ and $|\Delta\tau_b|$ are the mean life average and difference between b and \bar{b} hadrons.

VALUE	DOCUMENT ID	TECN	COMMENT
-0.001±0.012±0.008	30 ABBIENDI	99J	OPAL $e^+ e^- \rightarrow Z$

³⁰ Data analyzed using both the jet charge and the charge of secondary vertex in the opposite hemisphere.

\bar{b} PRODUCTION FRACTIONS AND DECAY MODES

The branching fraction measurements are for an admixture of B mesons and baryons at energies above the $\Upsilon(4S)$. Only the highest energy results (LEP, Tevatron, $S\bar{p}S$) are used in the branching fraction averages. In the following, we assume that the production fractions are the same at the LEP and at the Tevatron.

For inclusive branching fractions, *e.g.*, $B \rightarrow D^\pm$ anything, the values usually are multiplicities, not branching fractions. They can be greater than one.

The modes below are listed for a \bar{b} initial state. b modes are their charge conjugates. Reactions indicate the weak decay vertex and do not include mixing.

PRODUCTION FRACTIONS

The production fractions for weakly decaying b -hadrons at high energy have been calculated from the best values of mean lives, mixing parameters, and branching fractions in this edition by the Heavy Flavor Averaging Group (HFAG) as described in the note “ B^0 - \bar{B}^0 Mixing” in the B^0 Particle Listings. The production fractions in b -hadronic Z decay or $p\bar{p}$ collisions at the Tevatron are also listed at the end of the section. Values assume

$$\begin{aligned} B(\bar{b} \rightarrow B^+) &= B(\bar{b} \rightarrow B^0) \\ B(\bar{b} \rightarrow B^+) + B(\bar{b} \rightarrow B^0) + B(\bar{b} \rightarrow B_s^0) + B(b \rightarrow b\text{-baryon}) &= 100\%. \end{aligned}$$

The correlation coefficients between production fractions are also reported:

$$\begin{aligned} \text{cor}(B_s^0, b\text{-baryon}) &= -0.067 \\ \text{cor}(B_s^0, B^\pm = B^0) &= -0.476 \\ \text{cor}(b\text{-baryon}, B^\pm = B^0) &= -0.846. \end{aligned}$$

The notation for production fractions varies in the literature (f_d , d_{B^0} , $f(b \rightarrow \bar{B}^0)$, $\text{Br}(b \rightarrow \bar{B}^0)$). We use our own branching fraction notation here, $B(\bar{b} \rightarrow B^0)$.

Note these production fractions are b -hadronization fractions, not the conventional branching fractions of b -quark to a B -hadron, which may have considerable dependence on the initial and final state kinematic and production environment.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 B^+	$(40.3 \pm 1.1)\%$	
Γ_2 B^0	$(40.3 \pm 1.1)\%$	
Γ_3 B_s^0	$(11.0 \pm 1.2)\%$	
Γ_4 $b\text{-baryon}$	$(8.3 \pm 2.0)\%$	
Γ_5 B_c	—	

DECAY MODES

Semileptonic and leptonic modes

Γ_6	ν anything	$(23.1 \pm 1.5)\%$	
Γ_7	$\ell^+ \nu_\ell$ anything	[a] $(10.69 \pm 0.22)\%$	
Γ_8	$e^+ \nu_e$ anything	$(10.86 \pm 0.35)\%$	
Γ_9	$\mu^+ \nu_\mu$ anything	$(10.95^{+ 0.29}_{- 0.25})\%$	
Γ_{10}	$D^- \ell^+ \nu_\ell$ anything	[a] $(2.27 \pm 0.35)\%$	S=1.7
Γ_{11}	$D^- \pi^+ \ell^+ \nu_\ell$ anything	$(4.9 \pm 1.9) \times 10^{-3}$	
Γ_{12}	$D^- \pi^- \ell^+ \nu_\ell$ anything	$(2.6 \pm 1.6) \times 10^{-3}$	
Γ_{13}	$\bar{D}^0 \ell^+ \nu_\ell$ anything	[a] $(6.85 \pm 0.35)\%$	
Γ_{14}	$\bar{D}^0 \pi^- \ell^+ \nu_\ell$ anything	$(1.07 \pm 0.27)\%$	
Γ_{15}	$\bar{D}^0 \pi^+ \ell^+ \nu_\ell$ anything	$(2.3 \pm 1.6) \times 10^{-3}$	
Γ_{16}	$D^{*-} \ell^+ \nu_\ell$ anything	[a] $(2.75 \pm 0.19)\%$	

Γ_{17}	$D^{*-} \pi^- \ell^+ \nu_\ell$ anything	$(6 \pm 7) \times 10^{-4}$
Γ_{18}	$D^{*-} \pi^+ \ell^+ \nu_\ell$ anything	$(4.8 \pm 1.0) \times 10^{-3}$
Γ_{19}	$\overline{D}_j^0 \ell^+ \nu_\ell$ anything \times $B(\overline{D}_j^0 \rightarrow D^{*+} \pi^-)$	[a,b] $(2.6 \pm 0.9) \times 10^{-3}$
Γ_{20}	$D_j^- \ell^+ \nu_\ell$ anything \times $B(D_j^- \rightarrow D^0 \pi^-)$	[a,b] $(7.0 \pm 2.3) \times 10^{-3}$
Γ_{21}	$\overline{D}_2^*(2460)^0 \ell^+ \nu_\ell$ anything $\times B(\overline{D}_2^*(2460)^0 \rightarrow$ $D^{*-} \pi^+)$	$< 1.4 \times 10^{-3}$ CL=90%
Γ_{22}	$D_2^*(2460)^- \ell^+ \nu_\ell$ anything $\times B(D_2^*(2460)^- \rightarrow$ $D^0 \pi^-)$	$(4.2 \pm 1.5) \times 10^{-3}$
Γ_{23}	$\overline{D}_2^*(2460)^0 \ell^+ \nu_\ell$ anything $\times B(\overline{D}_2^*(2460)^0 \rightarrow$ $D^- \pi^+)$	$(1.6 \pm 0.8) \times 10^{-3}$
Γ_{24}	charmless $\ell \bar{\nu}_\ell$	[a] $(1.7 \pm 0.5) \times 10^{-3}$
Γ_{25}	$\tau^+ \nu_\tau$ anything	$(2.41 \pm 0.23)\%$
Γ_{26}	$D^{*-} \tau \nu_\tau$ anything	$(9 \pm 4) \times 10^{-3}$
Γ_{27}	$\overline{c} \rightarrow \ell^- \bar{\nu}_\ell$ anything	[a] $(8.02 \pm 0.19)\%$
Γ_{28}	$c \rightarrow \ell^+ \nu$ anything	$(1.6 \pm 0.4)\%$

Charmed meson and baryon modes

Γ_{29}	\overline{D}^0 anything	$(59.8 \pm 2.9)\%$
Γ_{30}	$D^0 D_s^\pm$ anything	[c] $(9.1 \pm 4.0) \%$
Γ_{31}	$D^\mp D_s^\pm$ anything	[c] $(4.0 \pm 2.3) \%$
Γ_{32}	$\overline{D}^0 D^0$ anything	[c] $(5.1 \pm 2.0) \%$
Γ_{33}	$D^0 D^\pm$ anything	[c] $(2.7 \pm 1.8) \%$
Γ_{34}	$D^\pm D^\mp$ anything	[c] $< 9 \times 10^{-3}$ CL=90%
Γ_{35}	D^0 anything	
Γ_{36}	D^+ anything	
Γ_{37}	D^- anything	$(23.3 \pm 1.7)\%$
Γ_{38}	$D^*(2010)^+$ anything	$(17.3 \pm 2.0)\%$
Γ_{39}	$D_1(2420)^0$ anything	$(5.0 \pm 1.5)\%$
Γ_{40}	$D^*(2010)^\mp D_s^\pm$ anything	[c] $(3.3 \pm 1.6) \%$
Γ_{41}	$D^0 D^*(2010)^\pm$ anything	[c] $(3.0 \pm 1.1) \%$
Γ_{42}	$D^*(2010)^\pm D^\mp$ anything	[c] $(2.5 \pm 1.2) \%$
Γ_{43}	$D^*(2010)^\pm D^*(2010)^\mp$ anything	[c] $(1.2 \pm 0.4) \%$

Γ_{44}	$\overline{D}D$ anything	(10 \pm 11) %
Γ_{45}	$D_2^*(2460)^0$ anything	(4.7 \pm 2.7) %
Γ_{46}	D_s^- anything	(14.7 \pm 2.1) %
Γ_{47}	D_s^+ anything	(10.1 \pm 3.1) %
Γ_{48}	Λ_c^+ anything	(9.7 \pm 2.9) %
Γ_{49}	\overline{c}/c anything	[d] (116.2 \pm 3.2) %

Charmonium modes

Γ_{50}	$J/\psi(1S)$ anything	(1.16 \pm 0.10) %
Γ_{51}	$\psi(2S)$ anything	(4.8 \pm 2.4) $\times 10^{-3}$
Γ_{52}	$\chi_{c1}(1P)$ anything	(1.4 \pm 0.4) %

K or K^* modes

Γ_{53}	$\overline{s}\gamma$	(3.1 \pm 1.1) $\times 10^{-4}$
Γ_{54}	$\overline{s}\overline{\nu}\nu$	< 6.4 $\times 10^{-4}$ CL=90%
Γ_{55}	K^\pm anything	(74 \pm 6) %
Γ_{56}	K_S^0 anything	(29.0 \pm 2.9) %

Pion modes

Γ_{57}	π^\pm anything	(397 \pm 21) %
Γ_{58}	π^0 anything	[d] (278 \pm 60) %
Γ_{59}	ϕ anything	(2.82 \pm 0.23) %

Baryon modes

Γ_{60}	p/\overline{p} anything	(13.1 \pm 1.1) %
---------------	---------------------------	----------------------

Other modes

Γ_{61}	charged anything	[d] (497 \pm 7) %
Γ_{62}	hadron $^+$ hadron $^-$	(1.7 \pm 1.0) $\times 10^{-5}$
Γ_{63}	charmless	(7 \pm 21) $\times 10^{-3}$

Baryon modes

Γ_{64}	$\Lambda/\overline{\Lambda}$ anything	(5.9 \pm 0.6) %
Γ_{65}	b -baryon anything	(10.2 \pm 2.8) %

$\Delta B = 1$ weak neutral current ($B1$) modes

Γ_{66}	$e^+ e^-$ anything	
Γ_{67}	$\mu^+ \mu^-$ anything	$B1 < 3.2 \times 10^{-4}$ CL=90%
Γ_{68}	$\nu\overline{\nu}$ anything	

[a] An ℓ indicates an e or a μ mode, not a sum over these modes.

[b] D_j represents an unresolved mixture of pseudoscalar and tensor D^{**} (P -wave) states.

[c] The value is for the sum of the charge states or particle/antiparticle states indicated.

[d] Inclusive branching fractions have a multiplicity definition and can be greater than 100%.

$B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE BRANCHING RATIOS

$\Gamma(B^+)/\Gamma_{\text{total}}$

Γ_1/Γ

“OUR EVALUATION” is an average using rescaled values of the data listed below and from the best values of mean lives, mixing parameters, and branching fractions in this edition by the Heavy Flavor Averaging Group (HFAG) as described at <http://www.slac.stanford.edu/xorg/hfag/>.

VALUE	DOCUMENT ID	TECN	COMMENT
0.403 ± 0.011 OUR EVALUATION			

0.4099 ± 0.0082 ± 0.0111	³¹ ABDALLAH	03K DLPH	$e^+ e^- \rightarrow Z$
---------------------------------	------------------------	----------	-------------------------

³¹ The analysis is based on a neural network, to estimate the charge of the weakly-decaying b hadron by distinguishing its decay products from particles produced at the primary vertex.

$\Gamma(B^+)/\Gamma(B^0)$

Γ_1/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
1.054 ± 0.018 ± 0.062	AALTONEN	08N CDF	$p\bar{p}$ at 1.96 TeV

$\Gamma(B_s^0)/[\Gamma(B^+) + \Gamma(B^0)]$

$\Gamma_3/(\Gamma_1+\Gamma_2)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.136 ± 0.017 OUR EVALUATION			

0.140 ± 0.020 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.
----------------------------------	---

0.128 ^{+0.011} _{-0.010} ± 0.011	³² AALTONEN	08N CDF	$p\bar{p}$ at 1.96 TeV
---	------------------------	---------	------------------------


0.213 ± 0.068	³³ AFFOLDER	00E CDF	$p\bar{p}$ at 1.8 TeV
---------------	------------------------	---------	-----------------------

0.21 ± 0.036 ^{+0.038} _{-0.030}	³⁴ ABE	99P CDF	$\bar{p}p$ at 1.8 TeV
--	-------------------	---------	-----------------------

³² AALTONEN 08N reports $[\Gamma(\bar{b} \rightarrow B_s^0)/[\Gamma(\bar{b} \rightarrow B^+) + \Gamma(\bar{b} \rightarrow B^0)]] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (5.76 \pm 0.18)^{+0.45}_{-0.42} \times 10^{-3}$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³³ AFFOLDER 00E uses several electron-charm final states in $b \rightarrow ce^- X$.

³⁴ ABE 99P uses the numbers of $K^*(892)^0$, $K^*(892)^+$, and $\phi(1020)$ events produced in association with the double semileptonic decays $b \rightarrow c\mu^- X$ with $c \rightarrow s\mu^+ X$.



$$\Gamma(b\text{-baryon})/\left[\Gamma(B^+) + \Gamma(B^0)\right] \quad \Gamma_4/(\Gamma_1+\Gamma_2)$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.103±0.027 OUR EVALUATION

0.29 ± 0.10 OUR AVERAGE

0.31 ± 0.11 ± 0.12 35 AALTONEN 09E CDF $p\bar{p}$ at 1.8 TeV

0.28 ± 0.09 ± 0.07 36 AALTONEN 08N CDF $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.118±0.042 37 AFFOLDER 00E CDF $p\bar{p}$ at 1.8 TeV

35 Errata to the measurement reported in AFFOLDER 00E using the p_T spectra from fully reconstructed B^0 and Λ_b decays.

36 AALTONEN 08N reports $[\Gamma(\bar{b} \rightarrow b\text{-baryon})/\left[\Gamma(\bar{b} \rightarrow B^+) + \Gamma(\bar{b} \rightarrow B^0)\right]] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = (14.1 \pm 0.6^{+5.3}_{-4.4}) \times 10^{-3}$ which we divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

37 AFFOLDER 00E uses several electron-charm final states in $b \rightarrow c e^- X$.

$$\Gamma(\nu\text{anything})/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.2308±0.0077±0.0124 38,39 ACCIARRI 96C L3 $e^+ e^- \rightarrow Z$

38 ACCIARRI 96C assumes relative b semileptonic decay rates $e:\mu:\tau$ of 1:1:0.25. Based on missing-energy spectrum.

39 Assumes Standard Model value for R_B .

$\Gamma(\ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$ Γ_7/Γ

"OUR EVALUATION" is an average of the data listed below, excluding all asymmetry measurements, performed by the LEP Electroweak Working Group as described in the "Note on the Z boson" in the Z Particle Listings.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.1069 ± 0.0022 OUR EVALUATION			
0.1064 ± 0.0016 OUR AVERAGE			
0.1070 $\pm 0.0010 \pm 0.0035$	40 HEISTER	02G ALEP	$e^+ e^- \rightarrow Z$
0.1070 $\pm 0.0008^{+0.0037}_{-0.0049}$	41 ABREU	01L DLPH	$e^+ e^- \rightarrow Z$
0.1083 $\pm 0.0010^{+0.0028}_{-0.0024}$	42 ABBIENDI	00E OPAL	$e^+ e^- \rightarrow Z$
0.1016 $\pm 0.0013 \pm 0.0030$	43 ACCIARRI	00 L3	$e^+ e^- \rightarrow Z$
0.1085 $\pm 0.0012 \pm 0.0047$	44,45 ACCIARRI	96C L3	$e^+ e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.1106 $\pm 0.0039 \pm 0.0022$	46 ABREU	95D DLPH	$e^+ e^- \rightarrow Z$
0.114 $\pm 0.003 \pm 0.004$	47 BUSKULIC	94G ALEP	$e^+ e^- \rightarrow Z$
0.100 $\pm 0.007 \pm 0.007$	48 ABREU	93C DLPH	$e^+ e^- \rightarrow Z$
0.105 $\pm 0.006 \pm 0.005$	49 AKERS	93B OPAL	Repl. by ABBIENDI 00E

40 Uses the combination of lepton transverse momentum spectrum and the correlation between the charge of the lepton and opposite jet charge. The first error is statistic and the second error is the total systematic error including the modeling.

41 The experimental systematic and model uncertainties are combined in quadrature.

42 ABBIENDI 00E result is determined by comparing the distribution of several kinematic variables of leptonic events in a lifetime tagged $Z \rightarrow b\bar{b}$ sample using artificial neural network techniques. The first error is statistic; the second error is the total systematic error.

43 ACCIARRI 00 result obtained from a combined fit of $R_B = \Gamma(Z \rightarrow b\bar{b})/\Gamma(Z \rightarrow \text{hadrons})$ and $B(b \rightarrow \ell\nu X)$, using double-tagging method.

44 ACCIARRI 96C result obtained by a fit to the single lepton spectrum.

45 Assumes Standard Model value for R_B .

46 ABREU 95D give systematic errors ± 0.0019 (model) and 0.0012 (R_c). We combine these in quadrature.

47 BUSKULIC 94G uses e and μ events. This value is from a global fit to the lepton p and p_T (relative to jet) spectra which also determines the b and c production fractions, the fragmentation functions, and the forward-backward asymmetries. This branching ratio depends primarily on the ratio of dileptons to single leptons at high p_T , but the lower p_T portion of the lepton spectrum is included in the global fit to reduce the model dependence. The model dependence is ± 0.0026 and is included in the systematic error.

48 ABREU 93C event count includes ee events. Combining ee , $\mu\mu$, and $e\mu$ events, they obtain $0.100 \pm 0.007 \pm 0.007$.

49 AKERS 93B analysis performed using single and dilepton events.

 $\Gamma(e^+ \nu_e \text{anything})/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.1086 ± 0.0035 OUR AVERAGE				
0.1078 $\pm 0.0008^{+0.0050}_{-0.0046}$	50 ABBIENDI	00E OPAL	$e^+ e^- \rightarrow Z$	
0.1089 $\pm 0.0020 \pm 0.0051$	51,52 ACCIARRI	96C L3	$e^+ e^- \rightarrow Z$	
0.107 $\pm 0.015 \pm 0.007$	260 53 ABREU	93C DLPH	$e^+ e^- \rightarrow Z$	
0.138 $\pm 0.032 \pm 0.008$	54 ADEVA	91C L3	$e^+ e^- \rightarrow Z$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.086 ± 0.027 ± 0.008	55 ABE	93E VNS	$E_{\text{cm}}^{\text{ee}} = 58 \text{ GeV}$
0.109 $^{+0.014}_{-0.013}$ ± 0.0055 2719	56 AKERS	93B OPAL	Repl. by ABBIENDI 00E
0.111 ± 0.028 ± 0.026	BEHREND	90D CELL	$E_{\text{cm}}^{\text{ee}} = 43 \text{ GeV}$
0.150 ± 0.011 ± 0.022	BEHREND	90D CELL	$E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$
0.112 ± 0.009 ± 0.011	ONG	88 MRK2	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
0.149 $^{+0.022}_{-0.019}$	PAL	86 DLCO	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
0.110 ± 0.018 ± 0.010	AIHARA	85 TPC	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
0.111 ± 0.034 ± 0.040	ALTHOFF	84J TASS	$E_{\text{cm}}^{\text{ee}} = 34.6 \text{ GeV}$
0.146 ± 0.028	KOOP	84 DLCO	Repl. by PAL 86
0.116 ± 0.021 ± 0.017	NELSON	83 MRK2	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

50 ABBIENDI 00E result is determined by comparing the distribution of several kinematic variables of leptonic events in a lifetime tagged $Z \rightarrow b\bar{b}$ sample using artificial neural network techniques. The first error is statistic; the second error is the total systematic error.

51 ACCIARRI 96C result obtained by a fit to the single lepton spectrum.

52 Assumes Standard Model value for R_B .

53 ABREU 93C event count includes ee events. Combining ee, $\mu\mu$, and e μ events, they obtain $0.100 \pm 0.007 \pm 0.007$.

54 ADEVA 91C measure the average $B(b \rightarrow eX)$ branching ratio using single and double tagged b enhanced Z events. Combining e and μ results, they obtain $0.113 \pm 0.010 \pm 0.006$. Constraining the initial number of b quarks by the Standard Model prediction ($378 \pm 3 \text{ MeV}$) for the decay of the Z into $b\bar{b}$, the electron result gives $0.112 \pm 0.004 \pm 0.008$. They obtain $0.119 \pm 0.003 \pm 0.006$ when e and μ results are combined. Used to measure the $b\bar{b}$ width itself, this electron result gives $370 \pm 12 \pm 24 \text{ MeV}$ and combined with the muon result gives $385 \pm 7 \pm 22 \text{ MeV}$.

55 ABE 93E experiment also measures forward-backward asymmetries and fragmentation functions for b and c.

56 AKERS 93B analysis performed using single and dilepton events.

$\Gamma(\mu^+ \nu_\mu \text{anything})/\Gamma_{\text{total}}$		Γ_9/Γ		
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1095 $^{+0.0029}_{-0.0025}$ OUR AVERAGE				
0.1096 ± 0.0008 $^{+0.0034}_{-0.0027}$	57 ABBIENDI	00E OPAL	$e^+ e^- \rightarrow Z$	
0.1082 ± 0.0015 ± 0.0059	58,59 ACCIARRI	96C L3	$e^+ e^- \rightarrow Z$	
0.110 ± 0.012 ± 0.007	656 ABREU	93C DLPH	$e^+ e^- \rightarrow Z$	
0.113 ± 0.012 ± 0.006	61 ADEVA	91C L3	$e^+ e^- \rightarrow Z$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.122 ± 0.006 ± 0.007	59 UENO	96 AMY	$e^+ e^- \text{ at } 57.9 \text{ GeV}$	
0.101 $^{+0.010}_{-0.009}$ ± 0.0055 4248	62 AKERS	93B OPAL	Repl. by ABBIENDI 00E	
0.104 ± 0.023 ± 0.016	BEHREND	90D CELL	$E_{\text{cm}}^{\text{ee}} = 43 \text{ GeV}$	
0.148 ± 0.010 ± 0.016	BEHREND	90D CELL	$E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$	
0.118 ± 0.012 ± 0.010	ONG	88 MRK2	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$	

0.117 $\pm 0.016 \pm 0.015$	BARTEL	87	JADE	$E_{\text{cm}}^{\text{ee}} = 34.6 \text{ GeV}$
0.114 $\pm 0.018 \pm 0.025$	BARTEL	85J	JADE	Repl. by BARTEL 87
0.117 $\pm 0.028 \pm 0.010$	ALTHOFF	84G	TASS	$E_{\text{cm}}^{\text{ee}} = 34.5 \text{ GeV}$
0.105 $\pm 0.015 \pm 0.013$	ADEVA	83B	MRKJ	$E_{\text{cm}}^{\text{ee}} = 33\text{--}38.5 \text{ GeV}$
0.155 $^{+0.054}_{-0.029}$	FERNANDEZ	83D	MAC	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

57 ABBIENDI 00E result is determined by comparing the distribution of several kinematic variables of leptonic events in a lifetime tagged $Z \rightarrow b\bar{b}$ sample using artificial neural network techniques. The first error is statistic; the second error is the total systematic error.

58 ACCIARRI 96C result obtained by a fit to the single lepton spectrum.

59 Assumes Standard Model value for R_B .

60 ABREU 93C event count includes $\mu\mu$ events. Combining ee , $\mu\mu$, and $e\mu$ events, they obtain $0.100 \pm 0.007 \pm 0.007$.

61 ADEVA 91C measure the average $B(b \rightarrow eX)$ branching ratio using single and double tagged b enhanced Z events. Combining e and μ results, they obtain $0.113 \pm 0.010 \pm 0.006$. Constraining the initial number of b quarks by the Standard Model prediction ($378 \pm 3 \text{ MeV}$) for the decay of the Z into $b\bar{b}$, the muon result gives $0.123 \pm 0.003 \pm 0.006$. They obtain $0.119 \pm 0.003 \pm 0.006$ when e and μ results are combined. Used to measure the $b\bar{b}$ width itself, this muon result gives $394 \pm 9 \pm 22 \text{ MeV}$ and combined with the electron result gives $385 \pm 7 \pm 22 \text{ MeV}$.

62 AKERS 93B analysis performed using single and dilepton events.

$\Gamma(D^- \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0227 ± 0.0035 OUR AVERAGE			Error includes scale factor of 1.7.
0.0272 $\pm 0.0028 \pm 0.0018$	63 ABREU	00R DLPH	$e^+ e^- \rightarrow Z$
0.0199 $\pm 0.0026 \pm 0.0004$	64 AKERS	95Q OPAL	$e^+ e^- \rightarrow Z$

63 ABREU 00R reports their experiment's uncertainties $\pm 0.0019 \pm 0.0016 \pm 0.0018$, where the first error is statistical, the second is systematic, and the third is the uncertainty due to the D branching fraction. We combine first two in quadrature.

64 AKERS 95Q reports $[\Gamma(\bar{b} \rightarrow D^- \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = (1.82 \pm 0.20 \pm 0.12) \times 10^{-3}$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.13 \pm 0.19) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^- \pi^+ \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0049 $\pm 0.0018 \pm 0.0007$	ABREU	00R DLPH	$e^+ e^- \rightarrow Z$

$\Gamma(D^- \pi^- \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0026 $\pm 0.0015 \pm 0.0004$	ABREU	00R DLPH	$e^+ e^- \rightarrow Z$

$\Gamma(\bar{D}^0 \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0685 ± 0.0035 OUR AVERAGE			
0.0704 $\pm 0.0040 \pm 0.0017$	65 ABREU	00R DLPH	$e^+ e^- \rightarrow Z$
0.065 $\pm 0.006 \pm 0.001$	66 AKERS	95Q OPAL	$e^+ e^- \rightarrow Z$

⁶⁵ ABREU 00R reports their experiment's uncertainties $\pm 0.0034 \pm 0.0036 \pm 0.0017$, where the first error is statistical, the second is systematic, and the third is the uncertainty due to the D branching fraction. We combine first two in quadrature.

⁶⁶ AKERS 95Q reports $[\Gamma(\bar{b} \rightarrow \bar{D}^0 \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = (2.52 \pm 0.14 \pm 0.17) \times 10^{-3}$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.87 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{D}^0 \pi^- \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0107 $\pm 0.0025 \pm 0.0011$	ABREU	00R	DLPH $e^+ e^- \rightarrow Z$

Γ_{14}/Γ

$\Gamma(\bar{D}^0 \pi^+ \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0023 $\pm 0.0015 \pm 0.0004$	ABREU	00R	DLPH $e^+ e^- \rightarrow Z$

Γ_{15}/Γ

$\Gamma(D^* - \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0275 ± 0.0019 OUR AVERAGE			

$0.0275 \pm 0.0021 \pm 0.0009$

⁶⁷ ABREU 00R

DLPH $e^+ e^- \rightarrow Z$

$0.0276 \pm 0.0027 \pm 0.0011$

⁶⁸ AKERS 95Q

OPAL $e^+ e^- \rightarrow Z$

Γ_{16}/Γ

⁶⁷ ABREU 00R reports their experiment's uncertainties $\pm 0.0017 \pm 0.0013 \pm 0.0009$, where the first error is statistical, the second is systematic, and the third is the uncertainty due to the D branching fraction. We combine first two in quadrature.

⁶⁸ AKERS 95Q reports $[\Gamma(\bar{b} \rightarrow D^* \ell^+ \nu_\ell X) \times B(D^{*+} \rightarrow D^0 \pi^+) \times B(D^0 \rightarrow K^- \pi^+)] = ((7.53 \pm 0.47 \pm 0.56) \times 10^{-4})$ and uses $B(D^{*+} \rightarrow D^0 \pi^+) = 0.681 \pm 0.013$ and $B(D^0 \rightarrow K^- \pi^+) = 0.0401 \pm 0.0014$ to obtain the above result. The first error is the experiments error and the second error is the systematic error from the D^{*+} and D^0 branching ratios.

$\Gamma(D^{*-} \pi^+ \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

Γ_{18}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0048 $\pm 0.0009 \pm 0.0005$	ABREU	00R	DLPH $e^+ e^- \rightarrow Z$

$\Gamma(D^{*-} \pi^- \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0006 $\pm 0.0007 \pm 0.0002$	ABREU	00R	DLPH $e^+ e^- \rightarrow Z$

$\Gamma(\bar{D}_j^0 \ell^+ \nu_\ell \text{anything} \times B(\bar{D}_j^0 \rightarrow D^{*+} \pi^-))/\Gamma_{\text{total}}$

Γ_{19}/Γ

D_j represents an unresolved mixture of pseudoscalar and tensor D^{**} (P -wave) states.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.64 $\pm 0.79 \pm 0.39$	ABBIENDI	03M	OPAL $e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.1 $\pm 1.3 \pm 1.3$	AKERS	95Q	OPAL Repl. by ABBIENDI 03M
-----------------------	-------	-----	----------------------------

$\Gamma(D_j^- \ell^+ \nu_\ell \text{anything} \times B(D_j^- \rightarrow D^0 \pi^-)) / \Gamma_{\text{total}}$ Γ_{20}/Γ

D_j represents an unresolved mixture of pseudoscalar and tensor D^{**} (P -wave) states.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$7.0 \pm 1.9 \pm 1.2$	AKERS	95Q OPAL	$e^+ e^- \rightarrow Z$

$\Gamma(\bar{D}_2^*(2460)^0 \ell^+ \nu_\ell \text{anything} \times B(\bar{D}_2^*(2460)^0 \rightarrow D^{*-} \pi^+)) / \Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	ABBIENDI	03M OPAL	$e^+ e^- \rightarrow Z$

$\Gamma(D_2^*(2460)^- \ell^+ \nu_\ell \text{anything} \times B(D_2^*(2460)^- \rightarrow D^0 \pi^-)) / \Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$4.2 \pm 1.3 \pm 0.7$	AKERS	95Q OPAL	$e^+ e^- \rightarrow Z$

$\Gamma(\bar{D}_2^*(2460)^0 \ell^+ \nu_\ell \text{anything} \times B(\bar{D}_2^*(2460)^0 \rightarrow D^- \pi^+)) / \Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.6 \pm 0.7 \pm 0.3$	AKERS	95Q OPAL	$e^+ e^- \rightarrow Z$

$\Gamma(\text{charmless } \ell \bar{\nu}_\ell) / \Gamma_{\text{total}}$ Γ_{24}/Γ

“OUR EVALUATION” is an average of the data listed below performed by the LEP Heavy Flavour Steering Group. The averaging procedure takes into account correlations between the measurements.

VALUE	DOCUMENT ID	TECN	COMMENT
0.00171 ± 0.00052 OUR EVALUATION			
0.0017 ± 0.0004 OUR AVERAGE			
$0.00163 \pm 0.00053 \begin{matrix} +0.00055 \\ -0.00062 \end{matrix}$	69 ABBIENDI	01R OPAL	$e^+ e^- \rightarrow Z$
$0.00157 \pm 0.00035 \pm 0.00055$	70 ABREU	00D DLPH	$e^+ e^- \rightarrow Z$
$0.00173 \pm 0.00055 \pm 0.00055$	71 BARATE	99G ALEP	$e^+ e^- \rightarrow Z$
$0.0033 \pm 0.0010 \pm 0.0017$	72 ACCIARRI	98K L3	$e^+ e^- \rightarrow Z$

⁶⁹ Obtained from the best fit of the MC simulated events to the data based on the $b \rightarrow X_u \ell \nu$ neutral network output distributions.

⁷⁰ ABREU 00D result obtained from a fit to the numbers of decays in $b \rightarrow u$ enriched and depleted samples and their lepton spectra, and assuming $|V_{cb}| = 0.0384 \pm 0.0033$ and $\tau_b = 1.564 \pm 0.014$ ps.

⁷¹ Uses lifetime tagged $b\bar{b}$ sample.

⁷² ACCIARRI 98K assumes $R_b = 0.2174 \pm 0.0009$ at Z decay.

$\Gamma(\tau^+ \nu_\tau \text{anything}) / \Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.41 ± 0.23 OUR AVERAGE				
$2.78 \pm 0.18 \pm 0.51$		73 ABBIENDI	01Q OPAL	$e^+ e^- \rightarrow Z$
$2.43 \pm 0.20 \pm 0.25$		74 BARATE	01E ALEP	$e^+ e^- \rightarrow Z$
$2.19 \pm 0.24 \pm 0.39$		75 ABREU	00C DLPH	$e^+ e^- \rightarrow Z$
$1.7 \pm 0.5 \pm 1.1$	76,77	ACCIARRI	96C L3	$e^+ e^- \rightarrow Z$
$2.4 \pm 0.7 \pm 0.8$	1032	78 ACCIARRI	94C L3	$e^+ e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.75 \pm 0.30 \pm 0.37$	405	79 BUSKULIC	95 ALEP	Repl. by BARATE 01E
$4.08 \pm 0.76 \pm 0.62$		BUSKULIC	93B ALEP	Repl. by BUSKULIC 95

⁷³ ABBIENDI 01Q uses a missing energy technique.

⁷⁴ The energy-flow and b -tagging algorithms were used.

⁷⁵ Uses the missing energy in $Z \rightarrow b\bar{b}$ decays without identifying leptons.

⁷⁶ ACCIARRI 96C result obtained from missing energy spectrum.

⁷⁷ Assumes Standard Model value for R_B .

⁷⁸ This is a direct result using tagged $b\bar{b}$ events at the Z , but species are not separated.

⁷⁹ BUSKULIC 95 uses missing-energy technique.

$\Gamma(D^{*-} \tau \nu_\tau \text{anything})/\Gamma_{\text{total}}$

Γ_{26}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(0.88 \pm 0.31 \pm 0.28) \times 10^{-2}$	80 BARATE	01E ALEP	$e^+ e^- \rightarrow Z$

80 The energy-flow and b -tagging algorithms were used.

$\Gamma(\bar{b} \rightarrow \bar{c} \rightarrow \ell^- \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}$

Γ_{27}/Γ

"OUR EVALUATION" is an average of the data listed below, excluding all asymmetry measurements, performed by the LEP Electroweak Working Group as described in the "Note on the Z boson" in the Z Particle Listings.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0802 ± 0.0019 OUR EVALUATION

0.0817 ± 0.0020 OUR AVERAGE

$0.0818 \pm 0.0015^{+0.0024}_{-0.0026}$	81 HEISTER	02G ALEP	$e^+ e^- \rightarrow Z$
---	------------	----------	-------------------------

$0.0798 \pm 0.0022^{+0.0025}_{-0.0029}$	82 ABREU	01L DLPH	$e^+ e^- \rightarrow Z$
---	----------	----------	-------------------------

$0.0840 \pm 0.0016^{+0.0039}_{-0.0036}$	83 ABBIENDI	00E OPAL	$e^+ e^- \rightarrow Z$
---	-------------	----------	-------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.0770 \pm 0.0097 \pm 0.0046$	84 ABREU	95D DLPH	$e^+ e^- \rightarrow Z$
--------------------------------	----------	----------	-------------------------

$0.082 \pm 0.003 \pm 0.012$	85 BUSKULIC	94G ALEP	$e^+ e^- \rightarrow Z$
-----------------------------	-------------	----------	-------------------------

$0.077 \pm 0.004 \pm 0.007$	86 AKERS	93B OPAL	Repl. by ABBIENDI 00E
-----------------------------	----------	----------	-----------------------

81 Uses the combination of lepton transverse momentum spectrum and the correlation between the charge of the lepton and opposite jet charge. The first error is statistic and the second error is the total systematic error including the modeling.

82 The experimental systematic and model uncertainties are combined in quadrature.

83 ABBIENDI 00E result is determined by comparing the distribution of several kinematic variables of leptonic events in a lifetime tagged $Z \rightarrow b\bar{b}$ sample using artificial neural network techniques. The first error is statistic; the second error is the total systematic error.

84 ABREU 95D give systematic errors ± 0.0033 (model) and 0.0032 (R_C). We combine these in quadrature. This result is from the same global fit as their $\Gamma(\bar{b} \rightarrow \ell^+ \nu_\ell X)$ data.

85 BUSKULIC 94G uses e and μ events. This value is from the same global fit as their $\Gamma(\bar{b} \rightarrow \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$ data.

86 AKERS 93B analysis performed using single and dilepton events.

$\Gamma(c \rightarrow \ell^+ \nu \text{anything})/\Gamma_{\text{total}}$

Γ_{28}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.0161 \pm 0.0020^{+0.0034}_{-0.0047}$	87 ABREU	01L DLPH	$e^+ e^- \rightarrow Z$

87 The experimental systematic and model uncertainties are combined in quadrature.

$\Gamma(\bar{D}^0 \text{anything})/\Gamma_{\text{total}}$

VALUE

0.598_{-0.008}^{+0.007}

DOCUMENT ID

88 BUSKULIC 96Y ALEP

TECN

ALEP

$e^+ e^- \rightarrow Z$

Γ_{29}/Γ

88 BUSKULIC 96Y reports $0.605 \pm 0.024 \pm 0.016$ from a measurement of $[\Gamma(\bar{b} \rightarrow \bar{D}^0 \text{anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)]$ assuming $B(D^0 \rightarrow K^- \pi^+) = 0.0383$, which we rescale to our best value $B(D^0 \rightarrow K^- \pi^+) = (3.87 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^0 D_s^\pm \text{anything})/\Gamma_{\text{total}}$

VALUE

0.091_{-0.018}^{+0.020}_{-0.022}^{+0.034}

DOCUMENT ID

89 BARATE 98Q ALEP

TECN

ALEP

$e^+ e^- \rightarrow Z$

Γ_{30}/Γ

89 The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(D^\mp D_s^\pm \text{anything})/\Gamma_{\text{total}}$

VALUE

0.040_{-0.014}^{+0.017}_{-0.011}^{+0.016}

DOCUMENT ID

90 BARATE 98Q ALEP

TECN

ALEP

$e^+ e^- \rightarrow Z$

Γ_{31}/Γ

90 The systematic error includes the uncertainties due to the charm branching ratios.

$[\Gamma(D^0 D_s^\pm \text{anything}) + \Gamma(D^\mp D_s^\pm \text{anything})]/\Gamma_{\text{total}}$ $(\Gamma_{30} + \Gamma_{31})/\Gamma$

VALUE

0.131_{-0.022}^{+0.026}_{-0.031}^{+0.048}

DOCUMENT ID

91 BARATE 98Q ALEP

TECN

ALEP

$e^+ e^- \rightarrow Z$

91 The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(\bar{D}^0 D^0 \text{anything})/\Gamma_{\text{total}}$

VALUE

0.051_{-0.014}^{+0.016}_{-0.011}^{+0.012}

DOCUMENT ID

92 BARATE 98Q ALEP

TECN

ALEP

$e^+ e^- \rightarrow Z$

Γ_{32}/Γ

92 The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(D^0 D^\pm \text{anything})/\Gamma_{\text{total}}$

VALUE

0.027_{-0.013}^{+0.015}_{-0.009}^{+0.010}

DOCUMENT ID

93 BARATE 98Q ALEP

TECN

ALEP

$e^+ e^- \rightarrow Z$

Γ_{33}/Γ

93 The systematic error includes the uncertainties due to the charm branching ratios.

$[\Gamma(\bar{D}^0 D^0 \text{anything}) + \Gamma(D^0 D^\pm \text{anything})]/\Gamma_{\text{total}}$ $(\Gamma_{32} + \Gamma_{33})/\Gamma$

VALUE

0.078_{-0.018}^{+0.020}_{-0.016}^{+0.018}

DOCUMENT ID

94 BARATE 98Q ALEP

TECN

ALEP

$e^+ e^- \rightarrow Z$

94 The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(D^\pm D^\mp \text{anything})/\Gamma_{\text{total}}$

VALUE

<0.009

CL%

90

DOCUMENT ID

BARATE 98Q ALEP

TECN

ALEP

$e^+ e^- \rightarrow Z$

Γ_{34}/Γ

$\Gamma(D^0 \text{anything}) + \Gamma(D^+ \text{anything})]/\Gamma_{\text{total}}$	$(\Gamma_{35} + \Gamma_{36})/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.093 ± 0.017 ± 0.014	95 ABDALLAH	03E DLPH	$e^+ e^- \rightarrow Z$

95 The second error is the total of systematic uncertainties including the branching fractions used in the measurement.

$\Gamma(D^- \text{anything})/\Gamma_{\text{total}}$	Γ_{37}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.233 ± 0.016 ± 0.005	96 BUSKULIC	96Y ALEP	$e^+ e^- \rightarrow Z$

96 BUSKULIC 96Y reports $0.234 \pm 0.013 \pm 0.010$ from a measurement of $[\Gamma(\bar{b} \rightarrow D^- \text{anything})/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)]$ assuming $B(D^+ \rightarrow K^- 2\pi^+) = 0.091$, which we rescale to our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.13 \pm 0.19) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^*(2010)^+ \text{anything})/\Gamma_{\text{total}}$	Γ_{38}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.173 ± 0.016 ± 0.012	97 ACKERSTAFF	98E OPAL	$e^+ e^- \rightarrow Z$

97 Uses lepton tags to select $Z \rightarrow b\bar{b}$ events.

$\Gamma(D_1(2420)^0 \text{anything})/\Gamma_{\text{total}}$	Γ_{39}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.050 ± 0.014 ± 0.006	98 ACKERSTAFF	97W OPAL	$e^+ e^- \rightarrow Z$

98 ACKERSTAFF 97W assumes $B(D_2^*(2460)^0 \rightarrow D^{*+} \pi^-) = 0.21 \pm 0.04$ and $\Gamma_{b\bar{b}}/\Gamma_{\text{hadrons}} = 0.216$ at Z decay.

$\Gamma(D^*(2010)^{\mp} D_s^{\pm} \text{anything})/\Gamma_{\text{total}}$	Γ_{40}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.033 $^{+0.010}_{-0.009}$ $^{+0.012}_{-0.009}$	99 BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

99 The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(D^0 D^*(2010)^{\pm} \text{anything})/\Gamma_{\text{total}}$	Γ_{41}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.030 $^{+0.009}_{-0.008}$ $^{+0.007}_{-0.005}$	100 BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

100 The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(D^*(2010)^{\pm} D^{\mp} \text{anything})/\Gamma_{\text{total}}$	Γ_{42}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.025 $^{+0.010}_{-0.009}$ $^{+0.006}_{-0.005}$	101 BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

101 The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(D^*(2010)^{\pm} D^*(2010)^{\mp} \text{anything})/\Gamma_{\text{total}}$	Γ_{43}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.012 $^{+0.004}_{-0.003}$ $^{+0.002}_{-0.002}$	102 BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

102 The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(\bar{D} \text{ anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{44}/Γ
0.10 ± 0.032 ± 0.107	103 ABBIENDI	04I OPAL	$e^+ e^- \rightarrow Z$	

103 Measurement performed using an inclusive identification of B mesons and the D candidates.

$\Gamma(D_2^*(2460)^0 \text{ anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{45}/Γ
0.047 ± 0.024 ± 0.013	104 ACKERSTAFF	97W OPAL	$e^+ e^- \rightarrow Z$	

104 ACKERSTAFF 97W assumes $B(D_2^*(2460)^0 \rightarrow D^+ \pi^-) = 0.21 \pm 0.04$ and $\Gamma_{b\bar{b}}/\Gamma_{\text{hadrons}} = 0.216$ at Z decay.

$\Gamma(D_s^- \text{ anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{46}/Γ
0.147 ± 0.017 ± 0.013	105 BUSKULIC	96Y ALEP	$e^+ e^- \rightarrow Z$	

105 BUSKULIC 96Y reports $0.183 \pm 0.019 \pm 0.009$ from a measurement of $[\Gamma(\bar{b} \rightarrow D_s^- \text{ anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^+ \text{ anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{47}/Γ
0.101 ± 0.010 ± 0.029	106 ABDALLAH	03E DLPH	$e^+ e^- \rightarrow Z$	

106 The second error is the total of systematic uncertainties including the branching fractions used in the measurement.

$\Gamma(b \rightarrow \Lambda_c^+ \text{ anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{48}/Γ
0.097 ± 0.013 ± 0.025	107 BUSKULIC	96Y ALEP	$e^+ e^- \rightarrow Z$	

107 BUSKULIC 96Y reports $0.110 \pm 0.014 \pm 0.006$ from a measurement of $[\Gamma(b \rightarrow \Lambda_c^+ \text{ anything})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow pK^-\pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.044$, which we rescale to our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{c}/c \text{ anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{49}/Γ
1.162 ± 0.032 OUR AVERAGE				
1.12 $^{+0.11}_{-0.10}$	108 ABBIENDI	04I OPAL	$e^+ e^- \rightarrow Z$	
1.166 $\pm 0.031 \pm 0.080$	109 ABREU	00 DLPH	$e^+ e^- \rightarrow Z$	
1.147 ± 0.041	110 ABREU	98D DLPH	$e^+ e^- \rightarrow Z$	
1.230 $\pm 0.036 \pm 0.065$	111 BUSKULIC	96Y ALEP	$e^+ e^- \rightarrow Z$	

- 108 Measurement performed using an inclusive identification of B mesons and the D candidates.
 109 Evaluated via summation of exclusive and inclusive channels.
 110 ABREU 98D results are extracted from a fit to the b -tagging probability distribution based on the impact parameter.
 111 BUSKULIC 96Y assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons, and PDG 96 branching ratios for charm decays. This is sum of their inclusive \bar{D}^0 , D^- , \bar{D}_s , and Λ_c branching ratios, corrected to include inclusive Ξ_c and charmonium.

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$

Γ_{50}/Γ

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.16±0.10 OUR AVERAGE					
1.12±0.12±0.10			112 ABREU	94P DLPH	$e^+ e^- \rightarrow Z$
1.16±0.16±0.14		121	113 ADRIANI	93J L3	$e^+ e^- \rightarrow Z$
1.21±0.13±0.08			BUSKULIC	92G ALEP	$e^+ e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.3 ± 0.2 ± 0.2			114 ADRIANI	92 L3	$e^+ e^- \rightarrow Z$
<4.9		90	MATTEUZZI	83 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
112 ABREU 94P is an inclusive measurement from b decays at the Z . Uses $J/\psi(1S) \rightarrow e^+ e^-$ and $\mu^+ \mu^-$ channels. Assumes $\Gamma(Z \rightarrow b\bar{b})/\Gamma_{\text{hadron}} = 0.22$.					
113 ADRIANI 93J is an inclusive measurement from b decays at the Z . Uses $J/\psi(1S) \rightarrow \mu^+ \mu^-$ and $J/\psi(1S) \rightarrow e^+ e^-$ channels.					
114 ADRIANI 92 measurement is an inclusive result for $B(Z \rightarrow J/\psi(1S)X) = (4.1 \pm 0.7 \pm 0.3) \times 10^{-3}$ which is used to extract the b -hadron contribution to $J/\psi(1S)$ production.					

$\Gamma(\psi(2S)\text{anything})/\Gamma_{\text{total}}$

Γ_{51}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0048±0.0022±0.0010	115 ABREU	94P DLPH	$e^+ e^- \rightarrow Z$
115 ABREU 94P is an inclusive measurement from b decays at the Z . Uses $\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-$, $J/\psi(1S) \rightarrow \mu^+ \mu^-$ channels. Assumes $\Gamma(Z \rightarrow b\bar{b})/\Gamma_{\text{hadron}} = 0.22$.			

$\Gamma(\chi_{c1}(1P)\text{anything})/\Gamma_{\text{total}}$

Γ_{52}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.014 ± 0.004 OUR AVERAGE				
0.0111 ^{+0.0057} _{-0.0050} ± 0.0005		116 ABREU	94P DLPH	$e^+ e^- \rightarrow Z$
0.019 ± 0.007 ± 0.001	19	117 ADRIANI	93J L3	$e^+ e^- \rightarrow Z$
116 ABREU 94P reports 0.014 ± 0.006 ^{+0.004} _{-0.002}				from a measurement of $[\Gamma(\bar{b} \rightarrow \chi_{c1}(1P)\text{anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.4 \pm 1.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes no $\chi_{c2}(1P)$ and $\Gamma(Z \rightarrow b\bar{b})/\Gamma_{\text{hadron}} = 0.22$.
117 ADRIANI 93J reports 0.024 ± 0.009 ± 0.002				from a measurement of $[\Gamma(\bar{b} \rightarrow \chi_{c1}(1P)\text{anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.4 \pm 1.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\chi_{c1}(1P)\text{anything})/\Gamma(J/\psi(1S)\text{anything})$ Γ_{52}/Γ_{50}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.92 \pm 0.82	121	118 ADRIANI	93J L3	$e^+ e^- \rightarrow Z$
118 ADRIANI 93J is a ratio of inclusive measurements from b decays at the Z using only the $J/\psi(1S) \rightarrow \mu^+ \mu^-$ channel since some systematics cancel.				

$\Gamma(\bar{s}\gamma)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
3.11 \pm 0.80 \pm 0.72		119 BARATE	98I ALEP	$e^+ e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 5.4	90	120 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
< 12	90	121 ADRIANI	93L L3	$e^+ e^- \rightarrow Z$
119 BARATE 98I uses lifetime tagged $Z \rightarrow b\bar{b}$ sample.				
120 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.				
121 ADRIANI 93L result is for $\bar{b} \rightarrow \bar{s}\gamma$ is performed inclusively.				

$\Gamma(\bar{s}\nu)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 6.4 $\times 10^{-4}$	90	122 BARATE	01E ALEP	$e^+ e^- \rightarrow Z$
122 The energy-flow and b -tagging algorithms were used.				

$\Gamma(K^\pm\text{anything})/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.74 \pm 0.06 OUR AVERAGE	BARATE	98V ALEP	$e^+ e^- \rightarrow Z$
0.72 \pm 0.02 \pm 0.06	ABREU	95C DLPH	$e^+ e^- \rightarrow Z$
0.88 \pm 0.05 \pm 0.18			

$\Gamma(K_S^0\text{anything})/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.290 \pm 0.011 \pm 0.027	ABREU	95C DLPH	$e^+ e^- \rightarrow Z$

$\Gamma(\pi^\pm\text{anything})/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
3.97 \pm 0.02 \pm 0.21	BARATE	98V ALEP	$e^+ e^- \rightarrow Z$

$\Gamma(\pi^0\text{anything})/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT	
2.78 \pm 0.15 \pm 0.60	123 ADAM	96 DLPH	$e^+ e^- \rightarrow Z$	
123 ADAM 96 measurement obtained from a fit to the rapidity distribution of π^0 's in $Z \rightarrow b\bar{b}$ events.				

$\Gamma(\phi\text{anything})/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0282 \pm 0.0013 \pm 0.0019	ABBIENDI	00Z OPAL	$e^+ e^- \rightarrow Z$

$\Gamma(p/\bar{p}\text{anything})/\Gamma_{\text{total}}$

VALUE

0.131±0.011 OUR AVERAGE

$0.131 \pm 0.004 \pm 0.011$

$0.141 \pm 0.018 \pm 0.056$

DOCUMENT ID TECN COMMENT

BARATE 98V ALEP $e^+ e^- \rightarrow Z$
ABREU 95C DLPH $e^+ e^- \rightarrow Z$

Γ_{60}/Γ

$\Gamma(\text{charged anything})/\Gamma_{\text{total}}$

VALUE

4.97±0.03±0.06

DOCUMENT ID TECN COMMENT

124 ABREU 98H DLPH $e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.84 \pm 0.04 \pm 0.38$

ABREU 95C DLPH Repl. by ABREU 98H

124 ABREU 98H measurement excludes the contribution from K^0 and Λ decay.

Γ_{61}/Γ

$\Gamma(\text{hadron}^+ \text{ hadron}^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})

1.7^{+1.0}_{-0.7}±0.2

DOCUMENT ID TECN COMMENT

125,126 BUSKULIC 96V ALEP $e^+ e^- \rightarrow Z$

125 BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

126 Average branching fraction of weakly decaying B hadrons into two long-lived charged hadrons, weighted by their production cross section and lifetimes.

Γ_{62}/Γ

$\Gamma(\text{charmless})/\Gamma_{\text{total}}$

VALUE

0.007±0.021

DOCUMENT ID TECN COMMENT

127 ABREU 98D DLPH $e^+ e^- \rightarrow Z$

127 ABREU 98D results are extracted from a fit to the b -tagging probability distribution based on the impact parameter. The expected hidden charm contribution of 0.026 ± 0.004 has been subtracted.

Γ_{63}/Γ

$\Gamma(\Lambda/\bar{\Lambda}\text{anything})/\Gamma_{\text{total}}$

VALUE

0.059 ± 0.006 OUR AVERAGE

$0.0587 \pm 0.0046 \pm 0.0048$

$0.059 \pm 0.007 \pm 0.009$

DOCUMENT ID TECN COMMENT

ACKERSTAFF 97N OPAL $e^+ e^- \rightarrow Z$

ABREU 95C DLPH $e^+ e^- \rightarrow Z$

Γ_{64}/Γ

$\Gamma(b\text{-baryon anything})/\Gamma_{\text{total}}$

VALUE

0.102±0.007±0.027

DOCUMENT ID TECN COMMENT

128 BARATE 98V ALEP $e^+ e^- \rightarrow Z$

128 BARATE 98V assumes $B(B_s \rightarrow pX) = 8 \pm 4\%$ and $B(b\text{-baryon} \rightarrow pX) = 58 \pm 6\%$.

Γ_{65}/Γ

$\Gamma(\mu^+ \mu^- \text{anything})/\Gamma_{\text{total}}$

Test for $\Delta B = 1$ weak neutral current.

VALUE

DOCUMENT ID TECN COMMENT

<3.2 × 10⁻⁴ 90 ABBOTT 98B D0 $p\bar{p}$ 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<5.0 \times 10^{-5}$ 90 129 ALBAJAR 91C UA1 $E_{\text{cm}}^{p\bar{p}} = 630$ GeV

<0.02 95 ALTHOFF 84G TASS $E_{\text{cm}}^{ee} = 34.5$ GeV

<0.007 95 ADEVA 83 MRKJ $E_{\text{cm}}^{ee} = 30\text{--}38$ GeV

<0.007 95 BARTEL 83B JADE $E_{\text{cm}}^{ee} = 33\text{--}37$ GeV

Γ_{67}/Γ

129 Both ABBOTT 98B and GLENN 98 claim that the efficiency quoted in ALBAJAR 91C was overestimated by a large factor.

$$[\Gamma(e^+ e^- \text{anything}) + \Gamma(\mu^+ \mu^- \text{anything})]/\Gamma_{\text{total}} \quad (\Gamma_{66} + \Gamma_{67})/\Gamma$$

Test for $\Delta B = 1$ weak neutral current.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.008	90	MATTEUZZI 83	MRK2	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

$$\Gamma(\nu\bar{\nu}\text{anything})/\Gamma_{\text{total}} \quad \Gamma_{68}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$<3.9 \times 10^{-4}$	130	GROSSMAN 96	RVUE $e^+ e^- \rightarrow Z$
130 GROSSMAN 96 limit is derived from the ALEPH BUSKULIC 95 limit $B(B^+ \rightarrow \tau^+ \nu_\tau) < 1.8 \times 10^{-3}$ at CL=90% using conservative simplifying assumptions.			

χ_b AT HIGH ENERGY

For a discussion of B - \bar{B} mixing, see the note on " B^0 - \bar{B}^0 Mixing" in the B^0 Particle Listings.

χ_b is the average B - \bar{B} mixing parameter at high-energy $\chi_b = f'_d \chi_d + f'_s \chi_s$ where f'_d and f'_s are the fractions of B^0 and B_s^0 hadrons in an unbiased sample of semileptonic b -hadron decays.

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.1284 ± 0.0069 OUR EVALUATION				
0.129 ± 0.004 OUR AVERAGE				
0.132 ± 0.001	± 0.024	131 ABAZOV	06S D0	$p\bar{p}$ at 1.96 TeV
0.152 ± 0.007	± 0.011	132 ACOSTA	04A CDF	$p\bar{p}$ at 1.8 TeV
0.1312 ± 0.0049	± 0.0042	133 ABBIENDI	03P OPAL	$e^+ e^- \rightarrow Z$
0.127 ± 0.013	± 0.006	134 ABREU	01L DLPH	$e^+ e^- \rightarrow Z$
0.1192 ± 0.0068	± 0.0051	135 ACCIARRI	99D L3	$e^+ e^- \rightarrow Z$
0.121 ± 0.016	± 0.006	136 ABREU	94J DLPH	$e^+ e^- \rightarrow Z$
0.114 ± 0.014	± 0.008	137 BUSKULIC	94G ALEP	$e^+ e^- \rightarrow Z$
0.129 ± 0.022		138 BUSKULIC	92B ALEP	$e^+ e^- \rightarrow Z$
0.176 ± 0.031	± 0.032	1112 ABE	91G CDF	$p\bar{p}$ 1.8 TeV
0.148 ± 0.029	± 0.017	140 ALBAJAR	91D UA1	$p\bar{p}$ 630 GeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.131 ± 0.020	± 0.016	141 ABE	97I CDF	Repl. by ACOSTA 04A
0.1107 ± 0.0062	± 0.0055	142 ALEXANDER	96 OPAL	Rep. by ABBIENDI 03P
0.136 ± 0.037	± 0.040	143 UENO	96 AMY	$e^+ e^-$ at 57.9 GeV
0.144 ± 0.014	+ 0.017 - 0.011	144 ABREU	94F DLPH	Sup. by ABREU 94J
0.131 ± 0.014		145 ABREU	94J DLPH	$e^+ e^- \rightarrow Z$
0.123 ± 0.012	± 0.008	ACCIARRI	94D L3	Repl. by ACCIARRI 99D

0.157	± 0.020	± 0.032		¹⁴⁶ ALBAJAR	94	UA1	$\sqrt{s} = 630$ GeV
0.121	$+0.044$ -0.040	± 0.017	1665	¹⁴⁷ ABREU	93C	DLPH	Sup. by ABREU 94J
0.143	$+0.022$ -0.021	± 0.007		¹⁴⁸ AKERS	93B	OPAL	Sup. by ALEXANDER 96
0.145	$+0.041$ -0.035	± 0.018		¹⁴⁹ ACTON	92C	OPAL	$e^+ e^- \rightarrow Z$
0.121	± 0.017	± 0.006		¹⁵⁰ ADEVA	92C	L3	Sup. by ACCIARRI 99D
0.132	± 0.22	$+0.015$ -0.012	823	¹⁵¹ DECAMP	91	ALEP	$e^+ e^- \rightarrow Z$
0.178	$+0.049$ -0.040	± 0.020		¹⁵² ADEVA	90P	L3	$e^+ e^- \rightarrow Z$
0.17	$+0.15$ -0.08			^{153,154} WEIR	90	MRK2	$e^+ e^-$ 29 GeV
0.21	$+0.29$ -0.15			¹⁵³ BAND	88	MAC	$E_{cm}^{ee} = 29$ GeV
>0.02 at 90% CL				¹⁵³ BAND	88	MAC	$E_{cm}^{ee} = 29$ GeV
0.121	± 0.047			^{153,155} ALBAJAR	87C	UA1	Repl. by ALBAJAR 91D
<0.12 at 90% CL				^{153,156} SCHAAD	85	MRK2	$E_{cm}^{ee} = 29$ GeV

131 Uses the dimuon charge asymmetry. Averaged over the mix of *b*-flavored hadrons.

132 Measurement performed using events containing a dimuon or an e/μ pair.

133 The average *B* mixing parameter is determined simultaneously with *b* and *c* forward-backward asymmetries in the fit.

134 The experimental systematic and model uncertainties are combined in quadrature.

135 ACCIARRI 99D uses maximum-likelihood fits to extract χ_b as well as the A_{FB}^b in $Z \rightarrow b\bar{b}$ events containing prompt leptons.

136 This ABREU 94J result is from 5182 $\ell\ell$ and 279 $\Lambda\ell$ events. The systematic error includes 0.004 for model dependence.

137 BUSKULIC 94G data analyzed using ee , $e\mu$, and $\mu\mu$ events.

138 BUSKULIC 92B uses a jet charge technique combined with electrons and muons.

139 ABE 91G measurement of χ is done with $e\mu$ and ee events.

140 ALBAJAR 91D measurement of χ is done with dimuons.

141 Uses di-muon events.

142 ALEXANDER 96 uses a maximum likelihood fit to simultaneously extract χ as well as the forward-backward asymmetries in $e^+e^- \rightarrow Z \rightarrow b\bar{b}$ and $c\bar{c}$.

143 UENO 96 extracted χ from the energy dependence of the forward-backward asymmetry.

144 ABREU 94F uses the average electric charge sum of the jets recoiling against a *b*-quark jet tagged by a high p_T muon. The result is for $\overline{\chi} = f_d \chi_d + 0.9 f_s \chi_s$.

145 This ABREU 94J result combines $\ell\ell$, $\Lambda\ell$, and jet-charge ℓ (ABREU 94F) analyses. It is for $\overline{\chi} = f_d \chi_d + 0.96 f_s \chi_s$.

146 ALBAJAR 94 uses dimuon events. Not independent of ALBAJAR 91D.

147 ABREU 93C data analyzed using ee , $e\mu$, and $\mu\mu$ events.

148 AKERS 93B analysis performed using dilepton events.

149 ACTON 92C uses electrons and muons. Superseded by AKERS 93B.

150 ADEVA 92C uses electrons and muons.

151 DECAMP 91 done with opposite and like-sign dileptons. Superseded by BUSKULIC 92B.

152 ADEVA 90P measurement uses ee , $\mu\mu$, and $e\mu$ events from 118k events at the Z . Superseded by ADEVA 92C.

153 These experiments are not in the average because the combination of B_s and B_d mesons which they see could differ from those at higher energy.

154 The WEIR 90 measurement supersedes the limit obtained in SCHAAD 85. The 90% CL are 0.06 and 0.38.

155 ALBAJAR 87C measured $\chi = (\overline{B}^0 \rightarrow B^0 \rightarrow \mu^+ X)$ divided by the average production weighted semileptonic branching fraction for *B* hadrons at 546 and 630 GeV.

156 Limit is average probability for hadron containing *B* quark to produce a positive lepton.

CP VIOLATION PARAMETERS in semileptonic b -hadron decays.

$\text{Re}(\epsilon_b) / (1 + |\epsilon_b|^2)$

CP impurity in semileptonic b -hadron decays.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$-2.39 \pm 0.63 \pm 0.37$	157 ABAZOV	10H D0	$p\bar{p}$ at 1.96 TeV
157 ABAZOV 10H reports a measurement of like-sign dimuon charge asymmetry of $A_{SL}^b = (-9.57 \pm 2.51 \pm 1.46) \times 10^{-3}$ in semileptonic b -hadron decays. Using the measured production ratio of B_d^0 and B_s^0 , and the asymmetry of B_d^0 $A_{SL}^d = (-4.7 \pm 4.6) \times 10^{-3}$ measured from B -factories, they obtain the asymmetry for B_s^0 as $A_{SL}^s = (-14.6 \pm 7.5) \times 10^{-3}$.			

B -HADRON PRODUCTION FRACTIONS IN HADRONIC Z DECAY

The production fractions of b -hadrons in hadronic Z decays have been calculated using the best values of mean lives, mixing parameters and branching fractions in this edition by the Heavy Flavor Averaging Group (HFAG) (see <http://www.slac.stanford.edu/xorg/hfag/>).

The values reported below assume:

$$f(\bar{b} \rightarrow B^+) = f(\bar{b} \rightarrow B^0)$$

$$f(\bar{b} \rightarrow B^+) + f(\bar{b} \rightarrow B^0) + f(\bar{b} \rightarrow B_s^0) + f(b \rightarrow b\text{-baryon}) = 1$$

The values are:

$$f(\bar{b} \rightarrow B^+) = f(\bar{b} \rightarrow B^0) = 0.403 \pm 0.009$$

$$f(\bar{b} \rightarrow B_s^0) = 0.105 \pm 0.009$$

$$f(b \rightarrow b\text{-baryon}) = 0.089 \pm 0.015$$

and their correlation coefficients are:

$$\text{cor}(B_s^0, b\text{-baryon}) = +0.022$$

$$\text{cor}(B_s^0, B^+ = B^0) = -0.528$$

$$\text{cor}(b\text{-baryon}, B^+ = B^0) = -0.861$$

as obtained using a time-integrated mixing parameter $\overline{\chi} = 0.1259 \pm 0.0042$ given by a fit to heavy quark quantities with asymmetries removed (see the note "The Z boson").

B -HADRON PRODUCTION FRACTIONS IN $p\bar{p}$ COLLISIONS AT Tevatron

The production fractions for b -hadrons in $p\bar{p}$ collisions at the Tevatron have been calculated from the best values of mean lifetimes, mixing parameters, and branching fractions in this edition by the Heavy Flavor Averaging Group (HFAG) (see <http://www.slac.stanford.edu/xorg/hfag/>).

The values reported below assume:

$$f(\bar{b} \rightarrow B^+) = f(\bar{b} \rightarrow B^0)$$

$$f(\bar{b} \rightarrow B^+) + f(\bar{b} \rightarrow B^0) + f(\bar{b} \rightarrow B_s^0) + f(b \rightarrow b\text{-baryon}) = 1$$

The values are:

$$f(\bar{b} \rightarrow B^+) = f(\bar{b} \rightarrow B^0) = 0.339 \pm 0.031$$

$$f(\bar{b} \rightarrow B_s^0) = 0.111 \pm 0.014$$

$$f(b \rightarrow b\text{-baryon}) = 0.211 \pm 0.014$$

and their correlation coefficients are:

$$\text{cor}(B_s^0, b\text{-baryon}) = -0.585$$

$$\text{cor}(B_s^0, B^+ = B^0) = +0.429$$

$$\text{cor}(b\text{-baryon}, B^+ = B^0) = -0.984$$

as obtained with the Tevatron average of time-integrated mixing parameter

$$\overline{\chi} = 0.147 \pm 0.011.$$

$B^\pm/B^0/B_s^0/b\text{-baryon}$ ADMIXTURE REFERENCES

ABAZOV Also	10H PRL 105 081801	V.M. Abazov <i>et al.</i>	(D0 Collab.) (D0 Collab.)
AALTONEN	09E PR D79 032001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	08N PR D77 072003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	06S PR D74 092001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBIENDI	04I EPJ C35 149	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABDALLAH	04E EPJ C33 307	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ACOSTA	04A PR D69 012002	D. Acosta <i>et al.</i>	(CDF Collab.)
ABBIENDI	03M EPJ C30 467	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	03P PL B577 18	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABDALLAH	03E PL B561 26	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	03K PL B576 29	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
HEISTER	02G EPJ C22 613	A. Heister <i>et al.</i>	(ALEPH Collab.)
ABBIENDI	01Q PL B520 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	01R EPJ C21 399	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	01L EPJ C20 455	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BARATE	01E EPJ C19 213	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABBIENDI	00E EPJ C13 225	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	00Z PL B492 13	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	00 EPJ C12 225	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00C PL B496 43	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00D PL B478 14	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00R PL B475 407	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	00 EPJ C13 47	M. Acciarri <i>et al.</i>	(L3 Collab.)
AFFOLDER	00E PRL 84 1663	T. Affolder <i>et al.</i>	(CDF Collab.)
ABBIENDI	99J EPJ C12 609	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABE	99P PR D60 092005	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	99D PL B448 152	M. Acciarri <i>et al.</i>	(L3 Collab.)
BARATE	99G EPJ C6 555	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABBOTT	98B PL B423 419	B. Abbott <i>et al.</i>	(D0 Collab.)
ABE	98B PR D57 5382	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	98D PL B426 193	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	98H PL B425 399	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	98 PL B416 220	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	98K PL B436 174	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	98E EPJ C1 439	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98I PL B429 169	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98Q EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98V EPJ C5 205	R. Barate <i>et al.</i>	(ALEPH Collab.)
GLENN	98 PRL 80 2289	S. Glenn <i>et al.</i>	(CLEO Collab.)
ABE	97I PR D55 2546	F. Abe <i>et al.</i>	(CDF Collab.)
ACKERSTAFF	97F ZPHY C73 397	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97N ZPHY C74 423	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97W ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ABREU	96E PL B377 195	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	96C ZPHY C71 379	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAM	96 ZPHY C69 561	W. Adam <i>et al.</i>	(DELPHI Collab.)
ADAM	96D ZPHY C72 207	W. Adam <i>et al.</i>	(DELPHI Collab.)
ALEXANDER	96 ZPHY C70 357	G. Alexander <i>et al.</i>	(OPAL Collab.)
BUSKULIC	96F PL B369 151	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96V PL B384 471	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96Y PL B388 648	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
GROSSMAN	96 NP B465 369	Y. Grossman, Z. Ligeti, E. Nardi	(REHO, CIT)
Also	NP B480 753 (erratum)	Y. Grossman, Z. Ligeti, E. Nardi	
PDG	96 PR D54 1	R. M. Barnett <i>et al.</i>	
UENO	96 PL B381 365	K. Ueno <i>et al.</i>	(AMY Collab.)
ABE,K	95B PRL 75 3624	K. Abe <i>et al.</i>	(SLD Collab.)
ABREU	95C PL B347 447	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95D ZPHY C66 323	P. Abreu <i>et al.</i>	(DELPHI Collab.)

ADAM	95	ZPHY C68 363	W. Adam <i>et al.</i>	(DELPHI Collab.)
AKERS	95Q	ZPHY C67 57	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	95	PL B343 444	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	94F	PL B322 459	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	94J	PL B332 488	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	94L	ZPHY C63 3	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	94P	PL B341 109	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	94C	PL B332 201	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	94D	PL B335 542	M. Acciarri <i>et al.</i>	(L3 Collab.)
ALBAJAR	94	ZPHY C61 41	C. Albajar <i>et al.</i>	(UA1 Collab.)
BUSKULIC	94G	ZPHY C62 179	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABE	93E	PL B313 288	K. Abe <i>et al.</i>	(VENUS Collab.)
ABE	93J	PRL 71 3421	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	93C	PL B301 145	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	93D	ZPHY C57 181	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	93G	PL B312 253	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	93C	PL B307 247	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ACTON	93L	ZPHY C60 217	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ADRIANI	93J	PL B317 467	O. Adriani <i>et al.</i>	(L3 Collab.)
ADRIANI	93K	PL B317 474	O. Adriani <i>et al.</i>	(L3 Collab.)
ADRIANI	93L	PL B317 637	O. Adriani <i>et al.</i>	(L3 Collab.)
AKERS	93B	ZPHY C60 199	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	93B	PL B298 479	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	93O	PL B314 459	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	92	ZPHY C53 567	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	92	PL B274 513	D.P. Acton <i>et al.</i>	(OPAL Collab.)
ACTON	92C	PL B276 379	D.P. Acton <i>et al.</i>	(OPAL Collab.)
ADEVA	92C	PL B288 395	B. Adeva <i>et al.</i>	(L3 Collab.)
ADRIANI	92	PL B288 412	O. Adriani <i>et al.</i>	(L3 Collab.)
BUSKULIC	92B	PL B284 177	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	92F	PL B295 174	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	92G	PL B295 396	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABE	91G	PRL 67 3351	F. Abe <i>et al.</i>	(CDF Collab.)
ADEVA	91C	PL B261 177	B. Adeva <i>et al.</i>	(L3 Collab.)
ADEVA	91H	PL B270 111	B. Adeva <i>et al.</i>	(L3 Collab.)
ALBAJAR	91C	PL B262 163	C. Albajar <i>et al.</i>	(UA1 Collab.)
ALBAJAR	91D	PL B262 171	C. Albajar <i>et al.</i>	(UA1 Collab.)
ALEXANDER	91G	PL B266 485	G. Alexander <i>et al.</i>	(OPAL Collab.)
DECAMP	91	PL B258 236	D. Decamp <i>et al.</i>	(ALEPH Collab.)
DECAMP	91C	PL B257 492	D. Decamp <i>et al.</i>	(ALEPH Collab.)
ADEVA	90P	PL B252 703	B. Adeva <i>et al.</i>	(L3 Collab.)
BEHREND	90D	ZPHY C47 333	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
HAGEMANN	90	ZPHY C48 401	J. Hagemann <i>et al.</i>	(JADE Collab.)
LYONS	90	PR D41 982	L. Lyons, A.J. Martin, D.H. Saxon	(OXF, BRIS+)
WEIR	90	PL B240 289	A.J. Weir <i>et al.</i>	(Mark II Collab.)
BRAUNSCH...	89B	ZPHY C44 1	R. Braunschweig <i>et al.</i>	(TASSO Collab.)
ONG	89	PRL 62 1236	R.A. Ong <i>et al.</i>	(Mark II Collab.)
BAND	88	PL B200 221	H.R. Band <i>et al.</i>	(MAC Collab.)
KLEM	88	PR D37 41	D.E. Klem <i>et al.</i>	(DELCO Collab.)
ONG	88	PRL 60 2587	R.A. Ong <i>et al.</i>	(Mark II Collab.)
ALBAJAR	87C	PL B186 247	C. Albajar <i>et al.</i>	(UA1 Collab.)
ASH	87	PRL 58 640	W.W. Ash <i>et al.</i>	(MAC Collab.)
BARTEL	87	ZPHY C33 339	W. Bartel <i>et al.</i>	(JADE Collab.)
BROM	87	PL B195 301	J.M. Brom <i>et al.</i>	(HRS Collab.)
PAL	86	PR D33 2708	T. Pal <i>et al.</i>	(DELCO Collab.)
AIHARA	85	ZPHY C27 39	H. Aihara <i>et al.</i>	(TPC Collab.)
BARTEL	85J	PL 163B 277	W. Bartel <i>et al.</i>	(JADE Collab.)
SCHAAD	85	PL 160B 188	T. Schaad <i>et al.</i>	(Mark II Collab.)
ALTHOFF	84G	ZPHY C22 219	M. Althoff <i>et al.</i>	(TASSO Collab.)
ALTHOFF	84J	PL 146B 443	M. Althoff <i>et al.</i>	(TASSO Collab.)
KOOP	84	PRL 52 970	D.E. Koop <i>et al.</i>	(DELCO Collab.)
ADEVA	83	PRL 50 799	B. Adeva <i>et al.</i>	(Mark-J Collab.)
ADEVA	83B	PRL 51 443	B. Adeva <i>et al.</i>	(Mark-J Collab.)
BARTEL	83B	PL 132B 241	W. Bartel <i>et al.</i>	(JADE Collab.)
FERNANDEZ	83D	PRL 50 2054	E. Fernandez <i>et al.</i>	(MAC Collab.)
MATTEUZZI	83	PL 129B 141	C. Matteuzzi <i>et al.</i>	(Mark II Collab.)
NELSON	83	PRL 50 1542	M.E. Nelson <i>et al.</i>	(Mark II Collab.)