

## **$a_1(1640)$**


$I^G(J^{PC}) = 1^-(1^{++})$

### OMMITTED FROM SUMMARY TABLE

Seen in the amplitude analysis of the  $3\pi^0$  system produced in  $\bar{p}p \rightarrow 4\pi^0$ . Possibly seen in the study of the hadronic structure in decay  $\tau \rightarrow 3\pi\nu_\tau$  (ABREU 98G and ASNER 00). Needs confirmation.

### **$a_1(1640)$ MASS**

| VALUE (MeV)  | EVTS  | DOCUMENT ID                         | TECN    | COMMENT                                      |
|--|-------|-------------------------------------|---------|--|
| <b><math>1647 \pm 22</math> OUR AVERAGE</b>  |       | Error includes scale factor of 1.4. |         | See the ideogram below.                      |
| $1630 \pm 20$  | 35280 | 1 BAKER                             | 03 SPEC | $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$ |
| $1714 \pm 9 \pm 36$  |       | CHUNG                               | 02 B852 | $18.3\pi^-p \rightarrow \pi^+\pi^-\pi^-p$    |
| $1640 \pm 12 \pm 30$   |       | BAKER                               | 99 SPEC | $1.94\bar{p}p \rightarrow 4\pi^0$            |
| <b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b> |       |                                     |         |  |
| $1670 \pm 90$  |       | BELLINI                             | 85 SPEC | $40\pi^-A \rightarrow \pi^-\pi^+\pi^-A$      |



<sup>1</sup> Using the  $a_1(1260)$  mass and width results of BOWLER 88.

## **$a_1(1640)$ WIDTH**

| VALUE (MeV)   | EVTS  | DOCUMENT ID        | TECN | COMMENT   |
|---|-------|--------------------|------|---|
| <b>254± 27 OUR AVERAGE</b>  |       |                    |      | Error includes scale factor of 1.1.               |
| 225± 30   | 35280 | <sup>2</sup> BAKER | 03   | SPEC $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$ |
| 308± 37±62  |       | CHUNG              | 02   | B852 18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$ |
| 300± 22±40  |       | BAKER              | 99   | SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$           |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |       |                    |      |   |
| 300±100   |       | BELLINI            | 85   | SPEC 40 $\pi^- A \rightarrow \pi^-\pi^+\pi^- A$   |

<sup>2</sup> Using the  $a_1(1260)$  mass and width results of BOWLER 88.

## **$a_1(1640)$ DECAY MODES**

| Mode                             | Fraction ( $\Gamma_i/\Gamma$ ) |
|----------------------------------|--------------------------------|
| $\Gamma_1 \pi\pi\pi$             | seen                           |
| $\Gamma_2 f_2(1270)\pi$          | seen                           |
| $\Gamma_3 \sigma\pi$             | seen                           |
| $\Gamma_4 \rho\pi S\text{-wave}$ | seen                           |
| $\Gamma_5 \rho\pi D\text{-wave}$ | seen                           |
| $\Gamma_6 \omega\pi\pi$          | seen                           |
| $\Gamma_7 f_1(1285)\pi$          | seen                           |
| $\Gamma_8 a_1(1260)\eta$         | not seen                       |

## **$a_1(1640)$ BRANCHING RATIOS**

### $\Gamma(f_2(1270)\pi)/\Gamma(\sigma\pi)$ $\Gamma_2/\Gamma_3$

| VALUE   | DOCUMENT ID | TECN | COMMENT                                 |
|---|-------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |      |   |
| 0.24±0.07   | BAKER       | 99   | SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$ |

### $\Gamma(\rho\pi D\text{-wave})/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma$

| VALUE   | DOCUMENT ID | TECN | COMMENT   |
|---|-------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |      |   |
| seen  | CHUNG       | 02   | B852 18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$ |
| seen  | AMELIN      | 95B  | VES 36 $\pi^- A \rightarrow \pi^+\pi^-\pi^- A$    |

### $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma$

| VALUE   | EVTS  | DOCUMENT ID        | TECN | COMMENT   |
|---|-------|--------------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |       |                    |      |   |
| seen  | 35280 | <sup>3</sup> BAKER | 03   | SPEC $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$ |

### $\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}}$ $\Gamma_7/\Gamma$

| VALUE   | DOCUMENT ID | TECN | COMMENT  |
|---|-------------|------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |      |  |
| not seen  | KUHN        | 04   | B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$    |
| seen  | LEE         | 94   | MPS2 18 $\pi^- p \rightarrow K^+\bar{K}^0\pi^-\pi^- p$ |

$\Gamma(a_1(1260)\eta)/\Gamma_{\text{total}}$

VALUE

**not seen**

DOCUMENT ID

KUHN

TECN

04 B852

COMMENT

18  $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

$\Gamma_8/\Gamma$

<sup>3</sup> Assuming the  $\omega\rho$  mechanism for the  $\omega\pi\pi$  state.

---

**a<sub>1</sub>(1640) REFERENCES**

---

|         |     |               |                           |                         |
|---------|-----|---------------|---------------------------|-------------------------|
| KUHN    | 04  | PL B595 109   | J. Kuhn <i>et al.</i>     | (BNL E852 Collab.)      |
| BAKER   | 03  | PL B563 140   | C.A. Baker <i>et al.</i>  |                         |
| CHUNG   | 02  | PR D65 072001 | S.U. Chung <i>et al.</i>  | (BNL E852 Collab.)      |
| ASNER   | 00  | PR D61 012002 | D.M. Asner <i>et al.</i>  | (CLEO Collab.)          |
| BAKER   | 99  | PL B449 114   | C.A. Baker <i>et al.</i>  |                         |
| ABREU   | 98G | PL B426 411   | P. Abreu <i>et al.</i>    | (DELPHI Collab.)        |
| AMELIN  | 95B | PL B356 595   | D.V. Amelin <i>et al.</i> | (SERP, TBIL)            |
| LEE     | 94  | PL B323 227   | J.H. Lee <i>et al.</i>    | (BNL, IND, KYUN, MASD+) |
| BOWLER  | 88  | PL B209 99    | M.G. Bowler               | (OXF)                   |
| BELLINI | 85  | SJNP 41 781   | D. Bellini <i>et al.</i>  |                         |

Translated from YAF 41 1223.

---