

$\eta'(958)$ $I^G(J^{PC}) = 0^+(0^-+)$ **$\eta'(958)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
957.78 ± 0.06 OUR AVERAGE				
957.793 ± 0.054 ± 0.036	3.9k	LIBBY 08	CLEO	$J/\psi \rightarrow \gamma\eta'$
957.9 ± 0.2 ± 0.6	4800	WURZINGER 96	SPEC	$1.68 pd \rightarrow {}^3\text{He}\eta'$
957.46 ± 0.33		DUANE 74	MMS	$\pi^- p \rightarrow n\text{MM}$
958.2 ± 0.5	1414	DANBURG 73	HBC	$2.2 K^- p \rightarrow \Lambda\eta'$
958 ± 1	400	JACOBS 73	HBC	$2.9 K^- p \rightarrow \Lambda\eta'$
956.1 ± 1.1	3415	¹ BASILE 71	CNTR	$1.6 \pi^- p \rightarrow n\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
957.5 ± 0.2		BAI 04J	BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
959 ± 1	630	² BELADIDZE 92C	VES	$36 \pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
958 ± 1	340	² ARMSTRONG 91B	OMEG	$300 pp \rightarrow pp\eta\pi^+\pi^-$
958.2 ± 0.4	622	² AUGUSTIN 90	DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
957.8 ± 0.2	2420	² AUGUSTIN 90	DM2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
956.3 ± 1.0	143	² GIDAL 87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$
957.4 ± 1.4	535	³ BASILE 71	CNTR	$1.6 \pi^- p \rightarrow n\eta'$
957 ± 1		RITTENBERG 69	HBC	$1.7\text{--}2.7 K^- p$

¹ Using all η' decays.² Systematic uncertainty not estimated.³ Using η' decays into neutrals. Not independent of the other listed BASILE 71 η' mass measurement. **$\eta'(958)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.199 ± 0.009 OUR FIT					
0.230 ± 0.021 OUR AVERAGE					
0.226 ± 0.017 ± 0.014	2300	CZERWINSKI 10	MMS		$pp \rightarrow pp\eta'$
0.40 ± 0.22	4800	WURZINGER 96	SPEC		$1.68 pd \rightarrow {}^3\text{He}\eta'$
0.28 ± 0.10	1000	BINNIE 79	MMS	0	$\pi^- p \rightarrow n\text{MM}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.20 ± 0.04		BAI 04J	BES2		$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

$\eta'(958)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \pi^+ \pi^- \eta$	(43.4 ± 0.7) %	
$\Gamma_2 \rho^0 \gamma$ (including non-resonant $\pi^+ \pi^- \gamma$)	(29.3 ± 0.6) %	
$\Gamma_3 \pi^0 \pi^0 \eta$	(21.6 ± 0.8) %	
$\Gamma_4 \omega \gamma$	(2.75 ± 0.22) %	
$\Gamma_5 \gamma \gamma$	(2.18 ± 0.08) %	
$\Gamma_6 3\pi^0$	(1.68 ± 0.22) $\times 10^{-3}$	
$\Gamma_7 \mu^+ \mu^- \gamma$	(1.07 ± 0.26) $\times 10^{-4}$	
$\Gamma_8 \pi^+ \pi^- \mu^+ \mu^-$	< 2.2 $\times 10^{-4}$	90%
$\Gamma_9 \pi^+ \pi^- \pi^0$	(3.6 $^{+1.1}_{-0.9}$) $\times 10^{-3}$	
$\Gamma_{10} \pi^0 \rho^0$	< 4 %	90%
$\Gamma_{11} 2(\pi^+ \pi^-)$	< 2.4 $\times 10^{-4}$	90%
$\Gamma_{12} \pi^+ \pi^- 2\pi^0$	< 2.6 $\times 10^{-3}$	90%
$\Gamma_{13} 2(\pi^+ \pi^-)$ neutrals	< 1 %	95%
$\Gamma_{14} 2(\pi^+ \pi^-) \pi^0$	< 1.9 $\times 10^{-3}$	90%
$\Gamma_{15} 2(\pi^+ \pi^-) 2\pi^0$	< 1 %	95%
$\Gamma_{16} 3(\pi^+ \pi^-)$	< 5 $\times 10^{-4}$	90%
$\Gamma_{17} \pi^+ \pi^- e^+ e^-$	(2.4 $^{+1.3}_{-1.0}$) $\times 10^{-3}$	
$\Gamma_{18} \gamma e^+ e^-$	< 9 $\times 10^{-4}$	90%
$\Gamma_{19} \pi^0 \gamma \gamma$	< 8 $\times 10^{-4}$	90%
$\Gamma_{20} 4\pi^0$	< 5 $\times 10^{-4}$	90%
$\Gamma_{21} e^+ e^-$	< 2.1 $\times 10^{-7}$	90%
Γ_{22} invisible	< 9 $\times 10^{-4}$	90%

Charge conjugation (C), Parity (P), Lepton family number (LF) violating modes

$\Gamma_{23} \pi^+ \pi^-$	P, CP	< 2.9	$\times 10^{-3}$	90%
$\Gamma_{24} \pi^0 \pi^0$	P, CP	< 1.0	$\times 10^{-3}$	90%
$\Gamma_{25} \pi^0 e^+ e^-$	C	[a] < 1.4	$\times 10^{-3}$	90%
$\Gamma_{26} \eta e^+ e^-$	C	[a] < 2.4	$\times 10^{-3}$	90%
$\Gamma_{27} 3\gamma$	C	< 1.0	$\times 10^{-4}$	90%
$\Gamma_{28} \mu^+ \mu^- \pi^0$	C	[a] < 6.0	$\times 10^{-5}$	90%
$\Gamma_{29} \mu^+ \mu^- \eta$	C	[a] < 1.5	$\times 10^{-5}$	90%
$\Gamma_{30} e \mu$	LF	< 4.7	$\times 10^{-4}$	90%

[a] C parity forbids this to occur as a single-photon process.

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, 2 combinations of partial widths obtained from integrated cross section, and 13 branching ratios uses 40 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 31.5$ for 32 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	1							
x_3	-76	-57						
x_4	-20	-24	6					
x_5	-31	-26	35	0				
x_6	-23	-17	28	1	10			
x_9	-1	-5	-7	-3	-4	-2		
x_{17}	-4	-6	-5	-2	-3	-2	-1	
Γ	26	5	-21	4	-72	-6	4	3
	x_1	x_2	x_3	x_4	x_5	x_6	x_9	x_{17}

	Mode	Rate (MeV)
Γ_1	$\pi^+ \pi^- \eta$	0.086 ± 0.004
Γ_2	$\rho^0 \gamma$ (including non-resonant $\pi^+ \pi^- \gamma$)	0.0583 ± 0.0028
Γ_3	$\pi^0 \pi^0 \eta$	0.0430 ± 0.0022
Γ_4	$\omega \gamma$	0.0055 ± 0.0005
Γ_5	$\gamma \gamma$	0.00434 ± 0.00013
Γ_6	$3\pi^0$	$(3.3 \pm 0.5) \times 10^{-4}$
Γ_9	$\pi^+ \pi^- \pi^0$	$(7.2 \pm 2.2) \times 10^{-4}$
Γ_{17}	$\pi^+ \pi^- e^+ e^-$	$(4.8 \pm 2.6) \times 10^{-4}$

$\eta'(958)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$		Γ_5
VALUE (keV)	EVTS	DOCUMENT ID
4.34 \pm 0.14 OUR FIT		TECN
4.28 \pm 0.19 OUR AVERAGE		COMMENT
4.17 \pm 0.10 \pm 0.27	2000	4 ACCIARRI 98Q L3 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \gamma$
4.53 \pm 0.29 \pm 0.51	266	KARCH 92 CBAL $e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$
3.61 \pm 0.13 \pm 0.48		5 BEHREND 91 CELL $e^+ e^- \rightarrow e^+ e^- \eta'(958)$
4.6 \pm 1.1 \pm 0.6	23	BARU 90 MD1 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \gamma$

$4.57 \pm 0.25 \pm 0.44$		BUTLER	90	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta'(958)$
$5.08 \pm 0.24 \pm 0.71$	547	⁶ ROE	90	ASP	$e^+ e^- \rightarrow e^+ e^- 2\gamma$
$3.8 \pm 0.7 \pm 0.6$	34	AIHARA	88C	TPC	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
$4.9 \pm 0.5 \pm 0.5$	136	⁷ WILLIAMS	88	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$4.7 \pm 0.6 \pm 0.9$	143	⁸ GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
4.0 ± 0.9		⁹ BARTEL	85E	JADE	$e^+ e^- \rightarrow e^+ e^- 2\gamma$

⁴ No non-resonant $\pi^+ \pi^-$ contribution found.

⁵ Reevaluated by us using $B(\eta' \rightarrow \rho(770)\gamma) = (30.2 \pm 1.3)\%$.

⁶ Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.

⁷ Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.

⁸ Superseded by BUTLER 90.

⁹ Systematic error not evaluated.

$\eta'(958) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $\gamma\gamma$ and with the total width is obtained from the integrated cross section into channel(i) in the $\gamma\gamma$ annihilation.

$$\Gamma(\gamma\gamma) \times \Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / \Gamma_{\text{total}} \quad \Gamma_5 \Gamma_2 / \Gamma$$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

1.27 ± 0.04 OUR FIT

1.26 ± 0.07 OUR AVERAGE Error includes scale factor of 1.2.

$1.09 \pm 0.04 \pm 0.13$		BEHREND	91	CELL	$e^+ e^- \rightarrow e^+ e^- \rho(770)^0 \gamma$
$1.35 \pm 0.09 \pm 0.21$		AIHARA	87	TPC	$e^+ e^- \rightarrow e^+ e^- \rho\gamma$
$1.13 \pm 0.04 \pm 0.13$	867	ALBRECHT	87B	ARG	$e^+ e^- \rightarrow e^+ e^- \rho\gamma$
$1.53 \pm 0.09 \pm 0.21$		ALTHOFF	84E	TASS	$e^+ e^- \rightarrow e^+ e^- \rho\gamma$
$1.14 \pm 0.08 \pm 0.11$	243	BERGER	84B	PLUT	$e^+ e^- \rightarrow e^+ e^- \rho\gamma$
$1.73 \pm 0.34 \pm 0.35$	95	JENNI	83	MRK2	$e^+ e^- \rightarrow e^+ e^- \rho\gamma$
$1.49 \pm 0.13 \pm 0.027$	213	BARTEL	82B	JADE	$e^+ e^- \rightarrow e^+ e^- \rho\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.85 \pm 0.31 \pm 0.24$ 43 BEHREND 83B CELL $e^+ e^- \rightarrow e^+ e^- \rho\gamma$

$$\Gamma(\gamma\gamma) \times \Gamma(\pi^0 \pi^0 \eta) / \Gamma_{\text{total}} \quad \Gamma_5 \Gamma_3 / \Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

0.94 ± 0.05 OUR FIT

0.92 ± 0.06 ± 0.11 10 KARCH 92 CBAL $e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.95 \pm 0.05 \pm 0.08$	¹¹ KARCH	90	CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$
$1.00 \pm 0.08 \pm 0.10$	11,12 ANTREASYAN	87	CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$

¹⁰ Reevaluated by us using $B(\eta \rightarrow \gamma\gamma) = (39.21 \pm 0.34)\%$. Supersedes ANTREASYAN 87 and KARCH 90.

¹¹ Superseded by KARCH 92.

¹² Using $BR(\eta \rightarrow 2\gamma) = (38.9 \pm 0.5)\%$.

$\eta'(958) \rightarrow \eta\pi\pi$ DECAY PARAMETERS

$$|\text{MATRIX ELEMENT}|^2 = |1 + \alpha Y|^2 + CX + DX^2$$

X and Y are Dalitz variables; α is complex and C , and D are real-valued. Parameters C and D are not necessarily equal to c and d , respectively, in the generalized parameterization following this one. May be different for $\eta'(958) \rightarrow \eta\pi^+\pi^-$ and $\eta'(958) \rightarrow \eta\pi^0\pi^0$ decays. Because of different initial assumptions and strong correlations of the parameters we do not average the parameters in the section below.

$\text{Re}(\alpha)$ decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$-0.033 \pm 0.005 \pm 0.003$	44k	¹³ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$-0.072 \pm 0.012 \pm 0.006$	7k	¹⁴ AMELIN	05A	VES $28\pi^-A \rightarrow \eta\pi^+\pi^-\pi^-A^*$
$-0.021 \pm 0.018 \pm 0.017$	6.7k	¹⁵ BRIERE	00	CLEO $10.6e^+e^- \rightarrow \eta\pi^+\pi^-X$
$-0.058 \pm 0.013 \pm 0.003$	5.4k	¹⁶ ALDE	86	GAM2 $38\pi^-p \rightarrow n\eta\pi^0\pi^0$
-0.08 ± 0.03		^{16,17} KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta\pi^+\pi^-$

¹³ See ABLIKIM 11 for the full correlation matrix.

¹⁴ Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

¹⁵ Assuming $\text{Im}(\alpha) = 0$, $C = 0$, and $D = 0$.

¹⁶ Assuming $C = 0$.

¹⁷ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

$\text{Im}(\alpha)$ decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$0.000 \pm 0.049 \pm 0.001$	44k	¹⁸ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$0.0 \pm 0.1 \pm 0.0$	7k	¹⁹ AMELIN	05A	VES $28\pi^-A \rightarrow \eta\pi^+\pi^-\pi^-A^*$
$-0.00 \pm 0.13 \pm 0.00$	5.4k	²⁰ ALDE	86	GAM2 $38\pi^-p \rightarrow n\eta\pi^0\pi^0$
0.0 ± 0.3		^{20,21} KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta\pi^+\pi^-$

¹⁸ See ABLIKIM 11 for the full correlation matrix.

¹⁹ Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

²⁰ Assuming $C = 0$.

²¹ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

C decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$+0.018 \pm 0.009 \pm 0.003$	44k	²² ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$0.020 \pm 0.018 \pm 0.004$	7k	²³ AMELIN	05A	VES $28\pi^-A \rightarrow \eta\pi^+\pi^-\pi^-A^*$

²² See ABLIKIM 11 for the full correlation matrix.

²³ Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

D decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.059±0.012±0.004	44k	²⁴ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.066±0.030±0.015	7k	²⁵ AMELIN	05A	VES $\pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
0.00 ± 0.03 ± 0.00	5.4k	²⁶ ALDE	86	GAM2 $38 \pi^- p \rightarrow n\eta\pi^0\pi^0$
0		^{26,27} KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta\pi^+\pi^-$

²⁴ See ABLIKIM 11 for the full correlation matrix.
²⁵ Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.
²⁶ Assuming $C = 0$.
²⁷ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

$\eta'(958) \rightarrow \eta\pi\pi$ DECAY PARAMETERS

$$|\text{MATRIX ELEMENT}|^2 \propto 1 + a Y + b Y^2 + c X + d X^2$$

X and Y are Dalitz variables and a , b , c , and d are real-valued parameters.

May be different for $\eta'(958) \rightarrow \eta\pi^+\pi^-$ and $\eta'(958) \rightarrow \eta\pi^0\pi^0$ decays.

We do not average measurements in the section below because parameter values from each experiment are strongly correlated.

a decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.047±0.011±0.003	44k	²⁸ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.066±0.016±0.003	15k	²⁹ BLIK	09	GAM4 $32.5 \pi^- p \rightarrow \eta' n$
-0.127±0.016±0.008	20k	³⁰ DOROFEEV	07	VES $27 \pi^- p \rightarrow \eta' n, \pi^- A \rightarrow \eta' \pi^- A^*$

²⁸ See ABLIKIM 11 for the full correlation matrix.

²⁹ From $\eta' \rightarrow \eta\pi^0\pi^0$ decay.

³⁰ From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

b decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.069±0.019±0.009	44k	³¹ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.063±0.028±0.004	15k	³² BLIK	09	GAM4 $32.5 \pi^- p \rightarrow \eta' n$
-0.106±0.028±0.014	20k	³³ DOROFEEV	07	VES $27 \pi^- p \rightarrow \eta' n, \pi^- A \rightarrow \eta' \pi^- A^*$

³¹ See ABLIKIM 11 for the full correlation matrix.

³² From $\eta' \rightarrow \eta\pi^0\pi^0$ decay.

³³ From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

c decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.019±0.011±0.003	44k	³⁴ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.107±0.096±0.003	15k	³⁵ BLIK	09	GAM4 $32.5\pi^- p \rightarrow \eta' n$
0.015±0.011±0.014	20k	³⁶ DOROFEEV	07	VES $27\pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

³⁴ See ABLIKIM 11 for the full correlation matrix.

³⁵ From $\eta' \rightarrow \eta\pi^0\pi^0$ decay.

³⁶ From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

d decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.073±0.012±0.003	44k	³⁷ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
0.018±0.078±0.006	15k	³⁸ BLIK	09	GAM4 $32.5\pi^- p \rightarrow \eta' n$
-0.082±0.017±0.008	20k	³⁹ DOROFEEV	07	VES $27\pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

³⁷ See ABLIKIM 11 for the full correlation matrix.

³⁸ From $\eta' \rightarrow \eta\pi^0\pi^0$ decay. If $c \equiv 0$ from Bose-Einstein symmetry, $d = -0.067 \pm 0.020 \pm 0.003$.

³⁹ From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

$\eta'(958)$ β PARAMETER |MATRIX ELEMENT|² = (1 + 2 βZ)

See the "Note on η Decay Parameters" in our 1994 edition Physical Review
D50 1173 (1994), p. 1454.

β decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.46±0.22 OUR AVERAGE				Error includes scale factor of 1.4.
-0.59±0.18	235	BLIK	08	GAMS $32\pi^- p \rightarrow \eta' n$
-0.1 ± 0.3		ALDE	87B	GAM2 $38\pi^- p \rightarrow n3\pi^0$

$\eta'(958)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.434±0.007 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.424±0.011±0.004 1.2k ⁴⁰ PEDLAR 09 CLEO $J/\psi \rightarrow \gamma\eta'$

⁴⁰ Not independent of other η' branching fractions and ratios in PEDLAR 09.

Γ_1/Γ

$\Gamma(\pi^+\pi^-\eta(\text{charged decay}))/\Gamma_{\text{total}}$ $0.286\Gamma_1/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.1240 ± 0.0020 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.123 ± 0.014	107	RITTENBERG 69	HBC	1.7–2.7 $K^- p$
0.10 ± 0.04	10	LONDON 66	HBC	$2.24 K^- p \rightarrow \Lambda 2\pi^+ 2\pi^- \pi^0$
0.07 ± 0.04	7	BADIER 65B	HBC	$3 K^- p$

 $\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))/\Gamma_{\text{total}}$ $0.714\Gamma_1/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.310 ± 0.005 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.314 ± 0.026	281	RITTENBERG 69	HBC	1.7–2.7 $K^- p$
-------------------	-----	---------------	-----	-----------------

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.293 ± 0.006 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.287 $\pm 0.007 \pm 0.004$	0.2k	⁴¹ PEDLAR 09	CLEO	$J/\psi \rightarrow \gamma\eta'$
0.329 ± 0.033	298	RITTENBERG 69	HBC	1.7–2.7 $K^- p$
0.2 ± 0.1	20	LONDON 66	HBC	$2.24 K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$
0.34 ± 0.09	35	BADIER 65B	HBC	$3 K^- p$

⁴¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi^+\pi^-\eta)$ Γ_2/Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.676 ± 0.017 OUR FIT			
0.683 ± 0.020 OUR AVERAGE			

0.677 $\pm 0.024 \pm 0.011$	PEDLAR 09	CLE3	$J/\psi \rightarrow \eta'\gamma$
0.69 ± 0.03	ABLIKIM 06E	BES2	$J/\psi \rightarrow \eta'\gamma$

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))$ $\Gamma_2/0.714\Gamma_1$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.947 ± 0.024 OUR FIT				
0.97 ± 0.09 OUR AVERAGE				

0.70 ± 0.22	AMSLER 04B	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\eta$
1.07 ± 0.17	BELADIDZE 92C	VES	$36 \pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
0.92 ± 0.14	DANBURG 73	HBC	$2.2 K^- p \rightarrow \Lambda X^0$
1.11 ± 0.18	JACOBS 73	HBC	$2.9 K^- p \rightarrow \Lambda X^0$

 $\Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.216 ± 0.008 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.235 $\pm 0.013 \pm 0.004$	3.2k	⁴² PEDLAR 09	CLEO	$J/\psi \rightarrow \gamma\eta'$
-----------------------------	------	-------------------------	------	----------------------------------

⁴² Not independent of other η' branching fractions and ratios in PEDLAR 09.

$\Gamma(\pi^0\pi^0\eta(3\pi^0\text{decay}))/\Gamma_{\text{total}}$

$0.321\Gamma_3/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0694 ± 0.0026 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.11 ± 0.06	4	BENSINGER	70	DBC $2.2\pi^+d$
-----------------	---	-----------	----	-----------------

$\Gamma(\pi^0\pi^0\eta)/\Gamma(\pi^+\pi^-\eta)$

Γ_3/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.498 ± 0.025 OUR FIT			

$0.555 \pm 0.043 \pm 0.013$	PEDLAR	09	CLE3 $J/\psi \rightarrow \eta'\gamma$
-----------------------------------------------	--------	----	---------------------------------------

$\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi\pi\eta)$

$\Gamma_2/(\Gamma_1+\Gamma_3)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.451 ± 0.012 OUR FIT			

$0.43 \pm 0.02 \pm 0.02$	BARBERIS	98C	OMEG $450\text{ pp} \rightarrow p_f\eta' p_s$
--------------------------------------------	----------	-----	-----------------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ± 0.15	DAVIS	68	HBC $5.5K^-p$
-----------------	-------	----	---------------

$\Gamma(\omega\gamma)/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0275 ± 0.0022 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0234 $\pm 0.0030 \pm 0.0004$	70	43 PEDLAR	09	CLEO $J/\psi \rightarrow \gamma\eta'$
--------------------------------	----	-----------	----	---------------------------------------

43 Not independent of other η' branching fractions and ratios in PEDLAR 09.

$\Gamma(\omega\gamma)/\Gamma(\pi^+\pi^-\eta)$

Γ_4/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.063 ± 0.005 OUR FIT				

$0.055 \pm 0.007 \pm 0.001$	PEDLAR	09	CLE3 $J/\psi \rightarrow \eta'\gamma$
-----------------------------------------------	--------	----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.068 ± 0.013	68	ZANFINO	77	ASPK $8.4\pi^-p$
-------------------	----	---------	----	------------------

$\Gamma(\omega\gamma)/\Gamma(\pi^0\pi^0\eta)$

Γ_4/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.127 ± 0.011 OUR FIT			

0.147 ± 0.016	ALDE	87B	GAM2 $38\pi^-p \rightarrow n4\gamma$
-------------------------------------	------	-----	--------------------------------------

$\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/[\Gamma(\pi^+\pi^-\eta) + \Gamma(\pi^0\pi^0\eta) + \Gamma(\omega\gamma)]$

$\Gamma_2/(\Gamma_1+\Gamma_3+\Gamma_4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.433 ± 0.012 OUR FIT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.25 ± 0.14	DAUBER	64	HBC $1.95K^-p$
-----------------	--------	----	----------------

$[\Gamma(\pi^0\pi^0\eta(\text{charged decay})) + \Gamma(\omega(\text{charged decay})\gamma)]/\Gamma_{\text{total}}$

$(0.286\Gamma_3+0.89\Gamma_4)/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0863 ± 0.0032 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.045 ± 0.029	42	RITTENBERG	69	HBC $1.7\text{--}2.7K^-p$
-------------------	----	------------	----	---------------------------

$\Gamma(\pi^+ \pi^- \text{ neutrals})/\Gamma_{\text{total}}$ $(0.714\Gamma_1 + 0.286\Gamma_3 + 0.89\Gamma_4)/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.396 ± 0.004 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.4 ± 0.1	39	LONDON	66	HBC $2.24 K^- p \rightarrow \Lambda \pi^+ \pi^- \text{ neutrals}$
0.35 ± 0.06	33	BADIER	65B	HBC $3 K^- p$

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.18 ± 0.08 OUR FIT				
2.00 ± 0.15 OUR AVERAGE				

1.98 ^{+0.31} _{-0.27} ± 0.07	114	⁴⁴ WICHT	08	BELL $B^\pm \rightarrow K^\pm \gamma\gamma$
2.00 ± 0.18		⁴⁵ STANTON	80	SPEC $8.45 \pi^- p \rightarrow n \pi^+ \pi^- 2\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.25 ± 0.16 ± 0.03	0.3k	⁴⁶ PEDLAR	09	CLEO $J/\psi \rightarrow \gamma\eta'$
1.8 ± 0.2	6000	⁴⁷ APEL	79	NICE $15-40 \pi^- p \rightarrow n 2\gamma$
2.5 ± 0.7		DUANE	74	MMS $\pi^- p \rightarrow n \text{MM}$
1.71 ± 0.33	68	DALPIAZ	72	CNTR $1.6 \pi^- p \rightarrow n X^0$
2.0 ^{+0.8} _{-0.6}	31	HARVEY	71	OSPK $3.65 \pi^- p \rightarrow n X^0$

⁴⁴ WICHT 08 reports $[\Gamma(\eta'(958) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta' K^+)] = (1.40^{+0.16+0.15}_{-0.15-0.12}) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta' K^+) = (7.06 \pm 0.25) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁵ Includes APEL 79 result.

⁴⁶ Not independent of other η' branching fractions and ratios in PEDLAR 09.

⁴⁷ Data is included in STANTON 80 evaluation.

$\Gamma(\gamma\gamma)/\Gamma(\pi^+ \pi^- \eta)$

Γ_5/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.0503 ± 0.0022 OUR FIT			
0.053 ± 0.004 ± 0.001	PEDLAR	09	CLE3 $J/\psi \rightarrow \eta' \gamma$

$\Gamma(\gamma\gamma)/\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma))$

Γ_5/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
0.0744 ± 0.0033 OUR FIT			
0.080 ± 0.008	ABLIKIM	06E	BES2 $J/\psi \rightarrow \eta' \gamma$

$\Gamma(\gamma\gamma)/\Gamma(\pi^0 \pi^0 \eta)$

Γ_5/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.101 ± 0.004 OUR FIT			
0.105 ± 0.010 OUR AVERAGE			
Error includes scale factor of 1.9.			
0.091 ± 0.009	AMSLER	93	CBAR $0.0 \bar{p}p$
0.112 ± 0.002 ± 0.006	ALDE	87B	GAM2 $38 \pi^- p \rightarrow n 2\gamma$

$\Gamma(\gamma\gamma)/\Gamma(\pi^0 \pi^0 \eta (\text{neutral decay}))$

$\Gamma_5/0.714\Gamma_3$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.141 ± 0.006 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.188 ± 0.058	16	APEL	72	OSPK $3.8 \pi^- p \rightarrow n X^0$

$\Gamma(\text{ neutrals})/\Gamma_{\text{total}}$

VALUE EVTS
0.179±0.006 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.185±0.022	535	BASILE	71	CNTR	1.6 $\pi^- p \rightarrow n X^0$
0.189±0.026	123	RITTENBERG	69	HBC	1.7–2.7 $K^- p$

(0.714 $\Gamma_3+0.09\Gamma_4+\Gamma_5)/\Gamma$

DOCUMENT ID TECN COMMENT

$\Gamma(3\pi^0)/\Gamma(\pi^0\pi^0\eta)$

VALUE (units 10^{-4}) EVTS
78±10 OUR FIT

78±10 OUR AVERAGE

86±19	235	BLIK	08	GAMS	32 $\pi^- p \rightarrow \eta' n$
74±15		ALDE	87B	GAM2	38 $\pi^- p \rightarrow n 6\gamma$
75±18		BINON	84	GAM2	30–40 $\pi^- p \rightarrow n 6\gamma$

Γ_6/Γ_3

$\Gamma(\mu^+\mu^-\gamma)/\Gamma(\gamma\gamma)$

VALUE (units 10^{-3}) EVTS
4.9±1.2

DOCUMENT ID TECN COMMENT

VIKTOROV	80	CNTR	25,33	$\pi^- p \rightarrow 2\mu\gamma$
----------	----	------	-------	----------------------------------

Γ_7/Γ_5

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4}) CL%

DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.4	90	48 NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
------	----	---------	----	------	----------------------------------

48 Not independent of measured value of Γ_8/Γ_1 from NAIK 09.

Γ_8/Γ

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\pi^+\pi^-\eta)$

VALUE (units 10^{-3}) CL%

DOCUMENT ID TECN COMMENT

<0.5	90	49 NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
------	----	---------	----	------	----------------------------------

49 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 1.3 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.

Γ_8/Γ_1

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-2}) CL%

DOCUMENT ID TECN COMMENT

0.36 $^{+0.11}_{-0.09}$ OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.37 $^{+0.11}_{-0.09} \pm 0.04$	50 NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
----------------------------------	---------	----	------	----------------------------------

<9	95	DANBURG	73	HBC	2.2 $K^- p \rightarrow \Lambda X^0$
----	----	---------	----	-----	-------------------------------------

<5	90	RITTENBERG	69	HBC	1.7–2.7 $K^- p$
----	----	------------	----	-----	-----------------

Γ_9/Γ

50 Not independent of measured value of Γ_9/Γ_1 from NAIK 09.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\eta)$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.3 ± 2.5 OUR FIT

8.25
 ± 2.49
 ± 0.04

20 51 NAIK

09 CLEO $J/\psi \rightarrow \gamma\eta'$

51 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (21^{+6}_{-5} \pm 2) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0\rho^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	RITTENBERG 65	HBC	$2.7 K^- p$

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.4	90	52 NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
<100	90	RITTENBERG 69	HBC	1.7–2.7	$K^- p$

52 Not independent of measured value of Γ_{11}/Γ_1 from NAIK 09.

$\Gamma(2(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.6	90	53 NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$

53 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 1.4 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<27	90	54 NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
-----	----	---------	----	------	----------------------------------

54 Not independent of measured value of Γ_{12}/Γ_1 from NAIK 09.

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi^+\pi^-\eta)$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	55 NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$

55 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 15 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.

$\Gamma(2(\pi^+\pi^-) \text{ neutrals})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	95	DANBURG 73	HBC	$2.2 K^- p \rightarrow \Lambda X^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.01	90	RITTENBERG 69	HBC	1.7–2.7 $K^- p$
-------	----	---------------	-----	-----------------

Γ_9/Γ_1

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.002	90	56 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<0.01	90	RITTENBERG 69	HBC	$1.7\text{--}2.7 K^- p$

56 Not independent of measured value of Γ_{14}/Γ_1 from NAIK 09.

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{14}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	57 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
57 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 11 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.				

$\Gamma(2(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	95	KALBFLEISCH 64B	HBC	$K^- p \rightarrow \Lambda 2(\pi^+\pi^-)+\text{MM}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.01	90	LONDON	66 HBC	Compilation

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.53	90	58 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<5	95	KALBFLEISCH 64B	HBC	$K^- p \rightarrow \Lambda 2(\pi^+\pi^-)$

58 Not independent of measured value of Γ_{16}/Γ_1 from NAIK 09.

$\Gamma(3(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ Γ_{16}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	59 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
59 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 3(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 3.0 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.				

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
$2.4^{+1.3}_{-1.0}$ OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.5^{+1.2}_{-0.9} \pm 0.5$	60 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<6	RITTENBERG 65	HBC	$2.7 K^- p$

60 Not independent of measured value of Γ_{17}/Γ_1 from NAIK 09.

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\pi^+\pi^-\eta)$

Γ_{17}/Γ_1

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	-------------	--------------------	-------------	----------------

5.6 $^{+3.0}_{-2.2}$ OUR FIT

5.50
 $^{+2.99}_{-2.29} \pm 0.03$

8 61 NAIK

09 CLEO $J/\psi \rightarrow \gamma\eta'$

61 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-e^+e^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (14^{+7}_{-5} \pm 3) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\gamma e^+e^-)/\Gamma_{\text{total}}$

Γ_{18}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

<0.9

90 BRIERE

00 CLEO $10.6 e^+e^-$

$\Gamma(\pi^0\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$

Γ_{19}/Γ_3

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

<37

90 ALDE 87B GAM2 $38 \pi^- p \rightarrow n4\gamma$

$\Gamma(4\pi^0)/\Gamma(\pi^0\pi^0\eta)$

Γ_{20}/Γ_3

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

<23

90 ALDE 87B GAM2 $38 \pi^- p \rightarrow n8\gamma$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$

Γ_{21}/Γ

<u>VALUE</u> (units 10^{-7})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

<2.1

90 VOROBIEV 88 ND $e^+e^- \rightarrow \pi^+\pi^-\eta$

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$

Γ_{22}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9.5

90 62 NAIK 09 CLEO $J/\psi \rightarrow \gamma\eta'$

62 Not independent of measured value of Γ_{22}/Γ_1 from NAIK 09.

$\Gamma(\text{invisible})/\Gamma(\gamma\gamma)$

Γ_{22}/Γ_5

<u>VALUE</u> (units 10^{-2})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.69

90 ABLIKIM 06Q BES $J/\psi \rightarrow \phi\eta'$

$\Gamma(\text{invisible})/\Gamma(\pi^+\pi^-\eta)$

Γ_{22}/Γ_1

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

<2.1

90 63 NAIK 09 CLEO $J/\psi \rightarrow \gamma\eta'$

63 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \text{invisible})/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 5.4 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.31 \times 10^{-2}$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 29	90	64 MORI	07A BELL	$\gamma\gamma \rightarrow \pi^+\pi^-$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
< 3.3	90	65 MORI	07A BELL	$\gamma\gamma \rightarrow \pi^+\pi^-$
<800	95	DANBURG	73 HBC	$2.2 K^- p \rightarrow \Lambda X^0$
<200	90	RITTENBERG	69 HBC	$1.7-2.7 K^- p$

⁶⁴ Taking into account interference with the $\gamma\gamma \rightarrow \pi^+\pi^-$ continuum.

⁶⁵ Without interference with the $\gamma\gamma \rightarrow \pi^+\pi^-$ continuum.

$\Gamma(\pi^0\pi^0)/\Gamma(\pi^0\pi^0\eta)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<45	90	ALDE	87B GAM2	$38 \pi^- p \rightarrow n 4\gamma$

$\Gamma(\pi^0e^+e^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.4	90	BRIERE	00 CLEO	$10.6 e^+e^-$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<13	90	RITTENBERG	65 HBC	$2.7 K^- p$

$\Gamma(\eta e^+e^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 2.4	90	BRIERE	00 CLEO	$10.6 e^+e^-$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<11	90	RITTENBERG	65 HBC	$2.7 K^- p$

$\Gamma(3\gamma)/\Gamma(\pi^0\pi^0\eta)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.6	90	ALDE	87B GAM2	$38 \pi^- p \rightarrow n 3\gamma$

$\Gamma(\mu^+\mu^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.0	90	DZHELYADIN	81 CNTR	$30 \pi^- p \rightarrow \eta' n$

$\Gamma(\mu^+\mu^-\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.5	90	DZHELYADIN	81 CNTR	$30 \pi^- p \rightarrow \eta' n$

$\Gamma(e\mu)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.7	90	BRIERE	00 CLEO	$10.6 e^+e^-$

Γ_{23}/Γ

Γ_{24}/Γ_3

Γ_{25}/Γ

Γ_{26}/Γ

Γ_{28}/Γ

Γ_{29}/Γ

Γ_{30}/Γ

$\eta'(958)$ C-NONCONSERVING DECAY PARAMETER

See the note on η decay parameters in the Stable Particle Particle Listings for definition of this parameter.

DECAY ASYMMETRY PARAMETER FOR $\pi^+ \pi^- \gamma$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.03 ± 0.04 OUR AVERAGE				
-0.019 ± 0.056		AIHARA 87	TPC	$2\gamma \rightarrow \pi^+ \pi^- \gamma$
-0.069 ± 0.078	295	GRIGORIAN 75	STRC	$2.1 \pi^- p$
0.00 ± 0.10	103	KALBFLEISCH 75	HBC	$2.18 K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.07 ± 0.08	152	RITTENBERG 65	HBC	$2.1-2.7 K^- p$

$\eta'(958)$ REFERENCES

ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BES III Collab.)
CZERWINSKI	10	PRL 105 122001	E. Czerwinski <i>et al.</i>	(COSY-11 Collab.)
BLIK	09	PAN 72 231	A.M. Blik <i>et al.</i>	(IHEP (Protvino))
		Translated from YAF 72 258.		
NAIK	09	PRL 102 061801	P. Naik <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
BLIK	08	PAN 71 2124	A. Blik <i>et al.</i>	(GAMS-4π Collab.)
		Translated from YAF 71 2161.		
LIBBY	08	PRL 101 182002	J. Libby <i>et al.</i>	(CLEO Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
DOROFEEV	07	PL B651 22	V. Dorofeev <i>et al.</i>	(VES Collab.)
MORI	07A	JPSJ 76 074102	T. Mori <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06Q	PRL 97 202002	M. Ablikim <i>et al.</i>	(BES Collab.)
AMELIN	05A	PAN 68 372	D.V. Amelin <i>et al.</i>	(VES Collab.)
		Translated from YAF 68 401.		
AMSLER	04B	EPJ C33 23	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
BRIERE	00	PRL 84 26	R. Briere <i>et al.</i>	(CLEO Collab.)
ACCIARRI	98Q	PL B418 399	M. Acciari <i>et al.</i>	(L3 Collab.)
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)
WURZINGER	96	PL B374 283	R. Wurzinger <i>et al.</i>	(BONN, ORSAY, SACL+)
PDG	94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)
AMSLER	93	ZPHY C58 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bityukov, G.V. Borisov	(SERP+)
		Translated from YAF 55 2748.		
KARCH	92	ZPHY C54 33	K. Karch <i>et al.</i>	(Crystal Ball Collab.)
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BEHREND	91	ZPHY C49 401	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BARU	90	ZPHY C48 581	S.E. Baru <i>et al.</i>	(MD-1 Collab.)
BUTLER	90	PR D42 1368	F. Butler <i>et al.</i>	(Mark II Collab.)
KARCH	90	PL B249 353	K. Karch <i>et al.</i>	(Crystal Ball Collab.)
ROE	90	PR D41 17	N.A. Roe <i>et al.</i>	(ASP Collab.)
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2γ Collab.)
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)
		Translated from YAF 48 436.		
WILLIAMS	88	PR D38 1365	D.A. Williams <i>et al.</i>	(Crystal Ball Collab.)
AIHARA	87	PR D35 2650	H. Aihara <i>et al.</i>	(TPC-2γ Collab.)
ALBRECHT	87B	PL B199 457	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALDE	87B	ZPHY C36 603	D.M. Alde <i>et al.</i>	(LANL, BELG, SERP, LAPP)
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)
ALDE	86	PL B177 115	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
BARTEL	85E	PL 160B 421	W. Bartel <i>et al.</i>	(JADE Collab.)
ALTHOFF	84E	PL 147B 487	M. Althoff <i>et al.</i>	(TASSO Collab.)
BERGER	84B	PL 142B 125	C. Berger	(PLUTO Collab.)
BINON	84	PL 140B 264	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP+)

BEHREND	83B	PL 125B 518 (erratum)	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
Also		PL 114B 378	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
JENNI	83	PR D27 1031	P. Jenni <i>et al.</i>	(SLAC, LBL)
BARTEL	82B	PL 113B 190	W. Bartel <i>et al.</i>	(JADE Collab.)
DZHELYADIN	81	PL 105B 239	R.I. Dzhelyadin <i>et al.</i>	(SERP)
STANTON	80	PL B92 353	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+)
VIKTOROV	80	SJNP 32 520	V.A. Viktorov <i>et al.</i>	(SERP)
		Translated from YAF 32	1005.	
APEL	79	PL 83B 131	W.D. Apel, K.H. Augenstein, E. Bertolucci	(KARLK+)
BINNIE	79	PL 83B 141	D.M. Binnie <i>et al.</i>	(LOIC)
ZANFINO	77	PRL 38 930	C. Zanfino <i>et al.</i>	(CARL, MCGI, OHIO+)
GRIGORIAN	75	NP B91 232	A. Grigorian <i>et al.</i>	(+)
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)
DUANE	74	PRL 32 425	A. Duane <i>et al.</i>	(LOIC, SHMP)
KALBFLEISCH	74	PR D10 916	G.R. Kalbfleisch	(BNL)
DANBURG	73	PR D8 3744	J.S. Danburg <i>et al.</i>	(BNL, MICH) JP
JACOBS	73	PR D8 18	S.M. Jacobs <i>et al.</i>	(BRAN, UMD, SYRA+) JP
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)
APEL	72	PL 40B 680	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)
DALPIAZ	72	PL 42B 377	P.F. Dalpiaz <i>et al.</i>	(CERN)
BASILE	71	NC 3A 371	M. Basile <i>et al.</i>	(CERN, BGNA, STRB)
HARVEY	71	PRL 27 885	E.H. Harvey <i>et al.</i>	(MINN, MICH)
BENSINGER	70	PL 33B 505	J.R. Bensinger <i>et al.</i>	(WISC)
RITTENBERG	69	Thesis UCRL 18863	A. Rittenberg	(LRL) I
DAVIS	68	PL 27B 532	R. Davis <i>et al.</i>	(NWES, ANL)
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IJP
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)
RITTENBERG	65	PRL 15 556	A. Rittenberg, G.R. Kalbfleisch	(LRL, BNL)
DAUBER	64	PRL 13 449	P.M. Dauber <i>et al.</i>	(UCLA) JP
KALBFLEISCH	64B	PRL 13 349	G.R. Kalbfleisch, O.I. Dahl, A. Rittenberg	(LRL) JP