

**$f_0(1710)$**

$I^G(J^{PC}) = 0^+(0^{++})$

See our mini-review in the 2004 edition of this *Review*, Physics Letters **B592** 1 (2004). See also the mini-review on scalar mesons under  $f_0(600)$  (see the index for the page number).

## $f_0(1710)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1720 \pm 6</math> OUR AVERAGE</b>				Error includes scale factor of 1.6. See the ideogram below.
$1701 \pm 5$	$\pm 9$	4k	1 CHEKANOV	$08$ ZEUS $e p \rightarrow K_S^0 K_S^0 X$
$1765 \pm 3$	$\pm 13$		ABLIKIM	$06V$ BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
$1760 \pm 15$	$\pm 15$		2 ABLIKIM	$05Q$ BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
$1738 \pm 30$			ABLIKIM	$04E$ BES2 $J/\psi \rightarrow \omega K^+ K^-$
$1740 \pm 4$	$\pm 10$		3 BAI	$03G$ BES $J/\psi \rightarrow \gamma K\bar{K}$
$1740 \pm 25$			3 BAI	$00A$ BES $J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
$1698 \pm 18$			4 BARBERIS	$00E$ $450 \text{ pp} \rightarrow p_f \eta \eta p_s$
$1710 \pm 12$	$\pm 11$		5 BARBERIS	$99D$ OMEG $450 \text{ pp} \rightarrow K^+ K^-, \pi^+ \pi^-$
$1710 \pm 25$			6 FRENCH	$99$ $300 \text{ pp} \rightarrow p_f(K^+ K^-)p_s$
$1707 \pm 10$			7 AUGUSTIN	$88$ DM2 $J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0 X$
$1698 \pm 15$			7 AUGUSTIN	$87$ DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
$1720 \pm 10$	$\pm 10$		8 BALTRUSAIT..87	$MRK3$ $J/\psi \rightarrow \gamma K^+ K^-$
$1742 \pm 15$			7 WILLIAMS	$84$ MPSF $200 \pi^- N \rightarrow 2K_S^0 X$
$1670 \pm 50$			BLOOM	$83$ CBAL $J/\psi \rightarrow \gamma 2\eta$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$1750 \pm 13$		AMSLER	$06$ CBAR	$1.64 \bar{p}p \rightarrow K^+ K^- \pi^0$
$1747 \pm 5$	$80k$	9,10 UMAN	$06$ E835	$5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
$1776 \pm 15$		VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$1790 \pm 40$		2 ABLIKIM	$05$ BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
$1670 \pm 20$		9 BINON	$05$ GAMS	$33 \pi^- p \rightarrow \eta \eta n$
$1726 \pm 7$	$74$	10 CHEKANOV	$04$ ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
$1732 \pm 15$		11 ANISOVICH	$03$ RVUE	
$1682 \pm 16$		TIKHOMIROV	$03$ SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
$1670 \pm 26$	$3651$	3,12 NICHTIU	$02$ OBLX	
$1770 \pm 12$		13,14 ANISOVICH	$99B$ SPEC	$0.6-1.2 p\bar{p} \rightarrow \eta \eta \pi^0$
$1730 \pm 15$		3 BARBERIS	$99$ OMEG	$450 \text{ pp} \rightarrow p_s p_f K^+ K^-$
$1750 \pm 20$		3 BARBERIS	$99B$ OMEG	$450 \text{ pp} \rightarrow p_s p_f \pi^+ \pi^-$
$1750 \pm 30$		15 ANISOVICH	$98B$ RVUE	Compilation
$1720 \pm 39$		BAI	$98H$ BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
$1775 \pm 1.5$	$57$	16 BARKOV	$98$	$\pi^- p \rightarrow K_S^0 K_S^0 n$
$1690 \pm 11$		17 ABREU	$96C$ DLPH	$Z^0 \rightarrow K^+ K^- + X$
$1696 \pm 5$	$\pm 9$	8 BAI	$96C$ BES	$J/\psi \rightarrow \gamma K^+ K^-$

$1781 \pm 8$	$\begin{array}{c} +10 \\ -31 \end{array}$	<sup>3</sup> BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
$1768 \pm 14$		BALOSHIN	95	SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 \chi$
$1750 \pm 15$		<sup>18</sup> BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
$1620 \pm 16$		<sup>8</sup> BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
$1748 \pm 10$		<sup>7</sup> ARMSTRONG	93C	E760	$\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
$\sim 1750$		BREAKSTONE	93	SFM	$p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$
$1744 \pm 15$		<sup>19</sup> ALDE	92D	GAM2	$38 \pi^- p \rightarrow \eta \eta n$
$1713 \pm 10$		<sup>20</sup> ARMSTRONG	89D	OMEG	$300 p p \rightarrow p p K^+ K^-$
$1706 \pm 10$		<sup>20</sup> ARMSTRONG	89D	OMEG	$300 p p \rightarrow p p K_S^0 K_S^0$
$1700 \pm 15$		<sup>8</sup> BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$1720 \pm 60$		<sup>3</sup> BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$1638 \pm 10$		<sup>21</sup> FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
$1690 \pm 4$		<sup>22</sup> FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
$1755 \pm 8$		<sup>23</sup> ALDE	86C	GAM2	$38 \pi^- p \rightarrow n 2\eta$
$1730 \pm 2$	$-10$	<sup>24</sup> LONGACRE	86	RVUE	$22 \pi^- p \rightarrow n 2K_S^0$
$1650 \pm 50$		BURKE	82	MRK2	$J/\psi \rightarrow \gamma 2\rho$
$1640 \pm 50$		<sup>25,26</sup> EDWARDS	82D	CBAL	$J/\psi \rightarrow \gamma 2\eta$
$1730 \pm 10$	$\pm 20$	<sup>27</sup> ETKIN	82C	MPS	$23 \pi^- p \rightarrow n 2K_S^0$

<sup>1</sup> In the SU(3) based model with a specific interference pattern of the  $f_2(1270)$ ,  $a_2^0(1320)$ , and  $f'_2(1525)$  mesons incoherently added to the  $f_0(1710)$  and non-resonant background.

<sup>2</sup> This state may be different from  $f_0(1710)$ , see CLOSE 05.

<sup>3</sup>  $J^P = 0^+$ .

<sup>4</sup> T-matrix pole.

<sup>5</sup> Supersedes BARBERIS 99 and BARBERIS 99B.

<sup>6</sup>  $J^P = 0^+$ , supersedes by ARMSTRONG 89D.

<sup>7</sup> No  $J^P C$  determination.

<sup>8</sup>  $J^P = 2^+$ .

<sup>9</sup> Breit-Wigner mass.

<sup>10</sup> Systematic errors not estimated.

<sup>11</sup> K-matrix pole, assuming  $J^P = 0^+$ , from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K \bar{K} n$ ,  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ ,  $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$ ,  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^0$ ,  $K^+ K_S^0 \pi^-$  at rest,  $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$ ,  $K_S^0 K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^-$  at rest.

<sup>12</sup> Decaying to  $f_0(1370) \pi \pi$ .

<sup>13</sup>  $J^P = 0^+$ .

<sup>14</sup> Not seen by AMSLER 02.

<sup>15</sup> T-matrix pole, assuming  $J^P = 0^+$

<sup>16</sup> No  $J^P C$  determination.

<sup>17</sup> No  $J^P C$  determination, width not determined.

<sup>18</sup> From a fit to the  $0^+$  partial wave.

<sup>19</sup> ALDE 92D combines all the GAMS-2000 data.

<sup>20</sup>  $J^P = 2^+$ , superseded by FRENCH 99.

<sup>21</sup> From an analysis ignoring interference with  $f'_2(1525)$ .

<sup>22</sup> From an analysis including interference with  $f'_2(1525)$ .


<sup>23</sup> Superseded by ALDE 92D.

<sup>24</sup> Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

<sup>25</sup>  $J^P = 2^+$  preferred.

<sup>26</sup> From fit neglecting nearby  $f'_2(1525)$ . Replaced by BLOOM 83.

27 Superseded by LONGACRE 86.



### $f_0(1710)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>135 ± 8 OUR AVERAGE</b>		Error includes scale factor of 1.1.		
100 ± 24 +7 -22	4k	28 CHEKANOV	08 ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
145 ± 8 ±69		ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
125 ± 25 +10 -15		29 ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
125 ± 20		ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+ K^-$
166 + 5 +15 - 8 -10	30 BAI		03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
120 + 50 - 40	30 BAI		00A BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
120 ± 26		31 BARBERIS	00E	$450 pp \rightarrow p_f \eta \eta p_s$
126 ± 16 ±18		32 BARBERIS	99D OMEG	$450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$
105 ± 34		33 FRENCH	99	$300 pp \rightarrow p_f(K^+ K^-) p_s$
166.4 ± 33.2		34 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0 X$
136 ± 28		34 AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
130 ± 20		35 BALTRUSAIT...	87 MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
57 ± 38		36 WILLIAMS	84 MPSF	$200 \pi^- N \rightarrow 2K_S^0 X$
160 ± 80		BLOOM	83 CBAL	$J/\psi \rightarrow \gamma 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

148	$\pm 40$	AMSLER	06	CBAR	$1.64 \bar{p}p \rightarrow K^+ K^- \pi^0$		
188	$\pm 13$	80k	29,37	UMAN	06	E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
250	$\pm 30$			Vladimirsk...	06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
270	$\pm 60$		38	Ablikim	05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
260	$\pm 50$		29	Binon	05	GAMS	$33 \pi^- p \rightarrow \eta\eta n$
38	$\pm 20$		74	Chekhanov	04	ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
144	$\pm 30$		39,40	Anisovich	03	RVUE	
320	$\pm 50$		40,41	Anisovich	03	RVUE	
102	$\pm 26$			Tikhomirov	03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
267	$\pm 44$	3651	30,42	Nichitiu	02	OBLX	
220	$\pm 40$		43,44	Anisovich	99B	SPEC	$0.6-1.2 p\bar{p} \rightarrow \eta\eta\pi^0$
100	$\pm 25$		30	Barberis	99	OMEG	$450 pp \rightarrow p_s p_f K^+ K^-$
160	$\pm 30$		30	Barberis	99B	OMEG	$450 pp \rightarrow p_s p_f \pi^+ \pi^-$
250	$\pm 140$		45	Anisovich	98B	RVUE	Compilation
30	$\pm 7$		46	Barkov	98		$\pi^- p \rightarrow K_S^0 K_S^0 n$
103	$\pm 18$	$\pm 30$	35	BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
85	$\pm 24$	$\pm 19$	30	BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
56	$\pm 19$			Baloshin	95	SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 X$
160	$\pm 40$		47	Bugg	95	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
160	$\pm 60$	$\pm 20$	35	Bugg	95	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
264	$\pm 25$		34	ARMSTRONG	93C	E760	$\bar{p}p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$
200	to 300			BREAKSTONE	93	SFM	$pp \rightarrow pp\pi^+\pi^-\pi^+\pi^-$
< 80	90% CL		48	ALDE	92D	GAM2	$38 \pi^- p \rightarrow \eta\eta N^*$
181	$\pm 30$		49	ARMSTRONG	89D	OMEG	$300 pp \rightarrow ppK^+ K^-$
104	$\pm 30$		49	ARMSTRONG	89D	OMEG	$300 pp \rightarrow ppK_S^0 K_S^0$
30	$\pm 20$		35	Bolonkin	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
350	$\pm 150$		30	Bolonkin	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
148	$\pm 17$		50	Falvard	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
184	$\pm 6$		51	Falvard	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
122	$\pm 74$	$\pm 15$	52	Longacre	86	RVUE	$22 \pi^- p \rightarrow n2K_S^0$
200	$\pm 100$			Burke	82	MRK2	$J/\psi \rightarrow \gamma 2\rho$
220	$\pm 100$	$\pm 70$	53,54	EDWARDS	82D	CBAL	$J/\psi \rightarrow \gamma 2\eta$
200	$\pm 156$	$\pm 9$	55	ETKIN	82B	MPS	$23 \pi^- p \rightarrow n2K_S^0$

28 In the SU(3) based model with a specific interference pattern of the  $f_2(1270)$ ,  $a_2^0(1320)$ , and  $f'_2(1525)$  mesons incoherently added to the  $f_0(1710)$  and non-resonant background.

29 Breit-Wigner width.

30  $J^P = 0^+$ .

31 T-matrix pole.

32 Supersedes Barberis 99 and Barberis 99B.

33  $J^P = 0^+$ , supersedes by ARMSTRONG 89D.

- 34 No  $J^{PC}$  determination.  
 35  $J^P = 2^+$ .  
 36 No  $J^{PC}$  determination.  
 37 Systematic errors not estimated.  
 38 This state may be different from  $f_0(1710)$ , see CLOSE 05.  
 39 (Solution I)  
 40 K-matrix pole, assuming  $J^P = 0^+$ , from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K\bar{K}n$ ,  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ ,  $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$ ,  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^0$ ,  $K^+ K_S^0 \pi^-$  at rest,  $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$ ,  $K_S^0 K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^-$  at rest.  
 41 (Solution I)  
 42 Decaying to  $f_0(1370)\pi\pi$ .  
 43  $J^P = 0^+$ .  
 44 Not seen by AMSLER 02.  
 45 T-matrix pole, assuming  $J^P = 0^+$   
 46 No  $J^{PC}$  determination.  
 47 From a fit to the  $0^+$  partial wave.  
 48 ALDE 92D combines all the GAMS-2000 data.  
 49  $J^P = 2^+$ , ( $0^+$  excluded).  
 50 From an analysis ignoring interference with  $f'_2(1525)$ .  
 51 From an analysis including interference with  $f'_2(1525)$ .  
 52 Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.  
 53  $J^P = 2^+$  preferred.  
 54 From fit neglecting nearby  $f'_2(1525)$ . Replaced by BLOOM 83.  
 55 From an amplitude analysis of the  $K_S^0 K_S^0$  system, superseded by LONGACRE 86.
- 

## $f_0(1710)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 K\bar{K}$	seen
$\Gamma_2 \eta\eta$	seen
$\Gamma_3 \pi\pi$	seen
$\Gamma_4 \gamma\gamma$	
$\Gamma_5 \omega\omega$	seen

---

## $f_0(1710) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1 \Gamma_4 / \Gamma$
VALUE (eV)	CL%
<110	95

56 BEHREND 89C CELL  $\gamma\gamma \rightarrow K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

	DOCUMENT ID	TECN	COMMENT
<480	ALBRECHT 90G ARG	$\gamma\gamma \rightarrow K^+ K^-$	
<280	ALTHOFF 85B TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$	

56 Assuming helicity 2.

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_4/\Gamma$			
<u>VALUE (keV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.82	95	57 BARATE	00E ALEP	$\gamma\gamma \rightarrow \pi^+\pi^-$
57 Assuming spin 0.				

## $f_0(1710)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.36 ± 0.12	ALBALADEJO 08	RVUE	
0.38 <sup>+0.09</sup> <sub>-0.19</sub>	58,59 LONGACRE 86	MPS	$22\pi^- p \rightarrow n2K_S^0$

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.22 ± 0.12	ALBALADEJO 08	RVUE	
0.18 <sup>+0.03</sup> <sub>-0.13</sub>	58,59 LONGACRE 86	RVUE	

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AMSLER 02	CBAR	$0.9\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
0.039 <sup>+0.002</sup> <sub>-0.024</sub>	58,59 LONGACRE 86	RVUE	

$\Gamma(\pi\pi)/\Gamma(K\bar{K})$	$\Gamma_3/\Gamma_1$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<b>0.41<sup>+0.11</sup><sub>-0.17</sub></b>		ABLIKIM 06V	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
0.32 ± 0.14		ALBALADEJO 08	RVUE	
< 0.11	95	60 ABLIKIM 04E	BES2	$J/\psi \rightarrow \omega K^+K^-$
5.8 <sup>+9.1</sup> <sub>-5.5</sub>		61 ANISOVICH 02D	SPEC	Combined fit
0.2 ± 0.024 ± 0.036		BARBERIS 99D	OMEG	$450\bar{p}p \rightarrow K^+K^-, \pi^+\pi^-$
0.39 ± 0.14		ARMSTRONG 91	OMEG	$300\bar{p}p \rightarrow pp\pi\pi, ppK\bar{K}$

$\Gamma(\eta\eta)/\Gamma(K\bar{K})$	$\Gamma_2/\Gamma_1$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<b>0.48 ± 0.15</b>		BARBERIS 00E		$450\bar{p}p \rightarrow p_f\eta\eta p_s$
0.46 <sup>+0.70</sup> <sub>-0.38</sub>		61 ANISOVICH 02D	SPEC	Combined fit
< 0.02	90	62 PROKOSHKIN 91	GA24	$300\pi^-p \rightarrow \pi^-p\eta\eta$

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$	$\Gamma_5/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$
58	From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2.			
59	Fit with constrained inelasticity.			
60	Using data from ABLIKIM 04A.			
61	From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$ , $\pi^0\eta\eta$ , $\pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0n$ , $\eta\eta n$ , $\eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K}n$ ) data.			
62	Combining results of GAM4 with those of ARMSTRONG 89D.			

## f<sub>0</sub>(1710) REFERENCES

ALBALADEJO	08	PRL 101 252002	M. Albaladejo, J.A. Oller	
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
Vladimirsky...	06	PAN 69 493	V.V. Vladimirsyky <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		
CLOSE	05	PR D71 094022	F.E. Close, Q. Zhao	
ABLIKIM	04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)
CHEKANOV	04	PL B578 33	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>	
		Translated from YAF 65 1583.		
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>	
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)
FRENCH	99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
		Translated from UFN 168 481.		
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARKOV	98	JETPL 68 764	B.P. Barkov <i>et al.</i>	
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)
		Translated from YAF 58 50.		
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BREAKSTONE	93	ZPHY C58 251	A.M. Breakstone <i>et al.</i>	(IOWA, CERN, DORT+)
ALDE	92D	PL B284 457	D.M. Alde <i>et al.</i>	(GAM2 Collab.)
Also		SJNP 54 451	D.M. Alde <i>et al.</i>	(GAM2 Collab.)
		Translated from YAF 54 745.		
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2, GAM4 Collab.)
		Translated from DANS 316 900.		

ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
ALDE	86C	PL B182 105	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)
WILLIAMS	84	PR D30 877	E.G.H. Williams <i>et al.</i>	(VAND, NDAM, TUFTS+)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
ETKIN	82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)