


**$\pi_2(1670)$**  $I^G(J^{PC}) = 1^-(2^-+)$  **$\pi_2(1670)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>1672.2 \pm 3.0</math> OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.			
1658 $\pm$ 3 $\pm$ 24 8	420k	ALEKSEEV	10	COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
1749 $\pm$ 10 $\pm$ 100	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
1676 $\pm$ 3 $\pm$ 8		<sup>1</sup> CHUNG	02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1685 $\pm$ 10 $\pm$ 30		<sup>2</sup> BARBERIS	01		$450 pp \rightarrow p_f 3\pi^0 p_s$
1687 $\pm$ 9 $\pm$ 15		AMELIN	99	VES	$37 \pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
1669 $\pm$ 4		BARBERIS	98B		$450 pp \rightarrow p_f \rho \pi p_s$
1670 $\pm$ 4		BARBERIS	98B		$450 pp \rightarrow p_f f_2(1270) \pi p_s$
1730 $\pm$ 20		<sup>3</sup> AMELIN	95B	VES	$36 \pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
1690 $\pm$ 14		<sup>4</sup> BERDNIKOV	94	VES	$37 \pi^- A \rightarrow K^+ K^- \pi^- A$
1710 $\pm$ 20	700	ANTIPOV	87	SIGM	$50 \pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
1676 $\pm$ 6		<sup>4</sup> EVANGELIS...	81	OMEG	$12 \pi^- p \rightarrow 3\pi p$
1657 $\pm$ 14		<sup>4,5</sup> DAUM	80D	SPEC	$63-94 \pi p \rightarrow 3\pi X$
1662 $\pm$ 10	2000	<sup>4</sup> BALTAY	77	HBC	$15 \pi^+ p \rightarrow p 3\pi$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
1742 $\pm$ 31 $\pm$ 49		ANTREASYAN	90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
1624 $\pm$ 21		<sup>1</sup> BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1622 $\pm$ 35		<sup>6</sup> BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1693 $\pm$ 28		<sup>7</sup> BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1710 $\pm$ 20		<sup>8</sup> DAUM	81B	SPEC	$63,94 \pi^- p$
1660 $\pm$ 10		<sup>4</sup> ASCOLI	73	HBC	$5-25 \pi^- p \rightarrow p \pi_2$

<sup>1</sup> From  $f_2(1270)\pi$  decay.<sup>2</sup> From a fit to the invariant mass distribution.<sup>3</sup> From a fit to  $J^{PC} = 2^-+ f_2(1270)\pi$ ,  $f_0(1370)\pi$  waves.<sup>4</sup> From a fit to  $J^P = 2^- S$ -wave  $f_2(1270)\pi$  partial wave.<sup>5</sup> Clear phase rotation seen in  $2^- S$ ,  $2^- P$ ,  $2^- D$  waves. We quote central value and spread of single-resonance fits to three channels.<sup>6</sup> From  $\rho \pi$  decay.<sup>7</sup> From  $\sigma \pi$  decay.<sup>8</sup> From a two-resonance fit to four  $2^- 0^+$  waves. This should not be averaged with all the single resonance fits.



### $\pi_2(1670)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>260± 9 OUR AVERAGE</b>		Error includes scale factor of 1.2.			
271± 9 <sup>+ 22</sup> <sub>- 24</sub>	420k	ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
408± 60±250	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
254± 3± 31		<sup>9</sup> CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
265± 30± 40		<sup>10</sup> BARBERIS	01		450 $pp \rightarrow p_f 3\pi^0 p_s$
168± 43± 53		AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
268± 15		BARBERIS	98B		450 $pp \rightarrow p_f \rho \pi p_s$
256± 15		BARBERIS	98B		450 $pp \rightarrow p_f f_2(1270) \pi p_s$
310± 20		<sup>11</sup> AMELIN	95B	VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
190± 50		<sup>12</sup> BERDNIKOV	94	VES	37 $\pi^- A \rightarrow K^+ K^- \pi^- A$
170± 80	700	ANTIPOV	87	SIGM	50 $\pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
260± 20		<sup>12</sup> EVANGELIS...	81	OMEG	12 $\pi^- p \rightarrow 3\pi p$
219± 20		<sup>12,13</sup> DAUM	80D	SPEC	63–94 $\pi p \rightarrow 3\pi X$
285± 60	2000	<sup>12</sup> BALTAY	77	HBC	15 $\pi^+ p \rightarrow p 3\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$236 \pm 49 \pm 36$	ANTREASYAN 90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
$304 \pm 22$	<sup>9</sup> BELLINI	85	SPEC $40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
$404 \pm 108$	<sup>14</sup> BELLINI	85	SPEC $40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
$330 \pm 90$	<sup>15</sup> BELLINI	85	SPEC $40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
$312 \pm 50$	<sup>16</sup> DAUM	81B	SPEC $63,94 \pi^- p$
$270 \pm 60$	<sup>12</sup> ASCOLI	73	HBC — $5-25 \pi^- p \rightarrow p \pi_2^-$

<sup>9</sup> From  $f_2(1270)\pi$  decay.

<sup>10</sup> From a fit to the invariant mass distribution.

<sup>11</sup> From a fit to  $J^{PC} = 2^{-+}$   $f_2(1270)\pi$ ,  $f_0(1370)\pi$  waves.

<sup>12</sup> From a fit to  $J^P = 2^- f_2(1270)\pi$  partial wave.

<sup>13</sup> Clear phase rotation seen in  $2^- S$ ,  $2^- P$ ,  $2^- D$  waves. We quote central value and spread of single-resonance fits to three channels.

<sup>14</sup> From  $\rho\pi$  decay.

<sup>15</sup> From  $\sigma\pi$  decay.

<sup>16</sup> From a two-resonance fit to four  $2^{-0^+}$  waves. This should not be averaged with all the single resonance fits.

## $\pi_2(1670)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1 3\pi$	(95.8 ± 1.4) %	
$\Gamma_2 \pi^+ \pi^- \pi^0$		
$\Gamma_3 \pi^0 \pi^0 \pi^0$		
$\Gamma_4 f_2(1270)\pi$	(56.3 ± 3.2) %	
$\Gamma_5 \rho\pi$	(31 ± 4) %	
$\Gamma_6 \sigma\pi$	(10.9 ± 3.4) %	
$\Gamma_7 (\pi\pi)_{S\text{-wave}}$	( 8.7 ± 3.4) %	
$\Gamma_8 K\bar{K}^*(892) + \text{c.c.}$	( 4.2 ± 1.4) %	
$\Gamma_9 \omega\rho$	( 2.7 ± 1.1) %	
$\Gamma_{10} \gamma\gamma$	< 2.8 × 10 <sup>-7</sup>	90%
$\Gamma_{11} \eta\pi$		
$\Gamma_{12} \pi^\pm 2\pi^+ 2\pi^-$		
$\Gamma_{13} \rho(1450)\pi$	< 3.6 × 10 <sup>-3</sup>	97.7%
$\Gamma_{14} b_1(1235)\pi$	< 1.9 × 10 <sup>-3</sup>	97.7%
$\Gamma_{15} \eta 3\pi$		
$\Gamma_{16} f_1(1285)\pi$	possibly seen	
$\Gamma_{17} a_2(1320)\pi$	not seen	

## CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 6 measurements and one constraint to determine 4 parameters. The overall fit has a  $\chi^2 = 1.9$  for 3 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_5$	-53		
$x_7$	-29	-59	
$x_8$	-8	-21	-9
	$x_4$	$x_5$	$x_7$

## $\pi_2(1670)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	$\Gamma_{10}$				
VALUE (keV)	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt;0.072</b>	90	17 ACCIARRI	97T	L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>					
<0.19	90	17 ALBRECHT	97B	ARG	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
1.41 $\pm 0.23 \pm 0.28$		ANTREASYAN	90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
0.8 $\pm 0.3 \pm 0.12$		18 BEHREND	90C	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
1.3 $\pm 0.3 \pm 0.2$		19 BEHREND	90C	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$

17 Decaying into  $f_2(1270)\pi$  and  $\rho\pi$ .

18 Constructive interference between  $f_2(1270)\pi, \rho\pi$  and background.

19 Incoherent Ansatz.

## $\pi_2(1670) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

$\Gamma(\pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$	$\Gamma_2 \Gamma_{10} / \Gamma$				
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.1</b>	95	20 SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

20 From analysis of L3 data at 183–209 GeV.

## $\pi_2(1670)$ BRANCHING RATIOS

$\Gamma(3\pi) / \Gamma_{\text{total}}$	$\Gamma_1 / \Gamma = (\Gamma_4 + \Gamma_5 + \Gamma_7) / \Gamma$		
VALUE	DOCUMENT ID		
<b>0.958 <math>\pm 0.014</math> OUR FIT</b>			

$$\Gamma(\pi^0 \pi^0 \pi^0)/\Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_3/\Gamma_2$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.29±0.03±0.05</b>	21 BARBERIS	01	$450 \text{ } pp \rightarrow p_f 3\pi^0 p_s$

$$\Gamma(\rho\pi)/0.565\Gamma(f_2(1270)\pi) \quad \Gamma_5/0.565\Gamma_4$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.97±0.09 OUR AVERAGE</b>			Error includes scale factor of 1.9.
0.76±0.07±0.10	CHUNG	02	$B852 \quad 18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1.01±0.05	BARBERIS	98B	$450 \text{ } pp \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$

$$\Gamma(\sigma\pi)/\Gamma(f_2(1270)\pi) \quad \Gamma_6/\Gamma_4$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.19±0.06 OUR AVERAGE</b>			
0.17±0.02±0.07	CHUNG	02	$B852 \quad 18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
0.24±0.10	22,23 BAKER	99	$SPEC \quad 1.94 \bar{p}p \rightarrow 4\pi^0$

$$\frac{1}{2}\Gamma(\rho\pi)/\Gamma(\pi^\pm \pi^\mp \pi^\pm) \quad \frac{1}{2}\Gamma_5/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<b>0.29±0.04 OUR FIT</b>				
<b>0.29±0.05</b>	24 DAUM	81B	SPEC	$63,94 \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.3	BARTSCH	68	HBC	$+ \quad 8 \pi^+ p \rightarrow 3\pi p$

$$0.565\Gamma(f_2(1270)\pi)/\Gamma(\pi^\pm \pi^\mp \pi^\pm) \quad 0.565\Gamma_4/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<b>0.604±0.035 OUR FIT</b>				
<b>0.60 ±0.05 OUR AVERAGE</b>				Error includes scale factor of 1.3.
0.61 ±0.04	24 DAUM	81B	SPEC	$63,94 \pi^- p$
0.76 +0.24 -0.34	ARMENISE	69	DBC	$+ \quad 5.1 \pi^+ d \rightarrow d 3\pi$
0.35 ±0.20	BALTAY	68	HBC	$+ \quad 7-8.5 \pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.59	BARTSCH	68	HBC	$+ \quad 8 \pi^+ p \rightarrow 3\pi p$

$$0.624\Gamma((\pi\pi)_{S\text{-wave}})/\Gamma(\pi^\pm \pi^\mp \pi^\pm) \quad 0.624\Gamma_7/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.10±0.04 OUR FIT</b>			
<b>0.10±0.05</b>	24 DAUM	81B	SPEC $63,94 \pi^- p$

$$\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma(f_2(1270)\pi) \quad \Gamma_8/\Gamma_4$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<b>0.075±0.025 OUR FIT</b>				
<b>0.075±0.025</b>	25 ARMSTRONG	82B	OMEG	$- \quad 16 \pi^- p \rightarrow K^+ K^- \pi^- p$

$$\Gamma(\omega\rho)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.027±0.004±0.010</b>	26 AMELIN	99	VES $37 \pi^- A \xrightarrow{\omega\pi^- \pi^0} A^*$

$\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$   
(All  $\eta$  decays.)

VALUE

**<0.09**

DOCUMENT ID

BALTAY 68 HBC + 7–8.5  $\pi^+ p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.10

DOCUMENT ID

CRENNELL 70 HBC – 6  $\pi^- p \rightarrow f_2\pi^- N$

$\Gamma_{11}/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$

$\Gamma(\pi^\pm 2\pi^+ 2\pi^-)/\Gamma(\pi^\pm\pi^+\pi^-)$

VALUE

**<0.10**

DOCUMENT ID

CRENNELL 70 HBC – 6  $\pi^- p \rightarrow f_2\pi^- N$

$\Gamma_{12}/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$

$\Gamma(\rho(1450)\pi)/\Gamma_{\text{total}}$

VALUE

**<0.0036**

CL%

97.7

DOCUMENT ID

AMELIN 99 VES 37  $\pi^- A \xrightarrow{\omega\pi^-\pi^0} A^*$

$\Gamma_{13}/\Gamma$

$\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$

VALUE

**<0.0019**

CL%

97.7

DOCUMENT ID

AMELIN 99 VES 37  $\pi^- A \xrightarrow{\omega\pi^-\pi^0} A^*$

$\Gamma_{14}/\Gamma$

$\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}}$

VALUE

**possibly seen**

EVTS

69k

DOCUMENT ID

KUHN 04 B852 18  $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

$\Gamma_{16}/\Gamma$

$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$

VALUE

**not seen**

EVTS

69k

DOCUMENT ID

KUHN 04 B852 18  $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

$\Gamma_{17}/\Gamma$

**D-wave/S-wave RATIO FOR  $\pi_2(1670) \rightarrow f_2(1270)\pi$**

VALUE

**-0.18±0.06**

DOCUMENT ID

22 BAKER 99 SPEC 1.94  $\bar{p}p \rightarrow 4\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.22±0.10

DOCUMENT ID

24 DAUM 81B SPEC 63,94  $\pi^- p$

**F-wave/P-wave RATIO FOR  $\pi_2(1670) \rightarrow \rho\pi$**

VALUE

**-0.72±0.07±0.14**

DOCUMENT ID

CHUNG 02 B852 18.3  $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$

21 Using BARBERIS 98B.

22 Using preliminary CBAR data.

23 With the  $\sigma\pi$  in  $L=2$  and the  $f_2(1270)\pi$  in  $L=0$ .

24 From a two-resonance fit to four  $2^-0^+$  waves.

25 From a partial-wave analysis of  $K^+K^-\pi^-$  system.

26 Normalized to the  $B(\pi_2(1670) \rightarrow f_2\pi)$ .

## $\pi_2(1670)$ REFERENCES

ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>	
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)
		Translated from YAF 62 487.		
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>	
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
BERDNIKOV	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)
ANTREASYAN	90	ZPHY C48 561	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BEHREND	90C	ZPHY C46 583	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
ANTIPOV	87	EPL 4 403	Y.M. Antipov <i>et al.</i>	(SERP, JINR, INRM+)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	
		Translated from YAF 41 1223.		
ARMSTRONG	82B	NP B202 1	T.A. Armstrong, B. Baccari	(AACH3, BARI, BONN+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
Also		NP B186 594	C. Evangelista	
DAUM	80D	PL 89B 285	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) JP
BALTAY	77	PRL 39 591	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP
ASCOLI	73	PR D7 669	G. Ascoli	(ILL, TNTO, GENO, HAMB, MILA+) JP
CRENNELL	70	PRL 24 781	D.J. Crennell <i>et al.</i>	(BNL)
ARMENISE	69	LNC 2 501	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE) I
BARTSCH	68	NP B7 345	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN) JP

---