

**$\rho_5(2350)$**

$I^G(J^{PC}) = 1^+(5^{--})$

### OMMITTED FROM SUMMARY TABLE

This entry was previously called  $U_1(2400)$ . See also  $\rho(2150)$ ,  $f_2(2150)$ ,  $\rho_3(2250)$ ,  $f_4(2300)$ .

### **$\rho_5(2350)$ MASS**

#### $\pi^- p \rightarrow \omega\pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>2330 \pm 35</math></b>	ALDE 95	GAM2	$38 \pi^- p \rightarrow \omega\pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$\sim 2303$	HASAN 94	RVUE		$\bar{p}p \rightarrow \pi\pi$
$\sim 2300$	<sup>1</sup> MARTIN 80B	RVUE		
$\sim 2250$	<sup>1</sup> MARTIN 80C	RVUE		
$\sim 2500$	<sup>2</sup> CARTER 78B	CNTR 0		$0.7\text{--}2.4 \bar{p}p \rightarrow$
$\sim 2480$	<sup>3</sup> CARTER 77	CNTR 0		$K^-\bar{K}^+ \quad 0.7\text{--}2.4 \bar{p}p \rightarrow \pi\pi$

#### **S-CHANNEL $\bar{N}N$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$2300 \pm 45$	<sup>4</sup> ANISOVICH 02	SPEC		$0.6\text{--}1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
$2295 \pm 30$	ANISOVICH 00J	SPEC		
$\sim 2380$	<sup>5</sup> CUTTS 78B	CNTR		$0.97\text{--}3 \bar{p}p \rightarrow \bar{N}N$
$2345 \pm 15$	<sup>5,6</sup> COUPLAND 77	CNTR 0		$0.7\text{--}2.4 \bar{p}p \rightarrow \bar{p}p$
$2359 \pm 2$	<sup>5,7</sup> ALSPECTOR 73	CNTR		$\bar{p}p$ S channel
$2350 \pm 10$	<sup>8</sup> ABRAMS 70	CNTR		S channel $\bar{N}N$
$2360 \pm 25$	<sup>9</sup> OH 70B	HDBC -0		$\bar{p}(pn), K^*K2\pi$

#### $\pi^- p \rightarrow K^+ K^- n$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$2307 \pm 6$	ALPER 80	CNTR 0		$62 \pi^- p \rightarrow K^+ K^- n$

<sup>1</sup>  $I(J^P) = 1(5^-)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ .<sup>2</sup>  $I = 0(1); J^P = 5^-$  from Barrelet-zero analysis.<sup>3</sup>  $I(J^P) = 1(5^-)$  from amplitude analysis.<sup>4</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.<sup>5</sup> Isospins 0 and 1 not separated.<sup>6</sup> From a fit to the total elastic cross section.<sup>7</sup> Referred to as  $U$  or  $U$  region by ALSPECTOR 73.<sup>8</sup> For  $I = 1 \bar{N}N$ .<sup>9</sup> No evidence for this bump seen in the  $\bar{p}p$  data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.

## $\rho_5(2350)$ WIDTH

### $\pi^- p \rightarrow \omega \pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>400±100</b>	ALDE	95	GAM2 38 $\pi^- p \rightarrow \omega \pi^0 n$

### $\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
~ 169	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 250	<sup>10</sup> MARTIN	80B	RVUE	
~ 300	<sup>10</sup> MARTIN	80C	RVUE	
~ 150	<sup>11</sup> CARTER	78B	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow K^- K^+$
~ 210	<sup>12</sup> CARTER	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

### S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
260± 75	<sup>13</sup> ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega \pi^0, \omega \eta \pi^0, \pi^+ \pi^-$
$235^{+ 65}_{- 40}$	ANISOVICH	00J	SPEC	
$135^{+ 150}_{- 65}$	<sup>14,15</sup> COUPLAND	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
$165^{+ 18}_{- 8}$	<sup>15</sup> ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
< 60	<sup>16</sup> OH	70B	HDBC –0	$\bar{p}(pn), K^* K 2\pi$
~ 140	ABRAMS	67C	CNTR	S channel $\bar{p}N$

### $\pi^- p \rightarrow K^+ K^- n$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
245±20	ALPER	80	CNTR 0	62 $\pi^- p \rightarrow K^+ K^- n$

<sup>10</sup>  $I(J^P) = 1(5^-)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ .

<sup>11</sup>  $I = 0(1)$ ;  $J^P = 5^-$  from Barrelet-zero analysis.

<sup>12</sup>  $I(J^P) = 1(5^-)$  from amplitude analysis.

<sup>13</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

<sup>14</sup> From a fit to the total elastic cross section.

<sup>15</sup> Isospins 0 and 1 not separated.

<sup>16</sup> No evidence for this bump seen in the  $\bar{p}p$  data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.

---

## **$\rho_5(2350)$ REFERENCES**

ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i> (GAMS Collab.) JP
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg (LOQM)
ALPER	80	PL 94B 422	B. Alper <i>et al.</i> (AMST, CERN, CRAC, MPIM+)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan (LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington (DURH) JP
CARTER	78B	NP B141 467	A.A. Carter (LOQM)
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i> (STON, WISC)
CARTER	77	PL 67B 117	A.A. Carter <i>et al.</i> (LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i> (LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i> (RUTG, UPNJ)
OH	73	NP B51 57	B.Y. Oh <i>et al.</i> (MSU)
CHAPMAN	71B	PR D4 1275	J.W. Chapman <i>et al.</i> (MICH)
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i> (BNL)
OH	70B	PRL 24 1257	B.Y. Oh <i>et al.</i> (MSU)
ABRAMS	67C	PRL 18 1209	R.J. Abrams <i>et al.</i> (BNL)

---