

$D_2^*(2460)^0$ 

$$I(J^P) = \frac{1}{2}(2^+)$$

$J^P = 2^+$  assignment strongly favored(ALBRECHT 89B, ALBRECHT 89H), natural parity confirmed by the helicity analysis(DEL-AMO-SANCHEZ 10P),

 **$D_2^*(2460)^0$  MASS**

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2462.6±0.7 OUR FIT</b>	Error includes scale factor of 1.3.			
<b>2461.8±0.8 OUR AVERAGE</b>	Error includes scale factor of 1.2.			
2462.2±0.1±0.8	243k	DEL-AMO-SA..10P	BABR	$e^+e^- \rightarrow D^+\pi^-X$
2460.4±1.2±2.2	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+\pi^-\pi^-$
2461.6±2.1±3.3		<sup>1</sup> ABE	04D BELL	$B^- \rightarrow D^+\pi^-\pi^-$
2464.5±1.1±1.9	5.8k	<sup>1</sup> LINK	04A FOCS	$\gamma A$
2465 ±3 ±3	486	AVERY	94C CLE2	$e^+e^- \rightarrow D^+\pi^-X$
2453 ±3 ±2	128	FRABETTI	94B E687	$\gamma Be \rightarrow D^+\pi^-X$
2461 ±3 ±1	440	AVERY	90 CLEO	$e^+e^- \rightarrow D^{*+}\pi^-X$
2455 ±3 ±5	337	ALBRECHT	89B ARG	$e^+e^- \rightarrow D^+\pi^-X$
2459 ±3 ±2	153	ANJOS	89C TPS	$\gamma N \rightarrow D^+\pi^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2469.1±3.7 <sup>+1.2</sup> <sub>-1.3</sub>	1560±230	<sup>2</sup> CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{(*)+}\pi^-X$
2463.3±0.6±0.8	20k	ABULENCIA	06A CDF	1900 $p\bar{p} \rightarrow D^+\pi^-X$
2461 ±6	126	<sup>3</sup> ABREU	98M DLPH	$e^+e^-$
2466 ±7	1	ASRATYAN	95 BEBC	53,40 $\nu(\bar{\nu}) \rightarrow pX,dX$

<sup>1</sup> Fit includes the contribution from  $D_0^*(2400)^0$ .

<sup>2</sup> Calculated using the mass difference  $m(D_2^{*0}) - m(D^{*+})_{PDG}$  reported below and  $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$  MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of  $^{+1.2}_{-1.3}$  MeV.

<sup>3</sup> No systematic error given.

 **$m_{D_2^{*0}} - m_{D^+}$** 

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>593.0±0.7 OUR FIT</b>	Error includes scale factor of 1.3.			
<b>593.9±0.6±0.5</b>	20k	ABULENCIA	06A CDF	1900 $p\bar{p} \rightarrow D^+\pi^-X$

$$m_{D_2^{*0}} - m_{D^{*+}}$$

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_s^{*0}$ ,  $D_1(2420)^0$ ,  $D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>452.3 ± 0.7 OUR FIT</b>	Error includes scale factor of 1.3.			
<b>458.8 ± 3.7<sup>+1.2</sup><sub>-1.3</sub></b>	1560 ± 230	CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$

### $D_2^*(2460)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>49.0 ± 1.4 OUR AVERAGE</b>	Error includes scale factor of 1.7. See the ideogram below.			
50.5 ± 0.6 ± 0.7	243k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^+ \pi^- X$
41.8 ± 2.5 ± 2.9	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$
49.2 ± 2.3 ± 1.3	20k	ABULENCIA	06A CDF	1900 $p\bar{p} \rightarrow D^+ \pi^- X$
45.6 ± 4.4 ± 6.7		<sup>4</sup> ABE	04D BELL	$B^- \rightarrow D^+ \pi^- \pi^-$
38.7 ± 5.3 ± 2.9	5.8k	<sup>4</sup> LINK	04A FOCS	$\gamma A$
28 <sup>+8</sup> <sub>-7</sub> ± 6	486	AVERY	94C CLE2	$e^+ e^- \rightarrow D^+ \pi^- X$
25 ± 10 ± 5	128	FRABETTI	94B E687	$\gamma Be \rightarrow D^+ \pi^- X$
20 <sup>+9</sup> <sub>-12</sub> <sup>+9</sup> <sub>-10</sub>	440	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
15 <sup>+13</sup> <sub>-10</sub> <sup>+5</sup> <sub>-10</sub>	337	ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- X$
20 ± 10 ± 5	153	ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- X$

<sup>4</sup> Fit includes the contribution from  $D_0^*(2400)^0$ .

WEIGHTED AVERAGE  
 $49.0 \pm 1.4$  (Error scaled by 1.7)



### $D_2^*(2460)^0$ DECAY MODES

$\bar{D}_2^*(2460)^0$  modes are charge conjugates of modes below.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^+ \pi^-$	seen
$\Gamma_2$ $D^*(2010)^+ \pi^-$	seen
$\Gamma_3$ $D^0 \pi^+ \pi^-$	not seen
$\Gamma_4$ $D^{*0} \pi^+ \pi^-$	not seen

### $D_2^*(2460)^0$ BRANCHING RATIOS

$\Gamma(D^+ \pi^-)/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	3.4k	AUBERT 09AB	BABR	$B^- \rightarrow D^+ \pi^- \pi^-$	
seen	337	ALBRECHT 89B	ARG	$e^+ e^- \rightarrow D^+ \pi^- X$	
seen		ANJOS 89C	TPS	$\gamma N \rightarrow D^+ \pi^- X$	

$\Gamma(D^*(2010)^+ \pi^-)/\Gamma_{\text{total}}$				$\Gamma_2/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	ACKERSTAFF 97W	OPAL	$e^+ e^- \rightarrow D^{*+} \pi^- X$	
seen	AVERY 90	CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$	
seen	ALBRECHT 89H	ARG	$e^+ e^- \rightarrow D^* \pi^- X$	

## $\Gamma(D^+\pi^-)/\Gamma(D^*(2010)^+\pi^-)$ $\Gamma_1/\Gamma_2$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.56±0.16 OUR AVERAGE</b>				
1.47±0.03±0.16	379k	DEL-AMO-SA..10P	BABR	$e^+e^- \rightarrow D^{(*)}\pi^- X$
2.8 ±0.8 $\begin{smallmatrix} +0.5 \\ -0.6 \end{smallmatrix}$	1560±230	CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{(*)}\pi^- X$
2.2 ±0.7 ±0.6		AVERY	94C CLE2	$e^+e^- \rightarrow D^{*+}\pi^- X$
2.3 ±0.8		AVERY	90 CLEO	$e^+e^-$
3.0 ±1.1 ±1.5		ALBRECHT	89H ARG	$e^+e^- \rightarrow D^*\pi^- X$
••• We do not use the following data for averages, fits, limits, etc. •••				
1.9 ±0.5		ABE	04D BELL	$B^- \rightarrow D^{(*)}\pi^-\pi^-$

## $\Gamma(D^+\pi^-)/[\Gamma(D^+\pi^-) + \Gamma(D^*(2010)^+\pi^-)]$ $\Gamma_1/(\Gamma_1+\Gamma_2)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
••• We do not use the following data for averages, fits, limits, etc. •••				
0.62±0.03±0.02	8414	<sup>5</sup> AUBERT	09Y BABR	$B^+ \rightarrow D_2^{*0}\ell^+\nu_\ell$
<sup>5</sup> Assuming $\Gamma(\Upsilon(4S) \rightarrow B^+B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths for charged and neutral $D_2^*$ mesons.				

## $D_2^*(2460)^0$ POLARIZATION AMPLITUDE $A_{D_2}$

A polarization amplitude  $A_{D_2}$  is a parameter that depends on the initial polarization of the  $D_2$ . For  $D_2$  decays the helicity angle,  $\theta_H$ , distribution varies like  $1 + A_{D_2} \cos(\theta_H)$ , where  $\theta_H$  is the angle in the  $D^*$  rest frame between the two pions emitted by the  $D_2 \rightarrow D^*\pi$  and  $D^* \rightarrow D\pi$ .

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
••• We do not use the following data for averages, fits, limits, etc. •••				
consistent with $-1$	243k	DEL-AMO-SA..10P	BABR	$e^+e^- \rightarrow D^+\pi^- X$
$-0.74 \begin{smallmatrix} +0.49 \\ -0.38 \end{smallmatrix}$		<sup>6</sup> AVERY	94C CLE2	$e^+e^- \rightarrow D^{*+}\pi^- X$
<sup>6</sup> Systematic uncertainties not estimated.				

## $D_2^*(2460)^0$ REFERENCES

DEL-AMO-SA...10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT 09AB	PR D79 112004	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHEKANOV 09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABULENCIA 06A	PR D73 051104	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABE 04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)
LINK 04A	PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)
ABREU 98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF 97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ASRATYAN 95	ZPHY C68 43	A.E. Asratyan <i>et al.</i>	(BIRM, BELG, CERN+)
AVERY 94C	PL B331 236	P. Avery <i>et al.</i>	(CLEO Collab.)
FRABETTI 94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AVERY 90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)
ALBRECHT 89B	PL B221 422	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP
ALBRECHT 89H	PL B232 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP
ANJOS 89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)