

X(1835)

$$I^G(J^{PC}) = ?^?(?^{-+})$$

OMITTED FROM SUMMARY TABLE

Could be a superposition of several states.

X(1835) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1835.7⁺₋ 5.0⁺₋ 3.2 OUR AVERAGE				
1836.5 ± 3.0 ⁺ ₋ 5.6 ⁺ ₋ 2.1	4265	¹ ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
1833.7 ± 6.1 ± 2.7	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1877.3 ± 6.3 ⁺ ₋ 3.4 ⁺ ₋ 7.4		² ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$
1837 ⁺ ₋ 10 ⁺ ₋ 12 ⁺ ₋ 9 ⁺ ₋ 7	231	^{3,4} ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p \bar{p}$
1831 ± 7		^{4,5} ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma p \bar{p}$
1859 ⁺ ₋ 3 ⁺ ₋ 10 ⁺ ₋ 5 ⁺ ₋ 25		⁴ BAI	03F BES2	$J/\psi \rightarrow \gamma p \bar{p}$

¹ From a fit of the $\pi^+ \pi^- \eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two unconfirmed states $\gamma X(2120)$, and $\gamma X(2370)$, for $M(p \bar{p}) < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0 \pi^+ \pi^- \eta'$.

² The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$. This state may be due also to $\eta_2(1870)$ or to a combination of $X(1835)$ and $\eta_2(1870)$.

³ From a fit of the $p \bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p \bar{p}$ phase space, for $M(p \bar{p}) < 2.85$ GeV.

⁴ Evidence for a threshold enhancement in the $p \bar{p}$ mass spectrum was also reported by ABE 02K, AUBERT, B 05L, and WANG 05A in $B^+ \rightarrow p \bar{p} K^+$, WANG 05A in $B^0 \rightarrow p \bar{p} K_S^0$, ABE 02W in $\bar{B}^0 \rightarrow p \bar{p} D^0$, and WEI 08 in $B^+ \rightarrow p \bar{p} \pi^+$ decays. Not seen by ATHAR 06 in $\Upsilon(1S) \rightarrow p \bar{p} \gamma$.

⁵ From the fit including final state interaction effects in isospin 0 S-wave according to SIBIRTSEV 05A. Systematic errors not estimated.

X(1835) WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
99 ± 50 OUR AVERAGE Error includes scale factor of 2.8.					
190 ± 9 ⁺ ₋ 38 ⁺ ₋ 36		4265	⁶ ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
67.7 ± 20.3 ± 7.7		264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
57 ± 12 ⁺ ₋ 19 ⁺ ₋ 4			⁷ ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$
0 ⁺ ₋ 44 ⁺ ₋ 0		231	^{8,9} ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p \bar{p}$
< 153	90		^{9,10} ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma p \bar{p}$
< 30			⁹ BAI	03F BES2	$J/\psi \rightarrow \gamma p \bar{p}$

- ⁶ From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two unconfirmed states $\gamma X(2120)$, and $\gamma X(2370)$, for $M(p\bar{p}) < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.
- ⁷ The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$. This state may be due also to $\eta_2(1870)$ or to a combination of $X(1835)$ and $\eta_2(1870)$.
- ⁸ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.
- ⁹ Evidence for a threshold enhancement in the $p\bar{p}$ mass spectrum was also reported by ABE 02K, AUBERT,B 05L, and WANG 05A in $B^+ \rightarrow p\bar{p}K^+$, WANG 05A in $B^0 \rightarrow p\bar{p}K_S^0$, ABE 02W in $\bar{B}^0 \rightarrow p\bar{p}D^0$, and WEI 08 in $B^+ \rightarrow p\bar{p}\pi^+$ decays. Not seen by ATHAR 06 in $\Upsilon(1S) \rightarrow p\bar{p}\gamma$.
- ¹⁰ From the fit including final state interaction effects in isospin 0 S -wave according to SIBIRTSEV 05A. Systematic errors not estimated.

X(1835) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $p\bar{p}$	seen
Γ_2 $\pi^+\pi^-\eta'$	seen

X(1835) BRANCHING RATIOS

$\Gamma(p\bar{p})/\Gamma(\pi^+\pi^-\eta')$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_2
	0.333	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$	

••• We do not use the following data for averages, fits, limits, etc. •••

X(1835) REFERENCES

ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
WEI	08	PL B659 80	J.-T. Wei <i>et al.</i>	(BELLE Collab.)
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT,B	05L	PR D72 051101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
ABE	02K	PRL 88 181803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02W	PRL 89 151802	K. Abe <i>et al.</i>	(BELLE Collab.)