

**$\eta_b(1S)$** 

$$I^G(J^{PC}) = 0^+(0^{-+})$$

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions. Observed in radiative decay of the  $\Upsilon(3S)$ , therefore  $C = +$ . **$\eta_b(1S)$  MASS**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9391.0 ± 2.8 OUR AVERAGE</b>				
9391.8 ± 6.6 ± 2.0	2.3 ± 0.5k	<sup>1</sup> BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
9394.2 <sup>+</sup> <sub>-</sub> 4.8 ± 2.0	13 ± 5k	<sup>1</sup> AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
9388.9 <sup>+</sup> <sub>-</sub> 3.1 ± 2.7	19 ± 3k	<sup>1</sup> AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9300 ± 20 ± 20		HEISTER	02D ALEP	181–209 $e^+ e^-$
<sup>1</sup> Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding $\gamma$ energy or mass difference measurements.				

 **$m_{\Upsilon(1S)} - m_{\eta_b}$** 

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>69.3 ± 2.8 OUR AVERAGE</b>				
68.5 ± 6.6 ± 2.0	2.3 ± 0.5k	<sup>2</sup> BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
66.1 <sup>+</sup> <sub>-</sub> 4.8 ± 2.0	13 ± 5k	<sup>2</sup> AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
71.4 <sup>+</sup> <sub>-</sub> 2.3 ± 2.7	19 ± 3k	<sup>2</sup> AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
<sup>2</sup> Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding $\gamma$ energy or mass measurements.				

 **$\gamma$  ENERGY IN  $\Upsilon(3S)$  DECAY**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>920.6<sup>+</sup><sub>-</sub> 2.8 ± 3.2 OUR AVERAGE</b>				
918.6 ± 6.0 ± 1.9	2.3 ± 0.5k	<sup>3</sup> BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
921.2 <sup>+</sup> <sub>-</sub> 2.1 ± 2.4	19 ± 3k	<sup>3</sup> AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
<sup>3</sup> Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.				

 **$\gamma$  ENERGY IN  $\Upsilon(2S)$  DECAY**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>609.3<sup>+</sup><sub>-</sub> 4.6 ± 4.5 ± 1.9</b>				
	13 ± 5k	<sup>4</sup> AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
<sup>4</sup> Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.				

## $\eta_b(1S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $3h^+ 3h^-$	not seen	
$\Gamma_2$ $2h^+ 2h^-$	not seen	
$\Gamma_3$ $4h^+ 4h^-$		
$\Gamma_4$ $\gamma\gamma$	not seen	
$\Gamma_5$ $\mu^+ \mu^-$	$<9 \times 10^{-3}$	90%
$\Gamma_6$ $\tau^+ \tau^-$	$<8\%$	90%

### $\eta_b(1S)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

#### $\Gamma(3h^+ 3h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_4/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<470	95	ABDALLAH	06	DLPH 161-209 $e^+ e^-$
<132	95	HEISTER	02D	ALEP 181-209 $e^+ e^-$

#### $\Gamma(2h^+ 2h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_4/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<190	95	ABDALLAH	06	DLPH 161-209 $e^+ e^-$
< 48	95	HEISTER	02D	ALEP 181-209 $e^+ e^-$

#### $\Gamma(4h^+ 4h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_4/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<660	95	ABDALLAH	06	DLPH 161-209 $e^+ e^-$

### $\eta_b(1S)$ BRANCHING RATIOS

#### $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9 \times 10^{-3}$	90	<sup>5</sup> AUBERT	09Z	BABR $e^+ e^- \rightarrow \Upsilon(2S, 3S) \rightarrow \gamma\eta_b$
<sup>5</sup> Obtained using $B(\Upsilon(2S) \rightarrow \gamma\eta_b) = (4.2^{+1.1}_{-1.0} \pm 0.9) \times 10^{-4}$ and $B(\Upsilon(3S) \rightarrow \gamma\eta_b) = (4.8 \pm 0.5 \pm 0.6) \times 10^{-4}$ . This limit is equivalent to $B(\eta_b \rightarrow \mu^+ \mu^-) = (-0.25 \pm 0.51 \pm 0.33)\%$ measurement.				

#### $\Gamma(\tau^+ \tau^-)/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-2}$	90	AUBERT	09P	BABR $e^+ e^- \rightarrow \gamma\tau^+ \tau^-$

## $\eta_b(1S)$ REFERENCES

BONVICINI	10	PR D81 031104R	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABDALLAH	06	PL B634 340	J.M. Abdallah <i>et al.</i>	(DELPHI Collab.)
HEISTER	02D	PL B530 56	A. Heister <i>et al.</i>	(ALEPH Collab.)

---