

$\psi(3770)$

$$J^{PC} = 0^{-}(1^{-}-)$$

 $\psi(3770)$ MASS (MeV)

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3773.15±0.33 OUR FIT				
3778.1 ±1.2 OUR AVERAGE				
3779.2	+1.8 -1.7	+0.6 -0.8	1	ANASHIN 12A KEDR $e^+e^- \rightarrow D\bar{D}$
3775.5	±2.4	±0.5	57	AUBERT 08B BABR $B \rightarrow D\bar{D}K$
3776	±5	±4	68	BRODZICKA 08 BELL $B^+ \rightarrow D^0\bar{D}^0K^+$
3778.8	±1.9	±0.9		AUBERT 07BE BABR $e^+e^- \rightarrow D\bar{D}\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3772.0	±1.9		2,3	ABLIKIM 08D BES2 $e^+e^- \rightarrow$ hadrons
3778.4	±3.0	±1.3	34	CHISTOV 04 BELL Sup. by BRODZICKA 08

¹ Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.

² Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

³ Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.


 $m_{\psi(3770)} - m_{\psi(2S)}$

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
87.04±0.33 OUR FIT			
86.6 ±0.7 OUR AVERAGE Error includes scale factor of 2.0. See the ideogram below.			
86.9	±0.4	4	ABLIKIM 07E BES2 $e^+e^- \rightarrow$ hadrons
86.7	±0.7		ABLIKIM 06L BES2 $e^+e^- \rightarrow$ hadrons
80	±2		SCHINDLER 80 MRK2 e^+e^-
86	±2	5	BACINO 78 DLCO e^+e^-
88	±3		RAPIDIS 77 LGW e^+e^-

⁴ BES-II $\psi(2S)$ mass subtracted (see ABLIKIM 06L).

⁵ SPEAR $\psi(2S)$ mass subtracted (see SCHINDLER 80).



$\psi(3770)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.2± 1.0 OUR FIT				
27.5± 0.9 OUR AVERAGE				
24.9 ^{+4.6+0.5} _{-4.0-1.1}		⁶ ANASHIN	12A KEDR	$e^+e^- \rightarrow D\bar{D}$
30.4± 8.5		^{7,8} ABLIKIM	08D BES2	$e^+e^- \rightarrow \text{hadrons}$
27 ±10 ±5	68	BRODZICKA	08 BELL	$B^+ \rightarrow D^0\bar{D}^0 K^+$
28.5± 1.2±0.2		⁸ ABLIKIM	07E BES2	$e^+e^- \rightarrow \text{hadrons}$
23.5± 3.7±0.9		AUBERT	07BE BABR	$e^+e^- \rightarrow D\bar{D}\gamma$
26.9± 2.4±0.3		⁸ ABLIKIM	06L BES2	$e^+e^- \rightarrow \text{hadrons}$
24 ± 5		⁸ SCHINDLER	80 MRK2	e^+e^-
24 ± 5		⁸ BACINO	78 DLCO	e^+e^-
28 ± 5		⁸ RAPIDIS	77 LGW	e^+e^-

⁶ Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.

⁷ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

⁸ Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.

$\psi(3770)$ DECAY MODES

In addition to the dominant decay mode to $D\bar{D}$, $\psi(3770)$ was found to decay into the final states containing the J/ψ (BAI 05, ADAM 06). ADAMS 06 and HUANG 06A searched for various decay modes with light hadrons and found a statistically significant signal for the decay to $\phi\eta$ only (ADAMS 06).

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $D\bar{D}$	(93 \pm 8 \pm 9) %	S=2.0
Γ_2 $D^0\bar{D}^0$	(52 \pm 5) %	S=2.0
Γ_3 D^+D^-	(41 \pm 4) %	S=2.0
Γ_4 $J/\psi\pi^+\pi^-$	(1.93 \pm 0.28) $\times 10^{-3}$	
Γ_5 $J/\psi\pi^0\pi^0$	(8.0 \pm 3.0) $\times 10^{-4}$	
Γ_6 $J/\psi\eta$	(9 \pm 4) $\times 10^{-4}$	
Γ_7 $J/\psi\pi^0$	< 2.8 $\times 10^{-4}$	CL=90%
Γ_8 e^+e^-	(9.6 \pm 0.7) $\times 10^{-6}$	S=1.3

Decays to light hadrons

Γ_9 $b_1(1235)\pi$	< 1.4 $\times 10^{-5}$	CL=90%
Γ_{10} $\phi\eta'$	< 7 $\times 10^{-4}$	CL=90%
Γ_{11} $\omega\eta'$	< 4 $\times 10^{-4}$	CL=90%
Γ_{12} $\rho^0\eta'$	< 6 $\times 10^{-4}$	CL=90%
Γ_{13} $\phi\eta$	(3.1 \pm 0.7) $\times 10^{-4}$	
Γ_{14} $\omega\eta$	< 1.4 $\times 10^{-5}$	CL=90%
Γ_{15} $\rho^0\eta$	< 5 $\times 10^{-4}$	CL=90%
Γ_{16} $\phi\pi^0$	< 3 $\times 10^{-5}$	CL=90%
Γ_{17} $\omega\pi^0$	< 6 $\times 10^{-4}$	CL=90%
Γ_{18} $\pi^+\pi^-\pi^0$	< 5 $\times 10^{-6}$	CL=90%
Γ_{19} $\rho\pi$	< 5 $\times 10^{-6}$	CL=90%
Γ_{20} $K^*(892)^+K^- + \text{c.c.}$	< 1.4 $\times 10^{-5}$	CL=90%
Γ_{21} $K^*(892)^0\bar{K}^0 + \text{c.c.}$	< 1.2 $\times 10^{-3}$	CL=90%
Γ_{22} $K_S^0 K_L^0$	< 1.2 $\times 10^{-5}$	CL=90%
Γ_{23} $2(\pi^+\pi^-)$	< 1.12 $\times 10^{-3}$	CL=90%
Γ_{24} $2(\pi^+\pi^-)\pi^0$	< 1.06 $\times 10^{-3}$	CL=90%
Γ_{25} $2(\pi^+\pi^-\pi^0)$	< 5.85 %	CL=90%
Γ_{26} $\omega\pi^+\pi^-$	< 6.0 $\times 10^{-4}$	CL=90%
Γ_{27} $3(\pi^+\pi^-)$	< 9.1 $\times 10^{-3}$	
Γ_{28} $3(\pi^+\pi^-)\pi^0$	< 1.37 %	
Γ_{29} $3(\pi^+\pi^-)2\pi^0$	< 11.74 %	CL=90%
Γ_{30} $\eta\pi^+\pi^-$	< 1.24 $\times 10^{-3}$	CL=90%
Γ_{31} $\pi^+\pi^-2\pi^0$	< 8.9 $\times 10^{-3}$	CL=90%
Γ_{32} $\rho^0\pi^+\pi^-$	< 6.9 $\times 10^{-3}$	CL=90%
Γ_{33} $\eta3\pi$	< 1.34 $\times 10^{-3}$	CL=90%
Γ_{34} $\eta2(\pi^+\pi^-)$	< 2.43 %	

Γ_{35}	$\eta\rho^0\pi^+\pi^-$	< 1.45	%	CL=90%
Γ_{36}	$\eta'3\pi$	< 2.44	$\times 10^{-3}$	CL=90%
Γ_{37}	$K^+K^-\pi^+\pi^-$	< 9.0	$\times 10^{-4}$	CL=90%
Γ_{38}	$\phi\pi^+\pi^-$	< 4.1	$\times 10^{-4}$	CL=90%
Γ_{39}	$K^+K^-2\pi^0$	< 4.2	$\times 10^{-3}$	CL=90%
Γ_{40}	$4(\pi^+\pi^-)$	< 1.67	%	CL=90%
Γ_{41}	$4(\pi^+\pi^-)\pi^0$	< 3.06	%	CL=90%
Γ_{42}	$\phi f_0(980)$	< 4.5	$\times 10^{-4}$	CL=90%
Γ_{43}	$K^+K^-\pi^+\pi^-\pi^0$	< 2.36	$\times 10^{-3}$	CL=90%
Γ_{44}	$K^+K^-\rho^0\pi^0$	< 8	$\times 10^{-4}$	CL=90%
Γ_{45}	$K^+K^-\rho^+\pi^-$	< 1.46	%	CL=90%
Γ_{46}	ωK^+K^-	< 3.4	$\times 10^{-4}$	CL=90%
Γ_{47}	$\phi\pi^+\pi^-\pi^0$	< 3.8	$\times 10^{-3}$	CL=90%
Γ_{48}	$K^{*0}K^-\pi^+\pi^0 + \text{c.c.}$	< 1.62	%	CL=90%
Γ_{49}	$K^{*+}K^-\pi^+\pi^- + \text{c.c.}$	< 3.23	%	CL=90%
Γ_{50}	$K^+K^-\pi^+\pi^-2\pi^0$	< 2.67	%	CL=90%
Γ_{51}	$K^+K^-2(\pi^+\pi^-)$	< 1.03	%	CL=90%
Γ_{52}	$K^+K^-2(\pi^+\pi^-)\pi^0$	< 3.60	%	CL=90%
Γ_{53}	ηK^+K^-	< 4.1	$\times 10^{-4}$	CL=90%
Γ_{54}	$\eta K^+K^-\pi^+\pi^-$	< 1.24	%	CL=90%
Γ_{55}	$\rho^0 K^+K^-$	< 5.0	$\times 10^{-3}$	CL=90%
Γ_{56}	$2(K^+K^-)$	< 6.0	$\times 10^{-4}$	CL=90%
Γ_{57}	ϕK^+K^-	< 7.5	$\times 10^{-4}$	CL=90%
Γ_{58}	$2(K^+K^-)\pi^0$	< 2.9	$\times 10^{-4}$	CL=90%
Γ_{59}	$2(K^+K^-)\pi^+\pi^-$	< 3.2	$\times 10^{-3}$	CL=90%
Γ_{60}	$K_S^0 K^-\pi^+$	< 3.2	$\times 10^{-3}$	CL=90%
Γ_{61}	$K_S^0 K^-\pi^+\pi^0$	< 1.33	%	CL=90%
Γ_{62}	$K_S^0 K^-\rho^+$	< 6.6	$\times 10^{-3}$	CL=90%
Γ_{63}	$K_S^0 K^-2\pi^+\pi^-$	< 8.7	$\times 10^{-3}$	CL=90%
Γ_{64}	$K_S^0 K^-\pi^+\rho^0$	< 1.6	%	CL=90%
Γ_{65}	$K_S^0 K^-\pi^+\eta$	< 1.3	%	CL=90%
Γ_{66}	$K_S^0 K^-2\pi^+\pi^-\pi^0$	< 4.18	%	CL=90%
Γ_{67}	$K_S^0 K^-2\pi^+\pi^-\eta$	< 4.8	%	CL=90%
Γ_{68}	$K_S^0 K^-\pi^+2(\pi^+\pi^-)$	< 1.22	%	CL=90%
Γ_{69}	$K_S^0 K^-\pi^+2\pi^0$	< 2.65	%	CL=90%
Γ_{70}	$K_S^0 K^-K^+K^-\pi^+$	< 4.9	$\times 10^{-3}$	CL=90%
Γ_{71}	$K_S^0 K^-K^+K^-\pi^+\pi^0$	< 3.0	%	CL=90%
Γ_{72}	$K_S^0 K^-K^+K^-\pi^+\eta$	< 2.2	%	CL=90%
Γ_{73}	$K^{*0}K^-\pi^+ + \text{c.c.}$	< 9.7	$\times 10^{-3}$	CL=90%
Γ_{74}	$\rho\bar{\rho}\pi^0$	< 1.2	$\times 10^{-3}$	
Γ_{75}	$\rho\bar{\rho}\pi^+\pi^-$	< 5.8	$\times 10^{-4}$	CL=90%
Γ_{76}	$\Lambda\bar{\Lambda}$	< 1.2	$\times 10^{-4}$	CL=90%
Γ_{77}	$\rho\bar{\rho}\pi^+\pi^-\pi^0$	< 1.85	$\times 10^{-3}$	CL=90%

Γ_{78}	$\omega p\bar{p}$	< 2.9	$\times 10^{-4}$	CL=90%
Γ_{79}	$\Lambda\bar{\Lambda}\pi^0$	< 1.2	$\times 10^{-3}$	CL=90%
Γ_{80}	$\rho\bar{\rho}2(\pi^+\pi^-)$	< 2.6	$\times 10^{-3}$	CL=90%
Γ_{81}	$\eta p\bar{p}$	< 5.4	$\times 10^{-4}$	CL=90%
Γ_{82}	$\eta p\bar{p}\pi^+\pi^-$	< 3.3	$\times 10^{-3}$	CL=90%
Γ_{83}	$\rho^0 p\bar{p}$	< 1.7	$\times 10^{-3}$	CL=90%
Γ_{84}	$\rho\bar{\rho}K^+K^-$	< 3.2	$\times 10^{-4}$	CL=90%
Γ_{85}	$\eta p\bar{p}K^+K^-$	< 6.9	$\times 10^{-3}$	CL=90%
Γ_{86}	$\pi^0 p\bar{p}K^+K^-$	< 1.2	$\times 10^{-3}$	CL=90%
Γ_{87}	$\phi p\bar{p}$	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{88}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	< 2.5	$\times 10^{-4}$	CL=90%
Γ_{89}	$\Lambda\bar{p}K^+$	< 2.8	$\times 10^{-4}$	CL=90%
Γ_{90}	$\Lambda\bar{p}K^+\pi^+\pi^-$	< 6.3	$\times 10^{-4}$	CL=90%

Radiative decays

Γ_{91}	$\gamma\chi_{c2}$	< 9	$\times 10^{-4}$	CL=90%
Γ_{92}	$\gamma\chi_{c1}$	$(2.9 \pm 0.6) \times 10^{-3}$		
Γ_{93}	$\gamma\chi_{c0}$	$(7.3 \pm 0.9) \times 10^{-3}$		
Γ_{94}	$\gamma\eta'$	< 1.8	$\times 10^{-4}$	CL=90%
Γ_{95}	$\gamma\eta$	< 1.5	$\times 10^{-4}$	CL=90%
Γ_{96}	$\gamma\pi^0$	< 2	$\times 10^{-4}$	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 3 branching ratios uses 23 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 20.0$ for 19 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3		98		
x_8		0	0	
Γ		0	0	-44
		x_2	x_3	x_8

	Mode	Rate (MeV)	Scale factor
Γ_2	$D^0\bar{D}^0$	14.1 ± 1.4	1.7
Γ_3	D^+D^-	11.2 ± 1.1	1.7
Γ_8	e^+e^-	$(2.62 \pm 0.18) \times 10^{-4}$	1.4

$\psi(3770)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$					Γ_8	
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT		
0.262±0.018 OUR FIT	Error includes scale factor of 1.4.					
0.256±0.016 OUR AVERAGE	Error includes scale factor of 1.2.					
0.154 ^{+0.079+0.021} _{-0.058-0.027}	9,10	ANASHIN	12A	KEDR	$e^+e^- \rightarrow D\bar{D}$	
0.22 ±0.05	11,12	ABLIKIM	08D	BES2	$e^+e^- \rightarrow$ hadrons	
0.277±0.011±0.013	12	ABLIKIM	07E	BES2	$e^+e^- \rightarrow$ hadrons	
0.203±0.003 ^{+0.041} _{-0.027}	1.4M 12,13	BESSION	06	CLEO	$e^+e^- \rightarrow$ hadrons	
0.276±0.050	12	SCHINDLER	80	MRK2	e^+e^-	
0.18 ±0.06	12	BACINO	78	DLCO	e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.414 ^{+0.072+0.093} _{-0.080-0.028}	10,14	ANASHIN	12A	KEDR	$e^+e^- \rightarrow D\bar{D}$	
0.37 ±0.09	15	RAPIDIS	77	LGW	e^+e^-	
⁹ Solution I of the two solutions.						
¹⁰ Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.						
¹¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.						
¹² Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.						
¹³ BESSION 06 (as corrected in BESSION 10) measure $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow$ hadrons) = $6.36 \pm 0.08^{+0.41}_{-0.30}$ nb at $\sqrt{s} = 3773 \pm 1$ MeV, and obtain Γ_{ee} from the Born-level cross section calculated using $\psi(3770)$ mass and width from our 2004 edition, PDG 04.						
¹⁴ Solution II of the two solutions.						
¹⁵ See also $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ below.						

 $\psi(3770)$ BRANCHING RATIOS

$\Gamma(D\bar{D})/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma = (\Gamma_2+\Gamma_3)/\Gamma$	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
0.93^{+0.08}_{-0.09} OUR FIT	Error includes scale factor of 2.0.					
0.93^{+0.08}_{-0.09} OUR AVERAGE	Error includes scale factor of 2.1.					
0.849±0.056±0.018	16	ABLIKIM	08B	BES2	$e^+e^- \rightarrow$ non- $D\bar{D}$	
1.033±0.014 ^{+0.048} _{-0.066}	1.427M 17	BESSION	06	CLEO	$e^+e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.866±0.050±0.036	18,19	ABLIKIM	07K	BES2	$e^+e^- \rightarrow$ non- $D\bar{D}$	
0.836±0.073±0.042	19	ABLIKIM	06L	BES2	$e^+e^- \rightarrow D\bar{D}$	
0.855±0.017±0.058	19,20	ABLIKIM	06N	BES2	$e^+e^- \rightarrow D\bar{D}$	

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.52 ± 0.05 OUR FIT	Error includes scale factor of 2.0.		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.467 ± 0.047 ± 0.023	ABLIKIM	06L BES2	$e^+e^- \rightarrow D^0\bar{D}^0$
0.499 ± 0.013 ± 0.038	²⁰ ABLIKIM	06N BES2	$e^+e^- \rightarrow D^0\bar{D}^0$

$\Gamma(D^+D^-)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.41 ± 0.04 OUR FIT	Error includes scale factor of 2.0.		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.369 ± 0.037 ± 0.028	ABLIKIM	06L BES2	$e^+e^- \rightarrow D^+D^-$
0.357 ± 0.011 ± 0.034	²⁰ ABLIKIM	06N BES2	$e^+e^- \rightarrow D^+D^-$

$\Gamma(D^0\bar{D}^0)/\Gamma(D^+D^-)$ Γ_2/Γ_3

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.260 ± 0.021 OUR FIT				
1.260 ± 0.021 OUR AVERAGE				
1.39 ± 0.31 ± 0.12		PAKHLOVA	08 BELL	10.6 $e^+e^- \rightarrow D\bar{D}\gamma$
1.78 ± 0.33 ± 0.24		AUBERT	07BE BABR	$e^+e^- \rightarrow D\bar{D}\gamma$
1.258 ± 0.016 ± 0.014		DOBBS	07 CLEO	$e^+e^- \rightarrow D\bar{D}$
1.27 ± 0.12 ± 0.08		ABLIKIM	06L BES2	$e^+e^- \rightarrow D\bar{D}$
2.43 ± 1.50 ± 0.43	34	²¹ CHISTOV	04 BELL	$B^+ \rightarrow \psi(3770)K^+$

$\Gamma(J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.93 ± 0.28 OUR AVERAGE				
1.89 ± 0.20 ± 0.20	231 ± 33	ADAM	06 CLEO	$e^+e^- \rightarrow \psi(3770)$
3.4 ± 1.4 ± 0.9	17.8 ± 4.8	BAI	05 BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(J/\psi\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.080 ± 0.025 ± 0.016	39 ± 14	ADAM	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
87 ± 33 ± 22	22 ± 10	ADAM	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<28	90	<10	ADAM	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.96 ± 0.07 OUR FIT	Error includes scale factor of 1.3.		
1.3 ± 0.2	RAPIDIS	77 LGW	e^+e^-

¹⁶ Neglecting interference.

¹⁷ Obtained by comparing a measurement of the total cross section (corrected in BESSON 10) with that of $D\bar{D}$ reported by CLEO in DOBBS 07.

¹⁸ Using $\sigma^{obs} = 7.07 \pm 0.58$ nb and neglecting interference.

¹⁹ Not independent of ABLIKIM 08B.

²⁰ From a measurement of $\sigma(e^+e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773$ MeV, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.

²¹ See ADLER 88C for older measurements of this quantity.

———— DECAYS TO LIGHT HADRONS ————

$\Gamma(b_1(1235)\pi)/\Gamma_{total}$ Γ_9/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	²² ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\phi\eta')/\Gamma_{total}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	²² ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\omega\eta')/\Gamma_{total}$ Γ_{11}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	²² ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho^0\eta')/\Gamma_{total}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	²² ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\phi\eta)/\Gamma_{total}$ Γ_{13}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.1 \pm 0.6 \pm 0.3$	²² ADAMS	06	CLEO $3.773 e^+e^- \rightarrow \phi\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	²³ ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$
-----	-----------------------	-----	--------------------------------------

$\Gamma(\omega\eta)/\Gamma_{total}$ Γ_{14}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	²² ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho^0\eta)/\Gamma_{total}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	²² ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\phi\pi^0)/\Gamma_{total}$ Γ_{16}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3	90	²² ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<50	²³ ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$
-----	-----------------------	-----	--------------------------------------

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6	90	22 ADAMS 06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	22,24 ADAMS 06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	22,24 ADAMS 06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(K^*(892)^+K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.4	90	22 ADAMS 06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.2	90	22 ADAMS 06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.2	90	25 CRONIN-HEN..06	CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	26 ABLIKIM 04F	BES	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------	-----	---------------------------------

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{23}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<11.2	90	27 HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<48		23 ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	--	----------------	------	---------------------------------

$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<10.6	90	27 HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<62		23 ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	--	----------------	------	---------------------------------

$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<58.5	90	305	ABLIKIM 08N	BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-4})	CL%		DOCUMENT ID	TECN	COMMENT
< 6.0	90	27	HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<55	90	23	ABLIKIM	07I	BES2 $3.77 e^+e^-$

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-4})			DOCUMENT ID	TECN	COMMENT
<91		23	ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-4})			DOCUMENT ID	TECN	COMMENT
<137		23	ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(3(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<117.4	90	59	ABLIKIM	08N	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-3})	CL%		DOCUMENT ID	TECN	COMMENT
<1.24	90	27	HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.3	90	23	ABLIKIM	10D	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8.9	90	218	ABLIKIM	08N	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-3})	CL%		DOCUMENT ID	TECN	COMMENT
<6.9	90	23	ABLIKIM	07F	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta3\pi)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-4})	CL%		DOCUMENT ID	TECN	COMMENT
<13.4	90	27	HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-4})			DOCUMENT ID	TECN	COMMENT
<243		23	ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-2})	CL%		DOCUMENT ID	TECN	COMMENT
<1.45	90	23	ABLIKIM	10D	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta' 3\pi)/\Gamma_{\text{total}}$					Γ_{36}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<24.4	90	27 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$	
$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{37}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 9.0	90	27 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<48		23 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$	
$\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{38}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 4.1	90	27 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<16		23 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$	
$\Gamma(K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$					Γ_{39}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.2	90	14	ABLIKIM	08N BES2	$e^+ e^- \rightarrow \psi(3770)$
$\Gamma(4(\pi^+ \pi^-))/\Gamma_{\text{total}}$					Γ_{40}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<16.7	90	23 ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$	
$\Gamma(4(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$					Γ_{41}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<30.6	90	23 ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$	
$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$					Γ_{42}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<4.5	90	27 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$	
$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{43}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 23.6	90	27 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<111		23 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$	
$\Gamma(K^+ K^- \rho^0 \pi^0)/\Gamma_{\text{total}}$					Γ_{44}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<8	90	23 ABLIKIM	07I BES2	$3.77 e^+ e^-$	
$\Gamma(K^+ K^- \rho^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{45}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<146	90	23 ABLIKIM	07I BES2	$3.77 e^+ e^-$	

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.4	90	²⁷ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<66	90	²³ ABLIKIM	07I	BES2 $e^+ e^-$

$\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<38	90	²³ ABLIKIM	07I	BES2 $e^+ e^-$

$\Gamma(K^{*0} K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<162	90	²³ ABLIKIM	07I	BES2 $e^+ e^-$

$\Gamma(K^{*+} K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<323	90	²³ ABLIKIM	07I	BES2 $e^+ e^-$

$\Gamma(K^+ K^- \pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<26.7	90	24	ABLIKIM	08N	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<10.3	90	²³ ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<36.0	90	²³ ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\eta K^+ K^-)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	²⁷ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<31	90	²³ ABLIKIM	10D	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\eta K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<1.24	90	²³ ABLIKIM	10D	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	²³ ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ Γ_{56}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 6.0	90	27 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<17		23 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$ Γ_{57}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 7.5	90	27 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<24		23 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(2(K^+ K^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{58}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 2.9	90	27 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<46		23 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(2(K^+ K^-)\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{59}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.2	90	23 ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{60}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.2	90	18	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{61}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<13.3	90	40	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \rho^+)/\Gamma_{\text{total}}$ Γ_{62}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.6	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- 2\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{63}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.7	90	39	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{64}/Γ

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.6	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+ \eta)/\Gamma_{\text{total}}$ Γ_{65}/Γ

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.3	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- 2\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<41.8	90	23	ABLIKIM	08M	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- 2\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<4.8	90	ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+ 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<12.2	90	4	ABLIKIM	08M	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+ 2\pi^0)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<26.5	90	17	ABLIKIM	08M	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- K^+ K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<4.9	90	ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- K^+ K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- K^+ K^- \pi^+ \eta)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K^{*0} K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<9.7	90	²³ ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
<12	²³ ABLIKIM	07B	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(p\bar{p}\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.8	90	²⁷ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<16		²³ ABLIKIM	07B	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	²⁷ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<4	90	²³ ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\rho\bar{\rho}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{77}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<18.5	90	27 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<73		23 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\omega\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{78}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 2.9	90	27 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<30	90	28 ABLIKIM	07I BES2	$3.77 e^+e^-$

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{79}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<12	90	23 ABLIKIM	07I BES2	$3.77 e^+e^-$

$\Gamma(\rho\bar{\rho}2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{80}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.6	90	23 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{81}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 5.4	90	27 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<11	90	23 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta\rho\bar{\rho}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{82}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.3	90	23 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho^0\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{83}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.7	90	23 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho\bar{\rho}K^+K^-)/\Gamma_{\text{total}}$ Γ_{84}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3.2	90	27 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<11		23 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta\rho\bar{\rho}K^+K^-)/\Gamma_{\text{total}}$ Γ_{85}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.9	90	23 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\pi^0 \rho \bar{\rho} K^+ K^-)/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	²³ ABLIKIM	10D	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\phi \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	²⁷ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9		²³ ABLIKIM	07B	BES2 $e^+ e^- \rightarrow \psi(3770)$
----	--	-----------------------	-----	---------------------------------------

$\Gamma(\Lambda \bar{\Lambda} \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.5	90	²⁷ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<39	90	²³ ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$
-----	----	-----------------------	-----	---------------------------------------

$\Gamma(\Lambda \bar{\rho} K^+)/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	²⁷ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\Lambda \bar{\rho} K^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6.3	90	²⁷ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$

²² Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

²³ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

²⁴ Data suggest possible destructive interference with continuum.

²⁵ Using $\sigma(e^+ e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = (6.38 \pm 0.08^{+0.41}_{-0.30})$ nb from BESSON 06 and $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6895 \pm 0.0014$.

²⁶ Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6860 \pm 0.0027$.

²⁷ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

²⁸ Using $\sigma^{obs} = 7.15 \pm 0.27 \pm 0.27$ nb and neglecting interference.

————— RADIATIVE DECAYS —————

$\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	²⁹ COAN	06A	CLEO $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma \gamma J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.0	90	³⁰ BRIERE	06	CLEO $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$
------	----	----------------------	----	---

$\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$					Γ_{92}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.9 \pm 0.5 \pm 0.4$		31 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}, \gamma\gamma J/\psi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$3.9 \pm 1.4 \pm 0.6$	54 ± 17	32 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
$2.8 \pm 0.5 \pm 0.4$	53 ± 10	29 COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	

$\Gamma(\gamma\chi_{c1})/\Gamma(J/\psi\pi^+\pi^-)$					Γ_{92}/Γ_4
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.49 \pm 0.31 \pm 0.26$	53 ± 10	33 COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	

$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$					Γ_{93}/Γ
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.3 \pm 0.7 \pm 0.6$		274 ± 27	34 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 44	90		29 COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

$\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c2})$					Γ_{93}/Γ_{91}
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
> 8	90	35 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770)$	

$\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c1})$					Γ_{93}/Γ_{92}
VALUE	DOCUMENT ID	TECN	COMMENT		
2.5 ± 0.6	35 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770)$		

$\Gamma(\gamma\eta')/\Gamma_{\text{total}}$					Γ_{94}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
< 1.8	90	36 PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$	

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$					Γ_{95}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
< 1.5	90	36 PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$	

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$					Γ_{96}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
< 2	90	PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$	

- ²⁹ Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+e^- \rightarrow D\bar{D})$ from HE 05 for $\sigma(e^+e^- \rightarrow \psi(3770))$.
- ³⁰ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = 9.22 \pm 0.11 \pm 0.46\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.
- ³¹ Averages the two measurements from COAN 06A and BRIERE 06.
- ³² Uses $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = 9.07 \pm 0.11 \pm 0.54\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.
- ³³ Using $B(\psi(3770) \rightarrow J/\psi\pi^+\pi^-) = (1.89 \pm 0.20 \pm 0.20) \times 10^{-3}$ from ADAM 06.
- ³⁴ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = 9.33 \pm 0.14 \pm 0.61\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.
- ³⁵ Not independent of other results in BRIERE 06.
- ³⁶ Assuming maximal destructive interference between $\psi(3770)$ and continuum sources.

$\psi(3770)$ REFERENCES

ANASHIN	12A	PL B711 292	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
ABLIKIM	10D	EPJ C66 11	M. Ablikim <i>et al.</i>	(BES II Collab.)
BESSON	10	PRL 104 159901E	D. Besson <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09C	EPJ C64 243	M. Ablikim <i>et al.</i>	(BES Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08M	PL B670 179	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08N	PL B670 184	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	08B	PR D77 011102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRODZICKA	08	PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08	PR D77 011103R	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07B	PL B650 111	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07E	PL B652 238	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07F	PL B656 30	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07I	EPJ C52 805	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07K	PR D76 122002	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	07BE	PR D76 111105R	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	07	PR D76 112001	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06N	PL B641 145	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAMS	06	PR D73 012002	G.S. Adams <i>et al.</i>	(CLEO Collab.)
BESSON	06	PRL 96 092002	D. Besson <i>et al.</i>	(CLEO Collab.)
Also		PRL 104 159901E	D. Besson <i>et al.</i>	(CLEO Collab.)
BRIERE	06	PR D74 031106R	R.A. Briere <i>et al.</i>	(CLEO Collab.)
COAN	06A	PRL 96 182002	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...	06	PR D74 012005	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
HUANG	06A	PRL 96 032003	G.S. Huang <i>et al.</i>	(CLEO Collab.)
BAI	05	PL B605 63	J.Z. Bai <i>et al.</i>	(BES Collab.)
HE	05	PRL 95 121801	Q. He <i>et al.</i>	(CLEO Collab.)
Also		PRL 96 199903 (errat.)	Q. He <i>et al.</i>	(CLEO Collab.)
ABLIKIM	04F	PR D70 077101	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ADLER	88C	PRL 60 89	J. Adler <i>et al.</i>	(Mark III Collab.)
SCHINDLER	80	PR D21 2716	R.H. Schindler <i>et al.</i>	(Mark II Collab.)
BACINO	78	PRL 40 671	W.J. Bacino <i>et al.</i>	(SLAC, UCLA, UCI)
RAPIDIS	77	PRL 39 526	P.A. Rapidis <i>et al.</i>	(LGW Collab.)
