

$\Upsilon(2S)$

$$J^{PC} = 0^{-}(1^{-}-)$$

$\Upsilon(2S)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.02326 ± 0.00031 OUR AVERAGE			
10.0235 ± 0.0005	¹ ARTAMONOV 00	MD1	$e^+e^- \rightarrow \text{hadrons}$
10.0231 ± 0.0004	BARBER 84	REDE	$e^+e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10.0236 ± 0.0005	^{2,3} BARU	86B REDE	$e^+e^- \rightarrow \text{hadrons}$
¹ Reanalysis of BARU 86B using new electron mass (COHEN 87).			
² Reanalysis of ARTAMONOV 84.			
³ Superseded by ARTAMONOV 00.			

$m\Upsilon(3S) - m\Upsilon(2S)$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
331.50 ± 0.02 ± 0.13	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

$\Upsilon(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID	COMMENT
31.98 ± 2.63 OUR EVALUATION		See the Note on "Width Determinations of the Υ States"

$\Upsilon(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\Upsilon(1S)\pi^+\pi^-$	(17.92 ± 0.26) %	
Γ_2 $\Upsilon(1S)\pi^0\pi^0$	(8.6 ± 0.4) %	
Γ_3 $\tau^+\tau^-$	(2.00 ± 0.21) %	
Γ_4 $\mu^+\mu^-$	(1.93 ± 0.17) %	S=2.2
Γ_5 e^+e^-	(1.91 ± 0.16) %	
Γ_6 $\Upsilon(1S)\pi^0$	< 1.8 × 10 ⁻⁴	CL=90%
Γ_7 $\Upsilon(1S)\eta$	(2.34 ± 0.31) × 10 ⁻⁴	
Γ_8 $J/\psi(1S)$ anything	< 6 × 10 ⁻³	CL=90%
Γ_9 \bar{d} anything	(3.4 ± 0.6) × 10 ⁻⁵	
Γ_{10} hadrons	(94 ± 11) %	
Γ_{11} ggg	(58.8 ± 1.2) %	
Γ_{12} γgg	(8.8 ± 1.1) %	

Radiative decays

Γ_{13}	$\gamma\chi_{b1}(1P)$		$(6.9 \pm 0.4) \%$		
Γ_{14}	$\gamma\chi_{b2}(1P)$		$(7.15 \pm 0.35) \%$		
Γ_{15}	$\gamma\chi_{b0}(1P)$		$(3.8 \pm 0.4) \%$		
Γ_{16}	$\gamma f_0(1710)$	< 5.9	$\times 10^{-4}$	CL=90%	
Γ_{17}	$\gamma f_2'(1525)$	< 5.3	$\times 10^{-4}$	CL=90%	
Γ_{18}	$\gamma f_2(1270)$	< 2.41	$\times 10^{-4}$	CL=90%	
Γ_{19}	$\gamma f_J(2220)$				
Γ_{20}	$\gamma\eta_c(1S)$	< 2.7	$\times 10^{-5}$	CL=90%	
Γ_{21}	$\gamma\chi_{c0}$	< 1.0	$\times 10^{-4}$	CL=90%	
Γ_{22}	$\gamma\chi_{c1}$	< 3.6	$\times 10^{-6}$	CL=90%	
Γ_{23}	$\gamma\chi_{c2}$	< 1.5	$\times 10^{-5}$	CL=90%	
Γ_{24}	$\gamma X(3872) \rightarrow \pi^+\pi^- J/\psi$	< 8	$\times 10^{-7}$	CL=90%	
Γ_{25}	$\gamma X(3872) \rightarrow \pi^+\pi^-\pi^0 J/\psi$	< 2.4	$\times 10^{-6}$	CL=90%	
Γ_{26}	$\gamma X(3915) \rightarrow \omega J/\psi$	< 2.8	$\times 10^{-6}$	CL=90%	
Γ_{27}	$\gamma X(4140) \rightarrow \phi J/\psi$	< 1.2	$\times 10^{-6}$	CL=90%	
Γ_{28}	$\gamma X(4350) \rightarrow \phi J/\psi$	< 1.3	$\times 10^{-6}$	CL=90%	
Γ_{29}	$\gamma\eta_b(1S)$		$(3.9 \pm 1.5) \times 10^{-4}$		
Γ_{30}	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 1.95	$\times 10^{-4}$	CL=95%	
Γ_{31}	$\gamma A^0 \rightarrow \gamma$ hadrons	< 8	$\times 10^{-5}$	CL=90%	

Lepton Family number (LF) violating modes

Γ_{32}	$e^\pm \tau^\mp$	LF	< 3.2	$\times 10^{-6}$	CL=90%
Γ_{33}	$\mu^\pm \tau^\mp$	LF	< 3.3	$\times 10^{-6}$	CL=90%

[a] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

$\mathcal{R}(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_4\Gamma_5/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
6.5±1.5±1.0	KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$	

$\Gamma(\mathcal{R}(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
105.4±1.0±4.2	11.8K	⁴ AUBERT	08BP BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$	
⁴ Using $B(\mathcal{R}(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\mathcal{R}(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.					

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{10}\Gamma_5/\Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
0.577±0.009 OUR AVERAGE					
0.581±0.004±0.009	⁵ ROSNER	06	CLEO	10.0 $e^+e^- \rightarrow$ hadrons	
0.552±0.031±0.017	⁵ BARU	96	MD1	$e^+e^- \rightarrow$ hadrons	
0.54 ±0.04 ±0.02	⁵ JAKUBOWSKI	88	CBAL	$e^+e^- \rightarrow$ hadrons	
0.58 ±0.03 ±0.04	⁶ GILES	84B	CLEO	$e^+e^- \rightarrow$ hadrons	

$0.60 \pm 0.12 \pm 0.07$	⁶ ALBRECHT	82	DASP	$e^+e^- \rightarrow$	hadrons
$0.54 \pm 0.07 \begin{smallmatrix} +0.09 \\ -0.05 \end{smallmatrix}$	⁶ NICZYPORUK	81C	LENA	$e^+e^- \rightarrow$	hadrons
0.41 ± 0.18	⁶ BOCK	80	CNTR	$e^+e^- \rightarrow$	hadrons

⁵ Radiative corrections evaluated following KURAEV 85.

⁶ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

$\Upsilon(2S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$

Γ_5

VALUE (keV)

DOCUMENT ID

0.612 ± 0.011 OUR EVALUATION

$\Upsilon(2S)$ BRANCHING RATIOS

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_1/Γ

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
17.92 ± 0.26 OUR AVERAGE				
$16.8 \pm 1.1 \pm 1.3$	906k	⁷ LEES	11C	BABR $e^+e^- \rightarrow \pi^+\pi^-X$
$17.80 \pm 0.05 \pm 0.37$	170k	⁸ LEES	11L	BABR $\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
$18.02 \pm 0.02 \pm 0.61$	851k	⁹ BHARI	09	CLEO $e^+e^- \rightarrow \pi^+\pi^-MM$
$17.22 \pm 0.17 \pm 0.75$	11.8K	¹⁰ AUBERT	08BP	BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
$19.2 \pm 0.2 \pm 1.0$	52.6k	¹¹ ALEXANDER	98	CLE2 $\pi^+\pi^-\ell^+\ell^-, \pi^+\pi^-MM$
$18.1 \pm 0.5 \pm 1.0$	11.6k	ALBRECHT	87	ARG $e^+e^- \rightarrow \pi^+\pi^-MM$
16.9 ± 4.0		GELPHMAN	85	CBAL $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
$19.1 \pm 1.2 \pm 0.6$		BESSON	84	CLEO $\pi^+\pi^-MM$
18.9 ± 2.6		FONSECA	84	CUSB $e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$
21 ± 7	7	NICZYPORUK	81B	LENA $e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$

⁷ LEES 11C reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁸ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

⁹ A weighted average of the inclusive and exclusive results.

¹⁰ Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$ and, $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.

¹¹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6 ± 0.4 OUR AVERAGE				
$8.43 \pm 0.16 \pm 0.42$	38k	¹² BHARI	09	CLEO $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
$9.2 \pm 0.6 \pm 0.8$	275	¹³ ALEXANDER	98	CLE2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
$9.5 \pm 1.9 \pm 1.9$	25	ALBRECHT	87	ARG $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
8.0 ± 1.5		GELPHMAN	85	CBAL $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
10.3 ± 2.3		FONSECA	84	CUSB $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

¹² Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

¹³ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_1

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.462±0.037 ¹⁴ BHARI 09 CLEO $e^+e^- \rightarrow \Upsilon(2S)$

¹⁴ Not independent of other values reported by BHARI 09.

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (units 10⁻²) EVTS DOCUMENT ID TECN COMMENT

2.00±0.21 OUR AVERAGE

2.00±0.12±0.18 22k ¹⁵ BESSON 07 CLEO $e^+e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+\tau^-$

1.7 ±1.5 ±0.6 HAAS 84B CLEO $e^+e^- \rightarrow \tau^+\tau^-$

¹⁵ BESSON 07 reports $[\Gamma(\Upsilon(2S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = 1.04 \pm 0.04 \pm 0.05$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE CL% EVTS DOCUMENT ID TECN COMMENT

0.0193±0.0017 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.

0.0203±0.0003±0.0008 120k ADAMS 05 CLEO $e^+e^- \rightarrow \mu^+\mu^-$

0.0122±0.0028±0.0019 ¹⁶ KOBEL 92 CBAL $e^+e^- \rightarrow \mu^+\mu^-$

0.0138±0.0025±0.0015 KAARSBERG 89 CSB2 $e^+e^- \rightarrow \mu^+\mu^-$

0.009 ±0.006 ±0.006 ¹⁷ ALBRECHT 85 ARG $e^+e^- \rightarrow \mu^+\mu^-$

0.018 ±0.008 ±0.005 HAAS 84B CLEO $e^+e^- \rightarrow \mu^+\mu^-$


• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.038 90 NICZYPORUK 81C LENA $e^+e^- \rightarrow \mu^+\mu^-$

¹⁶ Taking into account interference between the resonance and continuum.

¹⁷ Re-evaluated using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 0.026$.

WEIGHTED AVERAGE
 0.0193 ± 0.0017 (Error scaled by 2.2)



$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$

Γ_3/Γ_4

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.04 \pm 0.04 \pm 0.05$	22k	BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(2S)$

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{total}$

Γ_6/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.18	90	¹⁸ HE	08A	CLEO $e^+e^- \rightarrow l^+l^-\gamma\gamma$
<1.1	90	ALEXANDER	98	CLE2 $e^+e^- \rightarrow l^+l^-\gamma\gamma$
<8	90	LURZ	87	CBAL $e^+e^- \rightarrow l^+l^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •
¹⁸ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{total}$

Γ_7/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.34 ± 0.31 OUR AVERAGE					
$2.39 \pm 0.31 \pm 0.14$	112	¹⁹ LEES	11L	BABR	$\Upsilon(2S) \rightarrow l^+l^-\eta$
$2.1^{+0.7}_{-0.6} \pm 0.3$	14	²⁰ HE	08A	CLEO	$e^+e^- \rightarrow l^+l^-\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9	90	^{19,21} AUBERT	08BP	BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0 l^+l^-$
< 28	90	ALEXANDER	98	CLE2	$e^+e^- \rightarrow l^+l^-\eta$
< 50	90	ALBRECHT	87	ARG	$e^+e^- \rightarrow \pi^+\pi^-l^+l^-$ MM
< 70	90	LURZ	87	CBAL	$e^+e^- \rightarrow l^+l^-(\gamma\gamma, 3\pi^0)$
< 100	90	BESSON	84	CLEO	$e^+e^- \rightarrow \pi^+\pi^-l^+l^-$ MM
< 20	90	FONSECA	84	CUSB	$e^+e^- \rightarrow l^+l^-(\gamma\gamma, \pi^+\pi^-\pi^0)$

¹⁹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

²⁰ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

²¹ Using $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.

$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.35 \pm 0.17 \pm 0.08$		²² LEES	11L BABR	$\Upsilon(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$
<5.2	90	²³ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

²² Not independent of other values reported by LEES 11L.

²³ Not independent of other values reported by AUBERT 08BP.

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.006	90	MASCHMANN 90	CBAL	$e^+e^- \rightarrow \text{hadrons}$
-------------------------------	----	--------------	------	-------------------------------------

$\Gamma(\bar{d} \text{ anything})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$3.37 \pm 0.50 \pm 0.25$	58	ASNER	07 CLEO	$e^+e^- \rightarrow \bar{d}X$
--	----	-------	---------	-------------------------------

$\Gamma(ggg)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

58.8 ± 1.2	6M	²⁴ BESSON	06A CLEO	$\Upsilon(2S) \rightarrow \text{hadrons}$
----------------------------------	----	----------------------	----------	---

²⁴ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$ from BESSON 06A and PDG 08 values of $B(\pi^+\pi^-\Upsilon(1S)) = (18.1 \pm 0.4)\%$, $B(\pi^0\pi^0\Upsilon(1S)) = (8.6 \pm 0.4)\%$, $B(\mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.79 ± 1.05	100k	²⁵ BESSON	06A CLEO	$\Upsilon(2S) \rightarrow \gamma + \text{hadrons}$
-----------------------------------	------	----------------------	----------	--

²⁵ Calculated using BESSON 06A values of $\Gamma(\gamma gg)/\Gamma(ggg) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$ and $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

$\Gamma(\gamma gg)/\Gamma(ggg)$ Γ_{12}/Γ_{11}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$3.18 \pm 0.04 \pm 0.47$	6M	BESSON	06A CLEO	$\Upsilon(2S) \rightarrow (\gamma +) \text{hadrons}$
--	----	--------	----------	--

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$				Γ_{13}/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.069 ± 0.004 OUR AVERAGE				
0.0693 ± 0.0012 ± 0.0041	407k	ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$
0.069 ± 0.005 ± 0.009		EDWARDS	99 CLE2	$\Upsilon(2S) \rightarrow \gamma\chi(1P)$
0.091 ± 0.018 ± 0.022		ALBRECHT	85E ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$
0.065 ± 0.007 ± 0.012		NERNST	85 CBAL	$e^+e^- \rightarrow \gamma X$
0.080 ± 0.017 ± 0.016		HAAS	84 CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$
0.059 ± 0.014		KLOPFEN...	83 CUSB	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$				Γ_{14}/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0715 ± 0.0035 OUR AVERAGE				
0.0724 ± 0.0011 ± 0.0040	410k	ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$
0.074 ± 0.005 ± 0.008		EDWARDS	99 CLE2	$\Upsilon(2S) \rightarrow \gamma\chi(1P)$
0.098 ± 0.021 ± 0.024		ALBRECHT	85E ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$
0.058 ± 0.007 ± 0.010		NERNST	85 CBAL	$e^+e^- \rightarrow \gamma X$
0.102 ± 0.018 ± 0.021		HAAS	84 CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$
0.061 ± 0.014		KLOPFEN...	83 CUSB	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$				Γ_{15}/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.038 ± 0.004 OUR AVERAGE				
0.0375 ± 0.0012 ± 0.0047	198k	ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$
0.034 ± 0.005 ± 0.006		EDWARDS	99 CLE2	$\Upsilon(2S) \rightarrow \gamma\chi(1P)$
0.064 ± 0.014 ± 0.016		ALBRECHT	85E ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$
0.036 ± 0.008 ± 0.009		NERNST	85 CBAL	$e^+e^- \rightarrow \gamma X$
0.044 ± 0.023 ± 0.009		HAAS	84 CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.035 ± 0.014		KLOPFEN...	83 CUSB	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$				Γ_{16}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<59	90	²⁶ ALBRECHT	89 ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 5.9	90	²⁷ ALBRECHT	89 ARG	$\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-$
²⁶ Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$.				
²⁷ Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+\pi^-$.				

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$				Γ_{17}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<53	90	²⁸ ALBRECHT	89 ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
²⁸ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$.				

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$				Γ_{18}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<24.1	90	²⁹ ALBRECHT	89 ARG	$\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-$
²⁹ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.				

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.8	90	³⁰ ALBRECHT	89 ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
------	----	------------------------	--------	---

³⁰ Includes unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$.

 $\Gamma(\gamma \eta_b(1S))/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$3.9 \pm 1.1_{-0.9}^{+1.1}$	13 ± 5 k		³¹ AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------------	--------------	--	----------------------	-----------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90		LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$
-----	----	--	------	----------	-------------------------------------

< 8.4	90		³¹ BONVICINI	10 CLEO	$\Upsilon(2S) \rightarrow \gamma X$
-------	----	--	-------------------------	---------	-------------------------------------

< 5.1	90		³² ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$
-------	----	--	----------------------	---------	--------------------------------

³¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV.

³² Superseded by BONVICINI 10.

 $\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 2.7×10^{-5}	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$
------------------------	----	------	----------	-------------------------------------

 $\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 1.0×10^{-4}	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$
------------------------	----	------	----------	-------------------------------------

 $\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 3.6×10^{-6}	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$
------------------------	----	------	----------	-------------------------------------

 $\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 1.5×10^{-5}	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$
------------------------	----	------	----------	-------------------------------------

 $\Gamma(\gamma X(3872) \rightarrow \pi^+ \pi^- J/\psi)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 0.8×10^{-6}	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$
------------------------	----	------	----------	-------------------------------------

 $\Gamma(\gamma X(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 2.4×10^{-6}	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$
------------------------	----	------	----------	-------------------------------------

 $\Gamma(\gamma X(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 2.8×10^{-6}	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$
------------------------	----	------	----------	-------------------------------------

 $\Gamma(\gamma X(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 1.2×10^{-6}	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$
------------------------	----	------	----------	-------------------------------------

$\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-6}$	90	WANG	11B	BELL $\Upsilon(2S) \rightarrow \gamma X$

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ Γ_{30}/Γ
 (1.5 GeV < m_X < 5.0 GeV)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.95	95	ROSNER	07A	CLEO $e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma A^0 \rightarrow \gamma \text{ hadrons})/\Gamma_{\text{total}}$ Γ_{31}/Γ
 (0.3 GeV < m_{A^0} < 7 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-5}$	90	³³ LEES	11H	BABR $\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$

³³For a narrow scalar or pseudoscalar A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

LEPTON FAMILY NUMBER (LF) VIOLATING MODES

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.2	90	LEES	10B	BABR $e^+e^- \rightarrow e^\pm \tau^\mp$

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.3	90	LEES	10B	BABR $e^+e^- \rightarrow \mu^\pm \tau^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14.4	95	LOVE	08A	CLEO $e^+e^- \rightarrow \mu^\pm \tau^\mp$
---------	----	------	-----	--

$\Upsilon(2S)$ Cross-Particle Branching Ratios

$B(\Upsilon(2S) \rightarrow \pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \Upsilon(2S)X)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.78 \pm 0.02 \pm 0.11$	906k	LEES	11C	BABR $e^+e^- \rightarrow \pi^+\pi^- X$

$\Upsilon(2S)$ REFERENCES

LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104R	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)

ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GELPHMAN	85	PR D32 2893	D. Gelpman <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>	(DESY, ARGUS Collab.+)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)