

$N(1875) 3/2^-$

$$I(J^P) = \frac{1}{2}(3/2^-) \text{ Status: } ***$$

Before the 2012 *Review*, all the evidence for a $J^P = 3/2^-$ state with a mass above 1800 MeV was filed under a two-star $N(2080)$.

There is now evidence from ANISOVICH 12A for two $3/2^-$ states in this region, so we have split the older data (according to mass) between a three-star $N(1875)$ and a two-star $N(2120)$.

The latest GWU analysis (ARNDT 06) finds no evidence for this resonance.

 $N(1875)$ BREIT-WIGNER MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|---|
| 1820 to 1920 (≈ 1875) OUR ESTIMATE | | | |
| 1880 ± 20 | ANISOVICH | 12A | DPWA Multichannel |
| 1920 | BELL | 83 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 1880 ± 100 | ¹ CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| 1900 | SAXON | 80 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 1951 ± 27 | SHRESTHA | 12A | DPWA Multichannel |
| 2048 ± 65 | BATINIC | 10 | DPWA $\pi N \rightarrow N\pi, N\eta$ |
| 1946 ± 1 | PENNER | 02C | DPWA Multichannel |
| 1895 | MART | 00 | DPWA $\gamma p \rightarrow \Lambda K^+$ |
| 2003 ± 18 | VRANA | 00 | DPWA Multichannel |
| 1804 ± 55 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N \& N\pi\pi$ |
| 1880 | BAKER | 79 | DPWA $\pi^- p \rightarrow n\eta$ |

 $N(1875)$ BREIT-WIGNER WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|---|
| 160 to 320 (≈ 220) OUR ESTIMATE | | | |
| 200 ± 25 | ANISOVICH | 12A | DPWA Multichannel |
| 320 | BELL | 83 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 180 ± 60 | ¹ CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ (lower m) |
| 240 | SAXON | 80 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 500 ± 45 | SHRESTHA | 12A | DPWA Multichannel |
| 529 ± 128 | BATINIC | 10 | DPWA $\pi N \rightarrow N\pi, N\eta$ |
| 859 ± 7 | PENNER | 02C | DPWA Multichannel |
| 372 | MART | 00 | DPWA $\gamma p \rightarrow \Lambda K^+$ |
| 1070 ± 858 | VRANA | 00 | DPWA Multichannel |
| 450 ± 185 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N \& N\pi\pi$ |
| 87 | BAKER | 79 | DPWA $\pi^- p \rightarrow n\eta$ |

***N*(1875) POLE POSITION**

REAL PART

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|--|
| 1800 to 1950 OUR ESTIMATE | | | |
| 1860 ± 25 | ANISOVICH | 12A | DPWA Multichannel |
| 1880 ± 100 | ¹ CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ (lower <i>m</i>) |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 1975 | SHRESTHA | 12A | DPWA Multichannel |
| 1957 ± 49 | BATINIC | 10 | DPWA $\pi N \rightarrow N\pi, N\eta$ |
| 1824 | VRANA | 00 | DPWA Multichannel |
| not seen | ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |

– 2×IMAGINARY PART

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|--|
| 150 to 250 OUR ESTIMATE | | | |
| 200 ± 20 | ANISOVICH | 12A | DPWA Multichannel |
| 160 ± 80 | ¹ CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ (lower <i>m</i>) |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 495 | SHRESTHA | 12A | DPWA Multichannel |
| 467 ± 106 | BATINIC | 10 | DPWA $\pi N \rightarrow N\pi, N\eta$ |
| 614 | VRANA | 00 | DPWA Multichannel |
| not seen | ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |

***N*(1875) ELASTIC POLE RESIDUE**

MODULUS $|r|$

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|--|
| 2 to 10 OUR ESTIMATE | | | |
| 2.5 ± 1.0 | ANISOVICH | 12A | DPWA Multichannel |
| 10 ± 5 | ¹ CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ (lower <i>m</i>) |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 53 | BATINIC | 10 | DPWA $\pi N \rightarrow N\pi, N\eta$ |

PHASE θ

| <u>VALUE (°)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|--|
| 100 ± 80 | ¹ CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ (lower <i>m</i>) |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| – 65 | BATINIC | 10 | DPWA $\pi N \rightarrow N\pi, N\eta$ |

***N*(1875) INELASTIC POLE RESIDUE**

The “normalized residue” is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\pi \rightarrow N(1875) \rightarrow \Lambda K$

| <u>MODULUS (%)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|-------------------|
| 1.5 ± 0.5 | ANISOVICH | 12A | DPWA Multichannel |

Normalized residue in $N\pi \rightarrow N(1875) \rightarrow \Sigma K$

| <u>MODULUS (%)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|-------------------|
| 4 ± 2 | ANISOVICH | 12A | DPWA Multichannel |

Normalized residue in $N\pi \rightarrow N(1875) \rightarrow N\sigma$

| <u>MODULUS (%)</u> | <u>PHASE (°)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|------------------|--------------------|-------------|----------------|
| 8±3 | -170 ± 65 | ANISOVICH | 12A DPWA | Multichannel |

N(1875) DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor |
|---|--------------------------------|--------------|
| Γ_1 $N\pi$ | (12 ± 10) % | |
| Γ_2 $N\eta$ | (3.5 ± 3.5) % | 2.5 |
| Γ_3 $N\omega$ | (21 ± 7) % | |
| Γ_4 ΛK | | |
| Γ_5 ΣK | (7 ± 4) × 10 ⁻³ | |
| Γ_6 $N\pi\pi$ | | |
| Γ_7 $\Delta(1232)\pi$, S-wave | (40 ± 10) % | |
| Γ_8 $\Delta(1232)\pi$, D-wave | (17 ± 10) % | |
| Γ_9 $N\rho$, S=3/2, S-wave | (6 ± 6) % | |
| Γ_{10} $N(\pi\pi)_{S\text{-wave}}^{I=0}$ | (24 ± 24) % | |
| Γ_{11} $n\gamma$, helicity=1/2 | | |
| Γ_{12} $n\gamma$, helicity=3/2 | | |
| Γ_{13} $p\gamma$ | 0.008–0.016 % | |
| Γ_{14} $p\gamma$, helicity=1/2 | 0.006–0.010 % | |
| Γ_{15} $p\gamma$, helicity=3/2 | 0.002–0.006 % | |

N(1875) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

| <u>VALUE (%)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|--|
| 12±10 OUR ESTIMATE | | | |
| 3 ± 2 | ANISOVICH | 12A DPWA | Multichannel |
| 10 ± 4 | ¹ CUTKOSKY | 80 IPWA | $\pi N \rightarrow \pi N$ (lower m) |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 7 ± 2 | SHRESTHA | 12A DPWA | Multichannel |
| 17 ± 7 | BATINIC | 10 DPWA | $\pi N \rightarrow N\pi, N\eta$ |
| 12 ± 2 | PENNER | 02C DPWA | Multichannel |
| 13 ± 3 | VRANA | 00 DPWA | Multichannel |
| 23 ± 3 | MANLEY | 92 IPWA | $\pi N \rightarrow \pi N \& N\pi\pi$ |

$\Gamma(N\eta)/\Gamma_{\text{total}}$ Γ_2/Γ

| <u>VALUE (%)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-------------------------------------|
| 3.5±3.5 OUR AVERAGE | | | Error includes scale factor of 2.5. |
| 7 ± 2 | PENNER | 02C DPWA | Multichannel |
| 0 ± 2 | VRANA | 00 DPWA | Multichannel |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 8 ± 3 | BATINIC | 10 DPWA | $\pi N \rightarrow N\pi, N\eta$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1875) \rightarrow N\eta$ $(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|----------------------------------|
| 6 ±4 OUR ESTIMATE | | | |
| 5 ±2 | ANISOVICH | 12A | DPWA Multichannel |
| 6.5 | BAKER | 79 | DPWA $\pi^- p \rightarrow n\eta$ |

$\Gamma(N\omega) / \Gamma_{\text{total}}$ Γ_3 / Γ

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--------------|-------------|------|-------------------|
| 21 ±7 | PENNER | 02C | DPWA Multichannel |

$\Gamma(\Lambda K) / \Gamma_{\text{total}}$ Γ_4 / Γ

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|-------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.2 ±0.2 | PENNER | 02C | DPWA Multichannel |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1875) \rightarrow \Lambda K$ $(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|--|
| 4 ±2 OUR ESTIMATE | | | |
| 4 ±2 | ANISOVICH | 12A | DPWA Multichannel |
| 4 | BELL | 83 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |
| 3 | SAXON | 80 | DPWA $\pi^- p \rightarrow \Lambda K^0$ |

$\Gamma(\Sigma K) / \Gamma_{\text{total}}$ Γ_5 / Γ

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------------|-------------|------|-------------------|
| 0.7 ±0.4 | PENNER | 02C | DPWA Multichannel |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1875) \rightarrow \Sigma K$ $(\Gamma_1 \Gamma_5)^{1/2} / \Gamma$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|--------------------|------|-----------------------------------|
| 1 to 10 OUR ESTIMATE | | | |
| 15 ±8 | ANISOVICH | 12A | DPWA Multichannel |
| 1.4 to 3.7 | ² DEANS | 75 | DPWA $\pi N \rightarrow \Sigma K$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1875) \rightarrow \Delta(1232)\pi, S\text{-wave}$ $(\Gamma_1 \Gamma_7)^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| −0.09 ±0.09 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N \& N\pi\pi$ |

$\Gamma(\Delta(1232)\pi, S\text{-wave}) / \Gamma_{\text{total}}$ Γ_7 / Γ

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|-------------------|
| 40 ±10 | VRANA | 00 | DPWA Multichannel |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 87 ± 3 | SHRESTHA | 12A | DPWA Multichannel |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1875) \rightarrow \Delta(1232)\pi, D\text{-wave}$ $(\Gamma_1 \Gamma_8)^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| +0.22 ±0.07 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N \& N\pi\pi$ |

$\Gamma(\Delta(1232)\pi, D\text{-wave})/\Gamma_{\text{total}} \quad \Gamma_8/\Gamma$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|-------------------|
| 17 ± 10 | VRANA | 00 | DPWA Multichannel |
| < 6 | SHRESTHA | 12A | DPWA Multichannel |

$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1875) \rightarrow N\rho, S=3/2, S\text{-wave} \quad (\Gamma_1 \Gamma_9)^{1/2}/\Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------|-------------|------|--|
| -0.24 ± 0.06 | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$ |

$\Gamma(N\rho, S=3/2, S\text{-wave})/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------|-------------|------|-------------------|
| 6 ± 6 | VRANA | 00 | DPWA Multichannel |
| < 5 | SHRESTHA | 12A | DPWA Multichannel |

$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1875) \rightarrow N(\pi\pi)_{S\text{-wave}}^{I=0} \quad (\Gamma_1 \Gamma_{10})^{1/2}/\Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------|-------------|------|--|
| $+0.25 \pm 0.06$ | MANLEY | 92 | IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$ |

$\Gamma(N(\pi\pi)_{S\text{-wave}}^{I=0})/\Gamma_{\text{total}} \quad \Gamma_{10}/\Gamma$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|-------------------|
| 24 ± 24 | VRANA | 00 | DPWA Multichannel |
| < 4 | SHRESTHA | 12A | DPWA Multichannel |

$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1875) \rightarrow N\eta \quad (\Gamma_{13} \Gamma_2)^{1/2}/\Gamma$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|-----------------------------------|
| 60 ± 12 | ANISOVICH | 12A | DPWA Multichannel |
| 0.37 | HICKS | 73 | MPWA $\gamma p \rightarrow p\eta$ |

N(1875) PHOTON DECAY AMPLITUDES

Papers on γN amplitudes predating 1981 may be found in our 2006 edition, *Journal of Physics*, G **33** 1 (2006).

$N(1875) \rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

| VALUE ($\text{GeV}^{-1/2}$) | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-------------|------|-----------------------------------|
| 0.018 ± 0.010 | ANISOVICH | 12A | DPWA Multichannel |
| -0.020 ± 0.008 | AWAJI | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| 0.007 ± 0.008 | SHRESTHA | 12A | DPWA Multichannel |
| 0.012 | PENNER | 02D | DPWA Multichannel |
| 0.026 ± 0.052 | DEVENISH | 74 | DPWA $\gamma N \rightarrow \pi N$ |

$N(1875) \rightarrow p\gamma$, helicity-3/2 amplitude $A_{3/2}$

| <u>VALUE (GeV^{-1/2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-----------------------------------|
| -0.009 ± 0.005 | ANISOVICH | 12A | DPWA Multichannel |
| 0.017 ± 0.011 | AWAJI | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 0.043 ± 0.022 | SHRESTHA | 12A | DPWA Multichannel |
| -0.010 | PENNER | 02D | DPWA Multichannel |
| 0.128 ± 0.057 | DEVENISH | 74 | DPWA $\gamma N \rightarrow \pi N$ |

$N(1875) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

| <u>VALUE (GeV^{-1/2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-----------------------------------|
| 0.007 ± 0.013 | AWAJI | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 0.055 ± 0.021 | SHRESTHA | 12A | DPWA Multichannel |
| 0.023 | PENNER | 02D | DPWA Multichannel |
| 0.053 ± 0.083 | DEVENISH | 74 | DPWA $\gamma N \rightarrow \pi N$ |

$N(1875) \rightarrow n\gamma$, helicity-3/2 amplitude $A_{3/2}$

| <u>VALUE (GeV^{-1/2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-----------------------------------|
| -0.053 ± 0.034 | AWAJI | 81 | DPWA $\gamma N \rightarrow \pi N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| -0.085 ± 0.031 | SHRESTHA | 12A | DPWA Multichannel |
| -0.009 | PENNER | 02D | DPWA Multichannel |
| 0.100 ± 0.141 | DEVENISH | 74 | DPWA $\gamma N \rightarrow \pi N$ |

$N(1875) \quad \gamma p \rightarrow \Lambda K^+$ AMPLITUDES

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1875) \rightarrow \Lambda K^+$ (E_{2-} amplitude)

| <u>VALUE (units 10⁻³)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 2.29 ^{+0.7} _{-0.2} | MART | 00 | DPWA $\gamma p \rightarrow \Lambda K^+$ |
| 5.5 ± 0.3 | WORKMAN | 90 | DPWA |
| 4.09 | TANABE | 89 | DPWA |

$p\gamma \rightarrow N(1875) \rightarrow \Lambda K^+$ phase angle θ (E_{2-} amplitude)

| <u>VALUE (degrees)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| -48 ± 5 | WORKMAN | 90 | DPWA |
| -35.9 | TANABE | 89 | DPWA |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1875) \rightarrow \Lambda K^+$ (M_{2-} amplitude)

| <u>VALUE (units 10⁻³)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| -6.7 ± 0.2 | WORKMAN | 90 | DPWA |
| -4.09 | TANABE | 89 | DPWA |

N(1875) FOOTNOTES

- ¹ CUTKOSKY 80 finds a lower mass D_{13} resonance, as well as one in this region. Both are listed here.
² The range given for DEANS 75 is from the four best solutions. Disagrees with $\pi^+ p \rightarrow \Sigma^+ K^+$ data of WINNIK 77 around 1920 MeV.

N(1875) REFERENCES

For early references, see Physics Letters **111B** 1 (1982).

| | | | | |
|-----------|-----|------------------|---|-----------------------|
| ANISOVICH | 12A | EPJ A48 15 | A.V. Anisovich <i>et al.</i> | (BONN, PNPI) |
| SHRESTHA | 12A | PR C86 055203 | M. Shrestha, D.M. Manley | (KSU) |
| BATINIC | 10 | PR C82 038203 | M. Batinic <i>et al.</i> | (ZAGR) |
| ARNDT | 06 | PR C74 045205 | R.A. Arndt <i>et al.</i> | (GWU) |
| PDG | 06 | JPG 33 1 | W.-M. Yao <i>et al.</i> | (PDG Collab.) |
| PENNER | 02C | PR C66 055211 | G. Penner, U. Mosel | (GIES) |
| PENNER | 02D | PR C66 055212 | G. Penner, U. Mosel | (GIES) |
| MART | 00 | PR C61 012201 | T. Mart, C. Bennhold | |
| VRANA | 00 | PRPL 328 181 | T.P. Vrana, S.A. Dytman,, T.-S.H. Lee | (PITT+) |
| MANLEY | 92 | PR D45 4002 | D.M. Manley, E.M. Saleski | (KSA) IJP |
| Also | | PR D30 904 | D.M. Manley <i>et al.</i> | (VPI) |
| ARNDT | 91 | PR D43 2131 | R.A. Arndt <i>et al.</i> | (VPI, TELE) IJP |
| WORKMAN | 90 | PR C42 781 | R.L. Workman | (VPI) |
| TANABE | 89 | PR C39 741 | H. Tanabe, M. Kohno, C. Bennhold | (MANZ) |
| Also | | NC 102A 193 | M. Kohno, H. Tanabe, C. Bennhold | (MANZ) |
| BELL | 83 | NP B222 389 | K.W. Bell <i>et al.</i> | (RL) IJP |
| AWAJI | 81 | Bonn Conf. 352 | N. Awaji, R. Kajikawa | (NAGO) |
| Also | | NP B197 365 | K. Fujii <i>et al.</i> | (NAGO) |
| CUTKOSKY | 80 | Toronto Conf. 19 | R.E. Cutkosky <i>et al.</i> | (CMU, LBL) IJP |
| Also | | PR D20 2839 | R.E. Cutkosky <i>et al.</i> | (CMU, LBL) IJP |
| SAXON | 80 | NP B162 522 | D.H. Saxon <i>et al.</i> | (RHEL, BRIS) IJP |
| BAKER | 79 | NP B156 93 | R.D. Baker <i>et al.</i> | (RHEL) IJP |
| WINNIK | 77 | NP B128 66 | M. Winnik <i>et al.</i> | (HAIF) I |
| DEANS | 75 | NP B96 90 | S.R. Deans <i>et al.</i> | (SFLA, ALAH) IJP |
| DEVENISH | 74 | PL 52B 227 | R.C.E. Devenish, D.H. Lyth, W.A. Rankin | (DESY+) IJP |
| HICKS | 73 | PR D7 2614 | H.R. Hicks <i>et al.</i> | (CMU, ORNL, SFLA) IJP |