



$$I(J^P) = 0(\frac{1}{2}^+) \text{ Status: } ****$$

The parity of the Λ_c^+ is defined to be positive (as are the parities of the proton, neutron, and Λ). The quark content is udc . Results of an analysis of $pK^-\pi^+$ decays (JEZABEK 92) are consistent with $J = 1/2$. Nobody doubts that the spin is indeed $1/2$.

The only new measurements since our 2010 Review are of limits on rare or forbidden $\Lambda_c^+ \rightarrow p\ell^+\ell^-$ and $\bar{p}\ell^+\ell^+$ modes.

We have omitted some results that have been superseded by later experiments. The omitted results may be found in earlier editions.

Λ_c^+ MASS

Our value in 2004, 2284.9 ± 0.6 MeV, was the average of the measurements now filed below as "not used." The BABAR measurement is so much better that we use it alone. Note that it is about 2.6 (old) standard deviations above the 2004 value.

The fit also includes $\Sigma_c - \Lambda_c^+$ and $\Lambda_c^{*+} - \Lambda_c^+$ mass-difference measurements, but this doesn't affect the Λ_c^+ mass. The new (in 2006) Λ_c^+ mass simply pushes all those other masses higher.


| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|------|-----------------------|----------|-------------------------------------------------------------------------|
| 2286.46 ± 0.14 OUR FIT | | | | |
| 2286.46 ± 0.14 | 4891 | ¹ AUBERT,B | 05s BABR | $\Lambda K_S^0 K^+$ and $\Sigma^0 K_S^0 K^+$ |
| • • • | | | | We do not use the following data for averages, fits, limits, etc. • • • |
| 2284.7 ± 0.6 ± 0.7 | 1134 | AVERY | 91 CLEO | Six modes |
| 2281.7 ± 2.7 ± 2.6 | 29 | ALVAREZ | 90B NA14 | $pK^-\pi^+$ |
| 2285.8 ± 0.6 ± 1.2 | 101 | BARLAG | 89 NA32 | $pK^-\pi^+$ |
| 2284.7 ± 2.3 ± 0.5 | 5 | AGUILAR-... | 88B LEBC | $pK^-\pi^+$ |
| 2283.1 ± 1.7 ± 2.0 | 628 | ALBRECHT | 88C ARG | $pK^-\pi^+$, $p\bar{K}^0$, $\Lambda 3\pi$ |
| 2286.2 ± 1.7 ± 0.7 | 97 | ANJOS | 88B E691 | $pK^-\pi^+$ |
| 2281 ± 3 | 2 | JONES | 87 HBC | $pK^-\pi^+$ |
| 2283 ± 3 | 3 | BOSETTI | 82 HBC | $pK^-\pi^+$ |
| 2290 ± 3 | 1 | CALICCHIO | 80 HYBR | $pK^-\pi^+$ |

¹ AUBERT,B 05s uses low-Q $\Lambda K_S^0 K^+$ and $\Sigma^0 K_S^0 K^+$ decays to minimize systematic errors. The error above includes systematic as well as statistical errors. Many cross checks and adjustments to properties of the BABAR detector, as well as the large number of clean events, make this by far the best measurement of the Λ_c^+ mass.

Λ_c^+ MEAN LIFE

Measurements with an error $\geq 100 \times 10^{-15}$ s or with fewer than 20 events have been omitted from the Listings.

| <u>VALUE (10^{-15} s)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------------------------------------------------------|--------------------|-------------------------------------|-------------|-------------------------------------------------------------|
| 200 ± 6 | OUR AVERAGE | Error includes scale factor of 1.6. | | See the ideogram below. |
| 204.6 ± 3.4 ± 2.5 | 8034 | LINK | 02C | FOCS $pK^- \pi^+$ |
| 198.1 ± 7.0 ± 5.6 | 1630 | KUSHNIR... | 01 | SELX $\Lambda_c^+ \rightarrow pK^- \pi^+$ |
| 179.6 ± 6.9 ± 4.4 | 4749 | MAHMOOD | 01 | CLE2 $e^+ e^- \approx \Upsilon(4S)$ |
| 215 ± 16 ± 8 | 1340 | FRABETTI | 93D | E687 $\gamma \text{Be}, \Lambda_c^+ \rightarrow pK^- \pi^+$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 180 ± 30 ± 30 | 29 | ALVAREZ | 90 | NA14 $\gamma, \Lambda_c^+ \rightarrow pK^- \pi^+$ |
| 200 ± 30 ± 30 | 90 | FRABETTI | 90 | E687 $\gamma \text{Be}, \Lambda_c^+ \rightarrow pK^- \pi^+$ |
| 196 ⁺²³ ₋₂₀ | 101 | BARLAG | 89 | NA32 $pK^- \pi^+ + \text{c.c.}$ |
| 220 ± 30 ± 20 | 97 | ANJOS | 88B | E691 $pK^- \pi^+ + \text{c.c.}$ |



Λ_c^+ DECAY MODES

Nearly all branching fractions of the Λ_c^+ are measured relative to the $pK^-\pi^+$ mode, but there are no model-independent measurements of this branching fraction. We explain how we arrive at our value of $B(\Lambda_c^+ \rightarrow pK^-\pi^+)$ in a Note at the beginning of the branching-ratio measurements, below. When this branching fraction is eventually well determined, all the other branching fractions will slide up or down proportionally as the true value differs from the value we use here.

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|----------------------------------------------------------------------------------------|----------------------------------------|-----------------------------------|
| Hadronic modes with a p: $S = -1$ final states | | |
| Γ_1 $p\bar{K}^0$ | (2.3 \pm 0.6) % | |
| Γ_2 $pK^-\pi^+$ | [a] (5.0 \pm 1.3) % | |
| Γ_3 $p\bar{K}^*(892)^0$ | [b] (1.6 \pm 0.5) % | |
| Γ_4 $\Delta(1232)^{++}K^-$ | (8.6 \pm 3.0) $\times 10^{-3}$ | |
| Γ_5 $\Lambda(1520)\pi^+$ | [b] (1.8 \pm 0.6) % | |
| Γ_6 $pK^-\pi^+$ nonresonant | (2.8 \pm 0.8) % | |
| Γ_7 $p\bar{K}^0\pi^0$ | (3.3 \pm 1.0) % | |
| Γ_8 $p\bar{K}^0\eta$ | (1.2 \pm 0.4) % | |
| Γ_9 $p\bar{K}^0\pi^+\pi^-$ | (2.6 \pm 0.7) % | |
| Γ_{10} $pK^-\pi^+\pi^0$ | (3.4 \pm 1.0) % | |
| Γ_{11} $pK^*(892)^-\pi^+$ | [b] (1.1 \pm 0.5) % | |
| Γ_{12} $p(K^-\pi^+)_{\text{nonresonant}}\pi^0$ | (3.6 \pm 1.2) % | |
| Γ_{13} $\Delta(1232)\bar{K}^*(892)$ | seen | |
| Γ_{14} $pK^-\pi^+\pi^+\pi^-$ | (1.1 \pm 0.8) $\times 10^{-3}$ | |
| Γ_{15} $pK^-\pi^+\pi^0\pi^0$ | (8 \pm 4) $\times 10^{-3}$ | |
| Γ_{16} $pK^-\pi^+3\pi^0$ | | |
| Hadronic modes with a p: $S = 0$ final states | | |
| Γ_{17} $p\pi^+\pi^-$ | (3.5 \pm 2.0) $\times 10^{-3}$ | |
| Γ_{18} $pf_0(980)$ | [b] (2.8 \pm 1.9) $\times 10^{-3}$ | |
| Γ_{19} $p\pi^+\pi^+\pi^-\pi^-$ | (1.8 \pm 1.2) $\times 10^{-3}$ | |
| Γ_{20} pK^+K^- | (7.7 \pm 3.5) $\times 10^{-4}$ | |
| Γ_{21} $p\phi$ | [b] (8.2 \pm 2.7) $\times 10^{-4}$ | |
| Γ_{22} pK^+K^- non- ϕ | (3.5 \pm 1.7) $\times 10^{-4}$ | |
| Hadronic modes with a hyperon: $S = -1$ final states | | |
| Γ_{23} $\Lambda\pi^+$ | (1.07 \pm 0.28) % | |
| Γ_{24} $\Lambda\pi^+\pi^0$ | (3.6 \pm 1.3) % | |
| Γ_{25} $\Lambda\rho^+$ | < 5 % | CL=95% |
| Γ_{26} $\Lambda\pi^+\pi^+\pi^-$ | (2.6 \pm 0.7) % | |
| Γ_{27} $\Sigma(1385)^+\pi^+\pi^-$, $\Sigma^{*+} \rightarrow$ $\Lambda\pi^+$ | (7 \pm 4) $\times 10^{-3}$ | |
| Γ_{28} $\Sigma(1385)^-\pi^+\pi^+$, $\Sigma^{*-} \rightarrow$ $\Lambda\pi^-$ | (5.5 \pm 1.7) $\times 10^{-3}$ | |

| | | | |
|---------------|-------------------------------------------------------------------|----------------------------------------|--------|
| Γ_{29} | $\Lambda\pi^+\rho^0$ | (1.1 \pm 0.5) % | |
| Γ_{30} | $\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+$ | (3.7 \pm 3.1) $\times 10^{-3}$ | |
| Γ_{31} | $\Lambda\pi^+\pi^+\pi^-\text{nonresonant}$ | < 8 $\times 10^{-3}$ | CL=90% |
| Γ_{32} | $\Lambda\pi^+\pi^+\pi^-\pi^0\text{total}$ | (1.8 \pm 0.8) % | |
| Γ_{33} | $\Lambda\pi^+\eta$ | [b] (1.8 \pm 0.6) % | |
| Γ_{34} | $\Sigma(1385)^+\eta$ | [b] (8.5 \pm 3.3) $\times 10^{-3}$ | |
| Γ_{35} | $\Lambda\pi^+\omega$ | [b] (1.2 \pm 0.5) % | |
| Γ_{36} | $\Lambda\pi^+\pi^+\pi^-\pi^0, \text{no } \eta \text{ or } \omega$ | < 7 $\times 10^{-3}$ | CL=90% |
| Γ_{37} | $\Lambda K^+\bar{K}^0$ | (4.7 \pm 1.5) $\times 10^{-3}$ | S=1.2 |
| Γ_{38} | $\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Lambda\bar{K}^0$ | (1.3 \pm 0.5) $\times 10^{-3}$ | |
| Γ_{39} | $\Sigma^0\pi^+$ | (1.05 \pm 0.28) % | |
| Γ_{40} | $\Sigma^+\pi^0$ | (1.00 \pm 0.34) % | |
| Γ_{41} | $\Sigma^+\eta$ | (5.5 \pm 2.3) $\times 10^{-3}$ | |
| Γ_{42} | $\Sigma^+\pi^+\pi^-$ | (3.6 \pm 1.0) % | |
| Γ_{43} | $\Sigma^+\rho^0$ | < 1.4 % | CL=95% |
| Γ_{44} | $\Sigma^-\pi^+\pi^+$ | (1.7 \pm 0.5) % | |
| Γ_{45} | $\Sigma^0\pi^+\pi^0$ | (1.8 \pm 0.8) % | |
| Γ_{46} | $\Sigma^0\pi^+\pi^+\pi^-$ | (8.3 \pm 3.1) $\times 10^{-3}$ | |
| Γ_{47} | $\Sigma^+\pi^+\pi^-\pi^0$ | — | |
| Γ_{48} | $\Sigma^+\omega$ | [b] (2.7 \pm 1.0) % | |
| Γ_{49} | $\Sigma^+ K^+ K^-$ | (2.8 \pm 0.8) $\times 10^{-3}$ | |
| Γ_{50} | $\Sigma^+ \phi$ | [b] (3.1 \pm 0.9) $\times 10^{-3}$ | |
| Γ_{51} | $\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Sigma^+ K^-$ | (8.1 \pm 3.0) $\times 10^{-4}$ | |
| Γ_{52} | $\Sigma^+ K^+ K^- \text{nonresonant}$ | < 6 $\times 10^{-4}$ | CL=90% |
| Γ_{53} | $\Xi^0 K^+$ | (3.9 \pm 1.4) $\times 10^{-3}$ | |
| Γ_{54} | $\Xi^- K^+ \pi^+$ | (5.1 \pm 1.4) $\times 10^{-3}$ | |
| Γ_{55} | $\Xi(1530)^0 K^+$ | [b] (2.6 \pm 1.0) $\times 10^{-3}$ | |

Hadronic modes with a hyperon: $S = 0$ final states

| | | | |
|---------------|----------------------------|----------------------------------------|--------|
| Γ_{56} | ΛK^+ | (5.0 \pm 1.6) $\times 10^{-4}$ | |
| Γ_{57} | $\Lambda K^+ \pi^+ \pi^-$ | < 4 $\times 10^{-4}$ | CL=90% |
| Γ_{58} | $\Sigma^0 K^+$ | (4.2 \pm 1.3) $\times 10^{-4}$ | |
| Γ_{59} | $\Sigma^0 K^+ \pi^+ \pi^-$ | < 2.1 $\times 10^{-4}$ | CL=90% |
| Γ_{60} | $\Sigma^+ K^+ \pi^-$ | (1.7 \pm 0.7) $\times 10^{-3}$ | |
| Γ_{61} | $\Sigma^+ K^*(892)^0$ | [b] (2.8 \pm 1.1) $\times 10^{-3}$ | |
| Γ_{62} | $\Sigma^- K^+ \pi^+$ | < 1.0 $\times 10^{-3}$ | CL=90% |

Doubly Cabibbo-suppressed modes

| | | | |
|---------------|---------------|------------------------|--------|
| Γ_{63} | $p K^+ \pi^-$ | < 2.3 $\times 10^{-4}$ | CL=90% |
|---------------|---------------|------------------------|--------|

Semileptonic modes

| | | |
|---------------|-------------------------|-------------------------|
| Γ_{64} | $\Lambda\ell^+\nu_\ell$ | [c] (2.0 \pm 0.6) % |
| Γ_{65} | $\Lambda e^+\nu_e$ | (2.1 \pm 0.6) % |
| Γ_{66} | $\Lambda\mu^+\nu_\mu$ | (2.0 \pm 0.7) % |

Inclusive modes

| | | | | |
|---------------|------------------------------|-----|----------------------|-------|
| Γ_{67} | e^+ anything | | $(4.5 \pm 1.7) \%$ | |
| Γ_{68} | $p e^+$ anything | | $(1.8 \pm 0.9) \%$ | |
| Γ_{69} | Λe^+ anything | | | |
| Γ_{70} | p anything | | $(50 \pm 16) \%$ | |
| Γ_{71} | p anything (no Λ) | | $(12 \pm 19) \%$ | |
| Γ_{72} | p hadrons | | | |
| Γ_{73} | n anything | | $(50 \pm 16) \%$ | |
| Γ_{74} | n anything (no Λ) | | $(29 \pm 17) \%$ | |
| Γ_{75} | Λ anything | | $(35 \pm 11) \%$ | S=1.4 |
| Γ_{76} | Σ^\pm anything | [d] | $(10 \pm 5) \%$ | |
| Γ_{77} | 3prongs | | $(24 \pm 8) \%$ | |

**$\Delta C = 1$ weak neutral current ($C1$) modes, or
Lepton Family number (LF), or Lepton number (L), or
Baryon number (B) violating modes**

| | | | | | |
|---------------|------------------------|--------|---------|------------------|--------|
| Γ_{78} | $p e^+ e^-$ | $C1$ | < 5.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{79} | $p \mu^+ \mu^-$ | $C1$ | < 4.4 | $\times 10^{-5}$ | CL=90% |
| Γ_{80} | $p e^+ \mu^-$ | LF | < 9.9 | $\times 10^{-6}$ | CL=90% |
| Γ_{81} | $p e^- \mu^+$ | LF | < 1.9 | $\times 10^{-5}$ | CL=90% |
| Γ_{82} | $\bar{p} 2e^+$ | L, B | < 2.7 | $\times 10^{-6}$ | CL=90% |
| Γ_{83} | $\bar{p} 2\mu^+$ | L, B | < 9.4 | $\times 10^{-6}$ | CL=90% |
| Γ_{84} | $\bar{p} e^+ \mu^+$ | L, B | < 1.6 | $\times 10^{-5}$ | CL=90% |
| Γ_{85} | $\Sigma^- \mu^+ \mu^+$ | L | < 7.0 | $\times 10^{-4}$ | CL=90% |

[a] See the note on " Λ_c^+ Branching Fractions" below.

[b] This branching fraction includes all the decay modes of the final-state resonance.

[c] An ℓ indicates an e or a μ mode, not a sum over these modes.

[d] The value is for the sum of the charge states or particle/antiparticle states indicated.

CONSTRAINED FIT INFORMATION

An overall fit to 18 branching ratios uses 33 measurements and one constraint to determine 12 parameters. The overall fit has a $\chi^2 = 15.5$ for 22 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | | | | | | | | |
|----------|-------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| x_{23} | 96 | | | | | | | | | |
| x_{26} | 97 | 93 | | | | | | | | |
| x_{37} | 82 | 83 | 80 | | | | | | | |
| x_{39} | 95 | 98 | 92 | 82 | | | | | | |
| x_{42} | 93 | 90 | 91 | 77 | 88 | | | | | |
| x_{44} | 82 | 79 | 80 | 68 | 78 | 80 | | | | |
| x_{46} | 69 | 66 | 70 | 57 | 66 | 65 | 57 | | | |
| x_{49} | 88 | 85 | 86 | 72 | 84 | 93 | 75 | 61 | | |
| x_{50} | 85 | 82 | 83 | 70 | 81 | 90 | 72 | 59 | 84 | |
| x_{54} | 93 | 96 | 90 | 80 | 94 | 87 | 77 | 64 | 82 | 79 |
| | x_2 | x_{23} | x_{26} | x_{37} | x_{39} | x_{42} | x_{44} | x_{46} | x_{49} | x_{50} |

A REVIEW GOES HERE – Check our WWW List of Reviews

Λ_c^+ BRANCHING RATIOS

————— Hadronic modes with a p : $S = -1$ final states —————

| $\Gamma(p\bar{K}^0)/\Gamma(pK^-\pi^+)$ | | | | | Γ_1/Γ_2 |
|----------------------------------------|------|-------------|------|---------|-----------------------------|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 0.47±0.04 OUR AVERAGE | | | | | |
| 0.46±0.02±0.04 | 1025 | ALAM | 98 | CLE2 | $e^+e^- \approx \gamma(4S)$ |
| 0.44±0.07±0.05 | 133 | AVERY | 91 | CLEO | e^+e^- 10.5 GeV |
| 0.55±0.17±0.14 | 45 | ANJOS | 90 | E691 | γ Be 70–260 GeV |
| 0.62±0.15±0.03 | 73 | ALBRECHT | 88c | ARG | e^+e^- 10 GeV |

| $\Gamma(pK^-\pi^+)/\Gamma_{\text{total}}$ | | | | | Γ_2/Γ |
|-------------------------------------------------------------|--|--|--|--|-------------------|
| See the note on “ Λ_c^+ Branching Fractions” above. | | | | | |

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------------------------------------------------|------|-------------------------|------|---------------------------------|
| 0.050±0.013 OUR FIT | | | | |
| 0.050±0.013 | | PDG | 02 | See note at top of ratios |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.050±0.005±0.012 | 1205 | ² JAFFE | 00 | CLE2 e^+e^- 10.52–10.58 GeV |
| 0.041±0.010 | | ^{3,4} ALBRECHT | 92o | ARG $e^+e^- \approx \gamma(4S)$ |
| 0.044±0.012 | | ^{3,5} CRAWFORD | 92 | CLEO e^+e^- 10.5 GeV |

² JAFFE 00 assumes that a \bar{D} meson and an antiproton in opposite hemispheres tags for a Λ_c^+ in the hemisphere of the \bar{p} . The fraction of such $\bar{D}\bar{p}$ events with a $\Lambda_c^+ \rightarrow pK^-\pi^+$ decay then gives the $pK^-\pi^+$ branching fraction. See the paper for assumptions, caveats, etc.

³ To extract $\Gamma(pK^-\pi^+)/\Gamma_{\text{total}}$, we use $B(\bar{B} \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (0.28 \pm 0.06)\%$, which is the average of measurements from ARGUS (ALBRECHT 88C) and CLEO (CRAWFORD 92).

⁴ ALBRECHT 920 measures $B(\bar{B} \rightarrow \Lambda_c^+ X) = (6.8 \pm 0.5 \pm 0.3)\%$.

⁵ CRAWFORD 92 measures $B(\bar{B} \rightarrow \Lambda_c^+ X) = (6.4 \pm 0.8 \pm 0.8)\%$.

$\Gamma(p\bar{K}^*(892)^0)/\Gamma(pK^-\pi^+)$

Γ_3/Γ_2

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------------------------------------------------|------|------------------------|------|------------------------------------|
| 0.31±0.04 OUR AVERAGE | | | | |
| 0.29±0.04±0.03 | | ⁶ AITALA 00 | E791 | $\pi^- N$, 500 GeV |
| 0.35 ^{+0.06} _{-0.07} ±0.03 | 39 | BOZEK 93 | NA32 | $\pi^- Cu$ 230 GeV |
| 0.42±0.24 | 12 | BASILE 81B | CNTR | $pp \rightarrow \Lambda_c^+ e^- X$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.35±0.11 | | BARLAG 90D | NA32 | See BOZEK 93 |

⁶ AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow pK^-\pi^+$ decays.

$\Gamma(\Delta(1232)^{++}K^-)/\Gamma(pK^-\pi^+)$

Γ_4/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------------------------------------------|------|------------------------|------|------------------------------------|
| 0.17±0.04 OUR AVERAGE Error includes scale factor of 1.1. | | | | |
| 0.18±0.03±0.03 | | ⁷ AITALA 00 | E791 | $\pi^- N$, 500 GeV |
| 0.12 ^{+0.04} _{-0.05} ±0.05 | 14 | BOZEK 93 | NA32 | $\pi^- Cu$ 230 GeV |
| 0.40±0.17 | 17 | BASILE 81B | CNTR | $pp \rightarrow \Lambda_c^+ e^- X$ |

⁷ AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow pK^-\pi^+$ decays.

$\Gamma(\Lambda(1520)\pi^+)/\Gamma(pK^-\pi^+)$

Γ_5/Γ_2

Unseen decay modes of the $\Lambda(1520)$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------------------|------|------------------------|------|---------------------|
| 0.35±0.08 OUR AVERAGE | | | | |
| 0.34±0.08±0.05 | | ⁸ AITALA 00 | E791 | $\pi^- N$, 500 GeV |
| 0.40 ^{+0.18} _{-0.13} ±0.09 | 12 | BOZEK 93 | NA32 | $\pi^- Cu$ 230 GeV |

⁸ AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow pK^-\pi^+$ decays.

$\Gamma(pK^-\pi^+ \text{ nonresonant})/\Gamma(pK^-\pi^+)$

Γ_6/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------------------|------|------------------------|------|---------------------|
| 0.55±0.06 OUR AVERAGE | | | | |
| 0.55±0.06±0.04 | | ⁹ AITALA 00 | E791 | $\pi^- N$, 500 GeV |
| 0.56 ^{+0.07} _{-0.09} ±0.05 | 71 | BOZEK 93 | NA32 | $\pi^- Cu$ 230 GeV |

⁹ AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow p K^- \pi^+$ decays.

$\Gamma(p\bar{K}^0\pi^0)/\Gamma(pK^-\pi^+)$ Γ_7/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|-------------------------------|
| 0.66±0.05±0.07 | 774 | ALAM | 98 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(p\bar{K}^0\eta)/\Gamma(pK^-\pi^+)$ Γ_8/Γ_2

Unseen decay modes of the η are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|-------------------------------|
| 0.25±0.04±0.04 | 57 | AMMAR | 95 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(p\bar{K}^0\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_9/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------|-------------|--------------------|-------------|-------------------------------|
| 0.51±0.06 OUR AVERAGE | | | | |
| 0.52±0.04±0.05 | 985 | ALAM | 98 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| 0.43±0.12±0.04 | 83 | AVERY | 91 CLEO | e^+e^- 10.5 GeV |
| 0.98±0.36±0.08 | 12 | BARLAG | 90D NA32 | π^- 230 GeV |

$\Gamma(pK^-\pi^+\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{10}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|-------------------------------|
| 0.67±0.04±0.11 | 2606 | ALAM | 98 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(pK^*(892)^-\pi^+)/\Gamma(p\bar{K}^0\pi^+\pi^-)$ Γ_{11}/Γ_9

Unseen decay modes of the $K^*(892)^-$ are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|-------------|--------------------|-------------|----------------|
| 0.44±0.14 | 17 | ALEEV | 94 BIS2 | nN 20–70 GeV |

$\Gamma(p(K^-\pi^+)_{\text{nonresonant}}\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{12}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|--------------------|
| 0.73±0.12±0.05 | 67 | BOZEK | 93 NA32 | π^- Cu 230 GeV |

$\Gamma(\Delta(1232)\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{13}/Γ

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|-------------|--------------------|-------------|----------------|
| seen | 35 | AMENDOLIA | 87 SPEC | γ Ge-Si |

$\Gamma(pK^-\pi^+\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{14}/Γ_2

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|-----------------|
| 0.022±0.015 | BARLAG 90D | NA32 | π^- 230 GeV |

$\Gamma(pK^-\pi^+\pi^0\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{15}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|--------------------|
| 0.16±0.07±0.03 | 15 | BOZEK | 93 NA32 | π^- Cu 230 GeV |

$\Gamma(pK^-\pi^+3\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{16}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------|-------------|--------------------|-------------|--------------------|
| ••• | | | | |
| 0.10±0.06±0.02 | 8 | BOZEK | 93 NA32 | π^- Cu 230 GeV |

————— **Hadronic modes with a p : $S = 0$ final states** —————

$\Gamma(p\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{17}/Γ_2

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|-----------------|
| 0.069±0.036 | BARLAG | 90D NA32 | π^- 230 GeV |

$\Gamma(p f_0(980))/\Gamma(pK^-\pi^+)$ Γ_{18}/Γ_2

Unseen decay modes of the $f_0(980)$ are included.

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|-----------------|
| 0.055±0.036 | BARLAG | 90D NA32 | π^- 230 GeV |

$\Gamma(p\pi^+\pi^+\pi^-\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{19}/Γ_2

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|-----------------|
| 0.036±0.023 | BARLAG | 90D NA32 | π^- 230 GeV |

$\Gamma(pK^+K^-)/\Gamma(pK^-\pi^+)$ Γ_{20}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------------------------------------------------------|-------------|--------------------|-------------|---------------------------------------|
| 0.015±0.006 OUR AVERAGE | | | | Error includes scale factor of 2.1. |
| 0.014±0.002±0.002 | 676 | ABE | 02C BELL | $e^+e^- \approx \Upsilon(4S)$ |
| 0.039±0.009±0.007 | 214 | ALEXANDER | 96C CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.096±0.029±0.010 | 30 | FRABETTI | 93H E687 | γ Be, \bar{E}_γ 220 GeV |
| 0.048±0.027 | | BARLAG | 90D NA32 | π^- 230 GeV |

$\Gamma(p\phi)/\Gamma(pK^-\pi^+)$ Γ_{21}/Γ_2

Unseen decay modes of the ϕ are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------------------------------------------------------|-------------|--------------------|-------------|-------------------------------------|
| 0.0164±0.0032 OUR AVERAGE | | | | Error includes scale factor of 1.2. |
| 0.015 ±0.002 ±0.002 | 345 | ABE | 02C BELL | $e^+e^- \approx \Upsilon(4S)$ |
| 0.024 ±0.006 ±0.003 | 54 | ALEXANDER | 96C CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.040 ±0.027 | | BARLAG | 90D NA32 | π^- 230 GeV |

$\Gamma(pK^+K^- \text{ non-}\phi)/\Gamma(pK^-\pi^+)$ Γ_{22}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|-------------------------------|
| 0.007±0.002±0.002 | 344 | ABE | 02C BELL | $e^+e^- \approx \Upsilon(4S)$ |

————— **Hadronic modes with a hyperon: $S = -1$ final states** —————

$\Gamma(\Lambda\pi^+)/\Gamma(pK^-\pi^+)$ Γ_{23}/Γ_2

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------------------------------------------------------|------------|-------------|--------------------|-------------|----------------------------------------------------|
| 0.214±0.016 OUR FIT | | | | | Error includes scale factor of 1.1. |
| 0.204±0.019 OUR AVERAGE | | | | | |
| 0.217±0.013±0.020 | | 750 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.18 ±0.03 ±0.04 | | | ALBRECHT | 92 ARG | $e^+e^- \approx 10.4$ GeV |
| 0.18 ±0.03 ±0.03 | | 87 | AVERY | 91 CLEO | e^+e^- 10.5 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <0.33 | | 90 | ANJOS | 90 E691 | γ Be 70–260 GeV |
| <0.16 | | 90 | ALBRECHT | 88C ARG | e^+e^- 10 GeV |

$\Gamma(\Lambda\pi^+\pi^0)/\Gamma(\rho K^-\pi^+)$ Γ_{24}/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|---------|---------------------------------------------|
| 0.73±0.09±0.16 | 464 | AVERY | 94 CLE2 | $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$ |

$\Gamma(\Lambda\rho^+)/\Gamma(\rho K^-\pi^+)$ Γ_{25}/Γ_2

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------|-----|-------------|---------|---------------------------------------------|
| <0.95 | 95 | AVERY | 94 CLE2 | $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$ |

$\Gamma(\Lambda\pi^+\pi^+\pi^-)/\Gamma(\rho K^-\pi^+)$ Γ_{26}/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------|----------|----------------------------------------------------|
| 0.525±0.032 OUR FIT | | | | |
| 0.522±0.032 OUR AVERAGE | | | | |
| 0.508±0.024±0.024 | 1356 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.65 ±0.11 ±0.12 | 289 | AVERY | 91 CLEO | e^+e^- 10.5 GeV |
| 0.82 ±0.29 ±0.27 | 44 | ANJOS | 90 E691 | γ Be 70–260 GeV |
| 0.94 ±0.41 ±0.13 | 10 | BARLAG | 90D NA32 | π^- 230 GeV |
| 0.61 ±0.16 ±0.04 | 105 | ALBRECHT | 88C ARG | e^+e^- 10 GeV |

$\Gamma(\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow \Lambda\pi^+)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_{27}/Γ_{26}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|----------|----------------------------------------------------|
| 0.28±0.10±0.08 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma(1385)^-\pi^+\pi^+, \Sigma^{*-} \rightarrow \Lambda\pi^-)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_{28}/Γ_{26}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|----------|----------------------------------------------------|
| 0.21±0.03±0.02 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Lambda\pi^+\rho^0)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_{29}/Γ_{26}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|----------|----------------------------------------------------|
| 0.40±0.12±0.12 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_{30}/Γ_{26}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|----------|----------------------------------------------------|
| 0.14±0.09±0.07 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Lambda\pi^+\pi^+\pi^- \text{ nonresonant})/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_{31}/Γ_{26}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------|-----|-------------|----------|----------------------------------------------------|
| <0.3 | 90 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\rho\bar{K}^0\pi^+\pi^-)/\Gamma(\Lambda\pi^+\pi^+\pi^-)$ Γ_9/Γ_{26}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------------------------------------------------|------|-------------|---------|--------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2.6±1.2 | | ALEEV | 96 SPEC | n nucleus, 50 GeV/ c |
| 4.3±1.2 | 130 | ALEEV | 84 BIS2 | n C 40–70 GeV |

$\Gamma(\Lambda\pi^+\pi^+\pi^-\pi^0 \text{ total})/\Gamma(\rho K^-\pi^+)$ Γ_{32}/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|------------------------------|------|-------------------------------|
| 0.36±0.09±0.09 | 50 | ¹⁰ CRONIN-HEN..03 | CLE3 | $e^+e^- \approx \Upsilon(4S)$ |

¹⁰ CRONIN-HENNESSY 03 finds this channel to be dominantly $\Lambda\eta\pi^+$ and $\Lambda\omega\pi^+$; see below.

$\Gamma(\Lambda\pi^+\eta)/\Gamma(\rho K^-\pi^+)$ Γ_{33}/Γ_2

Unseen decay modes of the η are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------|-------------|--------------------|-------------|-------------------------------|
| 0.36±0.07 OUR AVERAGE | | | | |
| 0.41±0.17±0.10 | 11 | CRONIN-HEN..03 | CLE3 | $e^+e^- \approx \Upsilon(4S)$ |
| 0.35±0.05±0.06 | 116 | AMMAR 95 | CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma(1385)^+\eta)/\Gamma(\rho K^-\pi^+)$ Γ_{34}/Γ_2

Unseen decay modes of the $\Sigma(1385)^+$ and η are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|-------------------------------|
| 0.17±0.04±0.03 | 54 | AMMAR 95 | CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda\pi^+\omega)/\Gamma(\rho K^-\pi^+)$ Γ_{35}/Γ_2

Unseen decay modes of the ω are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|-------------------------------|
| 0.24±0.06±0.06 | 32 | CRONIN-HEN..03 | CLE3 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda\pi^+\pi^+\pi^-\pi^0, \text{no } \eta \text{ or } \omega)/\Gamma(\rho K^-\pi^+)$ Γ_{36}/Γ_2

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|------------|--------------------|-------------|-------------------------------|
| <0.13 | 90 | CRONIN-HEN..03 | CLE3 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda K^+\bar{K}^0)/\Gamma(\rho K^-\pi^+)$ Γ_{37}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|--------------------|-------------|----------------------------------------------------|
| 0.093±0.018 OUR FIT | | | | Error includes scale factor of 1.7. |
| 0.131±0.020 OUR AVERAGE | | | | |
| 0.142±0.018±0.022 | 251 | LINK 05F | FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.12 ±0.02 ±0.02 | 59 | AMMAR 95 | CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Lambda\bar{K}^0)/\Gamma(\Lambda K^+\bar{K}^0)$ Γ_{38}/Γ_{37}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------|-------------|--------------------|-------------|----------------------------------------------------|
| 0.28±0.07 OUR AVERAGE | | | | |
| 0.32±0.10±0.04 | 84 ± 24 | LINK 05F | FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.26±0.08±0.03 | 93 | ABE 02C | BELL | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda K^+\bar{K}^0)/\Gamma(\Lambda\pi^+)$ Γ_{37}/Γ_{23}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|-------------|--------------------|-------------|-------------------------------------|
| 0.43 ±0.08 OUR FIT | | | | Error includes scale factor of 2.0. |
| 0.395±0.026±0.036 | 460 ± 30 | AUBERT 07U | BABR | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^0\pi^+)/\Gamma(\rho K^-\pi^+)$ Γ_{39}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------|-------------|--------------------|-------------|---------------------------------------------|
| 0.210±0.018 OUR FIT | | | | |
| 0.20 ±0.04 OUR AVERAGE | | | | |
| 0.21 ±0.02 ±0.04 | 196 | AVERY 94 | CLE2 | $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$ |
| 0.17 ±0.06 ±0.04 | | ALBRECHT 92 | ARG | $e^+e^- \approx 10.4$ GeV |

$\Gamma(\Sigma^0\pi^+)/\Gamma(\Lambda\pi^+)$ Γ_{39}/Γ_{23}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------|-------------|--------------------|-------------|----------------------------------------------------|
| 0.98 ±0.05 OUR FIT | | | | |
| 0.98 ±0.05 OUR AVERAGE | | | | |
| 0.977±0.015±0.051 | 33k | AUBERT 07U | BABR | $e^+e^- \approx \Upsilon(4S)$ |
| 1.09 ±0.11 ±0.19 | 750 | LINK 05F | FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma^+ \pi^0)/\Gamma(\rho K^- \pi^+)$ Γ_{40}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|--------------------------------|
| 0.20±0.03±0.03 | 93 | KUBOTA | 93 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^+ \eta)/\Gamma(\rho K^- \pi^+)$ Γ_{41}/Γ_2

Unseen decay modes of the η are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|--------------------------------|
| 0.11±0.03±0.02 | 26 | AMMAR | 95 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^+ \pi^+ \pi^-)/\Gamma(\rho K^- \pi^+)$ Γ_{42}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------------|-------------|--------------------|-------------|--------------------------------|
| 0.72±0.07 OUR FIT | | | | |
| 0.69±0.08 OUR AVERAGE | | | | |
| 0.72±0.14 | 47 ± 9 | VAZQUEZ-JA..08 | SELX | Σ^- nucleus, 600 GeV |
| 0.74±0.07±0.09 | 487 | KUBOTA | 93 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |
| 0.54 ^{+0.18} _{-0.15} | 11 | BARLAG | 92 NA32 | π^- Cu 230 GeV |

$\Gamma(\Sigma^+ \rho^0)/\Gamma(\rho K^- \pi^+)$ Γ_{43}/Γ_2

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|------------|--------------------|-------------|--------------------------------|
| <0.27 | 95 | KUBOTA | 93 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^- \pi^+ \pi^+)/\Gamma(\rho K^- \pi^+)$ Γ_{44}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|-------------|--------------------|-------------|-----------------------------|
| 0.33 ±0.06 OUR FIT | | | | |
| 0.314±0.067 | 30 ± 6 | VAZQUEZ-JA..08 | SELX | Σ^- nucleus, 600 GeV |

$\Gamma(\Sigma^- \pi^+ \pi^+)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{44}/Γ_{42}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|---------------------------------------|
| 0.46±0.09 OUR FIT | | | | |
| 0.53±0.15±0.07 | 56 | FRABETTI | 94E E687 | γ Be, \bar{E}_γ 220 GeV |

$\Gamma(\Sigma^0 \pi^+ \pi^0)/\Gamma(\rho K^- \pi^+)$ Γ_{45}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|----------------------------------------------|
| 0.36±0.09±0.10 | 117 | AVERY | 94 CLE2 | $e^+ e^- \approx \Upsilon(3S), \Upsilon(4S)$ |

$\Gamma(\Sigma^0 \pi^+ \pi^+ \pi^-)/\Gamma(\rho K^- \pi^+)$ Γ_{46}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|----------------------------------------------|
| 0.17±0.04 OUR FIT | | | | |
| 0.21±0.05±0.05 | 90 | AVERY | 94 CLE2 | $e^+ e^- \approx \Upsilon(3S), \Upsilon(4S)$ |

$\Gamma(\Sigma^0 \pi^+ \pi^+ \pi^-)/\Gamma(\Lambda \pi^+ \pi^+ \pi^-)$ Γ_{46}/Γ_{26}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|----------------------------------------------------|
| 0.31±0.08 OUR FIT | | | | |
| 0.26±0.06±0.09 | 480 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma^+ \omega)/\Gamma(\rho K^- \pi^+)$ Γ_{48}/Γ_2

Unseen decay modes of the ω are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|--------------------------------|
| 0.54±0.13±0.06 | 107 | KUBOTA | 93 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^+ K^+ K^-)/\Gamma(p K^- \pi^+)$ Γ_{49}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|-------------|--------------------|-------------|---------------------------------|
| 0.056±0.008 OUR FIT | | | | |
| 0.070±0.011±0.011 | 59 | AVERY | 93 | CLE2 $e^+ e^- \approx 10.5$ GeV |

$\Gamma(\Sigma^+ K^+ K^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{49}/Γ_{42}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|--------------------|-------------|------------------------------------------|
| 0.078±0.009 OUR FIT | | | | |
| 0.074±0.009 OUR AVERAGE | | | | |
| 0.076±0.007±0.009 | 246 | ABE | 02C | BELL $e^+ e^- \approx \Upsilon(4S)$ |
| 0.071±0.011±0.011 | 103 | LINK | 02G | FOCS γ nucleus, ≈ 180 GeV |

$\Gamma(\Sigma^+ \phi)/\Gamma(p K^- \pi^+)$ Γ_{50}/Γ_2

Unseen decay modes of the ϕ are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|-------------|--------------------|-------------|---------------------------------|
| 0.062±0.010 OUR FIT | | | | |
| 0.069±0.023±0.016 | 26 | AVERY | 93 | CLE2 $e^+ e^- \approx 10.5$ GeV |

$\Gamma(\Sigma^+ \phi)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{50}/Γ_{42}

Unseen decay modes of the ϕ are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|--------------------|-------------|------------------------------------------|
| 0.087±0.012 OUR FIT | | | | |
| 0.086±0.012 OUR AVERAGE | | | | |
| 0.085±0.012±0.012 | 129 | ABE | 02C | BELL $e^+ e^- \approx \Upsilon(4S)$ |
| 0.087±0.016±0.006 | 57 | LINK | 02G | FOCS γ nucleus, ≈ 180 GeV |

$\Gamma(\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Sigma^+ K^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{51}/Γ_{42}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|--------------------|-------------|------------------------------------------|
| 0.023±0.005 OUR AVERAGE | | | | |
| 0.023±0.005±0.005 | 75 | ABE | 02C | BELL $e^+ e^- \approx \Upsilon(4S)$ |
| 0.022±0.006±0.006 | 34 | LINK | 02G | FOCS γ nucleus, ≈ 180 GeV |

$\Gamma(\Sigma^+ K^+ K^- \text{ nonresonant})/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{52}/Γ_{42}


| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------------------------------------------------------|------------|--------------------|-------------|------------------------------------------|
| <0.018 | 90 | ABE | 02C | BELL $e^+ e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.028 | 90 | LINK | 02G | FOCS γ nucleus, ≈ 180 GeV |

$\Gamma(\Xi^0 K^+)/\Gamma(p K^- \pi^+)$ Γ_{53}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|---------------------------------|
| 0.078±0.013±0.013 | 56 | AVERY | 93 | CLE2 $e^+ e^- \approx 10.5$ GeV |

$\Gamma(\Xi^- K^+ \pi^+)/\Gamma(p K^- \pi^+)$ Γ_{54}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------------------------------------------------------|--------------------|-------------|---------------------------------|
| 0.102±0.010 OUR FIT | Error includes scale factor of 1.1. | | | |
| 0.098±0.021 OUR AVERAGE | Error includes scale factor of 1.3. See the ideogram below. | | | |
| 0.14 ±0.03 ±0.02 | 34 | ALBRECHT | 95B | ARG $e^+ e^- \approx 10.4$ GeV |
| 0.079±0.013±0.014 | 60 | AVERY | 93 | CLE2 $e^+ e^- \approx 10.5$ GeV |
| 0.15 ±0.04 ±0.03 | 30 | AVERY | 91 | CLEO $e^+ e^- 10.5$ GeV |



$\Gamma(\Xi(1530)^0 K^+)/\Gamma(p K^- \pi^+)$ Γ_{55}/Γ_2

Unseen decay modes of the $\Xi(1530)^0$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|--------------|------|----------------------------|
| 0.052±0.014 OUR AVERAGE | | | | |
| 0.05 ±0.02 ±0.01 | 11 | ALBRECHT 95B | ARG | $e^+ e^- \approx 10.4$ GeV |
| 0.053±0.016±0.010 | 24 | AVERY 93 | CLE2 | $e^+ e^- \approx 10.5$ GeV |

$\Gamma(\Xi^- K^+ \pi^+)/\Gamma(\Lambda \pi^+)$ Γ_{54}/Γ_{23}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----------|-------------|----------|------------------------------|
| 0.47 ±0.04 OUR FIT | | | | |
| 0.480±0.016±0.039 | 2665 ± 84 | AUBERT | 07U BABR | $e^+ e^- \approx \gamma(4S)$ |

————— **Hadronic modes with a hyperon: S = 0 final states** —————

$\Gamma(\Lambda K^+)/\Gamma(\Lambda \pi^+)$ Γ_{56}/Γ_{23}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------------|-------------|----------|-------------------------------------|
| 0.047±0.009 OUR AVERAGE | | | | Error includes scale factor of 1.8. |
| 0.044±0.004±0.003 | 1162 ± 101 | AUBERT | 07U BABR | $e^+ e^- \approx \gamma(4S)$ |
| 0.074±0.010±0.012 | 265 | ABE | 02C BELL | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Lambda K^+ \pi^+ \pi^-)/\Gamma(\Lambda \pi^+)$ Γ_{57}/Γ_{23}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-------------|----------|------------------------------|
| <4.1 × 10⁻² | 90 | AUBERT | 07U BABR | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Sigma^0 K^+)/\Gamma(\Sigma^0 \pi^+)$ Γ_{58}/Γ_{39}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|--------------------|-------------|--------------------------------|
| 0.040±0.006 OUR AVERAGE | | | | |
| 0.038±0.005±0.003 | 366 ± 52 | AUBERT | 07U BABR | $e^+ e^- \approx \Upsilon(4S)$ |
| 0.056±0.014±0.008 | 75 | ABE | 02C BELL | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^0 K^+ \pi^+ \pi^-)/\Gamma(\Sigma^0 \pi^+)$ Γ_{59}/Γ_{39}

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------|--------------------|-------------|--------------------------------|
| <2.0 × 10⁻² | 90 | AUBERT | 07U BABR | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^+ K^+ \pi^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{60}/Γ_{42}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|--------------------------------|
| 0.047±0.011±0.008 | 105 | ABE | 02C BELL | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^+ K^*(892)^0)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{61}/Γ_{42}

Unseen decay modes of the $K^*(892)^0$ are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|-------------------------------------|
| 0.078±0.018±0.013 | 49 | LINK | 02G FOCS | γ nucleus, ≈ 180 GeV |

$\Gamma(\Sigma^- K^+ \pi^+)/\Gamma(\Sigma^+ K^*(892)^0)$ Γ_{62}/Γ_{61}

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|------------|--------------------|-------------|-------------------------------------|
| <0.35 | 90 | LINK | 02G FOCS | γ nucleus, ≈ 180 GeV |

———— Doubly Cabibbo-suppressed modes ————

$\Gamma(\rho K^+ \pi^-)/\Gamma(\rho K^- \pi^+)$ Γ_{63}/Γ_2

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------|------------|--------------------|-------------|----------------------------------|
| <0.0046 | 90 | LINK | 05K FOCS | $R = (0.05 \pm 0.26 \pm 0.02)\%$ |

———— Semileptonic modes ————

$\Gamma(\Lambda \ell^+ \nu_\ell)/\Gamma(\rho K^- \pi^+)$ Γ_{64}/Γ_2

We average here the averages of the next two data blocks.

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>COMMENT</u> |
|------------------------------|--------------------|------------------------------------------------------------|
| 0.41±0.05 OUR AVERAGE | | |
| 0.42±0.07 | PDG 02 | Our $\Gamma(\Lambda e^+ \nu_e)/\Gamma(\rho K^- \pi^+)$ |
| 0.39±0.08 | PDG 02 | Our $\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma(\rho K^- \pi^+)$ |

$\Gamma(\Lambda e^+ \nu_e)/\Gamma(\rho K^- \pi^+)$ Γ_{65}/Γ_2

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------|--------------------|-------------|--------------------------------|
| 0.42±0.07 OUR AVERAGE | | | |
| 0.43±0.08 | 11,12 BERGFELD 94 | CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |
| 0.38±0.14 | 12,13 ALBRECHT 91G | ARG | $e^+ e^- \approx 10.4$ GeV |

¹¹ BERGFELD 94 measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (4.87 \pm 0.28 \pm 0.69)$ pb.

¹² To extract $\Gamma(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e)/\Gamma(\Lambda_c^+ \rightarrow \rho K^- \pi^+)$, we use $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c \rightarrow \rho K^- \pi^+) = (11.2 \pm 1.3)$ pb, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (AVERY 91).

¹³ ALBRECHT 91G measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (4.20 \pm 1.28 \pm 0.71)$ pb.

$\Gamma(\Lambda\mu^+\nu_\mu)/\Gamma(pK^-\pi^+)$

Γ_{66}/Γ_2

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|-------------|----------------------------------|
| 0.39±0.08 OUR AVERAGE | | | |
| 0.40±0.09 | 14,15 BERGFELD 94 | CLE2 | $e^+e^- \approx \mathcal{R}(4S)$ |
| 0.35±0.20 | 15,16 ALBRECHT 91G | ARG | $e^+e^- \approx 10.4$ GeV |
| <p>¹⁴ BERGFELD 94 measures $\sigma(e^+e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda\mu^+\nu_\mu) = (4.43 \pm 0.51 \pm 0.64)$ pb.</p> <p>¹⁵ To extract $\Gamma(\Lambda_c^+ \rightarrow \Lambda\mu^+\nu_\mu)/\Gamma(\Lambda_c^+ \rightarrow pK^-\pi^+)$, we use $\sigma(e^+e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (11.2 \pm 1.3)$ pb, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (AVERY 91).</p> <p>¹⁶ ALBRECHT 91G measures $\sigma(e^+e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda\mu^+\nu_\mu) = (3.91 \pm 2.02 \pm 0.90)$ pb.</p> | | | |

Inclusive modes

$\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}}$

Γ_{67}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|----------------------|
| 0.045±0.017 | VELLA 82 | MRK2 | e^+e^- 4.5–6.8 GeV |

$\Gamma(pe^+ \text{ anything})/\Gamma_{\text{total}}$

Γ_{68}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|------------------------|-------------|----------------------|
| 0.018±0.009 | ¹⁷ VELLA 82 | MRK2 | e^+e^- 4.5–6.8 GeV |

¹⁷ VELLA 82 includes protons from Λ decay.

$\Gamma(\Lambda e^+ \text{ anything})/\Gamma_{\text{total}}$

Γ_{69}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------------------------------------------------------|------------------------|-------------|----------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.011±0.008 | ¹⁸ VELLA 82 | MRK2 | e^+e^- 4.5–6.8 GeV |

¹⁸ VELLA 82 includes Λ 's from Σ^0 decay.

$\Gamma(p \text{ anything})/\Gamma_{\text{total}}$

Γ_{70}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|---------------------------|-------------|-------------------|
| 0.50±0.08±0.14 | ¹⁹ CRAWFORD 92 | CLEO | e^+e^- 10.5 GeV |

¹⁹ This CRAWFORD 92 value includes protons from Λ decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(p \text{ anything (no } \Lambda))/\Gamma_{\text{total}}$

Γ_{71}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|-------------|-------------------|
| 0.12±0.10±0.16 | CRAWFORD 92 | CLEO | e^+e^- 10.5 GeV |

$\Gamma(n \text{ anything})/\Gamma_{\text{total}}$

Γ_{73}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|---------------------------|-------------|-------------------|
| 0.50±0.08±0.14 | ²⁰ CRAWFORD 92 | CLEO | e^+e^- 10.5 GeV |

²⁰ This CRAWFORD 92 value includes neutrons from Λ decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(n \text{ anything (no } \Lambda))/\Gamma_{\text{total}}$

Γ_{74}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|-------------|-------------------|
| 0.29±0.09±0.15 | CRAWFORD 92 | CLEO | e^+e^- 10.5 GeV |


$\Gamma(\rho \text{ hadrons})/\Gamma_{\text{total}}$ **Γ_{72}/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|--------------------|-------------|------------------------|
| 0.41 ± 0.24 | ADAMOVICH 87 | EMUL | γA 20–70 GeV/c |

$\Gamma(\Lambda \text{ anything})/\Gamma_{\text{total}}$ **Γ_{75}/Γ**

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------------------------|-----------------|--------------------|-------------|-------------------------------------------------------------|
| 0.35 ± 0.11 OUR AVERAGE | | | | Error includes scale factor of 1.4. See the ideogram below. |
| $0.59 \pm 0.10 \pm 0.12$ | | CRAWFORD 92 | CLEO | $e^+ e^-$ 10.5 GeV |
| 0.49 ± 0.24 | | ADAMOVICH 87 | EMUL | γA 20–70 GeV/c |
| 0.23 ± 0.10 | 8 ²¹ | ABE 86 | HYBR | 20 GeV γp |

²¹ ABE 86 includes Λ 's from Σ^0 decay.



$\Gamma(\Sigma^\pm \text{ anything})/\Gamma_{\text{total}}$ **Γ_{76}/Γ**

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|-------------|--------------------|-------------|-------------------|
| 0.1 ± 0.05 | 5 | ABE 86 | HYBR | 20 GeV γp |

$\Gamma(3\text{prongs})/\Gamma_{\text{total}}$ **Γ_{77}/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------------------|--------------------|-------------|--------------------------------------|
| $0.24 \pm 0.07 \pm 0.04$ | KAYIS-TOPAK.03 | CHRS | ν_μ emulsion, $\bar{E}=27$ GeV |

————— Rare or forbidden modes —————

$\Gamma(p e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{78}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|---------------|--------------------|-------------|------------------------------|
| $<5.5 \times 10^{-6}$ | 90 | 4.0 ± 7.1 | LEES | 11G BABR | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(p \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{79}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|----------------|--------------------|-------------|------------------------------|
| $<44 \times 10^{-6}$ | 90 | 11.1 ± 5.6 | LEES | 11G BABR | $e^+ e^- \approx \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------------|----|---|--------|---------|--------------------------|
| $<3.4 \times 10^{-4}$ | 90 | 0 | KODAMA | 95 E653 | π^- emulsion 600 GeV |
|-----------------------|----|---|--------|---------|--------------------------|

$\Gamma(p e^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{80}/Γ**

A test of lepton family-number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|----------------|--------------------|-------------|------------------------------|
| $<9.9 \times 10^{-6}$ | 90 | -0.7 ± 3.0 | LEES | 11G BABR | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(p e^- \mu^+)/\Gamma_{\text{total}}$ **Γ_{81}/Γ**

A test of lepton family-number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|---------------|--------------------|-------------|------------------------------|
| $<19 \times 10^{-6}$ | 90 | 6.2 ± 4.9 | LEES | 11G BABR | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\bar{p} 2e^+)/\Gamma_{\text{total}}$ **Γ_{82}/Γ**

A test of lepton- and baryon-number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|----------------|--------------------|-------------|------------------------------|
| $<2.7 \times 10^{-6}$ | 90 | -1.5 ± 4.5 | LEES | 11G BABR | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\bar{p} 2\mu^+)/\Gamma_{\text{total}}$ **Γ_{83}/Γ**

A test of lepton- and baryon-number conservation and of lepton family-number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|---------------|--------------------|-------------|------------------------------|
| $<9.4 \times 10^{-6}$ | 90 | 0.0 ± 2.2 | LEES | 11G BABR | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\bar{p} e^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{84}/Γ**

A test of lepton- and baryon-number conservation and of lepton family-number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|----------------|--------------------|-------------|------------------------------|
| $<16 \times 10^{-6}$ | 90 | 10.1 ± 6.8 | LEES | 11G BABR | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Sigma^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{85}/Γ**

A test of lepton-number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|-------------|--------------------|-------------|--------------------------|
| $<7.0 \times 10^{-4}$ | 90 | 0 | KODAMA | 95 E653 | π^- emulsion 600 GeV |

Λ_c^+ DECAY PARAMETERS

See the note on “Baryon Decay Parameters” in the neutron Listings.

α FOR $\Lambda_c^+ \rightarrow \Lambda\pi^+$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------------------------|------|----------------------|------|---------------------------------------------------|
| -0.91 ± 0.15 OUR AVERAGE | | | | |
| $-0.78 \pm 0.16 \pm 0.19$ | | LINK | 06A | FOCS γ A, $\bar{E}_\gamma \approx 180$ GeV |
| $-0.94 \pm 0.21 \pm 0.12$ | 414 | ²² BISHAI | 95 | CLE2 $e^+e^- \approx \mathcal{T}(4S)$ |
| -0.96 ± 0.42 | | ALBRECHT | 92 | ARG $e^+e^- \approx 10.4$ GeV |
| -1.1 ± 0.4 | 86 | AVERY | 90B | CLEO $e^+e^- \approx 10.6$ GeV |

²² BISHAI 95 actually gives $\alpha = -0.94 \pm 0.21 \pm 0.12$, chopping the errors at the physical limit -1.0 . However, for $\alpha \approx -1.0$, some experiments should get unphysical values ($\alpha < -1.0$), and for averaging with other measurements such values (or errors that extend below -1.0) should *not* be chopped.

α FOR $\Lambda_c^+ \rightarrow \Sigma^+\pi^0$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------------------|------|-------------|------|---------------------------------------|
| $-0.45 \pm 0.31 \pm 0.06$ | 89 | BISHAI | 95 | CLE2 $e^+e^- \approx \mathcal{T}(4S)$ |

α FOR $\Lambda_c^+ \rightarrow \Lambda\ell^+\nu_\ell$

The experiments don't cover the complete (or same incomplete) $M(\Lambda\ell^+)$ range, but we average them together anyway.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------------------------|------|------------------------|------|---------------------------------------|
| -0.86 ± 0.04 OUR AVERAGE | | | | |
| $-0.86 \pm 0.03 \pm 0.02$ | 3201 | ²³ HINSON | 05 | CLEO $e^+e^- \approx \mathcal{T}(4S)$ |
| $-0.91 \pm 0.42 \pm 0.25$ | | ²⁴ ALBRECHT | 94B | ARG $e^+e^- \approx 10$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------------------------------------------|-----|------------------------|----|----------------------|
| $-0.82 \pm 0.09 \pm 0.06$ $-0.06 - 0.03$ | 700 | ²⁵ CRAWFORD | 95 | CLE2 See HINSON 05 |
| $-0.89 \pm 0.17 \pm 0.09$ $-0.11 - 0.05$ | 350 | ²⁶ BERGFELD | 94 | CLE2 See CRAWFORD 95 |

²³ HINSON 05 measures the form-factor ratio $R \equiv f_2/f_1$ for $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$ events to be $-0.31 \pm 0.05 \pm 0.04$ and the pole mass to be $2.21 \pm 0.08 \pm 0.14$ GeV/ c^2 , and from these calculates α , averaged over q^2 , where $\langle q^2 \rangle = 0.67$ (GeV/ c)².

²⁴ ALBRECHT 94B uses Λe^+ and $\Lambda \mu^+$ events in the mass range $1.85 < M(\Lambda\ell^+) < 2.20$ GeV.

²⁵ CRAWFORD 95 measures the form-factor ratio $R \equiv f_2/f_1$ for $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$ events to be $-0.25 \pm 0.14 \pm 0.08$ and from this calculates α , averaged over q^2 , to be the above.

²⁶ BERGFELD 94 uses Λe^+ events.

$\Lambda_c^+, \bar{\Lambda}_c^-$ CP-VIOLATING DECAY ASYMMETRIES

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda\pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}\pi^-$

This is zero if CP is conserved.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------------------|-------------|------|---------------------------------------------------|
| $-0.07 \pm 0.19 \pm 0.24$ | LINK | 06A | FOCS γ A, $\bar{E}_\gamma \approx 180$ GeV |

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} e^- \bar{\nu}_e$

This is zero if *CP* is conserved.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|------|-------------------------------------|
| 0.00 ± 0.03 ± 0.02 | HINSON | 05 | CLEO $e^+ e^- \approx \Upsilon(4S)$ |

Λ_c^+ REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1992 edition (Physical Review **D45**, 1 June, Part II) or in earlier editions.

| | | | | |
|----------------|-----|------------------------------|-----------------------------------|-----------------------------|
| LEES | 11G | PR D84 072006 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| VAZQUEZ-JA... | 08 | PL B666 299 | E. Vazquez-Jauregui <i>et al.</i> | (SELEX Collab.) |
| AUBERT | 07U | PR D75 052002 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| LINK | 06A | PL B634 165 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| AUBERT,B | 05S | PR D72 052006 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| HINSON | 05 | PRL 94 191801 | J.W. Hinson <i>et al.</i> | (CLEO Collab.) |
| LINK | 05F | PL B624 22 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| LINK | 05K | PL B624 166 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| CRONIN-HEN... | 03 | PR D67 012001 | D. Cronin-Hennessy <i>et al.</i> | (CLEO Collab.) |
| KAYIS-TOPAK... | 03 | PL B555 156 | A. Kayis-Topaksu <i>et al.</i> | (CERN CHORUS Collab.) |
| ABE | 02C | PL B524 33 | K. Abe <i>et al.</i> | (KEK BELLE Collab.) |
| LINK | 02C | PRL 88 161801 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| LINK | 02G | PL B540 25 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| PDG | 02 | PR D66 010001 | K. Hagiwara <i>et al.</i> | |
| KUSHNIR... | 01 | PRL 86 5243 | A. Kushnirenko <i>et al.</i> | (FNAL SELEX Collab.) |
| MAHMOOD | 01 | PRL 86 2232 | A.H. Mahmood <i>et al.</i> | (CLEO Collab.) |
| AITALA | 00 | PL B471 449 | E.M. Aitala <i>et al.</i> | (FNAL E791 Collab.) |
| JAFFE | 00 | PR D62 072005 | D.E. Jaffe <i>et al.</i> | (CLEO Collab.) |
| ALAM | 98 | PR D57 4467 | M.S. Alam <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 96E | PRPL 276 223 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALEEV | 96 | JINRRC 3-77 31 | A.N. Alev <i>et al.</i> | (Serpukhov EXCHARM Collab.) |
| ALEXANDER | 96C | PR D53 R1013 | J.P. Alexander <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 95B | PL B342 397 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| AMMAR | 95 | PRL 74 3534 | R. Ammar <i>et al.</i> | (CLEO Collab.) |
| BISHAI | 95 | PL B350 256 | M. Bishai <i>et al.</i> | (CLEO Collab.) |
| CRAWFORD | 95 | PRL 75 624 | G. Crawford <i>et al.</i> | (CLEO Collab.) |
| KODAMA | 95 | PL B345 85 | K. Kodama <i>et al.</i> | (FNAL E653 Collab.) |
| ALBRECHT | 94B | PL B326 320 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALEEV | 94 | PAN 57 1370 | A.N. Alev <i>et al.</i> | (Serpukhov BIS-2 Collab.) |
| | | Translated from YF 57 1443. | | |
| AVERY | 94 | PL B325 257 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| BERGFELD | 94 | PL B323 219 | T. Bergfeld <i>et al.</i> | (CLEO Collab.) |
| FRABETTI | 94E | PL B328 193 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| AVERY | 93 | PRL 71 2391 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| BOZEK | 93 | PL B312 247 | A. Bozek <i>et al.</i> | (CERN NA32 Collab.) |
| FRABETTI | 93D | PRL 70 1755 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| FRABETTI | 93H | PL B314 477 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| KUBOTA | 93 | PRL 71 3255 | Y. Kubota <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 92 | PL B274 239 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 92O | ZPHY C56 1 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BARLAG | 92 | PL B283 465 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| CRAWFORD | 92 | PR D45 752 | G. Crawford <i>et al.</i> | (CLEO Collab.) |
| JEZABEK | 92 | PL B286 175 | M. Jezabek, K. Rybicki, R. Rylko | (CRAC) |
| ALBRECHT | 91G | PL B269 234 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| AVERY | 91 | PR D43 3599 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| ALVAREZ | 90 | ZPHY C47 539 | M.P. Alvarez <i>et al.</i> | (CERN NA14/2 Collab.) |
| ALVAREZ | 90B | PL B246 256 | M.P. Alvarez <i>et al.</i> | (CERN NA14/2 Collab.) |
| ANJOS | 90 | PR D41 801 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| AVERY | 90B | PRL 65 2842 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| BARLAG | 90D | ZPHY C48 29 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| FRABETTI | 90 | PL B251 639 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| BARLAG | 89 | PL B218 374 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| AGUILAR... | 88B | ZPHY C40 321 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| Also | | PL B189 254 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| Also | | PL B199 462 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| Also | | SJNP 48 833 | M. Begalli <i>et al.</i> | (LEBC-EHS Collab.) |
| | | Translated from YAF 48 1310. | | |

| | | | | |
|-----------|-----|-----------------------------|------------------------------|---------------------------|
| ALBRECHT | 88C | PL B207 109 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ANJOS | 88B | PRL 60 1379 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| ADAMOVICH | 87 | EPL 4 887 | M.I. Adamovich <i>et al.</i> | (Photon Emulsion Collab.) |
| Also | | SJNP 46 447 | F. Viaggi <i>et al.</i> | (Photon Emulsion Collab.) |
| | | Translated from YAF 46 799. | | |
| AMENDOLIA | 87 | ZPHY C36 513 | S.R. Amendolia <i>et al.</i> | (CERN NA1 Collab.) |
| JONES | 87 | ZPHY C36 593 | G.T. Jones <i>et al.</i> | (CERN WA21 Collab.) |
| ABE | 86 | PR D33 1 | K. Abe <i>et al.</i> | |
| ALEEV | 84 | ZPHY C23 333 | A.N. Aleev <i>et al.</i> | (BIS-2 Collab.) |
| BOSETTI | 82 | PL 109B 234 | P.C. Bosetti <i>et al.</i> | (AACH3, BONN, CERN+) |
| VELLA | 82 | PRL 48 1515 | E. Vella <i>et al.</i> | (SLAC, LBL, UCB) |
| BASILE | 81B | NC 62A 14 | M. Basile <i>et al.</i> | (CERN, BGNA, PGIA, FRAS) |
| CALICCHIO | 80 | PL 93B 521 | M. Calicchio <i>et al.</i> | (BARI, BIRM, BRUX+) |

OTHER RELATED PAPERS

| | | | |
|-----------|----|---------------|---------------------------|
| MIGLIOZZI | 99 | PL B462 217 | P. Migliozi <i>et al.</i> |
| DUNIETZ | 98 | PR D58 094010 | I. Dunietz |
