



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ****$$

p MASS (atomic mass units u)

The mass is known much more precisely in u (atomic mass units) than in MeV. See the next data block.

VALUE (u)	DOCUMENT ID	TECN	COMMENT
1.007276466812 ± 0.000000000090	MOHR	12	RVUE 2010 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.00727646677 ± 0.00000000010	MOHR	08	RVUE 2006 CODATA value
1.00727646688 ± 0.00000000013	MOHR	05	RVUE 2002 CODATA value
1.00727646688 ± 0.00000000013	MOHR	99	RVUE 1998 CODATA value
1.007276470 ± 0.0000000012	COHEN	87	RVUE 1986 CODATA value

p MASS (MeV)

The mass is known much more precisely in u (atomic mass units) than in MeV. The conversion from u to MeV, $1 u = 931.494 061(21) \text{ MeV}/c^2$ (MOHR 12, the 2010 CODATA value), involves the relatively poorly known electronic charge.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
938.272046 ± 0.000021	MOHR	12	RVUE 2010 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
938.272013 ± 0.000023	MOHR	08	RVUE 2006 CODATA value
938.272029 ± 0.000080	MOHR	05	RVUE 2002 CODATA value
938.271998 ± 0.000038	MOHR	99	RVUE 1998 CODATA value
938.27231 ± 0.00028	COHEN	87	RVUE 1986 CODATA value
938.2796 ± 0.0027	COHEN	73	RVUE 1973 CODATA value

$$|m_p - m_{\bar{p}}|/m_p$$

A test of CPT invariance. Note that the comparison of the \bar{p} and p charge-to-mass ratio, given in the next data block, is much better determined.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2 × 10⁻⁹	90	¹ HORI	06	SPEC $\bar{p}e^- \text{He atom}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 1.0 × 10 ⁻⁸	90	¹ HORI	03	SPEC $\bar{p}e^- \text{}^4\text{He}, \bar{p}e^- \text{}^3\text{He}$
< 6 × 10 ⁻⁸	90	¹ HORI	01	SPEC $\bar{p}e^- \text{He atom}$
< 5 × 10 ⁻⁷		² TORII	99	SPEC $\bar{p}e^- \text{He atom}$

¹ HORI 01, HORI 03, and HORI 06 use the more-precisely-known constraint on the \bar{p} charge-to-mass ratio of GABRIELSE 99 (see below) to get their results. Their results are not independent of the HORI 01, HORI 03, and HORI 06 values for $|q_p + q_{\bar{p}}|/e$, below.

² TORII 99 uses the more-precisely-known constraint on the \bar{p} charge-to-mass ratio of GABRIELSE 95 (see below) to get this result. This is not independent of the TORII 99 value for $|q_p + q_{\bar{p}}|/e$, below.

\bar{p}/p CHARGE-TO-MASS RATIO, $|\frac{q_{\bar{p}}}{m_{\bar{p}}}|/(\frac{q_p}{m_p})$

A test of *CPT* invariance. Listed here are measurements involving the *inertial* masses. For a discussion of what may be inferred about the ratio of \bar{p} and p *gravitational* masses, see ERICSON 90; they obtain an upper bound of 10^{-6} – 10^{-7} for violation of the equivalence principle for \bar{p} 's.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.99999999991 ± 0.00000000009	GABRIELSE	99	TRAP Penning trap
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.0000000015 ± 0.0000000011	³ GABRIELSE	95	TRAP Penning trap
1.000000023 ± 0.000000042	⁴ GABRIELSE	90	TRAP Penning trap
³ Equation (2) of GABRIELSE 95 should read $M(\bar{p})/M(p) = 0.999\,999\,9985$ (11) (G. Gabrielse, private communication).			
⁴ GABRIELSE 90 also measures $m_{\bar{p}}/m_{e^-} = 1836.152660 \pm 0.000083$ and $m_p/m_{e^-} = 1836.152680 \pm 0.000088$. Both are completely consistent with the 1986 CODATA (COHEN 87) value for m_p/m_{e^-} of 1836.152701 ± 0.000037 .			

$$\left(\left|\frac{q_{\bar{p}}}{m_{\bar{p}}}\right| - \frac{q_p}{m_p}\right) / \frac{q_p}{m_p}$$

A test of *CPT* invariance. Taken from the \bar{p}/p charge-to-mass ratio, above.

<u>VALUE</u>	<u>DOCUMENT ID</u>
$(-9 \pm 9) \times 10^{-11}$ OUR EVALUATION	

$$|q_p + q_{\bar{p}}|/e$$

A test of *CPT* invariance. Note that the comparison of the \bar{p} and p charge-to-mass ratios given above is much better determined. See also a similar test involving the electron.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2 \times 10^{-9}$	90	⁵ HORI	06	SPEC $\bar{p}e^-$ He atom
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 1.0 \times 10^{-8}$	90	⁵ HORI	03	SPEC $\bar{p}e^-$ ⁴ He, $\bar{p}e^-$ ³ He
$< 6 \times 10^{-8}$	90	⁵ HORI	01	SPEC $\bar{p}e^-$ He atom
$< 5 \times 10^{-7}$		⁶ TORII	99	SPEC $\bar{p}e^-$ He atom
$< 2 \times 10^{-5}$		⁷ HUGHES	92	RVUE

⁵ HORI 01, HORI 03, and HORI 06 use the more-precisely-known constraint on the \bar{p} charge-to-mass ratio of GABRIELSE 99 (see above) to get their results. Their results are not independent of the HORI 01, HORI 03, and HORI 06 values for $|m_p - m_{\bar{p}}|/m_p$, above.

⁶ TORII 99 uses the more-precisely-known constraint on the \bar{p} charge-to-mass ratio of GABRIELSE 95 (see above) to get this result. This is not independent of the TORII 99 value for $|m_p - m_{\bar{p}}|/m_p$, above.

⁷ HUGHES 92 uses recent measurements of Rydberg-energy and cyclotron-frequency ratios.

$$|q_p + q_e|/e$$

See BRESSI 11 for a summary of experiments on the neutrality of matter.
See also “*n* CHARGE” in the neutron Listings.

VALUE	DOCUMENT ID	TECN	COMMENT
<1 × 10⁻²¹	⁸ BRESSI	11	Neutrality of SF ₆
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<3.2 × 10 ⁻²⁰	⁹ SENGUPTA	00	binary pulsar
<0.8 × 10 ⁻²¹	MARINELLI	84	Magnetic levitation
<1.0 × 10 ⁻²¹	⁸ DYLLA	73	Neutrality of SF ₆
⁸ BRESSI 11 uses the method of DYLLA 73 but finds serious errors in that experiment that greatly reduce its accuracy. The BRESSI 11 limit assumes that $n \rightarrow p e^- \nu_e$ conserves charge. Thus the limit applies equally to the charge of the neutron.			
⁹ SENGUPTA 00 uses the difference between the observed rate of rotational energy loss by the binary pulsar PSR B1913+16 and the rate predicted by general relativity to set this limit. See the paper for assumptions.			

ρ MAGNETIC MOMENT

See the “Note on Baryon Magnetic Moments” in the Λ Listings.

VALUE (μ_N)	DOCUMENT ID	TECN	COMMENT
2.792847356 ± 0.000000023	MOHR	12	RVUE 2010 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.792847356 ± 0.000000023	MOHR	08	RVUE 2006 CODATA value
2.792847351 ± 0.000000028	MOHR	05	RVUE 2002 CODATA value
2.792847337 ± 0.000000029	MOHR	99	RVUE 1998 CODATA value
2.792847386 ± 0.000000063	COHEN	87	RVUE 1986 CODATA value
2.7928456 ± 0.0000011	COHEN	73	RVUE 1973 CODATA value

\bar{p} MAGNETIC MOMENT

A few early results have been omitted.

VALUE (μ_N)	DOCUMENT ID	TECN	COMMENT
−2.792845 ± 0.000012	DISCIACCA	13	TRAP Single \bar{p} , Penning trap
• • • We do not use the following data for averages, fits, limits, etc. • • •			
−2.7862 ± 0.0083	PASK	09	CNTR \bar{p} He ⁺ hyperfine structure
−2.8005 ± 0.0090	KREISSL	88	CNTR \bar{p} ²⁰⁸ Pb 11→10 X-ray
−2.817 ± 0.048	ROBERTS	78	CNTR
−2.791 ± 0.021	HU	75	CNTR Exotic atoms

$$(\mu_p + \mu_{\bar{p}}) / \mu_p$$

A test of *CPT* invariance.

VALUE (units 10 ⁻⁶)	DOCUMENT ID	TECN	COMMENT
0 ± 5	DISCIACCA	13	TRAP Single \bar{p} , Penning trap

p ELECTRIC DIPOLE MOMENT

A nonzero value is forbidden by both T invariance and P invariance.

<u>VALUE (10^{-23} ecm)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 0.54		¹⁰ DMITRIEV 03		Uses ¹⁹⁹ Hg atom EDM
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
– 3.7 ± 6.3		CHO 89	NMR	TI F molecules
< 400		DZUBA 85	THEO	Uses ¹²⁹ Xe moment
130 ± 200		¹¹ WILKENING 84		
900 ± 1400		¹² WILKENING 84		
700 ± 900	1G	HARRISON 69	MBR	Molecular beam
¹⁰ DMITRIEV 03 calculates this limit from the limit on the electric dipole moment of the ¹⁹⁹ Hg atom.				
¹¹ This WILKENING 84 value includes a finite-size effect and a magnetic effect.				
¹² This WILKENING 84 value is more cautious than the other and excludes the finite-size effect, which relies on uncertain nuclear integrals.				

 p ELECTRIC POLARIZABILITY α_p

For a very complete review of the “polarizability of the nucleon and Compton scattering,” see SCHUMACHER 05. His recommended values for the proton are $\alpha_p = (12.0 \pm 0.6) \times 10^{-4} \text{ fm}^3$ and $\beta_p = (1.9 \mp 0.6) \times 10^{-4} \text{ fm}^3$, almost exactly our averages.

<u>VALUE (10^{-4} fm^3)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.2 ± 0.4 OUR AVERAGE			
10.65 ± 0.35 ± 0.36	MCGOVERN 13	RVUE	χ EFT + Compton scattering
12.1 ± 1.1 ± 0.5	¹³ BEANE 03		EFT + γp
11.82 ± 0.98 ^{+0.52} _{−0.98}	¹⁴ BLANPIED 01	LEGS	$p(\vec{\gamma}, \gamma)$, $p(\vec{\gamma}, \pi^0)$, $p(\vec{\gamma}, \pi^+)$
11.9 ± 0.5 ± 1.3	¹⁵ OLMOSDEL... 01	CNTR	γp Compton scattering
12.1 ± 0.8 ± 0.5	¹⁶ MACGIBBON 95	RVUE	global average
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
11.7 ± 0.8 ± 0.7	¹⁷ BARANOV 01	RVUE	Global average
12.5 ± 0.6 ± 0.9	MACGIBBON 95	CNTR	γp Compton scattering
9.8 ± 0.4 ± 1.1	HALLIN 93	CNTR	γp Compton scattering
10.62 ^{+1.25+1.07} _{−1.19−1.03}	ZIEGER 92	CNTR	γp Compton scattering
10.9 ± 2.2 ± 1.3	¹⁸ FEDERSPIEL 91	CNTR	γp Compton scattering
¹³ BEANE 03 uses effective field theory and low-energy γp and γd Compton-scattering data. It also gets for the isoscalar polarizabilities (see the erratum) $\alpha_N = (13.0 \pm 1.9+3.9−1.5) \times 10^{-4} \text{ fm}^3$ and $\beta_N = (-1.8 \pm 1.9+2.1−0.9) \times 10^{-4} \text{ fm}^3$.			
¹⁴ BLANPIED 01 gives $\alpha_p + \beta_p$ and $\alpha_p - \beta_p$. The separate α_p and β_p are provided to us by A. Sandorfi. The first error above is statistics plus systematics; the second is from the model.			
¹⁵ This OLMOSDELEON 01 result uses the TAPS data alone, and does not use the (re-evaluated) sum-rule constraint that $\alpha + \beta = (13.8 \pm 0.4) \times 10^{-4} \text{ fm}^3$. See the paper for a discussion.			

- ¹⁶ MACGIBBON 95 combine the results of ZIEGER 92, FEDERSPIEL 91, and their own experiment to get a “global average” in which model errors and systematic errors are treated in a consistent way. See MACGIBBON 95 for a discussion.
- ¹⁷ BARANOV 01 combines the results of 10 experiments from 1958 through 1995 to get a global average that takes into account both systematic and model errors and does not use the theoretical constraint on the sum $\alpha_p + \beta_p$.
- ¹⁸ FEDERSPIEL 91 obtains for the (static) electric polarizability α_p , defined in terms of the induced electric dipole moment by $\mathbf{D} = 4\pi\epsilon_0\alpha_p\mathbf{E}$, the value $(7.0 \pm 2.2 \pm 1.3) \times 10^{-4} \text{ fm}^3$.

p MAGNETIC POLARIZABILITY β_p

The electric and magnetic polarizabilities are subject to a dispersion sum-rule constraint $\bar{\alpha} + \bar{\beta} = (14.2 \pm 0.5) \times 10^{-4} \text{ fm}^3$. Errors here are anticorrelated with those on $\bar{\alpha}_p$ due to this constraint.

VALUE (10^{-4} fm^3)	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.4	OUR AVERAGE		Error includes scale factor of 1.2.
3.15 ± 0.35 ± 0.36	MCGOVERN	13	RVUE χ EFT + Compton scattering
3.4 ± 1.1 ± 0.1	¹⁹ BEANE	03	EFT + γp
1.43 ± 0.98 ^{+0.52} / _{-0.98}	²⁰ BLANPIED	01	LEGS $p(\vec{\gamma}, \gamma)$, $p(\vec{\gamma}, \pi^0)$, $p(\vec{\gamma}, \pi^+)$
1.2 ± 0.7 ± 0.5	²¹ OLMOSDEL...	01	CNTR γp Compton scattering
2.1 ± 0.8 ± 0.5	²² MACGIBBON	95	RVUE global average
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2.3 ± 0.9 ± 0.7	²³ BARANOV	01	RVUE Global average
1.7 ± 0.6 ± 0.9	MACGIBBON	95	CNTR γp Compton scattering
4.4 ± 0.4 ± 1.1	HALLIN	93	CNTR γp Compton scattering
3.58 ^{+1.19 +1.03} / _{-1.25 -1.07}	ZIEGER	92	CNTR γp Compton scattering
3.3 ± 2.2 ± 1.3	FEDERSPIEL	91	CNTR γp Compton scattering

- ¹⁹ BEANE 03 uses effective field theory and low-energy γp and γd Compton-scattering data. It also gets for the isoscalar polarizabilities (see the erratum) $\alpha_N = (13.0 \pm 1.9 ^{+3.9}/_{-1.5}) \times 10^{-4} \text{ fm}^3$ and $\beta_N = (-1.8 \pm 1.9 ^{+2.1}/_{-0.9}) \times 10^{-4} \text{ fm}^3$.
- ²⁰ BLANPIED 01 gives $\alpha_p + \beta_p$ and $\alpha_p - \beta_p$. The separate α_p and β_p are provided to us by A. Sandorfi. The first error above is statistics plus systematics; the second is from the model.
- ²¹ This OLMOSDELEON 01 result uses the TAPS data alone, and does not use the (re-evaluated) sum-rule constraint that $\alpha + \beta = (13.8 \pm 0.4) \times 10^{-4} \text{ fm}^3$. See the paper for a discussion.
- ²² MACGIBBON 95 combine the results of ZIEGER 92, FEDERSPIEL 91, and their own experiment to get a “global average” in which model errors and systematic errors are treated in a consistent way. See MACGIBBON 95 for a discussion.
- ²³ BARANOV 01 combines the results of 10 experiments from 1958 through 1995 to get a global average that takes into account both systematic and model errors and does not use the theoretical constraint on the sum $\alpha_p + \beta_p$.

p CHARGE RADIUS

This is the rms electric charge radius, $\sqrt{\langle r_E^2 \rangle}$.

Most measurements of the radius of the proton involve electron-proton interactions, and most of the more recent values agree with one another. The most precise of these is $r_p = 0.879(8)$ fm (BERNAUER 10). The CODATA 10 value (MOHR 12), obtained from the electronic results, is 0.8775(51). However, a measurement using muonic hydrogen finds $r_p = 0.84087(39)$ fm (ANTOIGNINI 13), which is 13 times more precise and seven standard deviations (using the CODATA 10 error) from the electronic results.

Since POHL 10 (the first μp result), there has been a lot of discussion about the disagreement, especially concerning the modeling of muonic hydrogen. Here is an incomplete list of papers: DERUJULA 10, CLOET 11, DISTLER 11, DERUJULA 11, ARRINGTON 11, BERNAUER 11, and HILL 11.

Until the difference between the ep and μp values is understood, it does not make sense to average the values together. For the present, we give both values. It is up to workers in this field to solve this puzzle.

<u>VALUE (fm)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.84087 ± 0.00026 ± 0.00029	ANTOIGNINI 13	LASR	μp -atom Lamb shift
0.8775 ± 0.0051	MOHR 12	RVUE	2010 CODATA, ep data
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.879 ± 0.005 ± 0.006	BERNAUER 10	SPEC	$ep \rightarrow ep$ form factor
0.912 ± 0.009 ± 0.007	BORISYUK 10		reanalyzes old ep data
0.871 ± 0.009 ± 0.003	HILL 10		z-expansion reanalysis
0.84184 ± 0.00036 ± 0.00056	POHL 10	LASR	See ANTOIGNINI 13
0.8768 ± 0.0069	MOHR 08	RVUE	2006 CODATA value
0.844 +0.008 -0.004	BELUSHKIN 07		Dispersion analysis
0.897 ± 0.018	BLUNDEN 05		SICK 03 + 2γ correction
0.8750 ± 0.0068	MOHR 05	RVUE	2002 CODATA value
0.895 ± 0.010 ± 0.013	SICK 03		$ep \rightarrow ep$ reanalysis
0.830 ± 0.040 ± 0.040	²⁴ ESCHRICH 01		$ep \rightarrow ep$
0.883 ± 0.014	MELNIKOV 00		1S Lamb Shift in H
0.880 ± 0.015	ROSENFELDR.00		ep + Coul. corrections
0.847 ± 0.008	MERGELL 96		ep + disp. relations
0.877 ± 0.024	WONG 94		reanalysis of Mainz ep data
0.865 ± 0.020	MCCORD 91		$ep \rightarrow ep$
0.862 ± 0.012	SIMON 80		$ep \rightarrow ep$
0.880 ± 0.030	BORKOWSKI 74		$ep \rightarrow ep$
0.810 ± 0.020	AKIMOV 72		$ep \rightarrow ep$
0.800 ± 0.025	FREREJACQ... 66		$ep \rightarrow ep$ (CH ₂ tgt.)
0.805 ± 0.011	HAND 63		$ep \rightarrow ep$

²⁴ESCHRICH 01 actually gives $\langle r^2 \rangle = (0.69 \pm 0.06 \pm 0.06)$ fm².

p MAGNETIC RADIUS

This is the rms magnetic radius, $\sqrt{\langle r_M^2 \rangle}$.

VALUE (fm)	DOCUMENT ID	TECN	COMMENT
0.777±0.013±0.010	BERNAUER	10 SPEC	$e p \rightarrow e p$ form factor
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.876±0.010±0.016	BORISYUK	10	reanalyzes old $e p \rightarrow e p$ data
0.854±0.005	BELUSHKIN	07	Dispersion analysis

p MEAN LIFE

A test of baryon conservation. See the “p Partial Mean Lives” section below for limits for identified final states. The limits here are to “anything” or are for “disappearance” modes of a bound proton (p) or (n). See also the 3ν modes in the “Partial Mean Lives” section. Table 1 of BACK 03 is a nice summary.

LIMIT (years)	PARTICLE	CL%	DOCUMENT ID	TECN	COMMENT
>5.8 × 10²⁹	n	90	25 ARAKI	06 KLND	$n \rightarrow$ invisible
>2.1 × 10²⁹	p	90	26 AHMED	04 SNO	$p \rightarrow$ invisible
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
>1.9 × 10 ²⁹	n	90	26 AHMED	04 SNO	$n \rightarrow$ invisible
>1.8 × 10 ²⁵	n	90	27 BACK	03 BORX	
>1.1 × 10 ²⁶	p	90	27 BACK	03 BORX	
>3.5 × 10 ²⁸	p	90	28 ZDESENKO	03	$p \rightarrow$ invisible
>1 × 10 ²⁸	p	90	29 AHMAD	02 SNO	$p \rightarrow$ invisible
>4 × 10 ²³	p	95	TRETYAK	01	$d \rightarrow n + ?$
>1.9 × 10 ²⁴	p	90	30 BERNABEI	00B DAMA	
>1.6 × 10 ²⁵	p, n		31,32 EVANS	77	
>3 × 10 ²³	p		32 DIX	70 CNTR	
>3 × 10 ²³	p, n		32,33 FLEROV	58	

²⁵ ARAKI 06 looks for signs of de-excitation of the residual nucleus after disappearance of a neutron from the s shell of ^{12}C .

²⁶ AHMED 04 looks for γ rays from the de-excitation of a residual $^{15}\text{O}^*$ or $^{15}\text{N}^*$ following the disappearance of a neutron or proton in ^{16}O .

²⁷ BACK 03 looks for decays of unstable nuclides left after N decays of parent ^{12}C , ^{13}C , ^{16}O nuclei. These are “invisible channel” limits.

²⁸ ZDESENKO 03 gets this limit on proton disappearance in deuterium by analyzing SNO data in AHMAD 02.

²⁹ AHMAD 02 (see its footnote 7) looks for neutrons left behind after the disappearance of the proton in deuterons.

³⁰ BERNABEI 00B looks for the decay of a $^{128}_{53}\text{I}$ nucleus following the disappearance of a proton in the otherwise-stable $^{129}_{54}\text{Xe}$ nucleus.

³¹ EVANS 77 looks for the daughter nuclide ^{129}Xe from possible ^{130}Te decays in ancient Te ore samples.

³² This mean-life limit has been obtained from a half-life limit by dividing the latter by $\ln(2) = 0.693$.

³³ FLEROV 58 looks for the spontaneous fission of a ^{232}Th nucleus after the disappearance of one of its nucleons.

\bar{p} MEAN LIFE

Of the two astrophysical limits here, that of GEER 00D involves considerably more refinements in its modeling. The other limits come from direct observations of stored antiprotons. See also “ \bar{p} Partial Mean Lives” after “ p Partial Mean Lives,” below, for exclusive-mode limits. The best (lifetime/branching fraction) limit there is 7×10^5 years, for $\bar{p} \rightarrow e^- \gamma$. We advance only the exclusive-mode limits to our Summary Tables.

<u>LIMIT</u> (years)	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$>8 \times 10^5$	90		³⁴ GEER	00D	\bar{p}/p ratio, cosmic rays
>0.28			GABRIELSE	90 TRAP	Penning trap
>0.08	90	1	BELL	79 CNTR	Storage ring
$>1 \times 10^7$			GOLDEN	79 SPEC	\bar{p}/p ratio, cosmic rays
$>3.7 \times 10^{-3}$			BREGMAN	78 CNTR	Storage ring
³⁴ GEER 00D uses agreement between a model of galactic \bar{p} production and propagation and the observed \bar{p}/p cosmic-ray spectrum to set this limit.					

p DECAY MODES

See the “Note on Nucleon Decay” in our 1994 edition (Phys. Rev. **D50**, 1173) for a short review.

The “partial mean life” limits tabulated here are the limits on τ/B_i , where τ is the total mean life and B_i is the branching fraction for the mode in question. For N decays, p and n indicate proton and neutron partial lifetimes.

Mode	Partial mean life (10^{30} years)	Confidence level
Antilepton + meson		
$\tau_1 \quad N \rightarrow e^+ \pi$	$> 2000 (n), > 8200 (p)$	90%
$\tau_2 \quad N \rightarrow \mu^+ \pi$	$> 1000 (n), > 6600 (p)$	90%
$\tau_3 \quad N \rightarrow \nu \pi$	$> 112 (n), > 16 (p)$	90%
$\tau_4 \quad p \rightarrow e^+ \eta$	> 4200	90%
$\tau_5 \quad p \rightarrow \mu^+ \eta$	> 1300	90%
$\tau_6 \quad n \rightarrow \nu \eta$	> 158	90%
$\tau_7 \quad N \rightarrow e^+ \rho$	$> 217 (n), > 710 (p)$	90%
$\tau_8 \quad N \rightarrow \mu^+ \rho$	$> 228 (n), > 160 (p)$	90%
$\tau_9 \quad N \rightarrow \nu \rho$	$> 19 (n), > 162 (p)$	90%
$\tau_{10} \quad p \rightarrow e^+ \omega$	> 320	90%
$\tau_{11} \quad p \rightarrow \mu^+ \omega$	> 780	90%
$\tau_{12} \quad n \rightarrow \nu \omega$	> 108	90%
$\tau_{13} \quad N \rightarrow e^+ K$	$> 17 (n), > 1000 (p)$	90%
$\tau_{14} \quad p \rightarrow e^+ K_S^0$		

τ_{15}	$p \rightarrow e^+ K_L^0$		
τ_{16}	$N \rightarrow \mu^+ K$	> 26 (<i>n</i>), > 1600 (<i>p</i>)	90%
τ_{17}	$p \rightarrow \mu^+ K_S^0$		
τ_{18}	$p \rightarrow \mu^+ K_L^0$		
τ_{19}	$N \rightarrow \nu K$	> 86 (<i>n</i>), > 2300 (<i>p</i>)	90%
τ_{20}	$n \rightarrow \nu K_S^0$	> 260	90%
τ_{21}	$p \rightarrow e^+ K^*(892)^0$	> 84	90%
τ_{22}	$N \rightarrow \nu K^*(892)$	> 78 (<i>n</i>), > 51 (<i>p</i>)	90%

Antilepton + mesons

τ_{23}	$p \rightarrow e^+ \pi^+ \pi^-$	> 82	90%
τ_{24}	$p \rightarrow e^+ \pi^0 \pi^0$	> 147	90%
τ_{25}	$n \rightarrow e^+ \pi^- \pi^0$	> 52	90%
τ_{26}	$p \rightarrow \mu^+ \pi^+ \pi^-$	> 133	90%
τ_{27}	$p \rightarrow \mu^+ \pi^0 \pi^0$	> 101	90%
τ_{28}	$n \rightarrow \mu^+ \pi^- \pi^0$	> 74	90%
τ_{29}	$n \rightarrow e^+ K^0 \pi^-$	> 18	90%

Lepton + meson

τ_{30}	$n \rightarrow e^- \pi^+$	> 65	90%
τ_{31}	$n \rightarrow \mu^- \pi^+$	> 49	90%
τ_{32}	$n \rightarrow e^- \rho^+$	> 62	90%
τ_{33}	$n \rightarrow \mu^- \rho^+$	> 7	90%
τ_{34}	$n \rightarrow e^- K^+$	> 32	90%
τ_{35}	$n \rightarrow \mu^- K^+$	> 57	90%

Lepton + mesons

τ_{36}	$p \rightarrow e^- \pi^+ \pi^+$	> 30	90%
τ_{37}	$n \rightarrow e^- \pi^+ \pi^0$	> 29	90%
τ_{38}	$p \rightarrow \mu^- \pi^+ \pi^+$	> 17	90%
τ_{39}	$n \rightarrow \mu^- \pi^+ \pi^0$	> 34	90%
τ_{40}	$p \rightarrow e^- \pi^+ K^+$	> 75	90%
τ_{41}	$p \rightarrow \mu^- \pi^+ K^+$	> 245	90%

Antilepton + photon(s)

τ_{42}	$p \rightarrow e^+ \gamma$	> 670	90%
τ_{43}	$p \rightarrow \mu^+ \gamma$	> 478	90%
τ_{44}	$n \rightarrow \nu \gamma$	> 28	90%
τ_{45}	$p \rightarrow e^+ \gamma \gamma$	> 100	90%
τ_{46}	$n \rightarrow \nu \gamma \gamma$	> 219	90%

Three (or more) leptons

τ_{47}	$p \rightarrow e^+ e^+ e^-$	> 793	90%
τ_{48}	$p \rightarrow e^+ \mu^+ \mu^-$	> 359	90%
τ_{49}	$p \rightarrow e^+ \nu \nu$	> 17	90%
τ_{50}	$n \rightarrow e^+ e^- \nu$	> 257	90%
τ_{51}	$n \rightarrow \mu^+ e^- \nu$	> 83	90%
τ_{52}	$n \rightarrow \mu^+ \mu^- \nu$	> 79	90%
τ_{53}	$p \rightarrow \mu^+ e^+ e^-$	> 529	90%
τ_{54}	$p \rightarrow \mu^+ \mu^+ \mu^-$	> 675	90%
τ_{55}	$p \rightarrow \mu^+ \nu \nu$	> 21	90%
τ_{56}	$p \rightarrow e^- \mu^+ \mu^+$	> 6	90%
τ_{57}	$n \rightarrow 3\nu$	> 0.0005	90%
τ_{58}	$n \rightarrow 5\nu$		

Inclusive modes

τ_{59}	$N \rightarrow e^+$ anything	> 0.6 (n, p)	90%
τ_{60}	$N \rightarrow \mu^+$ anything	> 12 (n, p)	90%
τ_{61}	$N \rightarrow \nu$ anything		
τ_{62}	$N \rightarrow e^+ \pi^0$ anything	> 0.6 (n, p)	90%
τ_{63}	$N \rightarrow 2$ bodies, ν -free		

$\Delta B = 2$ dinucleon modes

The following are lifetime limits per iron nucleus.

τ_{64}	$pp \rightarrow \pi^+ \pi^+$	> 0.7	90%
τ_{65}	$pn \rightarrow \pi^+ \pi^0$	> 2	90%
τ_{66}	$nn \rightarrow \pi^+ \pi^-$	> 0.7	90%
τ_{67}	$nn \rightarrow \pi^0 \pi^0$	> 3.4	90%
τ_{68}	$pp \rightarrow e^+ e^+$	> 5.8	90%
τ_{69}	$pp \rightarrow e^+ \mu^+$	> 3.6	90%
τ_{70}	$pp \rightarrow \mu^+ \mu^+$	> 1.7	90%
τ_{71}	$pn \rightarrow e^+ \bar{\nu}$	> 2.8	90%
τ_{72}	$pn \rightarrow \mu^+ \bar{\nu}$	> 1.6	90%
τ_{73}	$nn \rightarrow \nu_e \bar{\nu}_e$	> 1.4	90%
τ_{74}	$nn \rightarrow \nu_\mu \bar{\nu}_\mu$	> 1.4	90%
τ_{75}	$pn \rightarrow$ invisible	> 0.000021	90%
τ_{76}	$pp \rightarrow$ invisible	> 0.00005	90%

\bar{p} DECAY MODES

	Mode	Partial mean life (years)	Confidence level
τ_{77}	$\bar{p} \rightarrow e^- \gamma$	> 7×10^5	90%
τ_{78}	$\bar{p} \rightarrow \mu^- \gamma$	> 5×10^4	90%
τ_{79}	$\bar{p} \rightarrow e^- \pi^0$	> 4×10^5	90%
τ_{80}	$\bar{p} \rightarrow \mu^- \pi^0$	> 5×10^4	90%

τ_{81}	$\bar{p} \rightarrow e^- \eta$	$> 2 \times 10^4$	90%
τ_{82}	$\bar{p} \rightarrow \mu^- \eta$	$> 8 \times 10^3$	90%
τ_{83}	$\bar{p} \rightarrow e^- K_S^0$	> 900	90%
τ_{84}	$\bar{p} \rightarrow \mu^- K_S^0$	$> 4 \times 10^3$	90%
τ_{85}	$\bar{p} \rightarrow e^- K_L^0$	$> 9 \times 10^3$	90%
τ_{86}	$\bar{p} \rightarrow \mu^- K_L^0$	$> 7 \times 10^3$	90%
τ_{87}	$\bar{p} \rightarrow e^- \gamma \gamma$	$> 2 \times 10^4$	90%
τ_{88}	$\bar{p} \rightarrow \mu^- \gamma \gamma$	$> 2 \times 10^4$	90%
τ_{89}	$\bar{p} \rightarrow e^- \rho$		
τ_{90}	$\bar{p} \rightarrow e^- \omega$	> 200	90%
τ_{91}	$\bar{p} \rightarrow e^- K^*(892)^0$		

p PARTIAL MEAN LIVES

The “partial mean life” limits tabulated here are the limits on τ/B_i , where τ is the total mean life for the proton and B_i is the branching fraction for the mode in question.

Decaying particle: p = proton, n = bound neutron. The same event may appear under more than one partial decay mode. Background estimates may be accurate to a factor of two.

Antilepton + meson

$\tau(N \rightarrow e^+ \pi)$

τ_1

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>2000	n	90	0	0.27	NISHINO 12	SKAM
>8200	p	90	0	0.3	NISHINO 09	SKAM
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
> 540	p	90	0	0.2	MCGREW 99	IMB3
> 158	n	90	3	5	MCGREW 99	IMB3
>1600	p	90	0	0.1	SHIOZAWA 98	SKAM
> 70	p	90	0	0.5	BERGER 91	FREJ
> 70	n	90	0	≤ 0.1	BERGER 91	FREJ
> 550	p	90	0	0.7	³⁵ BECKER-SZ... 90	IMB3
> 260	p	90	0	<0.04	HIRATA 89C	KAMI
> 130	n	90	0	<0.2	HIRATA 89C	KAMI
> 310	p	90	0	0.6	SEIDEL 88	IMB
> 100	n	90	0	1.6	SEIDEL 88	IMB
> 1.3	n	90	0		BARTELT 87	SOUD
> 1.3	p	90	0		BARTELT 87	SOUD
> 250	p	90	0	0.3	HAINES 86	IMB
> 31	n	90	8	9	HAINES 86	IMB
> 64	p	90	0	<0.4	ARISAKA 85	KAMI
> 26	n	90	0	<0.7	ARISAKA 85	KAMI
> 82	p (free)	90	0	0.2	BLEWITT 85	IMB
> 250	p	90	0	0.2	BLEWITT 85	IMB

> 25	<i>n</i>	90	4 4	PARK	85	IMB
> 15	<i>p, n</i>	90	0	BATTISTONI	84	NUSX
> 0.5	<i>p</i>	90	1 0.3	³⁶ BARTELT	83	SOUD
> 0.5	<i>n</i>	90	1 0.3	³⁶ BARTELT	83	SOUD
> 5.8	<i>p</i>	90	2	³⁷ KRISHNA...	82	KOLR
> 5.8	<i>n</i>	90	2	³⁷ KRISHNA...	82	KOLR
> 0.1	<i>n</i>	90		³⁸ GURR	67	CNTR

³⁵ This BECKER-SZENDY 90 result includes data from SEIDEL 88.

³⁶ Limit based on zero events.

³⁷ We have calculated 90% CL limit from 1 confined event.

³⁸ We have converted half-life to 90% CL mean life.

$\tau(N \rightarrow \mu^+ \pi)$

T2

<u>LIMIT</u> (10 ³⁰ years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>1000	<i>n</i>	90	1	0.43	NISHINO 12	SKAM
>6600	<i>p</i>	90	0	0.3	NISHINO 09	SKAM

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 473	<i>p</i>	90	0 0.6	MCGREW	99	IMB3
> 90	<i>n</i>	90	1 1.9	MCGREW	99	IMB3
> 81	<i>p</i>	90	0 0.2	BERGER	91	FREJ
> 35	<i>n</i>	90	1 1.0	BERGER	91	FREJ
> 230	<i>p</i>	90	0 <0.07	HIRATA	89C	KAMI
> 100	<i>n</i>	90	0 <0.2	HIRATA	89C	KAMI
> 270	<i>p</i>	90	0 0.5	SEIDEL	88	IMB
> 63	<i>n</i>	90	0 0.5	SEIDEL	88	IMB
> 76	<i>p</i>	90	2 1	HAINES	86	IMB
> 23	<i>n</i>	90	8 7	HAINES	86	IMB
> 46	<i>p</i>	90	0 <0.7	ARISAKA	85	KAMI
> 20	<i>n</i>	90	0 <0.4	ARISAKA	85	KAMI
> 59	<i>p</i> (free)	90	0 0.2	BLEWITT	85	IMB
> 100	<i>p</i>	90	1 0.4	BLEWITT	85	IMB
> 38	<i>n</i>	90	1 4	PARK	85	IMB
> 10	<i>p, n</i>	90	0	BATTISTONI	84	NUSX
> 1.3	<i>p, n</i>	90	0	ALEKSEEV	81	BAKS

$\tau(N \rightarrow \nu \pi)$

T3

<u>LIMIT</u> (10 ³⁰ years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
> 16	<i>p</i>	90	6	6.7	WALL 00B	SOU2
>112	<i>n</i>	90	6	6.6	MCGREW 99	IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 39	<i>n</i>	90	4 3.8	WALL	00B	SOU2
> 10	<i>p</i>	90	15 20.3	MCGREW	99	IMB3
> 13	<i>n</i>	90	1 1.2	BERGER	89	FREJ
> 10	<i>p</i>	90	11 14	BERGER	89	FREJ
> 25	<i>p</i>	90	32 32.8	³⁹ HIRATA	89C	KAMI
>100	<i>n</i>	90	1 3	HIRATA	89C	KAMI
> 6	<i>n</i>	90	73 60	HAINES	86	IMB
> 2	<i>p</i>	90	16 13	KAJITA	86	KAMI

> 40	<i>n</i>	90	0	1	KAJITA	86	KAMI
> 7	<i>n</i>	90	28	19	PARK	85	IMB
> 7	<i>n</i>	90	0		BATTISTONI	84	NUSX
> 2	<i>p</i>	90	≤	3	BATTISTONI	84	NUSX
> 5.8	<i>p</i>	90	1		⁴⁰ KRISHNA...	82	KOLR
> 0.3	<i>p</i>	90	2		⁴¹ CHERRY	81	HOME
> 0.1	<i>p</i>	90			⁴² GURR	67	CNTR

³⁹ In estimating the background, this HIRATA 89C limit (as opposed to the later limits of WALL 00B and MCGREW 99) does not take into account present understanding that the flux of ν_μ originating in the upper atmosphere is depleted. Doing so would reduce the background and thus also would reduce the limit here.

⁴⁰ We have calculated 90% CL limit from 1 confined event.

⁴¹ We have converted 2 possible events to 90% CL limit.

⁴² We have converted half-life to 90% CL mean life.

$\tau(p \rightarrow e^+ \eta)$

T4

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>4200	<i>p</i>	90	0	0.44	NISHINO	12 SKAM

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 81	<i>p</i>	90	1	1.7	WALL	00B	SOU2
> 313	<i>p</i>	90	0	0.2	MCGREW	99	IMB3
> 44	<i>p</i>	90	0	0.1	BERGER	91	FREJ
> 140	<i>p</i>	90	0	<0.04	HIRATA	89C	KAMI
> 100	<i>p</i>	90	0	0.6	SEIDEL	88	IMB
> 200	<i>p</i>	90	5	3.3	HAINES	86	IMB
> 64	<i>p</i>	90	0	<0.8	ARISAKA	85	KAMI
> 64	<i>p</i> (free)	90	5	6.5	BLEWITT	85	IMB
> 200	<i>p</i>	90	5	4.7	BLEWITT	85	IMB
> 1.2	<i>p</i>	90	2		⁴³ CHERRY	81	HOME

⁴³ We have converted 2 possible events to 90% CL limit.

$\tau(p \rightarrow \mu^+ \eta)$

T5

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>1300	<i>p</i>	90	2	0.49	NISHINO	12 SKAM

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 89	<i>p</i>	90	0	1.6	WALL	00B	SOU2
> 126	<i>p</i>	90	3	2.8	MCGREW	99	IMB3
> 26	<i>p</i>	90	1	0.8	BERGER	91	FREJ
> 69	<i>p</i>	90	1	<0.08	HIRATA	89C	KAMI
> 1.3	<i>p</i>	90	0	0.7	PHILLIPS	89	HPW
> 34	<i>p</i>	90	1	1.5	SEIDEL	88	IMB
> 46	<i>p</i>	90	7	6	HAINES	86	IMB
> 26	<i>p</i>	90	1	<0.8	ARISAKA	85	KAMI
> 17	<i>p</i> (free)	90	6	6	BLEWITT	85	IMB
> 46	<i>p</i>	90	7	8	BLEWITT	85	IMB

$\tau(n \rightarrow \nu\eta)$

T6

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>158	<i>n</i>	90	0	1.2	MCGREW	99 IMB3
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
> 71	<i>n</i>	90	2	3.7	WALL	00B SOU2
> 29	<i>n</i>	90	0	0.9	BERGER	89 FREJ
> 54	<i>n</i>	90	2	0.9	HIRATA	89C KAMI
> 16	<i>n</i>	90	3	2.1	SEIDEL	88 IMB
> 25	<i>n</i>	90	7	6	HAINES	86 IMB
> 30	<i>n</i>	90	0	0.4	KAJITA	86 KAMI
> 18	<i>n</i>	90	4	3	PARK	85 IMB
> 0.6	<i>n</i>	90	2		⁴⁴ CHERRY	81 HOME

⁴⁴We have converted 2 possible events to 90% CL limit.

$\tau(N \rightarrow e^+ \rho)$

T7

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>710	<i>p</i>	90	0	0.35	NISHINO	12 SKAM
>217	<i>n</i>	90	4	4.8	MCGREW	99 IMB3
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
> 70	<i>n</i>	90	1	0.38	NISHINO	12 SKAM
> 29	<i>p</i>	90	0	2.2	BERGER	91 FREJ
> 41	<i>n</i>	90	0	1.4	BERGER	91 FREJ
> 75	<i>p</i>	90	2	2.7	HIRATA	89C KAMI
> 58	<i>n</i>	90	0	1.9	HIRATA	89C KAMI
> 38	<i>n</i>	90	2	4.1	SEIDEL	88 IMB
> 1.2	<i>p</i>	90	0		BARTELT	87 SOUD
> 1.5	<i>n</i>	90	0		BARTELT	87 SOUD
> 17	<i>p</i>	90	7	7	HAINES	86 IMB
> 14	<i>n</i>	90	9	4	HAINES	86 IMB
> 12	<i>p</i>	90	0	<1.2	ARISAKA	85 KAMI
> 6	<i>n</i>	90	2	<1	ARISAKA	85 KAMI
> 6.7	<i>p</i> (free)	90	6	6	BLEWITT	85 IMB
> 17	<i>p</i>	90	7	7	BLEWITT	85 IMB
> 12	<i>n</i>	90	4	2	PARK	85 IMB
> 0.6	<i>n</i>	90	1	0.3	⁴⁵ BARTELT	83 SOUD
> 0.5	<i>p</i>	90	1	0.3	⁴⁵ BARTELT	83 SOUD
> 9.8	<i>p</i>	90	1		⁴⁶ KRISHNA...	82 KOLR
> 0.8	<i>p</i>	90	2		⁴⁷ CHERRY	81 HOME

⁴⁵Limit based on zero events.

⁴⁶We have calculated 90% CL limit from 0 confined events.

⁴⁷We have converted 2 possible events to 90% CL limit.

$\tau(N \rightarrow \mu^+ \rho)$

T8

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>160	<i>p</i>	90	1	0.42	NISHINO	12 SKAM
>228	<i>n</i>	90	3	9.5	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 36	<i>n</i>	90	0	0.29	NISHINO	12	SKAM
> 12	<i>p</i>	90	0	0.5	BERGER	91	FREJ
> 22	<i>n</i>	90	0	1.1	BERGER	91	FREJ
>110	<i>p</i>	90	0	1.7	HIRATA	89C	KAMI
> 23	<i>n</i>	90	1	1.8	HIRATA	89C	KAMI
> 4.3	<i>p</i>	90	0	0.7	PHILLIPS	89	HPW
> 30	<i>p</i>	90	0	0.5	SEIDEL	88	IMB
> 11	<i>n</i>	90	1	1.1	SEIDEL	88	IMB
> 16	<i>p</i>	90	4	4.5	HAINES	86	IMB
> 7	<i>n</i>	90	6	5	HAINES	86	IMB
> 12	<i>p</i>	90	0	<0.7	ARISAKA	85	KAMI
> 5	<i>n</i>	90	1	<1.2	ARISAKA	85	KAMI
> 5.5	<i>p</i> (free)	90	4	5	BLEWITT	85	IMB
> 16	<i>p</i>	90	4	5	BLEWITT	85	IMB
> 9	<i>n</i>	90	1	2	PARK	85	IMB

$\tau(N \rightarrow \nu \rho)$

79

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>162	<i>p</i>	90	18	21.7	MCGREW	99 IMB3
> 19	<i>n</i>	90	0	0.5	SEIDEL	88 IMB

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 9	<i>n</i>	90	4	2.4	BERGER	89	FREJ
> 24	<i>p</i>	90	0	0.9	BERGER	89	FREJ
> 27	<i>p</i>	90	5	1.5	HIRATA	89C	KAMI
> 13	<i>n</i>	90	4	3.6	HIRATA	89C	KAMI
> 13	<i>p</i>	90	1	1.1	SEIDEL	88	IMB
> 8	<i>p</i>	90	6	5	HAINES	86	IMB
> 2	<i>n</i>	90	15	10	HAINES	86	IMB
> 11	<i>p</i>	90	2	1	KAJITA	86	KAMI
> 4	<i>n</i>	90	2	2	KAJITA	86	KAMI
> 4.1	<i>p</i> (free)	90	6	7	BLEWITT	85	IMB
> 8.4	<i>p</i>	90	6	5	BLEWITT	85	IMB
> 2	<i>n</i>	90	7	3	PARK	85	IMB
> 0.9	<i>p</i>	90	2		⁴⁸ CHERRY	81	HOME
> 0.6	<i>n</i>	90	2		⁴⁸ CHERRY	81	HOME

⁴⁸We have converted 2 possible events to 90% CL limit.

$\tau(p \rightarrow e^+ \omega)$

710

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>320	<i>p</i>	90	1	0.53	NISHINO	12 SKAM

• • • We do not use the following data for averages, fits, limits, etc. • • •

>107	<i>p</i>	90	7	10.8	MCGREW	99	IMB3
> 17	<i>p</i>	90	0	1.1	BERGER	91	FREJ
> 45	<i>p</i>	90	2	1.45	HIRATA	89C	KAMI
> 26	<i>p</i>	90	1	1.0	SEIDEL	88	IMB
> 1.5	<i>p</i>	90	0		BARTELT	87	SOUD
> 37	<i>p</i>	90	6	5.3	HAINES	86	IMB

> 25	p	90	1	<1.4	ARISAKA	85	KAMI
> 12	p (free)	90	6	7.5	BLEWITT	85	IMB
> 37	p	90	6	5.7	BLEWITT	85	IMB
> 0.6	p	90	1	0.3	⁴⁹ BARTELT	83	SOUD
> 9.8	p	90	1		⁵⁰ KRISHNA...	82	KOLR
> 2.8	p	90	2		⁵¹ CHERRY	81	HOME

⁴⁹ Limit based on zero events.

⁵⁰ We have calculated 90% CL limit from 0 confined events.

⁵¹ We have converted 2 possible events to 90% CL limit.

$\tau(p \rightarrow \mu^+ \omega)$

τ_{11}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>780	p	90	0	0.48	NISHINO	12 SKAM

• • • We do not use the following data for averages, fits, limits, etc. • • •

>117	p	90	11	12.1	MCGREW	99	IMB3
> 11	p	90	0	1.0	BERGER	91	FREJ
> 57	p	90	2	1.9	HIRATA	89C	KAMI
> 4.4	p	90	0	0.7	PHILLIPS	89	HPW
> 10	p	90	2	1.3	SEIDEL	88	IMB
> 23	p	90	2	1	HAINES	86	IMB
> 6.5	p (free)	90	9	8.7	BLEWITT	85	IMB
> 23	p	90	8	7	BLEWITT	85	IMB

$\tau(n \rightarrow \nu \omega)$

τ_{12}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	
>108	n	90	12	22.5	MCGREW	99	IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 17	n	90	1	0.7	BERGER	89	FREJ
> 43	n	90	3	2.7	HIRATA	89C	KAMI
> 6	n	90	2	1.3	SEIDEL	88	IMB
> 12	n	90	6	6	HAINES	86	IMB
> 18	n	90	2	2	KAJITA	86	KAMI
> 16	n	90	1	2	PARK	85	IMB
> 2.0	n	90	2		⁵² CHERRY	81	HOME

⁵² We have converted 2 possible events to 90% CL limit.

$\tau(N \rightarrow e^+ K)$

τ_{13}

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	
>1000	p	90	6	4.7	KOBAYASHI	05	SKAM
> 17	n	90	35	29.4	MCGREW	99	IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 85	p	90	3	4.9	WALL	00	SOU2
> 31	p	90	23	25.2	MCGREW	99	IMB3
> 60	p	90	0		BERGER	91	FREJ
> 150	p	90	0	<0.27	HIRATA	89C	KAMI
> 70	p	90	0	1.8	SEIDEL	88	IMB
> 77	p	90	5	4.5	HAINES	86	IMB

> 38	p	90	0	<0.8	ARISAKA	85	KAMI
> 24	p (free)	90	7	8.5	BLEWITT	85	IMB
> 77	p	90	5	4	BLEWITT	85	IMB
> 1.3	p	90	0		ALEKSEEV	81	BAKS
> 1.3	n	90	0		ALEKSEEV	81	BAKS

$\tau(p \rightarrow e^+ K_S^0)$ **τ_{14}**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
------------------------------------	-----------------	------------	-------------	-----------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>120	p	90	1	1.3	WALL	00	SOU2
> 76	p	90	0	0.5	BERGER	91	FREJ

$\tau(p \rightarrow e^+ K_L^0)$ **τ_{15}**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
------------------------------------	-----------------	------------	-------------	-----------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>51	p	90	2	3.5	WALL	00	SOU2
>44	p	90	0	≤ 0.1	BERGER	91	FREJ

$\tau(N \rightarrow \mu^+ K)$ **τ_{16}**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
------------------------------------	-----------------	------------	-------------	-----------------	--------------------	-------------

>1600	p	90	13	13.2	REGIS	12	SKAM
> 26	n	90	20	28.4	MCGREW	99	IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>1300	p	90	3	3.9	KOBAYASHI	05	SKAM
> 120	p	90	0	<1.2	WALL	00	SOU2
> 120	p	90	4	7.2	MCGREW	99	IMB3
> 54	p	90	0		BERGER	91	FREJ
> 120	p	90	1	0.4	HIRATA	89C	KAMI
> 3.0	p	90	0	0.7	PHILLIPS	89	HPW
> 19	p	90	3	2.5	SEIDEL	88	IMB
> 1.5	p	90	0		⁵³ BARTELT	87	SOUD
> 1.1	n	90	0		BARTELT	87	SOUD
> 40	p	90	7	6	HAINES	86	IMB
> 19	p	90	1	<1.1	ARISAKA	85	KAMI
> 6.7	p (free)	90	11	13	BLEWITT	85	IMB
> 40	p	90	7	8	BLEWITT	85	IMB
> 6	p	90	1		BATTISTONI	84	NUSX
> 0.6	p	90	0		⁵⁴ BARTELT	83	SOUD
> 0.4	n	90	0		⁵⁴ BARTELT	83	SOUD
> 5.8	p	90	2		⁵⁵ KRISHNA...	82	KOLR
> 2.0	p	90	0		CHERRY	81	HOME
> 0.2	n	90			⁵⁶ GURR	67	CNTR

⁵³BARTELT 87 limit applies to $p \rightarrow \mu^+ K_S^0$.

⁵⁴Limit based on zero events.

⁵⁵We have calculated 90% CL limit from 1 confined event.

⁵⁶We have converted half-life to 90% CL mean life.

$\tau(p \rightarrow \mu^+ K_S^0)$

τ_{17}

LIMIT
(10^{30} years) PARTICLE CL% EVTS BKGD EST DOCUMENT ID TECN

• • • We do not use the following data for averages, fits, limits, etc. • • •

>150	p	90	0	<0.8	WALL	00	SOU2
> 64	p	90	0	1.2	BERGER	91	FREJ

$\tau(p \rightarrow \mu^+ K_L^0)$

τ_{18}

LIMIT
(10^{30} years) PARTICLE CL% EVTS BKGD EST DOCUMENT ID TECN

• • • We do not use the following data for averages, fits, limits, etc. • • •

>83	p	90	0	0.4	WALL	00	SOU2
>44	p	90	0	≤ 0.1	BERGER	91	FREJ

$\tau(N \rightarrow \nu K)$

τ_{19}

LIMIT
(10^{30} years) PARTICLE CL% EVTS BKGD EST DOCUMENT ID TECN

>2300	p	90	0	1.3	KOBAYASHI	05	SKAM
> 86	n	90	0	2.4	HIRATA	89C	KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 26	n	90	16	9.1	WALL	00	SOU2
> 670	p	90			HAYATO	99	SKAM
> 151	p	90	15	21.4	MCGREW	99	IMB3
> 30	n	90	34	34.1	MCGREW	99	IMB3
> 43	p	90	1	1.54	⁵⁷ ALLISON	98	SOU2
> 15	n	90	1	1.8	BERGER	89	FREJ
> 15	p	90	1	1.8	BERGER	89	FREJ
> 100	p	90	9	7.3	HIRATA	89C	KAMI
> 0.28	p	90	0	0.7	PHILLIPS	89	HPW
> 0.3	p	90	0		BARTELT	87	SOUD
> 0.75	n	90	0		⁵⁸ BARTELT	87	SOUD
> 10	p	90	6	5	HAINES	86	IMB
> 15	n	90	3	5	HAINES	86	IMB
> 28	p	90	3	3	KAJITA	86	KAMI
> 32	n	90	0	1.4	KAJITA	86	KAMI
> 1.8	p (free)	90	6	11	BLEWITT	85	IMB
> 9.6	p	90	6	5	BLEWITT	85	IMB
> 10	n	90	2	2	PARK	85	IMB
> 5	n	90	0		BATTISTONI	84	NUSX
> 2	p	90	0		BATTISTONI	84	NUSX
> 0.3	n	90	0		⁵⁹ BARTELT	83	SOUD
> 0.1	p	90	0		⁵⁹ BARTELT	83	SOUD
> 5.8	p	90	1		⁶⁰ KRISHNA...	82	KOLR
> 0.3	n	90	2		⁶¹ CHERRY	81	HOME

⁵⁷ This ALLISON 98 limit is with no background subtraction; with subtraction the limit becomes $> 46 \times 10^{30}$ years.

⁵⁸ BARTELT 87 limit applies to $n \rightarrow \nu K_S^0$.

⁵⁹ Limit based on zero events.

⁶⁰ We have calculated 90% CL limit from 1 confined event.

⁶¹ We have converted 2 possible events to 90% CL limit.

$\tau(n \rightarrow \nu K_S^0)$

T20

<u>LIMIT</u> (10 ³⁰ years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>260	n	90	34	30	⁶² KOBAYASHI 05	SKAM

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 51	n	90	16	9.1	WALL	00	SOU2
------	---	----	----	-----	------	----	------

⁶²We have doubled the $n \rightarrow \nu K^0$ limit given in KOBAYASHI 05 to obtain this $n \rightarrow \nu K_S^0$ limit.

$\tau(p \rightarrow e^+ K^*(892)^0)$

T21

<u>LIMIT</u> (10 ³⁰ years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	
>84	p	90	38	52.0	MCGREW	99	IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>10	p	90	0	0.8	BERGER	91	FREJ
>52	p	90	2	1.55	HIRATA	89C	KAMI
>10	p	90	1	<1	ARISAKA	85	KAMI

$\tau(N \rightarrow \nu K^*(892))$

T22

<u>LIMIT</u> (10 ³⁰ years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	
>51	p	90	7	9.1	MCGREW	99	IMB3
>78	n	90	40	50	MCGREW	99	IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>22	n	90	0	2.1	BERGER	89	FREJ
>17	p	90	0	2.4	BERGER	89	FREJ
>20	p	90	5	2.1	HIRATA	89C	KAMI
>21	n	90	4	2.4	HIRATA	89C	KAMI
>10	p	90	7	6	HAINES	86	IMB
> 5	n	90	8	7	HAINES	86	IMB
> 8	p	90	3	2	KAJITA	86	KAMI
> 6	n	90	2	1.6	KAJITA	86	KAMI
> 5.8	p (free)	90	10	16	BLEWITT	85	IMB
> 9.6	p	90	7	6	BLEWITT	85	IMB
> 7	n	90	1	4	PARK	85	IMB
> 2.1	p	90	1		⁶³ BATTISTONI	82	NUSX

⁶³We have converted 1 possible event to 90% CL limit.

————— **Antilepton + mesons** —————

$\tau(p \rightarrow e^+ \pi^+ \pi^-)$

T23

<u>LIMIT</u> (10 ³⁰ years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	
>82	p	90	16	23.1	MCGREW	99	IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>21	p	90	0	2.2	BERGER	91	FREJ
-----	---	----	---	-----	--------	----	------

$\tau(p \rightarrow e^+ \pi^0 \pi^0)$ **T24**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>147	p	90	2	0.8	MCGREW 99	IMB3
••• We do not use the following data for averages, fits, limits, etc. •••						
> 38	p	90	1	0.5	BERGER 91	FREJ

$\tau(n \rightarrow e^+ \pi^- \pi^0)$ **T25**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>52	n	90	38	34.2	MCGREW 99	IMB3
••• We do not use the following data for averages, fits, limits, etc. •••						
>32	n	90	1	0.8	BERGER 91	FREJ

$\tau(p \rightarrow \mu^+ \pi^+ \pi^-)$ **T26**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>133	p	90	25	38.0	MCGREW 99	IMB3
••• We do not use the following data for averages, fits, limits, etc. •••						
> 17	p	90	1	2.6	BERGER 91	FREJ
> 3.3	p	90	0	0.7	PHILLIPS 89	HPW

$\tau(p \rightarrow \mu^+ \pi^0 \pi^0)$ **T27**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>101	p	90	3	1.6	MCGREW 99	IMB3
••• We do not use the following data for averages, fits, limits, etc. •••						
> 33	p	90	1	0.9	BERGER 91	FREJ

$\tau(n \rightarrow \mu^+ \pi^- \pi^0)$ **T28**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>74	n	90	17	20.8	MCGREW 99	IMB3
••• We do not use the following data for averages, fits, limits, etc. •••						
>33	n	90	0	1.1	BERGER 91	FREJ

$\tau(n \rightarrow e^+ K^0 \pi^-)$ **T29**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>18	n	90	1	0.2	BERGER 91	FREJ

————— **Lepton + meson** —————

$\tau(n \rightarrow e^- \pi^+)$ **T30**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>65	n	90	0	1.6	SEIDEL 88	IMB
••• We do not use the following data for averages, fits, limits, etc. •••						
>55	n	90	0	1.09	BERGER 91B	FREJ
>16	n	90	9	7	HAINES 86	IMB
>25	n	90	2	4	PARK 85	IMB

$\tau(n \rightarrow \mu^- \pi^+)$ **T31**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>49	<i>n</i>	90	0	0.5	SEIDEL 88	IMB
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
>33	<i>n</i>	90	0	1.40	BERGER 91B	FREJ
> 2.7	<i>n</i>	90	0	0.7	PHILLIPS 89	HPW
>25	<i>n</i>	90	7	6	HAINES 86	IMB
>27	<i>n</i>	90	2	3	PARK 85	IMB

$\tau(n \rightarrow e^- \rho^+)$ **T32**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>62	<i>n</i>	90	2	4.1	SEIDEL 88	IMB
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
>12	<i>n</i>	90	13	6	HAINES 86	IMB
>12	<i>n</i>	90	5	3	PARK 85	IMB

$\tau(n \rightarrow \mu^- \rho^+)$ **T33**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>7	<i>n</i>	90	1	1.1	SEIDEL 88	IMB
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
>2.6	<i>n</i>	90	0	0.7	PHILLIPS 89	HPW
>9	<i>n</i>	90	7	5	HAINES 86	IMB
>9	<i>n</i>	90	2	2	PARK 85	IMB

$\tau(n \rightarrow e^- K^+)$ **T34**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>32	<i>n</i>	90	3	2.96	BERGER 91B	FREJ
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
> 0.23	<i>n</i>	90	0	0.7	PHILLIPS 89	HPW

$\tau(n \rightarrow \mu^- K^+)$ **T35**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>57	<i>n</i>	90	0	2.18	BERGER 91B	FREJ
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
> 4.7	<i>n</i>	90	0	0.7	PHILLIPS 89	HPW

————— **Lepton + mesons** —————

$\tau(p \rightarrow e^- \pi^+ \pi^+)$ **T36**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>30	<i>p</i>	90	1	2.50	BERGER 91B	FREJ
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
> 2.0	<i>p</i>	90	0	0.7	PHILLIPS 89	HPW

$\tau(n \rightarrow e^- \pi^+ \pi^0)$ **T37**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>29	<i>n</i>	90	1	0.78	BERGER	91B FREJ

$\tau(p \rightarrow \mu^- \pi^+ \pi^+)$ **T38**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>17	<i>p</i>	90	1	1.72	BERGER	91B FREJ

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 7.8	<i>p</i>	90	0	0.7	PHILLIPS	89 HPW
-------	----------	----	---	-----	----------	--------

$\tau(n \rightarrow \mu^- \pi^+ \pi^0)$ **T39**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>34	<i>n</i>	90	0	0.78	BERGER	91B FREJ

$\tau(p \rightarrow e^- \pi^+ K^+)$ **T40**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>75	<i>p</i>	90	81	127.2	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>20	<i>p</i>	90	3	2.50	BERGER	91B FREJ
-----	----------	----	---	------	--------	----------

$\tau(p \rightarrow \mu^- \pi^+ K^+)$ **T41**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>245	<i>p</i>	90	3	4.0	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 5	<i>p</i>	90	2	0.78	BERGER	91B FREJ
-----	----------	----	---	------	--------	----------

————— **Antilepton + photon(s)** —————

$\tau(p \rightarrow e^+ \gamma)$ **T42**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>670	<i>p</i>	90	0	0.1	MCGREW	99 IMB3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>133	<i>p</i>	90	0	0.3	BERGER	91 FREJ
>460	<i>p</i>	90	0	0.6	SEIDEL	88 IMB
>360	<i>p</i>	90	0	0.3	HAINES	86 IMB
> 87	<i>p</i> (free)	90	0	0.2	BLEWITT	85 IMB
>360	<i>p</i>	90	0	0.2	BLEWITT	85 IMB
> 0.1	<i>p</i>	90			⁶⁴ GURR	67 CNTR

⁶⁴We have converted half-life to 90% CL mean life.

$\tau(p \rightarrow \mu^+ \gamma)$

T43

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>478	p	90	0	0.1	MCGREW	99 IMB3
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>155	p	90	0	0.1	BERGER	91 FREJ
>380	p	90	0	0.5	SEIDEL	88 IMB
> 97	p	90	3	2	HAINES	86 IMB
> 61	p (free)	90	0	0.2	BLEWITT	85 IMB
>280	p	90	0	0.6	BLEWITT	85 IMB
> 0.3	p	90			⁶⁵ GURR	67 CNTR

⁶⁵We have converted half-life to 90% CL mean life.

$\tau(n \rightarrow \nu \gamma)$

T44

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>28	n	90	163	144.7	MCGREW	99 IMB3
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>24	n	90	10	6.86	BERGER	91B FREJ
> 9	n	90	73	60	HAINES	86 IMB
>11	n	90	28	19	PARK	85 IMB

$\tau(p \rightarrow e^+ \gamma \gamma)$

T45

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>100	p	90	1	0.8	BERGER	91 FREJ

$\tau(n \rightarrow \nu \gamma \gamma)$

T46

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>219	n	90	5	7.5	MCGREW	99 IMB3

————— **Three (or more) leptons** —————

$\tau(p \rightarrow e^+ e^+ e^-)$

T47

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>793	p	90	0	0.5	MCGREW	99 IMB3
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>147	p	90	0	0.1	BERGER	91 FREJ
>510	p	90	0	0.3	HAINES	86 IMB
> 89	p (free)	90	0	0.5	BLEWITT	85 IMB
>510	p	90	0	0.7	BLEWITT	85 IMB

$\tau(p \rightarrow e^+ \mu^+ \mu^-)$

T48

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>359	p	90	1	0.9	MCGREW	99 IMB3
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 81	p	90	0	0.16	BERGER	91 FREJ
> 5.0	p	90	0	0.7	PHILLIPS	89 HPW

$\tau(p \rightarrow e^+ \nu \nu)$

T49

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>17	p	90	152	153.7	MCGREW	99 IMB3
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
>11	p	90	11	6.08	BERGER	91B FREJ

$\tau(n \rightarrow e^+ e^- \nu)$

T50

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>257	n	90	5	7.5	MCGREW	99 IMB3
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
> 74	n	90	0	< 0.1	BERGER	91B FREJ
> 45	n	90	5	5	HAINES	86 IMB
> 26	n	90	4	3	PARK	85 IMB

$\tau(n \rightarrow \mu^+ e^- \nu)$

T51

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>83	n	90	25	29.4	MCGREW	99 IMB3
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
>47	n	90	0	< 0.1	BERGER	91B FREJ

$\tau(n \rightarrow \mu^+ \mu^- \nu)$

T52

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>79	n	90	100	145	MCGREW	99 IMB3
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
>42	n	90	0	1.4	BERGER	91B FREJ
> 5.1	n	90	0	0.7	PHILLIPS	89 HPW
>16	n	90	14	7	HAINES	86 IMB
>19	n	90	4	7	PARK	85 IMB

$\tau(p \rightarrow \mu^+ e^+ e^-)$

T53

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>529	p	90	0	1.0	MCGREW	99 IMB3
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
> 91	p	90	0	≤ 0.1	BERGER	91 FREJ

$\tau(p \rightarrow \mu^+ \mu^+ \mu^-)$

T54

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>675	p	90	0	0.3	MCGREW	99 IMB3
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
>119	p	90	0	0.2	BERGER	91 FREJ
> 10.5	p	90	0	0.7	PHILLIPS	89 HPW
>190	p	90	1	0.1	HAINES	86 IMB
> 44	p (free)	90	1	0.7	BLEWITT	85 IMB
>190	p	90	1	0.9	BLEWITT	85 IMB
> 2.1	p	90	1		⁶⁶ BATTISTONI	82 NUSX

⁶⁶We have converted 1 possible event to 90% CL limit.

$\tau(p \rightarrow \mu^+ \nu \nu)$ **T55**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>21	p	90	7	11.23	BERGER	91B FREJ

$\tau(p \rightarrow e^- \mu^+ \mu^+)$ **T56**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>6.0	p	90	0	0.7	PHILLIPS	89 HPW

$\tau(n \rightarrow 3\nu)$ **T57**

See also the “to anything” and “disappearance” limits for bound nucleons in the “ p Mean Life” data block just in front of the list of possible p decay modes. Such modes could of course be to three (or five) neutrinos, and the limits are stronger, but we do not repeat them here.

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>0.00049	n	90	2	2	⁶⁷ SUZUKI	93B KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.0023	n	90			⁶⁸ GLICENSTEIN	97 KAMI
>0.00003	n	90	11	6.1	⁶⁹ BERGER	91B FREJ
>0.00012	n	90	7	11.2	⁶⁹ BERGER	91B FREJ
>0.0005	n	90	0		LEARNED	79 RVUE

⁶⁷The SUZUKI 93B limit applies to any of $\nu_e \nu_e \bar{\nu}_e$, $\nu_\mu \nu_\mu \bar{\nu}_\mu$, or $\nu_\tau \nu_\tau \bar{\nu}_\tau$.

⁶⁸GLICENSTEIN 97 uses Kamioka data and the idea that the disappearance of the neutron's magnetic moment should produce radiation.

⁶⁹The first BERGER 91B limit is for $n \rightarrow \nu_e \nu_e \bar{\nu}_e$, the second is for $n \rightarrow \nu_\mu \nu_\mu \bar{\nu}_\mu$.

$\tau(n \rightarrow 5\nu)$ **T58**

See the note on $\tau(n \rightarrow 3\nu)$ on the previous data block.

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
------------------------------------	-----------------	------------	-------------	-----------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.0017	n	90			⁷⁰ GLICENSTEIN	97 KAMI
---------	-----	----	--	--	---------------------------	---------

⁷⁰GLICENSTEIN 97 uses Kamioka data and the idea that the disappearance of the neutron's magnetic moment should produce radiation.

————— Inclusive modes —————

$\tau(N \rightarrow e^+ \text{ anything})$ **T59**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>0.6	p, n	90			⁷¹ LEARNED	79 RVUE

⁷¹The electron may be primary or secondary.

$\tau(N \rightarrow \mu^+ \text{anything})$ **T60**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>12	p, n	90	2		^{72,73} CHERRY	81 HOME
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 1.8	p, n	90			⁷³ COWSIK	80 CNTR
> 6	p, n	90			⁷³ LEARNED	79 RVUE

⁷² We have converted 2 possible events to 90% CL limit.

⁷³ The muon may be primary or secondary.

$\tau(N \rightarrow \nu \text{anything})$ **T61**

Anything = π, ρ, K , etc.

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>0.0002	p, n	90	0		LEARNED	79 RVUE

$\tau(N \rightarrow e^+ \pi^0 \text{anything})$ **T62**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
>0.6	p, n	90	0		LEARNED	79 RVUE

$\tau(N \rightarrow 2 \text{ bodies}, \nu\text{-free})$ **T63**

<u>LIMIT</u> (10^{30} years)	<u>PARTICLE</u>	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>1.3	p, n	90	0		ALEKSEEV	81 BAKS

———— **$\Delta B = 2$ dinucleon modes** ————

$\tau(pp \rightarrow \pi^+ \pi^+)$ **T64**

<u>LIMIT</u> (10^{30} years)	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>0.7	90	4	2.34	BERGER	91B FREJ	τ per iron nucleus

$\tau(pn \rightarrow \pi^+ \pi^0)$ **T65**

<u>LIMIT</u> (10^{30} years)	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>2.0	90	0	0.31	BERGER	91B FREJ	τ per iron nucleus

$\tau(nn \rightarrow \pi^+ \pi^-)$ **T66**

<u>LIMIT</u> (10^{30} years)	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>0.7	90	4	2.18	BERGER	91B FREJ	τ per iron nucleus

$\tau(nn \rightarrow \pi^0 \pi^0)$ **T67**

<u>LIMIT</u> (10^{30} years)	<u>CL%</u>	<u>EVTS</u>	<u>BKGD EST</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>3.4	90	0	0.78	BERGER	91B FREJ	τ per iron nucleus

$\tau(pp \rightarrow e^+ e^+)$ **T68**

LIMIT (10^{30} years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>5.8	90	0	<0.1	BERGER	91B FREJ	τ per iron nucleus

$\tau(pp \rightarrow e^+ \mu^+)$ **T69**

LIMIT (10^{30} years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>3.6	90	0	<0.1	BERGER	91B FREJ	τ per iron nucleus

$\tau(pp \rightarrow \mu^+ \mu^+)$ **T70**

LIMIT (10^{30} years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>1.7	90	0	0.62	BERGER	91B FREJ	τ per iron nucleus

$\tau(pn \rightarrow e^+ \bar{\nu})$ **T71**

LIMIT (10^{30} years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>2.8	90	5	9.67	BERGER	91B FREJ	τ per iron nucleus

$\tau(pn \rightarrow \mu^+ \bar{\nu})$ **T72**

LIMIT (10^{30} years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>1.6	90	4	4.37	BERGER	91B FREJ	τ per iron nucleus

$\tau(nn \rightarrow \nu_e \bar{\nu}_e)$ **T73**

We include "invisible" modes here.

LIMIT (10^{30} years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>1.4	90			⁷⁴ ARAKI	06 KLND	$nn \rightarrow$ invisible
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
>0.000042	90			⁷⁵ TRETAYAK	04 CNTR	$nn \rightarrow$ invisible
>0.000049	90			⁷⁶ BACK	03 BORX	$nn \rightarrow$ invisible
>0.000012	90			⁷⁷ BERNABEI	00B DAMA	$nn \rightarrow$ invisible
>0.000012	90	5	9.7	BERGER	91B FREJ	τ per iron nucleus

⁷⁴ ARAKI 06 looks for signs of de-excitation of the residual nucleus after disappearance of two neutrons from the s shell of ^{12}C .

⁷⁵ TRETAYAK 04 uses data from an old Homestake-mine radiochemical experiment on limits for invisible decays of ^{39}K to ^{37}Ar .

⁷⁶ BACK 03 looks for decays of unstable nuclides left after NN decays of parent ^{12}C , ^{13}C , ^{16}O nuclei. These are "invisible channel" limits.

⁷⁷ BERNABEI 00B looks for the decay of a $^{127}_{54}\text{Xe}$ nucleus following the disappearance of an nn pair in the otherwise-stable $^{129}_{54}\text{Xe}$ nucleus. The limit here applies as well to $nn \rightarrow \nu_\mu \bar{\nu}_\mu$, $nn \rightarrow \nu_\tau \bar{\nu}_\tau$, or any "disappearance" mode.

$\tau(nn \rightarrow \nu_\mu \bar{\nu}_\mu)$ **T74**

See the preceding data block. "Invisible modes" would include any multi-neutrino mode.

LIMIT (10^{30} years)	CL%	EVTS	BKGD EST	CL%	DOCUMENT ID	TECN	COMMENT
>1.4	(CL = 90%) OUR LIMIT						

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

>0.000006	90	4	4.4	BERGER	91B FREJ	τ per iron nucleus
-----------	----	---	-----	--------	----------	-------------------------

$\tau(pn \rightarrow \text{invisible})$

T75

This violates charge conservation as well as baryon number conservation.

VALUE (10^{30} years)	CL%	DOCUMENT ID	TECN
>0.000021	90	⁷⁸ TRETYAK 04	CNTR

⁷⁸TRETYAK 04 uses data from an old Homestake-mine radiochemical experiment on limits for invisible decays of ³⁹K to ³⁷Ar.

$\tau(pp \rightarrow \text{invisible})$

T76

This violates charge conservation as well as baryon number conservation.

LIMIT (10^{30} years)	CL%	EVTS	BKGD EST	CL%	DOCUMENT ID	TECN
>0.00005				90	⁷⁹ BACK 03	BORX

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.0000055 90 ⁸⁰BERNABEI 00B DAMA

⁷⁹BACK 03 looks for decays of unstable nuclides left after *NN* decays of parent ¹²C, ¹³C, ¹⁶O nuclei. These are "invisible channel" limits.

⁸⁰BERNABEI 00B looks for the decay of a ¹²⁷₅₂Te nucleus following the disappearance of a *pp* pair in the otherwise-stable ¹²⁹₅₄Xe nucleus.

\bar{p} PARTIAL MEAN LIVES

The "partial mean life" limits tabulated here are the limits on $\bar{\tau}/B_i$, where $\bar{\tau}$ is the total mean life for the antiproton and B_i is the branching fraction for the mode in question.

$\tau(\bar{p} \rightarrow e^- \gamma)$

T77

VALUE (years)	CL%	DOCUMENT ID	TECN	COMMENT
> 7×10^5	90	GEER 00	APEX	8.9 GeV/ <i>c</i> \bar{p} beam
>1848	95	GEER 94	CALO	8.9 GeV/ <i>c</i> \bar{p} beam

$\tau(\bar{p} \rightarrow \mu^- \gamma)$

T78

VALUE (years)	CL%	DOCUMENT ID	TECN	COMMENT
> 5×10^4	90	GEER 00	APEX	8.9 GeV/ <i>c</i> \bar{p} beam
> 5.0×10^4	90	HU 98B	APEX	8.9 GeV/ <i>c</i> \bar{p} beam

$\tau(\bar{p} \rightarrow e^- \pi^0)$

T79

VALUE (years)	CL%	DOCUMENT ID	TECN	COMMENT
> 4×10^5	90	GEER 00	APEX	8.9 GeV/ <i>c</i> \bar{p} beam
>554	95	GEER 94	CALO	8.9 GeV/ <i>c</i> \bar{p} beam

$\tau(\bar{p} \rightarrow \mu^- \pi^0)$

T80

VALUE (years)	CL%	DOCUMENT ID	TECN	COMMENT
> 5×10^4	90	GEER 00	APEX	8.9 GeV/ <i>c</i> \bar{p} beam
> 4.8×10^4	90	HU 98B	APEX	8.9 GeV/ <i>c</i> \bar{p} beam

$\tau(\bar{p} \rightarrow e^- \eta)$ **T81**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
> 2 × 10⁴	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>171	95	GEER	94	CALO 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow \mu^- \eta)$ **T82**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>8 × 10³	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>7.9 × 10 ³	90	HU	98B	APEX 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow e^- K_S^0)$ **T83**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>900	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 29	95	GEER	94	CALO 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow \mu^- K_S^0)$ **T84**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>4 × 10³	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>4.3 × 10 ³	90	HU	98B	APEX 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow e^- K_L^0)$ **T85**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>9 × 10³	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>9	95	GEER	94	CALO 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow \mu^- K_L^0)$ **T86**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>7 × 10³	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>6.5 × 10 ³	90	HU	98B	APEX 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow e^- \gamma \gamma)$ **T87**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>2 × 10⁴	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow \mu^- \gamma \gamma)$ **T88**

<u>VALUE (years)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>2 × 10⁴	90	GEER	00	APEX 8.9 GeV/c \bar{p} beam
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>2.3 × 10 ⁴	90	HU	98B	APEX 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow e^- \rho)$ 789

VALUE (years)	CL%	DOCUMENT ID	TECN	COMMENT
---------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>200 90 ⁸¹ GEER 00 APEX 8.9 GeV/c \bar{p} beam

⁸¹ This GEER 00 measurement has been withdrawn; see GEER 00C.

$\tau(\bar{p} \rightarrow e^- \omega)$ 790

VALUE (years)	CL%	DOCUMENT ID	TECN	COMMENT
---------------	-----	-------------	------	---------

>200 90 GEER 00 APEX 8.9 GeV/c \bar{p} beam

$\tau(\bar{p} \rightarrow e^- K^*(892)^0)$ 791

VALUE (years)	CL%	DOCUMENT ID	TECN	COMMENT
---------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>1 × 10³ 90 ⁸² GEER 00 APEX 8.9 GeV/c \bar{p} beam

⁸² This GEER 00 measurement has been withdrawn; see GEER 00C.

p REFERENCES

ANTOGNINI	13	SCI 339 417	A. Antognini <i>et al.</i>	(MPI, ETH, UPMC+)
DISCIACCA	13	PRL 110 130801	J. DiSciacca <i>et al.</i>	(ATRAP Collab.)
MCGOVERN	13	EPJ A49 12	J.A. McGovern, D.R. Phillips, H.W. Griesshammer	
MOHR	12	RMP 84 1527	P.J. Mohr, B.N. Taylor, D.B. Newell	(NIST)
NISHINO	12	PR D85 112001	H. Nishino <i>et al.</i>	(Super Kamiokande Collab.)
REGIS	12	PR D86 012006	C. Regis <i>et al.</i>	(Super-Kamiokande Collab.)
ARRINGTON	11	PRL 107 119101	J. Arrington	(ANL)
BERNAUER	11	PRL 107 119102	J.C. Bernauer <i>et al.</i>	(MAMI A1 Collab.)
BRESSI	11	PR A83 052101	G. Bressi <i>et al.</i>	(LEGN, PAVII, PADO, TRST+)
CLOET	11	PR C83 012201	I.C. Cloet, G.A. Miller	(WASH)
DERUJULA	11	PL B697 26	A. de Rujula	(MADE, BOST, CERN)
DISTLER	11	PL B696 343	M.O. Distler, J.C. Bernauer, T. Walcher	(MANZ)
HILL	11	PRL 107 160402	R.J. Hill, G. Paz	(EFI)
BERNAUER	10	PRL 105 242001	J.C. Bernauer <i>et al.</i>	(MAMI A1 Collab.)
BORISYUK	10	NP A843 59	D. Borisyuk	(KIEV)
DERUJULA	10	PL B693 555	A. De Rujula	(MADU, CERN)
HILL	10	PR D82 113005	R.J. Hill, G. Paz	(CHIC)
POHL	10	NAT 466 213	R. Pohl <i>et al.</i>	(MPIQ, ENSP, COIM, +)
NISHINO	09	PRL 102 141801	H. Nishino <i>et al.</i>	(Super Kamiokande Collab.)
PASK	09	PL B678 55	T. Pask <i>et al.</i>	(Stefan Meyer Inst., Vienna, TOKY+)
MOHR	08	RMP 80 633	P.J. Mohr, B.N. Taylor, D.B. Newell	(NIST)
BELUSHKIN	07	PR C75 035202	M.A. Belushkin, H.W. Hammer, U.-G. Meissner	(BONN+)
ARAKI	06	PRL 96 101802	T. Araki <i>et al.</i>	(KamLAND Collab.)
HORI	06	PRL 96 243401	M. Hori <i>et al.</i>	(CERN, TOKYO+)
BLUNDEN	05	PR C72 057601	P.G. Blunden, I. Sick	(MANI, BASL)
KOBAYASHI	05	PR D72 052007	K. Kobayashi <i>et al.</i>	(Super-Kamiokande Collab.)
MOHR	05	RMP 77 1	P.J. Mohr, B.N. Taylor	(NIST)
SCHUMACHER	05	PPNP 55 567	M. Schumacher	(GOET)
AHMED	04	PRL 92 102004	S.N. Ahmed <i>et al.</i>	(SNO Collab.)
TRETYAK	04	JETPL 79 106	V.I. Tretyak, V.Yu. Denisov, Yu.G. Zdesenko	(KIEV)
BACK	03	PL B563 23	Translated from ZETFP 79 136. H.O. Back <i>et al.</i>	(BOREXINO Collab.)
BEANE	03	PL B567 200	S.R. Beane <i>et al.</i>	
Also		PL B607 320 (errata)	S.R. Beane <i>et al.</i>	
DMITRIEV	03	PRL 91 212303	V.F. Dmitriev, R.A. Senkov	(NOVO)
HORI	03	PRL 91 123401	M. Hori <i>et al.</i>	(CERN ASACUSA Collab.)
SICK	03	PL B576 62	I. Sick	(BASL)
ZDESENKO	03	PL B553 135	Yu.G. Zdesenko, V.I. Tretyak	(KIEV)
AHMAD	02	PRL 89 011301	Q.R. Ahmad <i>et al.</i>	(SNO Collab.)
BARANOV	01	PPN 32 376	P.S. Baranov <i>et al.</i>	
			Translated from FECAY 32 699.	

BLANPIED	01	PR C64 025203	G. Blanpied <i>et al.</i>	(BNL LEGS Collab.)
ESCHRICH	01	PL B522 233	I. Eschrich <i>et al.</i>	(FNAL SELEX Collab.)
HORI	01	PRL 87 093401	M. Hori <i>et al.</i>	(CERN ASACUSA Collab.)
OLMOSDEL...	01	EPJ A10 207	V. Olmos de Leon <i>et al.</i>	(MAMI TAPS Collab.)
TRETYAK	01	PL B505 59	V.I. Tretyak, Yu.G. Zdesenko	(KIEV)
BERNABEI	00B	PL B493 12	R. Bernabei <i>et al.</i>	(Gran Sasso DAMA Collab.)
GEER	00	PRL 84 590	S. Geer <i>et al.</i>	(FNAL APEX Collab.)
Also		PR D62 052004	S. Geer <i>et al.</i>	(FNAL APEX Collab.)
Also		PRL 85 3546 (errata)	S. Geer <i>et al.</i>	(FNAL APEX Collab.)
GEER	00C	PRL 85 3546 (errata)	S. Geer <i>et al.</i>	(FNAL APEX Collab.)
GEER	00D	APJ 532 648	S.H. Geer, D.C. Kennedy	
MELNIKOV	00	PRL 84 1673	K. Melnikov <i>et al.</i>	(SLAC, KARL)
ROSENFELDR...	00	PL B479 381	R. Rosenfelder	
SENGUPTA	00	PL B484 275	S. Sengupta	
WALL	00	PR D61 072004	D. Wall <i>et al.</i>	(Soudan-2 Collab.)
WALL	00B	PR D62 092003	D. Wall <i>et al.</i>	(Soudan-2 Collab.)
GABRIELSE	99	PRL 82 3198	G. Gabrielse <i>et al.</i>	
HAYATO	99	PRL 83 1529	Y. Hayato <i>et al.</i>	(Super-Kamiokande Collab.)
MCGREW	99	PR D59 052004	C. McGrew <i>et al.</i>	(IMB-3 Collab.)
MOHR	99	JPCRD 28 1713	P.J. Mohr, B.N. Taylor	(NIST)
Also		RMP 72 351	P.J. Mohr, B.N. Taylor	(NIST)
TORII	99	PR A59 223	H.A. Torii <i>et al.</i>	(CERN PS-205 Collab.)
ALLISON	98	PL B427 217	W.W.M. Allison <i>et al.</i>	(Soudan-2 Collab.)
HU	98B	PR D58 111101	M. Hu <i>et al.</i>	(FNAL APEX Collab.)
SHIOZAWA	98	PRL 81 3319	M. Shiozawa <i>et al.</i>	(Super-Kamiokande Collab.)
GLICENSTEIN	97	PL B411 326	J.F. Glicenstein	(SACL)
MERGELL	96	NP A596 367	P. Mergell <i>et al.</i>	(MANZ, BONN)
GABRIELSE	95	PRL 74 3544	G. Gabrielse <i>et al.</i>	(HARV, MANZ, SEOUL)
MACGIBBON	95	PR C52 2097	B.E. MacGibbon <i>et al.</i>	(ILL, SASK, INRM)
GEER	94	PRL 72 1596	S. Geer <i>et al.</i>	(FNAL, UCLA, PSU)
WONG	94	IJMP E3 821	C.W. Wong	(UCLA)
HALLIN	93	PR C48 1497	E.L. Hallin <i>et al.</i>	(SASK, BOST, ILL)
SUZUKI	93B	PL B311 357	Y. Suzuki <i>et al.</i>	(KAMIOKANDE Collab.)
HUGHES	92	PRL 69 578	R.J. Hughes, B.I. Deutch	(LANL, AARH)
ZIEGER	92	PL B278 34	A. Zieger <i>et al.</i>	(MPCM)
Also		PL B281 417 (erratum)	A. Zieger <i>et al.</i>	(MPCM)
BERGER	91	ZPHY C50 385	C. Berger <i>et al.</i>	(FREJUS Collab.)
BERGER	91B	PL B269 227	C. Berger <i>et al.</i>	(FREJUS Collab.)
FEDERSPIEL	91	PRL 67 1511	F.J. Federspiel <i>et al.</i>	(ILL)
MCCORD	91	NIM B56/57 496	M. McCord <i>et al.</i>	
BECKER-SZ...	90	PR D42 2974	R.A. Becker-Szendy <i>et al.</i>	(IMB-3 Collab.)
ERICSON	90	EPL 11 295	T.E.O. Ericson, A. Richter	(CERN, DARM)
GABRIELSE	90	PRL 65 1317	G. Gabrielse <i>et al.</i>	(HARV, MANZ, WASH+)
BERGER	89	NP B313 509	C. Berger <i>et al.</i>	(FREJUS Collab.)
CHO	89	PRL 63 2559	D. Cho, K. Sangster, E.A. Hinds	(YALE)
HIRATA	89C	PL B220 308	K.S. Hirata <i>et al.</i>	(Kamiokande Collab.)
PHILLIPS	89	PL B224 348	T.J. Phillips <i>et al.</i>	(HPW Collab.)
KREISSL	88	ZPHY C37 557	A. Kreissl <i>et al.</i>	(CERN PS176 Collab.)
SEIDEL	88	PRL 61 2522	S. Seidel <i>et al.</i>	(IMB Collab.)
BARTELT	87	PR D36 1990	J.E. Bartelt <i>et al.</i>	(Soudan Collab.)
Also		PR D40 1701 (erratum)	J.E. Bartelt <i>et al.</i>	(Soudan Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HAINES	86	PRL 57 1986	T.J. Haines <i>et al.</i>	(IMB Collab.)
KAJITA	86	JPSJ 55 711	T. Kajita <i>et al.</i>	(Kamiokande Collab.)
ARISAKA	85	JPSJ 54 3213	K. Arisaka <i>et al.</i>	(Kamiokande Collab.)
BLEWITT	85	PRL 55 2114	G.B. Blewitt <i>et al.</i>	(IMB Collab.)
DZUBA	85	PL 154B 93	V.A. Dzuba, V.V. Flambaum, P.G. Silvestrov	(NOVO)
PARK	85	PRL 54 22	H.S. Park <i>et al.</i>	(IMB Collab.)
BATTISTONI	84	PL 133B 454	G. Battistoni <i>et al.</i>	(NUSEX Collab.)
MARINELLI	84	PL 137B 439	M. Marinelli, G. Morpurgo	(GENO)
WILKENING	84	PR A29 425	D.A. Wilkening, N.F. Ramsey, D.J. Larson	(HARV+)
BARTELT	83	PRL 50 651	J.E. Bartelt <i>et al.</i>	(MINN, ANL)
BATTISTONI	82	PL 118B 461	G. Battistoni <i>et al.</i>	(NUSEX Collab.)
KRISHNA...	82	PL 115B 349	M.R. Krishnaswamy <i>et al.</i>	(TATA, OSKC+)
ALEKSEEV	81	JETPL 33 651	E.N. Alekseev <i>et al.</i>	(PNPI)

Translated from ZETFP 33 664.

CHERRY	81	PRL 47 1507	M.L. Cherry <i>et al.</i>	(PENN, BNL)
COWSIK	80	PR D22 2204	R. Cowsik, V.S. Narasimham	(TATA)
SIMON	80	NP A333 381	G.G. Simon <i>et al.</i>	
BELL	79	PL 86B 215	M. Bell <i>et al.</i>	(CERN)
GOLDEN	79	PRL 43 1196	R.L. Golden <i>et al.</i>	(NASA, PSLL)
LEARNED	79	PRL 43 907	J.G. Learned, F. Reines, A. Soni	(UCI)
BREGMAN	78	PL 78B 174	M. Bregman <i>et al.</i>	(CERN)
ROBERTS	78	PR D17 358	B.L. Roberts	(WILL, RHEL)
EVANS	77	SCI 197 989	J.C. Evans Jr., R.I. Steinberg	(BNL, PENN)
HU	75	NP A254 403	E. Hu <i>et al.</i>	(COLU, YALE)
BORKOWSKI	74	NP A222 269	F. Borkowski <i>et al.</i>	
COHEN	73	JPCRD 2 664	E.R. Cohen, B.N. Taylor	(RISC, NBS)
DYLLA	73	PR A7 1224	H.F. Dylla, J.G. King	(MIT)
AKIMOV	72	JETP 35 651	Yu.K. Akimov <i>et al.</i>	(YERE)
		Translated from ZETF 62 1231.		
DIX	70	Thesis Case	F.W. Dix	(CASE)
HARRISON	69	PRL 22 1263	G.E. Harrison, P.G.H. Sandars, S.J. Wright	(OXF)
GURR	67	PR 158 1321	H.S. Gurr <i>et al.</i>	(CASE, WITW)
FREREJACQ...	66	PR 141 1308	D. Frerejacque <i>et al.</i>	
HAND	63	RMP 35 335	L.N. Hand <i>et al.</i>	
FLEROV	58	DOKL 3 79	G.N. Flerov <i>et al.</i>	(ASCI)
