

$\pi_2(2100)$

$$I^G(J^{PC}) = 1^-(2^{-+})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

$\pi_2(2100)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2090 ± 29 OUR AVERAGE			
2090 ± 30	¹ AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
2100 ± 150	² DAUM	81B CNTR	63,94 $\pi^- p \rightarrow 3\pi X$

¹ From a fit to $J^{PC} = 2^{-+} f_2(1270)\pi, (\pi\pi)_S\pi$ waves.

² From a two-resonance fit to four 2^{-0^+} waves.

$\pi_2(2100)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
625 ± 50 OUR AVERAGE Error includes scale factor of 1.2.			
520 ± 100	³ AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
651 ± 50	⁴ DAUM	81B CNTR	63,94 $\pi^- p \rightarrow 3\pi X$

³ From a fit to $J^{PC} = 2^{-+} f_2(1270)\pi, (\pi\pi)_S\pi$ waves.

⁴ From a two-resonance fit to four 2^{-0^+} waves.

$\pi_2(2100)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 3π	seen
Γ_2 $\rho\pi$	seen
Γ_3 $f_2(1270)\pi$	seen
Γ_4 $(\pi\pi)_S\pi$	seen

$\pi_2(2100)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(3\pi)$	Γ_2/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.19 ± 0.05	⁵ DAUM 81B CNTR 63,94 $\pi^- p$
$\Gamma(f_2(1270)\pi)/\Gamma(3\pi)$	Γ_3/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.36 ± 0.09	⁵ DAUM 81B CNTR 63,94 $\pi^- p$
$\Gamma((\pi\pi)_S\pi)/\Gamma(3\pi)$	Γ_4/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.45 ± 0.07	⁵ DAUM 81B CNTR 63,94 $\pi^- p$

***D*-wave/*S*-wave RATIO FOR $\pi_2(2100) \rightarrow f_2(1270)\pi$**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.39 ± 0.23	⁵ DAUM	81B	CNTR 63,94 $\pi^- p$

⁵From a two-resonance fit to four 2^-0^+ waves.

$\pi_2(2100)$ REFERENCES

AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
