

B^\pm/B^0 ADMIXTURE

B DECAY MODES

The branching fraction measurements are for an admixture of B mesons at the $\Upsilon(4S)$. The values quoted assume that $B(\Upsilon(4S) \rightarrow B\bar{B}) = 100\%$.

For inclusive branching fractions, e.g., $B \rightarrow D^\pm$ anything, the treatment of multiple D 's in the final state must be defined. One possibility would be to count the number of events with one-or-more D 's and divide by the total number of B 's. Another possibility would be to count the total number of D 's and divide by the total number of B 's, which is the definition of average multiplicity. The two definitions are identical if only one D is allowed in the final state. Even though the “one-or-more” definition seems sensible, for practical reasons inclusive branching fractions are almost always measured using the multiplicity definition. For heavy final state particles, authors call their results inclusive branching fractions while for light particles some authors call their results multiplicities. In the B sections, we list all results as inclusive branching fractions, adopting a multiplicity definition. This means that inclusive branching fractions can exceed 100% and that inclusive partial widths can exceed total widths, just as inclusive cross sections can exceed total cross section.

\bar{B} modes are charge conjugates of the modes below. Reactions indicate the weak decay vertex and do not include mixing.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Semileptonic and leptonic modes		
$\Gamma_1 e^+ \nu_e$ anything	[a] (10.86 \pm 0.16) %	
$\Gamma_2 \bar{p} e^+ \nu_e$ anything	< 5.9 $\times 10^{-4}$ CL=90%	
$\Gamma_3 \mu^+ \nu_\mu$ anything	[a] (10.86 \pm 0.16) %	
$\Gamma_4 \ell^+ \nu_\ell$ anything	[a,b] (10.86 \pm 0.16) %	
$\Gamma_5 D^- \ell^+ \nu_\ell$ anything	[b] (2.8 \pm 0.9) %	
$\Gamma_6 \bar{D}^0 \ell^+ \nu_\ell$ anything	[b] (7.3 \pm 1.5) %	
$\Gamma_7 \bar{D} \ell^+ \nu_\ell$	(2.42 \pm 0.12) %	
$\Gamma_8 \bar{D} \tau^+ \nu_\tau$	(1.07 \pm 0.18) %	
$\Gamma_9 D^{*-} \ell^+ \nu_\ell$ anything	[c] (6.7 \pm 1.3) $\times 10^{-3}$	
$\Gamma_{10} D^{*0} \ell^+ \nu_\ell$ anything		
$\Gamma_{11} D^* \ell^+ \nu_\ell$	[d] (4.95 \pm 0.11) %	
$\Gamma_{12} D^* \tau^+ \nu_\tau$	(1.64 \pm 0.15) %	
$\Gamma_{13} \bar{D}^{**} \ell^+ \nu_\ell$	[b,e] (2.7 \pm 0.7) %	
$\Gamma_{14} \bar{D}_1(2420) \ell^+ \nu_\ell$ anything	(3.8 \pm 1.3) $\times 10^{-3}$	S=2.4
$\Gamma_{15} D \pi \ell^+ \nu_\ell$ anything + $D^* \pi \ell^+ \nu_\ell$ anything	(2.6 \pm 0.5) %	S=1.5
$\Gamma_{16} D \pi \ell^+ \nu_\ell$ anything	(1.5 \pm 0.6) %	

Γ_{17}	$D^* \pi \ell^+ \nu_\ell$ anything	(1.9 \pm 0.4) %
Γ_{18}	$\overline{D}_2^*(2460) \ell^+ \nu_\ell$ anything	(4.4 \pm 1.6) $\times 10^{-3}$
Γ_{19}	$D^{*-} \pi^+ \ell^+ \nu_\ell$ anything	(1.00 \pm 0.34) %
Γ_{20}	$D_s^- \ell^+ \nu_\ell$ anything	[b] < 7 $\times 10^{-3}$ CL=90%
Γ_{21}	$D_s^- \ell^+ \nu_\ell K^+$ anything	[b] < 5 $\times 10^{-3}$ CL=90%
Γ_{22}	$D_s^- \ell^+ \nu_\ell K^0$ anything	[b] < 7 $\times 10^{-3}$ CL=90%
Γ_{23}	$X_c \ell^+ \nu_\ell$	(10.65 \pm 0.16) %
Γ_{24}	$X_u \ell^+ \nu_\ell$	(2.14 \pm 0.31) $\times 10^{-3}$
Γ_{25}	$K^+ \ell^+ \nu_\ell$ anything	[b] (6.3 \pm 0.6) %
Γ_{26}	$K^- \ell^+ \nu_\ell$ anything	[b] (10 \pm 4) $\times 10^{-3}$
Γ_{27}	$K^0 / \overline{K}^0 \ell^+ \nu_\ell$ anything	[b] (4.6 \pm 0.5) %

 D , D^* , or D_s modes

Γ_{28}	D^\pm anything	(23.7 \pm 1.3) %
Γ_{29}	D^0 / \overline{D}^0 anything	(62.7 \pm 2.9) %
Γ_{30}	$D^*(2010)^\pm$ anything	(22.5 \pm 1.5) %
Γ_{31}	$D^*(2007)^0$ anything	(26.0 \pm 2.7) %
Γ_{32}	D_s^\pm anything	[f] (8.3 \pm 0.8) %
Γ_{33}	$D_s^{*\pm}$ anything	(6.3 \pm 1.0) %
Γ_{34}	$D_s^{*\pm} \overline{D}^{(*)}$	(3.4 \pm 0.6) %
Γ_{35}	$\overline{D} D_{s0}(2317)$	
Γ_{36}	$\overline{D} D_{sJ}(2457)$	
Γ_{37}	$D^{(*)} \overline{D}^{(*)} K^0 + D^{(*)} \overline{D}^{(*)} K^\pm$ [f,g]	(7.1 \pm 2.7) %
Γ_{38}	$b \rightarrow c \overline{c} s$	(22 \pm 4) %
Γ_{39}	$D_s^{(*)} \overline{D}^{(*)}$	[f,g] (3.9 \pm 0.4) %
Γ_{40}	$D^* D^*(2010)^\pm$	[f] < 5.9 $\times 10^{-3}$ CL=90%
Γ_{41}	$D D^*(2010)^\pm + D^* D^\pm$	[f] < 5.5 $\times 10^{-3}$ CL=90%
Γ_{42}	$D D^\pm$	[f] < 3.1 $\times 10^{-3}$ CL=90%
Γ_{43}	$D_s^{(*)\pm} \overline{D}^{(*)} X(n\pi^\pm)$	[f,g] (9 \pm 5) %
Γ_{44}	$D^*(2010) \gamma$	< 1.1 $\times 10^{-3}$ CL=90%
Γ_{45}	$D_s^+ \pi^-$, $D_s^{*+} \pi^-$, $D_s^+ \rho^-$, $D_s^{*+} \rho^-$, $D_s^+ \pi^0$, $D_s^{*+} \pi^0$, $D_s^+ \eta$, $D_s^{*+} \eta$, $D_s^+ \rho^0$, $D_s^{*+} \rho^0$, $D_s^+ \omega$, $D_s^{*+} \omega$	[f] < 4 $\times 10^{-4}$ CL=90%
Γ_{46}	$D_{s1}(2536)^+$ anything	< 9.5 $\times 10^{-3}$ CL=90%

Charmonium modes

Γ_{47}	$J/\psi(1S)$ anything	(1.094 \pm 0.032) %
Γ_{48}	$J/\psi(1S)$ (direct) anything	(7.8 \pm 0.4) $\times 10^{-3}$
Γ_{49}	$\psi(2S)$ anything	(3.07 \pm 0.21) $\times 10^{-3}$
Γ_{50}	$\chi_{c1}(1P)$ anything	(3.86 \pm 0.27) $\times 10^{-3}$
Γ_{51}	$\chi_{c1}(1P)$ (direct) anything	(3.24 \pm 0.25) $\times 10^{-3}$

Γ_{52}	$\chi_{c2}(1P)$ anything	(1.4 \pm 0.4) $\times 10^{-3}$	S=1.9
Γ_{53}	$\chi_{c2}(1P)$ (direct) anything	(1.65 \pm 0.31) $\times 10^{-3}$	
Γ_{54}	$\eta_c(1S)$ anything	< 9 $\times 10^{-3}$	CL=90%
Γ_{55}	$KX(3872) \times B(X \rightarrow D^0 \bar{D}^0 \pi^0)$	(1.2 \pm 0.4) $\times 10^{-4}$	
Γ_{56}	$KX(3872) \times B(X \rightarrow D^{*0} D^0)$	(8.0 \pm 2.2) $\times 10^{-5}$	
Γ_{57}	$KX(3940) \times B(X \rightarrow D^{*0} D^0)$	< 6.7 $\times 10^{-5}$	CL=90%
Γ_{58}	$K\chi_{c0}(2P), \chi_{c0} \rightarrow \omega J/\psi$	[h] (7.1 \pm 3.4) $\times 10^{-5}$	

 K or K^* modes

Γ_{59}	K^\pm anything	[f] (78.9 \pm 2.5) %	
Γ_{60}	K^+ anything	(66 \pm 5) %	
Γ_{61}	K^- anything	(13 \pm 4) %	
Γ_{62}	K^0 / \bar{K}^0 anything	[f] (64 \pm 4) %	
Γ_{63}	$K^*(892)^\pm$ anything	(18 \pm 6) %	
Γ_{64}	$K^*(892)^0 / \bar{K}^*(892)^0$ anything	[f] (14.6 \pm 2.6) %	
Γ_{65}	$K^*(892)\gamma$	(4.2 \pm 0.6) $\times 10^{-5}$	
Γ_{66}	$\eta K\gamma$	(8.5 \pm 1.8) $\times 10^{-6}$	
Γ_{67}	$K_1(1400)\gamma$	< 1.27 $\times 10^{-4}$	CL=90%
Γ_{68}	$K_2^*(1430)\gamma$	(1.7 \pm 0.6) $\times 10^{-5}$	
Γ_{69}	$K_2(1770)\gamma$	< 1.2 $\times 10^{-3}$	CL=90%
Γ_{70}	$K_3^*(1780)\gamma$	< 3.7 $\times 10^{-5}$	CL=90%
Γ_{71}	$K_4^*(2045)\gamma$	< 1.0 $\times 10^{-3}$	CL=90%
Γ_{72}	$K\eta'(958)$	(8.3 \pm 1.1) $\times 10^{-5}$	
Γ_{73}	$K^*(892)\eta'(958)$	(4.1 \pm 1.1) $\times 10^{-6}$	
Γ_{74}	$K\eta$	< 5.2 $\times 10^{-6}$	CL=90%
Γ_{75}	$K^*(892)\eta$	(1.8 \pm 0.5) $\times 10^{-5}$	
Γ_{76}	$K\phi\phi$	(2.3 \pm 0.9) $\times 10^{-6}$	
Γ_{77}	$\bar{b} \rightarrow \bar{s}\gamma$	(3.40 \pm 0.21) $\times 10^{-4}$	
Γ_{78}	$\bar{b} \rightarrow \bar{d}\gamma$	(9.2 \pm 3.0) $\times 10^{-6}$	
Γ_{79}	$\bar{b} \rightarrow \bar{s}$ gluon	< 6.8 %	CL=90%
Γ_{80}	η anything	(2.6 \pm 0.5) $\times 10^{-4}$	
Γ_{81}	η' anything	(4.2 \pm 0.9) $\times 10^{-4}$	
Γ_{82}	K^+ gluon (charmless)	< 1.87 $\times 10^{-4}$	CL=90%
Γ_{83}	K^0 gluon (charmless)	(1.9 \pm 0.7) $\times 10^{-4}$	

Light unflavored meson modes

Γ_{84}	$\rho\gamma$	(1.39 \pm 0.25) $\times 10^{-6}$	S=1.2
Γ_{85}	$\rho/\omega\gamma$	(1.30 \pm 0.23) $\times 10^{-6}$	S=1.2
Γ_{86}	π^\pm anything	[f,i] (358 \pm 7) %	
Γ_{87}	π^0 anything	(235 \pm 11) %	
Γ_{88}	η anything	(17.6 \pm 1.6) %	

Γ_{89}	ρ^0 anything	(21 \pm 5) %
Γ_{90}	ω anything	< 81 % CL=90%
Γ_{91}	ϕ anything	(3.43 \pm 0.12) %
Γ_{92}	$\phi K^*(892)$	< 2.2 $\times 10^{-5}$ CL=90%
Γ_{93}	$\bar{b} \rightarrow \bar{d}$ gluon	
Γ_{94}	π^+ gluon (charmless)	(3.7 \pm 0.8) $\times 10^{-4}$

Baryon modes

Γ_{95}	$\Lambda_c^+ / \bar{\Lambda}_c^-$ anything	(4.5 \pm 1.2) %
Γ_{96}	Λ_c^+ anything	< 1.7 % CL=90%
Γ_{97}	$\bar{\Lambda}_c^-$ anything	< 9 % CL=90%
Γ_{98}	$\bar{\Lambda}_c^- \ell^+$ anything	< 1.1 $\times 10^{-3}$ CL=90%
Γ_{99}	$\bar{\Lambda}_c^- e^+$ anything	< 2.3 $\times 10^{-3}$ CL=90%
Γ_{100}	$\bar{\Lambda}_c^- \mu^+$ anything	< - 1.8 $\times 10^{-3}$ CL=90%
Γ_{101}	$\bar{\Lambda}_c^- p$ anything	(2.6 \pm 0.8) %
Γ_{102}	$\bar{\Lambda}_c^- p e^+ \nu_e$	< 1.0 $\times 10^{-3}$ CL=90%
Γ_{103}	$\bar{\Sigma}_c^{--}$ anything	(4.2 \pm 2.4) $\times 10^{-3}$
Γ_{104}	$\bar{\Sigma}_c^-$ anything	< 9.6 $\times 10^{-3}$ CL=90%
Γ_{105}	$\bar{\Sigma}_c^0$ anything	(4.6 \pm 2.4) $\times 10^{-3}$
Γ_{106}	$\bar{\Sigma}_c^0 N$ ($N = p$ or n)	< 1.5 $\times 10^{-3}$ CL=90%
Γ_{107}	Ξ_c^0 anything	(1.93 \pm 0.30) $\times 10^{-4}$ S=1.1
	$\times B(\Xi_c^0 \rightarrow \Xi^- \pi^+)$	
Γ_{108}	Ξ_c^+ anything	(4.5 \pm 1.3) $\times 10^{-4}$
	$\times B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+)$	
Γ_{109}	p/\bar{p} anything	[f] (8.0 \pm 0.4) %
Γ_{110}	p/\bar{p} (direct) anything	[f] (5.5 \pm 0.5) %
Γ_{111}	$\Lambda/\bar{\Lambda}$ anything	[f] (4.0 \pm 0.5) %
Γ_{112}	Λ anything	
Γ_{113}	$\bar{\Lambda}$ anything	
Γ_{114}	$\Xi^-/\bar{\Xi}^+$ anything	[f] (2.7 \pm 0.6) $\times 10^{-3}$
Γ_{115}	baryons anything	(6.8 \pm 0.6) %
Γ_{116}	$p\bar{p}$ anything	(2.47 \pm 0.23) %
Γ_{117}	$\Lambda\bar{p}/\bar{\Lambda}p$ anything	[f] (2.5 \pm 0.4) %
Γ_{118}	$\Lambda\bar{\Lambda}$ anything	< 5 $\times 10^{-3}$ CL=90%

**Lepton Family number (*LF*) violating modes or
 $\Delta B = 1$ weak neutral current (*B1*) modes**

Γ_{119}	$s e^+ e^-$	<i>B1</i> (4.7 \pm 1.3) $\times 10^{-6}$
Γ_{120}	$s \mu^+ \mu^-$	<i>B1</i> (4.3 \pm 1.2) $\times 10^{-6}$
Γ_{121}	$s \ell^+ \ell^-$	<i>B1</i> [b] (4.5 \pm 1.0) $\times 10^{-6}$
Γ_{122}	$\pi \ell^+ \ell^-$	<i>B1</i> < 5.9 $\times 10^{-8}$ CL=90%
Γ_{123}	$\pi e^+ e^-$	< 1.10 $\times 10^{-7}$ CL=90%
Γ_{124}	$\pi \mu^+ \mu^-$	< 5.0 $\times 10^{-8}$ CL=90%

Γ_{125}	$K e^+ e^-$	<i>B1</i>	(4.4 \pm 0.6) $\times 10^{-7}$		
Γ_{126}	$K^*(892) e^+ e^-$	<i>B1</i>	(1.19 \pm 0.20) $\times 10^{-6}$	S=1.2	
Γ_{127}	$K \mu^+ \mu^-$	<i>B1</i>	(4.4 \pm 0.4) $\times 10^{-7}$		
Γ_{128}	$K^*(892) \mu^+ \mu^-$	<i>B1</i>	(1.06 \pm 0.09) $\times 10^{-6}$		
Γ_{129}	$K \ell^+ \ell^-$	<i>B1</i>	(4.8 \pm 0.4) $\times 10^{-7}$		
Γ_{130}	$K^*(892) \ell^+ \ell^-$	<i>B1</i>	(1.05 \pm 0.10) $\times 10^{-6}$		
Γ_{131}	$K \nu \bar{\nu}$	<i>B1</i>	< 1.7 $\times 10^{-5}$	CL=90%	
Γ_{132}	$K^* \nu \bar{\nu}$	<i>B1</i>	< 7.6 $\times 10^{-5}$	CL=90%	
Γ_{133}	$s e^\pm \mu^\mp$	<i>LF</i>	[f] < 2.2 $\times 10^{-5}$	CL=90%	
Γ_{134}	$\pi e^\pm \mu^\mp$	<i>LF</i>	< 9.2 $\times 10^{-8}$	CL=90%	
Γ_{135}	$\rho e^\pm \mu^\mp$	<i>LF</i>	< 3.2 $\times 10^{-6}$	CL=90%	
Γ_{136}	$K e^\pm \mu^\mp$	<i>LF</i>	< 3.8 $\times 10^{-8}$	CL=90%	
Γ_{137}	$K^*(892) e^\pm \mu^\mp$	<i>LF</i>	< 5.1 $\times 10^{-7}$	CL=90%	

[a] These values are model dependent.

[b] An ℓ indicates an e or a μ mode, not a sum over these modes.

[c] Here “anything” means at least one particle observed.

[d] This is a $B(B^0 \rightarrow D^{*-} \ell^+ \nu_\ell)$ value.

[e] D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances.

[f] The value is for the sum of the charge states or particle/antiparticle states indicated.

[g] $D^{(*)}\bar{D}^{(*)}$ stands for the sum of $D^*\bar{D}^*$, $D^*\bar{D}$, $D\bar{D}^*$, and $D\bar{D}$.

[h] $X(3915)$ denotes a near-threshold enhancement in the $\omega J/\psi$ mass spectrum.

[i] Inclusive branching fractions have a multiplicity definition and can be greater than 100%.

B^\pm/B^0 ADMIXTURE BRANCHING RATIOS

$\Gamma(e^+ \nu_e \text{anything})/\Gamma_{\text{total}}$

These branching fraction values are model dependent.

Γ_1/Γ

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

VALUE	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

0.1086 \pm 0.0016 OUR EVALUATION

0.1044 \pm 0.0025 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

0.1028 \pm 0.0018 \pm 0.0024	¹ URQUIJO	07	BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.0996 \pm 0.0019 \pm 0.0032	² AUBERT,B	06Y	BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.1091 \pm 0.0009 \pm 0.0024	³ MAHMOOD	04	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.097 \pm 0.005 \pm 0.004	⁴ ALBRECHT	93H	ARG	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1085 \pm 0.0021 \pm 0.0036	⁵ OKABE	05	BELL	Repl. by URQUIJO 07
0.1083 \pm 0.0016 \pm 0.0006	⁶ AUBERT	04X	BABR	Repl. by AUBERT,B 06Y
0.1036 \pm 0.0006 \pm 0.0023	⁷ AUBERT,B	04A	BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.1087 \pm 0.0018 \pm 0.0030	⁸ AUBERT	03	BABR	Repl. by AUBERT 04X
0.109 \pm 0.0012 \pm 0.0049	⁹ ABE	02Y	BELL	Repl. by OKABE 05
0.1049 \pm 0.0017 \pm 0.0043	¹⁰ BARISH	96B	CLE2	Repl. by MAHMOOD 04
0.100 \pm 0.004 \pm 0.003	¹¹ YANAGISAWA	91	CSB2	$e^+ e^- \rightarrow \gamma(4S)$
0.103 \pm 0.006 \pm 0.002	¹² ALBRECHT	90H	ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.117 \pm 0.004 \pm 0.010	¹³ WACHS	89	CBAL	Direct e at $\gamma(4S)$
0.120 \pm 0.007 \pm 0.005	CHEN	84	CLEO	Direct e at $\gamma(4S)$
0.132 \pm 0.008 \pm 0.014	¹⁴ KLOPFEN...	83B	CUSB	Direct e at $\gamma(4S)$

¹ URQUIJO 07 report a measurement of $(10.07 \pm 0.18 \pm 0.21)\%$ for the partial branching fraction of $B \rightarrow e\nu_e X_c$ decay with electron energy above 0.6 GeV. We converted the result to $B \rightarrow e\nu_e X$ branching fraction.

² The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame. The best precision on the ratio is achieved for a momentum threshold of 1.0 GeV: $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.074 \pm 0.041 \pm 0.026$.

³ Uses charge and angular correlations in $\gamma(4S)$ events with a high-momentum lepton and an additional electron.

⁴ ALBRECHT 93H analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.

⁵ The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame, and their ratio of $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.08 \pm 0.05 \pm 0.02$.

⁶ The semileptonic branching ratio, $|V_{cb}|$ and other heavy-quark parameters are determined from a simultaneous fit to moments of the hadronic-mass and lepton-energy distribution.

⁷ Uses the high-momentum lepton tag method and requires the electron energy above 0.6 GeV.

⁸ Uses the high-momentum lepton tag method. They also report $|V_{cb}| = 0.0423 \pm 0.0007(\text{exp}) \pm 0.0020(\text{theo.})$.

⁹ Uses the high-momentum lepton tag method. ABE 02Y also reports $|V_{cb}| = 0.0408 \pm 0.0010(\text{exp}) \pm 0.0025(\text{theo.})$. The second error is due to uncertainties of theoretical inputs.


¹⁰ BARISH 96B analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.

¹¹ YANAGISAWA 91 also measures an average semileptonic branching ratio at the $\gamma(5S)$ of 9.6–10.5% depending on assumptions about the relative production of different B meson species.

¹² ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.099 ± 0.006 is obtained using ISGUR 89B.

¹³ Using data above $p(e) = 2.4$ GeV, WACHS 89 determine $\sigma(B \rightarrow e\nu \text{ up}) / \sigma(B \rightarrow e\nu \text{ charm}) < 0.065$ at 90% CL.

¹⁴ Ratio $\sigma(b \rightarrow e\nu \text{ up}) / \sigma(b \rightarrow e\nu \text{ charm}) < 0.055$ at CL = 90%.



$\Gamma(\bar{p} e^+ \nu_e \text{ anything})/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.9 \times 10^{-4}$	90	¹ ADAM	03B	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0016	90	ALBRECHT	90H	$e^+ e^- \rightarrow \gamma(4S)$

¹ Based on $V-A$ model.

$\Gamma(\mu^+ \nu_\mu \text{ anything})/\Gamma_{\text{total}}$

Γ_3/Γ

These branching fraction values are model dependent.

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

VALUE	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

0.1086±0.0016 OUR EVALUATION

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.100 ± 0.006 ± 0.002	¹ ALBRECHT	90H	ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.108 ± 0.006 ± 0.01	CHEN	84	CLEO	Direct μ at $\gamma(4S)$
0.112 ± 0.009 ± 0.01	LEVMAN	84	CUSB	Direct μ at $\gamma(4S)$

¹ ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.097 ± 0.006 is obtained using ISGUR 89B.

$\Gamma(\ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

Γ_4/Γ


These branching fraction values are model dependent.

“OUR EVALUATION” is an average using rescaled values of the data listed below.

The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1086 ± 0.0016 OUR EVALUATION			
0.1044 ± 0.0025 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.5. See the ideogram below.			
$0.108 \pm 0.002 \pm 0.0056$	¹ HENDERSON 92 CLEO $e^+ e^- \rightarrow \gamma(4S)$		• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ HENDERSON 92 measurement employs e and μ . The systematic error contains 0.004 in quadrature from model dependence. The authors average a variation of the Isgur, Scora, Grinstein, and Wise model with that of the Altarelli-Cabibbo-Corbò-Maiani-Martinelli model for semileptonic decays to correct the acceptance.



$\Gamma(D^- \ell^+ \nu_\ell \text{anything})/\Gamma(\ell^+ \nu_\ell \text{anything})$

Γ_5/Γ_4

$\ell = e \text{ or } \mu$.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.26 \pm 0.07 \pm 0.04$			

¹ FULTON 91 uses $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.1 \pm 1.3 \pm 0.4)\%$ as measured by MARK III.

$\Gamma(\overline{D^0}\ell^+\nu_\ell\text{anything})/\Gamma(\ell^+\nu_\ell\text{anything})$
 $\ell = e \text{ or } \mu.$
 Γ_6/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
0.67±0.09±0.10	1 FULTON 91	CLEO	$e^+e^- \rightarrow \gamma(4S)$

¹ FULTON 91 uses $B(D^0 \rightarrow K^-\pi^+) = (4.2 \pm 0.4 \pm 0.4)\%$ as measured by MARK III.

 $\Gamma(\overline{D}\ell^+\nu_\ell)/\Gamma(\ell^+\nu_\ell\text{anything})$
 Γ_7/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
0.223±0.006±0.009	1 AUBERT 10	BABR	$e^+e^- \rightarrow \gamma(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

 $\Gamma(\overline{D}\tau^+\nu_\tau)/\Gamma(\overline{D}\ell^+\nu_\ell)$
 Γ_8/Γ_7

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
44.0 ± 5.8±4.2	1,2 LEES 12D	BABR	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.16±11.7±5.2 1 AUBERT 08N BABR Repl. by LEES 12D

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² Uses $\tau^+ \rightarrow e^+\nu_e\bar{\nu}_\tau$ and $\tau^+ \rightarrow \mu^+\nu_\mu\bar{\nu}_\tau$ and e^+ or μ^+ as ℓ^+ . Obtained from simultaneous fit to B^+ and B^0 assuming isospin symmetry.

 $\Gamma(D^{*-}\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$
 Γ_9/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
0.67±0.08±0.10	ABDALLAH 04D	DLPH	$e^+e^- \rightarrow Z^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.3 ± 0.1 1 BARISH 95 CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹ BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*+} \rightarrow D^0\pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$.

 $\Gamma(D^{*0}\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$
 Γ_{10}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

0.6±0.6±0.1 1 BARISH 95 CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹ BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$, $B(D^{*+} \rightarrow D^0\pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$, $B(D^{*0} \rightarrow D^0\pi^0) = (63.6 \pm 2.3 \pm 3.3)\%$.

 $\Gamma(D^{*+}\tau^+\nu_\tau)/\Gamma(D^{*+}\ell^+\nu_\ell)$
 Γ_{12}/Γ_{11}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
33.2±2.4±1.8	1 LEES 12D	BABR	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

29.7±5.6±1.8 2 AUBERT 08N BABR Repl. by LEES 12D

¹ Uses $\tau^+ \rightarrow e^+\nu_e\bar{\nu}_\tau$ and $\tau^+ \rightarrow \mu^+\nu_\mu\bar{\nu}_\tau$ and e^+ or μ^+ as ℓ^+ . Obtained from simultaneous fit to B^+ and B^0 assuming isospin symmetry. Uses a fully reconstructed B meson as a tag on the recoil side.

² Uses a fully reconstructed B meson as a tag on the recoil side. The results are normalized to the B^+ decay rate.

$\Gamma(\overline{D}^{**}\ell^+\nu_\ell)/\Gamma_{\text{total}}$ Γ_{13}/Γ

D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances. $\ell = e$ or μ , not sum over e and μ modes.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.027±0.005±0.005	63	1 ALBRECHT	93	ARG	$e^+e^- \rightarrow \gamma(4S)$
<0.028	95	2 BARISH	95	CLE2	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ALBRECHT 93 assumes the GISW model to correct for unseen modes. Using the BHKT model, the result becomes $0.023 \pm 0.006 \pm 0.004$. Assumes $B(D^{*+} \rightarrow D^0\pi^+) = 68.1\%$, $B(D^0 \rightarrow K^-\pi^+) = 3.65\%$, $B(D^0 \rightarrow K^-\pi^+\pi^-\pi^+) = 7.5\%$. We have taken their average e and μ value.

² BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$, assume all nonresonant channels are zero, and use GISW model for relative abundances of D^{**} states.

 $\Gamma(\overline{D}_1(2420)\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0038±0.0013 OUR AVERAGE			Error includes scale factor of 2.4.
0.0033±0.0006	1 ABAZOV	050	$D^0 p\bar{p}$ at 1.96 TeV
0.0074±0.0016	2 BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$
seen	3 BUSKULIC	95B ALEP	Repl. by BUSKULIC 97B

¹ Assumes $B(D_1 \rightarrow D^*\pi) = 1$, $B(D_1 \rightarrow D^*\pi^\pm) = 2/3$, and $B(b \rightarrow B) = 0.397$.

² BUSKULIC 97B assumes $B(D_1(2420) \rightarrow D^*\pi) = 1$, $B(D_1(2420) \rightarrow D^*\pi^\pm) = 2/3$, and $B(b \rightarrow B) = 0.378 \pm 0.022$.

³ BUSKULIC 95B reports $f_B \times B(B \rightarrow \overline{D}_1(2420)^0\ell^+\nu_\ell\text{anything}) \times B(\overline{D}_1(2420)^0 \rightarrow \overline{D}^*(2010)^-\pi^+) = (2.04 \pm 0.58 \pm 0.34)10^{-3}$, where f_B is the production fraction for a single B charge state.

 $[\Gamma(D\pi\ell^+\nu_\ell\text{anything}) + \Gamma(D^*\pi\ell^+\nu_\ell\text{anything})]/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.026 ± 0.005 OUR AVERAGE			Error includes scale factor of 1.5.
0.0340±0.0052±0.0032	1 ABREU	00R DLPFH	$e^+e^- \rightarrow Z$
0.0226±0.0029±0.0033	2 BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$

¹ Assumes no contribution from B_s and b baryons. Further assumes contributions from single pion ($D\pi$ and $D^*\pi$) states only, allowing isospin conservation to relate the relative π^0 and π^+ rates.

² BUSKULIC 97B assumes $B(b \rightarrow B) = 0.378 \pm 0.022$ and uses isospin invariance by assuming that all observed $D^0\pi^+$, $D^{*0}\pi^+$, $D^+\pi^-$, and $D^{*+}\pi^-$ are from D^{**} states. A correction has been applied to account for the production of B_s^0 and Λ_b^0 .

 $\Gamma(D\pi\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0154±0.0061	ABREU	00R DLPFH	$e^+e^- \rightarrow Z$

 $\Gamma(D^*\pi\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0186±0.0038	ABREU	00R DLPFH	$e^+e^- \rightarrow Z$

$\Gamma(X_c \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_{23}/Γ

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1065±0.0016 OUR EVALUATION			
0.1058±0.0015 OUR AVERAGE			
0.1064±0.0017±0.0006	¹ AUBERT	10A	BABR $e^+ e^- \rightarrow \gamma(4S)$
0.1044±0.0019±0.0022	² URQUIJO	07	BELL $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.1061±0.0016±0.0006	³ AUBERT	04X	BABR Repl. by AUBERT 10A
1 Obtained from a combined fit to the moments of observed spectra in inclusive $B \rightarrow X_c \ell^+ \nu_\ell$ decay.			
2 Measured the independent B^+ and B^0 partial branching fractions with electron energy above 0.4 GeV.			
3 The semileptonic branching ratio, $ V_{cb} $ and other heavy-quark parameters are determined from a simultaneous fit to moments of the hadronic-mass and lepton-energy distribution.			

 $\Gamma(X_u \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_{24}/Γ

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.14 ±0.31 OUR EVALUATION			
2.01 ±0.15 ±0.25	¹ LEES	12R	BABR $e^+ e^- \rightarrow \gamma(4S)$
2.27 ±0.26 ±0.37	² AUBERT	06H	BABR $e^+ e^- \rightarrow \gamma(4S)$
2.53 ±0.24 ±0.24	³ AUBERT,B	05X	BABR $e^+ e^- \rightarrow \gamma(4S)$
2.80 ±0.52 ±0.41	⁴ LIMOSANI	05	BELL $e^+ e^- \rightarrow \gamma(4S)$
1.77 ±0.29 ±0.38	⁵ BORNHEIM	02	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.963±0.173±0.159	⁶ URQUIJO	10	BELL $e^+ e^- \rightarrow \gamma(4S)$
1.18 ±0.09 ±0.07	⁷ AUBERT	08AS	BABR Repl. by LEES 12R
2.24 ±0.27 ±0.47	^{8,9} AUBERT	04I	BABR Repl. by AUBERT,B 05X

1 Measures several partial branching fractions in different phase space regions. The most precise result on the full branching fraction is obtained in the region for lepton momentum in B rest frame $p_\ell^* > 1$ GeV/c, where the measured partial branching fraction is $\Delta B = (1.80 \pm 0.13 \pm 0.15) \times 10^{-3}$. The acceptance in that region is reported in a private communication by the Authors to be 0.894. The corresponding $|V_{ub}|$ from the BLNP method is $(4.28 \pm 0.15 \pm 0.18 \pm 0.19) \times 10^{-3}$, where the last uncertainty comes from theoretical prediction.

2 Obtained from the partial rate $\Delta B = (0.572 \pm 0.041 \pm 0.065) \times 10^{-3}$ for the electron momentum interval of 2.0–2.6 GeV/c based on BLNP method.

3 Determined from the partial rate $\Delta B = (4.41 \pm 0.42 \pm 0.42) \times 10^{-4}$ measured for electron energy > 2 GeV and hadronic mass squared < 3.5 GeV 2 , and calculated acceptance 0.174 in that region. The V_{ub} is measured as $(4.41 \pm 0.30^{+0.65}_{-0.47} \pm 0.28) \times 10^{-3}$.

- ⁴ Uses electrons in the momentum interval 1.9–2.6 GeV/c in the center-of-mass frame. The V_{ub} is found to be $(5.08 \pm 0.47^{+0.49}_{-0.48}) \times 10^{-3}$.
- ⁵ BORNHEIM 02 uses the observed yield of leptons from semileptonic B decays in the end-point momentum interval 2.2–2.6 GeV/c with recent CLEO-2 data on $B \rightarrow X_s \gamma$. The V_{ub} is found to be $(4.08 \pm 0.34 \pm 0.53) \times 10^{-3}$.
- ⁶ Uses a multivariate analysis method and requires lepton momentum in the B rest frame, $p_l^* B > 1.0$ GeV/c.
- ⁷ Measures several partial branching fractions in different phase space regions. The most precise result is obtained in the region for hadronic mass $M_X < 1.55$ GeV/c², and is $\Delta B = (1.18 \pm 0.09 \pm 0.07) \times 10^{-3}$. The corresponding $|V_{ub}|$ from the BLNP method is $(4.27 \pm 0.16 \pm 0.13 \pm 0.30) \times 10^{-3}$, where the last uncertainty comes from the theoretical prediction of the partial rate in the given phase-space region.
- ⁸ Used BaBar measurement of Semileptonic branching fraction $B(B \rightarrow X \ell \nu_\ell) = (10.87 \pm 0.18 \pm 0.30)\%$ to convert the ratio of rates to branching fraction.
- ⁹ The third error includes the systematics and theoretical errors summed in quadrature.

$\Gamma(X_u \ell^+ \nu_\ell)/\Gamma(\ell^+ \nu_\ell \text{anything})$

Γ_{24}/Γ_4

ℓ denotes e or μ , not the sum. These experiments measure this ratio in very limited momentum intervals.

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.06±0.25±0.42			¹ AUBERT	04I BABR	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
			² ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$
107			³ BARTEL	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
77			⁴ ALBRECHT	91C ARG	$e^+ e^- \rightarrow \gamma(4S)$
41			⁵ ALBRECHT	90 ARG	$e^+ e^- \rightarrow \gamma(4S)$
76			⁶ FULTON	90 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
<4.0	90		⁷ BEHRENDS	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
<4.0	90		CHEN	84 CLEO	Direct e at $\gamma(4S)$
<5.5	90		KLOPFEN...	83B CUSB	Direct e at $\gamma(4S)$

¹ The third error includes the systematics and theoretical errors summed in quadrature.

² ALBRECHT 94C find $\Gamma(b \rightarrow c)/\Gamma(b \rightarrow \text{all}) = 0.99 \pm 0.02 \pm 0.04$.

³ BARTEL 93B (CLEO II) measures an excess of $107 \pm 15 \pm 11$ leptons in the lepton momentum interval 2.3–2.6 GeV/c which is attributed to $b \rightarrow u \ell \nu_\ell$. This corresponds to a model-dependent partial branching ratio ΔB_{ub} between $(1.15 \pm 0.16 \pm 0.15) \times 10^{-4}$, as evaluated using the KS model (KOERNER 88), and $(1.54 \pm 0.22 \pm 0.20) \times 10^{-4}$ using the ACCMM model (ARTUSO 93). The corresponding values of $|V_{ub}|/|V_{cb}|$ are 0.056 ± 0.006 and 0.076 ± 0.008 , respectively.

⁴ ALBRECHT 91C result supersedes ALBRECHT 90. Two events are fully reconstructed providing evidence for the $b \rightarrow u$ transition. Using the model of ALTARELLI 82, they obtain $|V_{ub}/V_{cb}| = 0.11 \pm 0.012$ from 77 leptons in the 2.3–2.6 GeV momentum range.

⁵ ALBRECHT 90 observes 41 ± 10 excess e and μ (lepton) events in the momentum interval $p = 2.3$ –2.6 GeV signaling the presence of the $b \rightarrow u$ transition. The events correspond to a model-dependent measurement of $|V_{ub}/V_{cb}| = 0.10 \pm 0.01$.

⁶ FULTON 90 observe 76 ± 20 excess e and μ (lepton) events in the momentum interval $p = 2.4$ –2.6 GeV signaling the presence of the $b \rightarrow u$ transition. The average branching ratio, $(1.8 \pm 0.4 \pm 0.3) \times 10^{-4}$, corresponds to a model-dependent measurement of approximately $|V_{ub}/V_{cb}| = 0.1$ using $B(b \rightarrow c \ell \nu) = 10.2 \pm 0.2 \pm 0.7\%$.

⁷ The quoted possible limits range from 0.018 to 0.04 for the ratio, depending on which model or momentum range is chosen. We select the most conservative limit they have calculated. This corresponds to a limit on $|V_{ub}|/|V_{cb}| < 0.20$. While the endpoint technique employed is more robust than their previous results in CHEN 84, these results do not provide a numerical improvement in the limit.

¹ GIBBONS 97B reports $[\Gamma(B \rightarrow D^\pm \text{anything})/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = 0.0216 \pm 0.0008 \pm 0.00082$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.13 \pm 0.19) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BORTOLETTO 92 reports $[\Gamma(B \rightarrow D^\pm \text{anything})/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = 0.0226 \pm 0.0030 \pm 0.0018$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.13 \pm 0.19) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 91H reports $[\Gamma(B \rightarrow D^\pm \text{anything})/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = 0.0209 \pm 0.0027 \pm 0.0040$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.13 \pm 0.19) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ BORTOLETTO 87 reports $[\Gamma(B \rightarrow D^\pm \text{anything})/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = 0.019 \pm 0.004 \pm 0.002$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.13 \pm 0.19) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^0/\bar{D}^0 \text{anything})/\Gamma_{\text{total}}$

Γ_{29}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.627 ± 0.029 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.647 ± 0.025	^{+0.007} _{-0.008}	1 GIBBONS	97B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.60	± 0.05	2 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.50	± 0.08	3 ALBRECHT	91H ARG	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.54	± 0.07	21k	4 BORTOLETTO87	CLEO $e^+ e^- \rightarrow \gamma(4S)$
0.62	± 0.19	± 0.01	5 GREEN	83 CLEO Repl. by BORTOLETTO 87


¹ GIBBONS 97B reports $[\Gamma(B \rightarrow D^0/\bar{D}^0 \text{anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = 0.0251 \pm 0.0006 \pm 0.00075$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.88 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BORTOLETTO 92 reports $[\Gamma(B \rightarrow D^0/\bar{D}^0 \text{anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = 0.0233 \pm 0.0012 \pm 0.0014$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.88 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 91H reports $[\Gamma(B \rightarrow D^0/\bar{D}^0 \text{anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = 0.0194 \pm 0.0015 \pm 0.0025$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.88 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ BORTOLETTO 87 reports $[\Gamma(B \rightarrow D^0/\bar{D}^0 \text{anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = 0.0210 \pm 0.0015 \pm 0.0021$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.88 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ GREEN 83 reports $[\Gamma(B \rightarrow D^0/\bar{D}^0 \text{anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = 0.024 \pm 0.006 \pm 0.004$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.88 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(D^*(2010)^{\pm} \text{anything})/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.225±0.015 OUR AVERAGE				
0.247±0.019±0.01	1	GIBBONS 97B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.205±0.019±0.007	2	ALBRECHT 96D	ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.230±0.028±0.009	3	BORTOLETTO 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.283±0.053±0.002	4	ALBRECHT 91H	ARG	Sup. by ALBRECHT 96D
0.22 ± 0.04 ± 0.07 - 0.04	5200	5 BORTOLETTO 87	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.27 ± 0.06 ± 0.08 - 0.06	510	6 CSORNA 85	CLEO	Repl. by BORTOLETTO 87

¹ GIBBONS 97B reports $B(B \rightarrow D^*(2010)^+ \text{anything}) = 0.239 \pm 0.015 \pm 0.014 \pm 0.009$ using CLEO measured D and D^* branching fractions. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ALBRECHT 96D reports $B(B \rightarrow D^*(2010)^+ \text{anything}) = 0.196 \pm 0.019$ using CLEO measured $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.681 \pm 0.01 \pm 0.013$, $B(D^0 \rightarrow K^- \pi^+) = 0.0401 \pm 0.0014$, $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-) = 0.081 \pm 0.005$. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ BORTOLETTO 92 reports $B(B \rightarrow D^*(2010)^+ \text{anything}) = 0.25 \pm 0.03 \pm 0.04$ using MARK II $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$ and $B(D^0 \rightarrow K^- \pi^+) = 0.042 \pm 0.008$. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ALBRECHT 91H reports $0.348 \pm 0.060 \pm 0.035$ from a measurement of $[\Gamma(B \rightarrow D^*(2010)^{\pm} \text{anything})/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0 \pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.55 \pm 0.04$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Uses the PDG 90 $B(D^0 \rightarrow K^- \pi^+) = 0.0371 \pm 0.0025$.

⁵ BORTOLETTO 87 uses old MARK III (BALTRUSAITIS 86E) branching ratios $B(D^0 \rightarrow K^- \pi^+) = 0.056 \pm 0.004 \pm 0.003$ and also assumes $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.60^{+0.08}_{-0.15}$. The product branching ratio for $B(B \rightarrow D^*(2010)^+) B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ is $0.13 \pm 0.02 \pm 0.012$. Superseded by BORTOLETTO 92.

⁶ $V-A$ momentum spectrum used to extrapolate below $p = 1$ GeV. We correct the value assuming $B(D^0 \rightarrow K^- \pi^+) = 0.042 \pm 0.006$ and $B(D^{*+} \rightarrow D^0 \pi^+) = 0.6^{+0.08}_{-0.15}$. The product branching fraction is $B(B \rightarrow D^{*+} X) \cdot B(D^{*+} \rightarrow \pi^+ D^0) \cdot B(D^0 \rightarrow K^- \pi^+) = (68 \pm 15 \pm 9) \times 10^{-4}$.

$\Gamma(D^*(2007)^0 \text{anything})/\Gamma_{\text{total}}$

Γ_{31}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.260±0.023±0.015	¹ GIBBONS 97B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ GIBBONS 97B reports $B(B \rightarrow D^*(2007)^0 \text{anything}) = 0.247 \pm 0.012 \pm 0.018 \pm 0.018$ using CLEO measured D and D^* branching fractions. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^{\pm} \text{anything})/\Gamma_{\text{total}}$

Γ_{32}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.083±0.008 OUR AVERAGE				
0.089±0.010±0.008		¹ ARTUSO 05B	CLE2	$e^+ e^- \rightarrow \gamma(5S)$
0.087±0.005±0.008		² AUBERT 02G	BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.065±0.011±0.006		³ ALBRECHT 92G	ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.068±0.010±0.006	257	⁴ BORTOLETTO90	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.085±0.022±0.008		⁵ HAAS 86	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.094±0.007±0.008		⁶ GIBAUT 96	CLE2	Repl. by ARTUSO 05B
0.094±0.024±0.008		⁷ ALBRECHT 87H	ARG	$e^+ e^- \rightarrow \gamma(4S)$

¹ ARTUSO 05B reports $0.0905 \pm 0.0025 \pm 0.0140$ from a measurement of $[\Gamma(B \rightarrow D_s^{\pm} \text{anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.5) \times 10^{-2}$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AUBERT 02G reports $[\Gamma(B \rightarrow D_s^{\pm} \text{anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)] = 0.00393 \pm 0.00007 \pm 0.00021$ which we divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 92G reports $[\Gamma(B \rightarrow D_s^{\pm} \text{anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)] = 0.00292 \pm 0.00039 \pm 0.00031$ which we divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ BORTOLETTO 90 reports $[\Gamma(B \rightarrow D_s^\pm \text{anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.00306 \pm 0.00047$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ HAAS 86 reports $[\Gamma(B \rightarrow D_s^\pm \text{anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.0038 \pm 0.0010$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $64 \pm 22\%$ decays are 2-body.

⁶ GIBAUT 96 reports $0.1211 \pm 0.0039 \pm 0.0088$ from a measurement of $[\Gamma(B \rightarrow D_s^\pm \text{anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷ ALBRECHT 87H reports $[\Gamma(B \rightarrow D_s^\pm \text{anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.0042 \pm 0.0009 \pm 0.0006$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $46 \pm 16\%$ of $B \rightarrow D_s X$ decays are 2-body. Superseded by ALBRECHT 92G.

$\Gamma(D_s^{*\pm} \text{anything})/\Gamma_{\text{total}}$	Γ_{33}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.063±0.009±0.006	¹ AUBERT 02G BABR $e^+e^- \rightarrow \gamma(4S)$
¹ AUBERT 02G reports $[\Gamma(B \rightarrow D_s^{*\pm} \text{anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.00284 \pm 0.00029 \pm 0.00025$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	

$\Gamma(D_s^{*\pm} \bar{D}^{(*)})/\Gamma(D_s^{*\pm} \text{anything})$	Γ_{34}/Γ_{33}
Sum over modes	
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.533±0.037±0.037	AUBERT 02G BABR $e^+e^- \rightarrow \gamma(4S)$

$\Gamma(\bar{D} D_{s0}(2317))/\Gamma_{\text{total}}$	Γ_{35}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	¹ KROKOVNY 03B BELL $e^+e^- \rightarrow \gamma(4S)$

¹ The product branching ratio for $B(B \rightarrow \bar{D} D_{s0}(2317)^+) \times B(D_{s0}(2317)^+ \rightarrow D_s \pi^0)$ is measured to be $(8.5^{+2.1}_{-1.9} \pm 2.6) \times 10^{-4}$.

$\Gamma(\bar{D} D_{sJ}(2457))/\Gamma_{\text{total}}$	Γ_{36}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	¹ KROKOVNY 03B BELL $e^+e^- \rightarrow \gamma(4S)$

¹ The product branching ratio for $B(B \rightarrow \bar{D} D_{sJ}(2457)^+) \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0, D_s^+ \gamma)$ are measured to be $(17.8^{+4.5}_{-3.9} \pm 5.3) \times 10^{-4}$ and $(6.7^{+1.3}_{-1.2} \pm 2.0) \times 10^{-4}$, respectively.

$$\frac{\Gamma(D_s^+\pi^-, D_s^{*+}\pi^-, D_s^+\rho^-, D_s^{*+}\rho^-, D_s^+\pi^0, D_s^{*+}\pi^0, D_s^+\eta, D_s^{*+}\eta, D_s^+\rho^0, D_s^{*+}\rho^0, D_s^+\omega, D_s^{*+}\omega)/\Gamma_{\text{total}}}{\text{Sum over modes.}} \quad \Gamma_{45}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4 \times 10^{-4}$	90	¹ ALEXANDER 93B	CLE2	$e^+e^- \rightarrow \gamma(4S)$

¹ ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ from a measurement of $[\Gamma(B \rightarrow D_s^+\pi^-, D_s^{*+}\pi^-, D_s^+\rho^-, D_s^{*+}\rho^-, D_s^+\pi^0, D_s^{*+}\pi^0, D_s^+\eta, D_s^{*+}\eta, D_s^+\rho^0, D_s^{*+}\rho^0, D_s^+\omega, D_s^{*+}\omega)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$. This branching ratio limit provides a model-dependent upper limit $|V_{ub}|/|V_{cb}| < 0.16$ at CL=90%.

$$\frac{\Gamma(D_{s1}(2536)^+\text{anything})/\Gamma_{\text{total}}}{\Gamma_{46}/\Gamma} \quad \Gamma_{46}/\Gamma$$

$D_{s1}(2536)^+$ is the narrow P -wave D_s^+ meson with $J^P = 1^+$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0095	90	¹ BISHAI 98	CLE2	$e^+e^- \rightarrow \gamma(4S)$

¹ Assuming factorization, the decay constant $f_{D_{s1}^+}$ is at least a factor of 2.5 times smaller than $f_{D_s^+}$.

$$\frac{\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}}{\Gamma_{47}/\Gamma} \quad \Gamma_{47}/\Gamma$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.094 ± 0.032 OUR AVERAGE		Error includes scale factor of 1.1.		
1.057 ± 0.012 ± 0.040		¹ AUBERT 03F	BABR	$e^+e^- \rightarrow \gamma(4S)$
1.121 ± 0.013 ± 0.042		ANDERSON 02	CLE2	$e^+e^- \rightarrow \gamma(4S)$
1.29 ± 0.45 ± 0.01	27	² MASCHMANN 90	CBAL	$e^+e^- \rightarrow \gamma(4S)$
1.24 ± 0.27 ± 0.01	120	³ ALBRECHT 87D	ARG	$e^+e^- \rightarrow \gamma(4S)$
1.35 ± 0.24 ± 0.01	52	⁴ ALAM 86	CLEO	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.12 ± 0.06 ± 0.01	1489	⁵ BALEST 95B	CLE2	$e^+e^- \rightarrow \gamma(4S)$
1.4 ± 0.6 ± 0.5	7	⁶ ALBRECHT 85H	ARG	$e^+e^- \rightarrow \gamma(4S)$
1.1 ± 0.21 ± 0.23	46	⁷ HAAS 85	CLEO	Repl. by ALAM 86

¹ AUBERT 03F also reports the momentum distribution and helicity of $J/\psi \rightarrow \ell^+\ell^-$ in the $\gamma(4S)$ center-of-mass frame.

² MASCHMANN 90 reports $(1.12 \pm 0.33 \pm 0.25) \times 10^{-2}$ from a measurement of $[\Gamma(B \rightarrow J/\psi(1S)\text{anything})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 87D reports $(1.07 \pm 0.16 \pm 0.22) \times 10^{-2}$ from a measurement of $[\Gamma(B \rightarrow J/\psi(1S)\text{anything})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. ALBRECHT 87D find the branching ratio for J/ψ not from $\psi(2S)$ to be 0.0081 ± 0.0023 .

⁴ ALAM 86 reports $(1.09 \pm 0.16 \pm 0.21) \times 10^{-2}$ from a measurement of $[\Gamma(B \rightarrow J/\psi(1S)\text{anything})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)]$ assuming $B(J/\psi(1S) \rightarrow$

$\mu^+ \mu^-) = 0.074 \pm 0.012$, which we rescale to our best value $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ BALEST 95B reports $(1.12 \pm 0.04 \pm 0.06) \times 10^{-2}$ from a measurement of $[\Gamma(B \rightarrow J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+ e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0599 \pm 0.0025$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. They measure $J/\psi(1S) \rightarrow e^+ e^-$ and $\mu^+ \mu^-$ and use PDG 1994 values for the branching fractions. The rescaling is the same for either mode so we use $e^+ e^-$.

⁶ Statistical and systematic errors were added in quadrature. ALBRECHT 85H also report a CL = 90% limit of 0.007 for $B \rightarrow J/\psi(1S) + X$ where $m_X < 1 \text{ GeV}$.

⁷ Dimuon and dielectron events used.

$\Gamma(J/\psi(1S)(\text{direct anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{48}/Γ
0.0078 ±0.0004 OUR AVERAGE			Error includes scale factor of 1.1.	
0.00740±0.00023±0.00043	¹ AUBERT	03F BABR	$e^+ e^- \rightarrow \gamma(4S)$	
0.00813±0.00017±0.00037	² ANDERSON	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0080 ±0.0008	³ BALEST	95B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

¹ AUBERT 03F also reports the helicity of $J/\psi \rightarrow \ell^+ \ell^-$ produced directly in B decay.

² Also reports the measurement of $J/\psi \rightarrow \ell^+ \ell^-$ polarization produced directly from B decay.

³ BALEST 95B assume PDG 1994 values for sub mode branching ratios. $J/\psi(1S)$ mesons are reconstructed in $J/\psi(1S) \rightarrow e^+ e^-$ and $J/\psi(1S) \rightarrow \mu^+ \mu^-$. The $B \rightarrow J/\psi(1S)X$ branching ratio contains $J/\psi(1S)$ mesons directly from B decays and also from feeddown through $\psi(2S) \rightarrow J/\psi(1S)$, $\chi_{c1}(1P) \rightarrow J/\psi(1S)$, or $\chi_{c2}(1P) \rightarrow J/\psi(1S)$. Using the measured inclusive rates, BALEST 95B corrects for the feeddown and finds the $B \rightarrow J/\psi(1S)(\text{direct}) X$ branching ratio.

$\Gamma(\psi(2S)\text{anything})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{49}/Γ
0.00307±0.00021 OUR AVERAGE					
0.00297±0.00020±0.00020		AUBERT	03F BABR	$e^+ e^- \rightarrow \gamma(4S)$	
0.00316±0.00014±0.00028		¹ ANDERSON	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
0.0046 ±0.0017 ±0.0011	8	ALBRECHT	87D ARG	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0034 ±0.0004 ±0.0003	240	² BALEST	95B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

¹ Also reports the measurement of $\psi(2S) \rightarrow \ell^+ \ell^-$ polarization produced directly from B decay.

² BALEST 95B assume PDG 1994 values for sub mode branching ratios. They find $B(B \rightarrow \psi(2S)X, \psi(2S) \rightarrow \ell^+ \ell^-) = 0.30 \pm 0.05 \pm 0.04$ and $B(B \rightarrow \psi(2S)X, \psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = 0.37 \pm 0.05 \pm 0.05$. Weighted average is quoted for $B(B \rightarrow \psi(2S)X)$.

$\Gamma(\chi_{c1}(1P)\text{anything})/\Gamma_{\text{total}}$	Γ_{50}/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.00386 ± 0.00027 OUR AVERAGE				
0.00367 ± 0.00035 ± 0.00044	AUBERT	03F	BABR	$e^+e^- \rightarrow \gamma(4S)$
0.00363 ± 0.00022 ± 0.00034	¹ ABE	02L	BELL	$e^+e^- \rightarrow \gamma(4S)$
0.00435 ± 0.00029 ± 0.00040	ANDERSON	02	CLE2	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0033 ± 0.0004 ± 0.0001	² CHEN	01	CLE2	$e^+e^- \rightarrow \gamma(4S)$
0.0040 ± 0.0006 ± 0.0004	³ BALEST	95B	CLE2	Repl. by CHEN 01
0.0105 ± 0.0035 ± 0.0025	⁴ ALBRECHT	92E	ARG	$e^+e^- \rightarrow \gamma(4S)$

¹ ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow e^+e^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.

² CHEN 01 reports $0.00414 \pm 0.00031 \pm 0.00040$ from a measurement of $[\Gamma(B \rightarrow \chi_{c1}(1P)\text{anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

³ BALEST 95B assume $B(\chi_{c1}(1P) \rightarrow J/\psi(1S)\gamma) = (27.3 \pm 1.6) \times 10^{-2}$, the PDG 1994 value. Fit to ψ -photon invariant mass distribution allows for a $\chi_{c1}(1P)$ and a $\chi_{c2}(1P)$ component.

⁴ ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production.

$\Gamma(\chi_{c1}(1P)(\text{direct})\text{anything})/\Gamma_{\text{total}}$	Γ_{51}/Γ			
VALUE	DOCUMENT ID	TECN	COMMENT	
0.00324 ± 0.00025 OUR AVERAGE				
0.00341 ± 0.00035 ± 0.00042	AUBERT	03F	BABR	$e^+e^- \rightarrow \gamma(4S)$
0.00332 ± 0.00022 ± 0.00034	¹ ABE	02L	BELL	$e^+e^- \rightarrow \gamma(4S)$
0.0031 ± 0.0004 ± 0.0001	² CHEN	01	CLE2	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0037 ± 0.0007	³ BALEST	95B	CLE2	Repl. by CHEN 01

¹ ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow e^+e^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.

² CHEN 01 reports $0.00383 \pm 0.00031 \pm 0.00040$ from a measurement of $[\Gamma(B \rightarrow \chi_{c1}(1P)(\text{direct})\text{anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

³ BALEST 95B assume PDG 1994 values. $J/\psi(1S)$ mesons are reconstructed in the e^+e^- and $\mu^+\mu^-$ modes. The $B \rightarrow \chi_{c1}(1P)X$ branching ratio contains $\chi_{c1}(1P)$ mesons directly from B decays and also from feeddown through $\psi(2S) \rightarrow \chi_{c1}(1P)\gamma$. Using the measured inclusive rates, BALEST 95B corrects for the feeddown and finds the $B \rightarrow \chi_{c1}(1P)(\text{direct})X$ branching ratio.

$\Gamma(\chi_{c2}(1P)\text{anything})/\Gamma_{\text{total}}$

Γ_{52}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
14 ± 4 OUR AVERAGE			Error includes scale factor of 1.9. See the ideogram below.		
21.0 ± 4.5 ± 3.1			AUBERT	03F BABR	$e^+ e^- \rightarrow \gamma(4S)$
18.0 $^{+2.3}_{-2.8}$ ± 2.6			¹ ABE	02L BELL	$e^+ e^- \rightarrow \gamma(4S)$
6.9 ± 3.5 ± 0.3			² CHEN	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$


• • • We do not use the following data for averages, fits, limits, etc. • • •

<38 90 35 ³ BALEST 95B CLE2 Repl. by CHEN 01

¹ ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+ \ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.

² CHEN 01 reports $(9.8 \pm 4.8 \pm 1.5) \times 10^{-4}$ from a measurement of $[\Gamma(B \rightarrow \chi_{c2}(1P)\text{anything})/\Gamma_{\text{total}}] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, which we rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

³ BALEST 95B assume $B(\chi_{c2}(1P) \rightarrow J/\psi(1S)\gamma) = (13.5 \pm 1.1) \times 10^{-2}$, the PDG 1994 value. $J/\psi(1S)$ mesons are reconstructed in the $e^+ e^-$ and $\mu^+ \mu^-$ modes, and PDG 1994 branching fractions are used. If interpreted as signal, the 35 ± 13 events correspond to $B(B \rightarrow \chi_{c2}(1P)X) = (0.25 \pm 0.10 \pm 0.03) \times 10^{-2}$.



¹ ALBRECHT 93I value is not independent of the sum of $B \rightarrow K^+$ anything and $B \rightarrow K^-$ anything ALBRECHT 94C values.

² Assuming $\Upsilon(4S) \rightarrow B\bar{B}$, a total of $3.38 \pm 0.34 \pm 0.68$ kaons per $\Upsilon(4S)$ decay is found (the second error is systematic). In the context of the standard B -decay model, this leads to a value for $(b\text{-quark} \rightarrow c\text{-quark})/(b\text{-quark} \rightarrow \text{all})$ of $1.09 \pm 0.33 \pm 0.13$.

³ GIANNINI 82 at CESR-CUSB observed 1.58 ± 0.35 K^0 per hadronic event much higher than 0.82 ± 0.10 below threshold. Consistent with predominant $b \rightarrow cX$ decay.

$\Gamma(K^+\text{ anything})/\Gamma_{\text{total}}$

Γ_{60}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.66 ±0.05	¹ ALBRECHT 94C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.620 \pm 0.013 \pm 0.038$	² ALBRECHT 94C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$0.66 \pm 0.05 \pm 0.07$	² ALAM 87B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Measurement relies on lepton-kaon correlations. It is for the weak decay vertex and does not include mixing of the neutral B meson. Mixing effects were corrected for by assuming a mixing parameter r of $(18.1 \pm 4.3)\%$.

² Measurement relies on lepton-kaon correlations. It includes production through mixing of the neutral B meson.

$\Gamma(K^-\text{ anything})/\Gamma_{\text{total}}$

Γ_{61}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.13 ±0.04	¹ ALBRECHT 94C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.165 \pm 0.011 \pm 0.036$	² ALBRECHT 94C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$0.19 \pm 0.05 \pm 0.02$	² ALAM 87B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Measurement relies on lepton-kaon correlations. It is for the weak decay vertex and does not include mixing of the neutral B meson. Mixing effects were corrected for by assuming a mixing parameter r of $(18.1 \pm 4.3)\%$.

² Measurement relies on lepton-kaon correlations. It includes production through mixing of the neutral B meson.

$\Gamma(K^0/\bar{K}^0\text{ anything})/\Gamma_{\text{total}}$

Γ_{62}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.64 ±0.04 OUR AVERAGE	¹ ALBRECHT 94C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
	ALAM 87B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 94C assume a K^0/\bar{K}^0 multiplicity twice that of K_S^0 .

$\Gamma(K^*(892)^{\pm}\text{ anything})/\Gamma_{\text{total}}$

Γ_{63}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.182±0.054±0.024	ALBRECHT 94J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)^0/\bar{K}^*(892)^0\text{ anything})/\Gamma_{\text{total}}$

Γ_{64}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.146±0.016±0.020	ALBRECHT 94J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)\gamma)/\Gamma_{\text{total}}$ Γ_{65}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.24 \pm 0.54 \pm 0.32$		¹ COAN 00	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<150	90	² LESIAK 92	CBAL	$e^+ e^- \rightarrow \gamma(4S)$
< 24	90	ALBRECHT 88H	ARG	$e^+ e^- \rightarrow \gamma(4S)$

¹ An average of $B(B^+ \rightarrow K^*(892)^+ \gamma)$ and $B(B^0 \rightarrow K^*(892)^0 \gamma)$ measurements reported in COAN 00 by assuming full correlated systematic errors.

² LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

 $\Gamma(\eta K\gamma)/\Gamma_{\text{total}}$ Γ_{66}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.5 \pm 1.3^{+1.2}_{-0.9}$		¹ NISHIDA 05	BELL	$e^+ e^- \rightarrow \gamma(4S)$

¹ $m_{\eta K} < 2.4 \text{ GeV}/c^2$

 $\Gamma(K_1(1400)\gamma)/\Gamma_{\text{total}}$ Γ_{67}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 12.7 \times 10^{-5}$	90	¹ COAN 00	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.6×10^{-3}	90	² LESIAK 92	CBAL	$e^+ e^- \rightarrow \gamma(4S)$
< 4.1×10^{-4}	90	ALBRECHT 88H	ARG	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

 $\Gamma(K_2^*(1430)\gamma)/\Gamma_{\text{total}}$ Γ_{68}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.66^{+0.59}_{-0.53} \pm 0.13$		¹ COAN 00	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<83	90	ALBRECHT 88H	ARG	$e^+ e^- \rightarrow \gamma(4S)$
-----	----	--------------	-----	----------------------------------

¹ COAN 00 obtains a fitted signal yield of $15.9^{+5.7}_{-5.2}$ events. A search for contamination by $K^*(1410)$ yielded a rate consistent with 0; the central value assumes no contamination.

 $\Gamma(K_2(1770)\gamma)/\Gamma_{\text{total}}$ Γ_{69}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.2 \times 10^{-3}$	90	¹ LESIAK 92	CBAL	$e^+ e^- \rightarrow \gamma(4S)$

¹ LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

$\Gamma(K_3^*(1780)\gamma)/\Gamma_{\text{total}}$				Γ_{70}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.7 \times 10^{-5}$	90	1 NISHIDA	05	BELL $e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<3.0 \times 10^{-3}$	90	ALBRECHT	88H	ARG $e^+ e^- \rightarrow \gamma(4S)$
¹ Uses $B(K_3^*(1780) \rightarrow \eta K) = 0.11^{+0.05}_{-0.04}$.				

$\Gamma(K_4^*(2045)\gamma)/\Gamma_{\text{total}}$				Γ_{71}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-3}$	90	1 LESIAK	92	CBAL $e^+ e^- \rightarrow \gamma(4S)$
¹ LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.				

$\Gamma(K\eta'(958))/\Gamma_{\text{total}}$				Γ_{72}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$(8.3^{+0.9}_{-0.8} \pm 0.7) \times 10^{-5}$	1	RICHICHI	00	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)\eta'(958))/\Gamma_{\text{total}}$				Γ_{73}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$4.1^{+1.0}_{-0.9} \pm 0.5$	1	AUBERT	07E	BABR $e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<22	90	1 RICHICHI	00	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

$\Gamma(K\eta)/\Gamma_{\text{total}}$				Γ_{74}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-6}$	90	1 RICHICHI	00	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)\eta)/\Gamma_{\text{total}}$				Γ_{75}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$(1.80^{+0.49}_{-0.43} \pm 0.18) \times 10^{-5}$	1	RICHICHI	00	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K\phi\phi)/\Gamma_{\text{total}}$				Γ_{76}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$2.3^{+0.9}_{-0.8} \pm 0.3$	1	HUANG	03	BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of charged and neutral B meson pairs and isospin symmetry.

\$\Gamma(\bar{b} \rightarrow \bar{s}\gamma)/\Gamma_{\text{total}}	\$\Gamma_{77}/\Gamma\$
---	------------------------

\$\Gamma(\bar{b} \rightarrow \bar{s}\gamma)/\Gamma_{\text{total}}	\$\Gamma_{77}/\Gamma\$		
3.40 ± 0.21 OUR AVERAGE			
3.52 ± 0.20 ± 0.51	^{1,2} LEES	12U BABR	$e^+ e^- \rightarrow \gamma(4S)$
3.32 ± 0.16 ± 0.31	^{1,3} LEES	12V BABR	$e^+ e^- \rightarrow \gamma(4S)$
3.47 ± 0.15 ± 0.40	^{1,4} LIMOSANI	09 BELL	$e^+ e^- \rightarrow \gamma(4S)$
3.90 ± 0.91 ± 0.64	^{1,5} AUBERT	080 BABR	$e^+ e^- \rightarrow \gamma(4S)$
3.36 ± 0.53 ^{+0.65} _{-0.68}	⁶ ABE	01F BELL	$e^+ e^- \rightarrow \gamma(4S)$
3.29 ± 0.44 ± 0.29	^{1,7} CHEN	01C CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.30 ± 0.08 ± 0.30	⁸ DEL-AMO-SA..10M	BABR	$e^+ e^- \rightarrow \gamma(4S)$
4.3 ± 0.3 ± 0.7	⁹ AUBERT	09U BABR	Repl. by DEL-AMO-SANCHEZ 10M
3.92 ± 0.31 ± 0.47	^{1,10} AUBERT,BE	06B BABR	Repl. by LEES 12V
3.49 ± 0.20 ^{+0.59} _{-0.46}	^{1,11} AUBERT,B	05R BABR	Repl. by LEES 12U
3.50 ± 0.32 ± 0.31	^{1,12} KOPPENBURG04	BELL	Repl. by LIMOSANI 09
2.32 ± 0.57 ± 0.35	ALAM	95 CLE2	Repl. by CHEN 01C

¹ We correct it to $E_\gamma > 1.6$ GeV using the method of BUCHMULLER 06 (average of three theoretical models).

² Reports $(3.29 \pm 0.19 \pm 0.48) \times 10^{-4}$ for $E_\gamma > 1.9$ GeV.

³ Reports $(3.21 \pm 0.15 \pm 0.29 \pm 0.08) \times 10^{-4}$ for $1.8 < E_\gamma < 2.8$ GeV, where the last systematic uncertainty is for model dependency. Results with other cutoffs are also reported.

⁴ The measurement reported is $(3.45 \pm 0.15 \pm 0.40) \times 10^{-4}$ for $E_\gamma > 1.7$ GeV.

⁵ Uses a fully reconstructed B meson as a tag on the recoil side. The measurement reported is $(3.66 \pm 0.85 \pm 0.60) \times 10^{-4}$ for $E_\gamma > 1.9$ GeV.

⁶ ABE 01F reports their systematic errors $(\pm 0.42^{+0.50}_{-0.54}) \times 10^{-4}$, where the second error is due to the theoretical uncertainty. We combine them in quadrature.

⁷ The measurement reported is $(3.21 \pm 0.43^{+0.32}_{-0.29}) \times 10^{-4}$ for $E_\gamma > 2.0$ GeV.

⁸ Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)}\gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range $0.5\text{--}2.0$ GeV/c 2 .

⁹ Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)}\gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range $0.6\text{--}1.8$ GeV/c 2 .

¹⁰ The measurement reported is $(3.67 \pm 0.29 \pm 0.45) \times 10^{-4}$ for $E_\gamma > 1.9$ GeV.

¹¹ The measurement reported is $(3.27 \pm 0.18^{+0.55}_{-0.42}) \times 10^{-4}$ for $E_\gamma > 1.9$ GeV.

¹² The measurement reported is $(3.55 \pm 0.32 \pm 0.32) \times 10^{-4}$ for $E_\gamma > 1.8$ GeV.

\$\Gamma(\bar{b} \rightarrow \bar{d}\gamma)/\Gamma_{\text{total}}	\$\Gamma_{78}/\Gamma\$
---	------------------------

\$\Gamma(\bar{b} \rightarrow \bar{d}\gamma)/\Gamma_{\text{total}}	\$\Gamma_{78}/\Gamma\$		
9.2 ± 2.0 ± 2.3			
1 DEL-AMO-SA..10M	¹ BABR	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
14 ± 5 ± 4	² AUBERT	09U BABR	Repl. by DEL-AMO-SANCHEZ 10M
¹ Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)}\gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range $0.5\text{--}2.0$ GeV/c 2 .			
² Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)}\gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range $0.6\text{--}1.8$ GeV/c 2 .			

$\Gamma(\bar{b} \rightarrow \bar{d}\gamma)/\Gamma(\bar{b} \rightarrow \bar{s}\gamma)$ Γ_{78}/Γ_{77}

VALUE	DOCUMENT ID	TECN	COMMENT
0.040±0.009±0.010	¹ DEL-AMO-SA..10M	BABR	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.033±0.013±0.009	² AUBERT	09U	BABR Repl. by DEL-AMO-SANCHEZ 10M
¹ Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)}\gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range 0.5–2.0 GeV/c ² . ² Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)}\gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range 0.6–1.8 GeV/c ² .			

 $\Gamma(\bar{b} \rightarrow \bar{s}\text{gluon})/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.068	90		¹ COAN	98	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.08		2	² ALBRECHT	95D	ARG $e^+ e^- \rightarrow \gamma(4S)$

¹ COAN 98 uses D - ℓ correlation.

² ALBRECHT 95D use full reconstruction of one B decay as tag. Two candidate events for charmless B decay can be interpreted as either $b \rightarrow s\text{gluon}$ or $b \rightarrow u$ transition. If interpreted as $b \rightarrow s\text{gluon}$ they find a branching ratio of ~ 0.026 or the upper limit quoted above. Result is highly model dependent.

 $\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
2.61±0.30^{+0.44}_{-0.74}		¹ NISHIMURA	10	BELL $e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
1.69±0.29 ^{+0.36} _{-0.62}		² NISHIMURA	10	BELL $e^+ e^- \rightarrow \gamma(4S)$
<4.4	90	³ BROWDER	98	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ Uses $B \rightarrow \eta X_s$ with $0.4 < m_{X_s} < 2.6$ GeV/c².

² Uses $B \rightarrow \eta X_s$ with $1.8 < m_{X_s} < 2.6$ GeV/c².

³ BROWDER 98 search for high momentum $B \rightarrow \eta X_s$ between 2.1 and 2.7 GeV/c.

 $\Gamma(\eta' \text{ anything})/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
4.2±0.9 OUR AVERAGE			
3.9±0.8±0.9	¹ AUBERT,B	04F	BABR $e^+ e^- \rightarrow \gamma(4S)$
4.6±1.1±0.6	² BONVICINI	03	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
6.2±1.6 ^{+1.3} _{-2.0}	³ BROWDER	98	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹AUBERT,B 04F reports branching ratio $B \rightarrow \eta' X_s$ for high momentum η' between 2.0 and 2.7 GeV/c in the $\Upsilon(4S)$ center-of-mass frame. X_s represents a recoil system consisting of a kaon and zero to four pions.

²BONVICINI 03 observed a signal of 61.2 ± 13.9 events in $B \rightarrow \eta' X_{nc}$ production for high momentum η' between 2.0 and 2.7 GeV/c in the $\Upsilon(4S)$ center-of-mass frame. The X_{nc} denotes “charmless” hadronic states recoiling against η' . The second error combines systematic and background subtraction uncertainties in quadrature.

³BROWDER 98 observed a signal of 39.0 ± 11.6 events in high momentum $B \rightarrow \eta' X_s$ production between 2.0 and 2.7 GeV/c. The branching fraction is based on the interpretation of $b \rightarrow sg$, where the last error includes additional uncertainties due to the color-suppressed $b \rightarrow$ backgrounds.

$\Gamma(K^+ \text{gluon (charmless)})/\Gamma_{\text{total}}$

Γ_{82}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.87	90	¹ DEL-AMO-SA..11	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ $B \rightarrow K^+ X$ with $m_X < 1.69$ GeV/c².

$\Gamma(K^0 \text{gluon (charmless)})/\Gamma_{\text{total}}$

Γ_{83}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$1.95^{+0.51}_{-0.45} \pm 0.50$	¹ DEL-AMO-SA..11	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ $B \rightarrow K^0 X$ with $m_X < 1.69$ GeV/c².

$\Gamma(\rho\gamma)/\Gamma_{\text{total}}$

Γ_{84}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.39 ± 0.25 OUR AVERAGE				Error includes scale factor of 1.2.
$1.73^{+0.34}_{-0.32} \pm 0.17$		^{1,2} AUBERT	08BH BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.21^{+0.24}_{-0.22} \pm 0.12$		^{1,2} TANIGUCHI	08 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.36^{+0.29}_{-0.27} \pm 0.10$		^{1,3} AUBERT	07L BABR	Repl. by AUBERT 08BH
< 1.9	90	^{1,3} AUBERT	04C BABR	Repl. by AUBERT 07L
< 14	90	^{1,4} COAN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes $\Gamma(B \rightarrow \rho\gamma) = \Gamma(B^+ \rightarrow \rho^+\gamma) = 2 \Gamma(B^0 \rightarrow \rho^0\gamma)$ and uses lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.071 \pm 0.009$.

³ Assumes $\Gamma(B \rightarrow \rho\gamma) = \Gamma(B^+ \rightarrow \rho^+\gamma) = 2 \Gamma(B^0 \rightarrow \rho^0\gamma)$ and uses lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.083 \pm 0.017$.

⁴ COAN 00 reports $B(B \rightarrow \rho\gamma)/B(B \rightarrow K^*(892)\gamma) < 0.32$ at 90%CL and scaled by the central value of $B(B \rightarrow K^*(892)\gamma) = (4.24 \pm 0.54 \pm 0.32) \times 10^{-5}$.

$\Gamma(\rho\gamma)/\Gamma(K^*(892)\gamma)$

Γ_{84}/Γ_{65}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$3.02^{+0.60+0.26}_{-0.55-0.28}$	TANIGUCHI 08	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\rho/\omega\gamma)/\Gamma_{\text{total}}$ Γ_{85}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.30 ± 0.23 OUR AVERAGE				Error includes scale factor of 1.2.
$1.63^{+0.30}_{-0.28} \pm 0.16$	1,2,3	AUBERT	08BH BABR	$e^+ e^- \rightarrow \gamma(4S)$
$1.14 \pm 0.20^{+0.10}_{-0.12}$	1,3	TANIGUCHI	08 BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.25^{+0.25}_{-0.24} \pm 0.09$	4	AUBERT	07L BABR	Repl. by AUBERT 08BH
$1.32^{+0.34}_{-0.31}^{+0.10}_{-0.09}$	4	MOHAPATRA	06 BELL	Repl. by TANIGUCHI 08
$0.6 \pm 0.3 \pm 0.1$	4	AUBERT	05 BABR	Repl. by AUBERT 07L
<1.4	90	4 MOHAPATRA	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes $\Gamma(B \rightarrow \rho\gamma) = \Gamma(B^+ \rightarrow \rho^+\gamma) = 2 \Gamma(B^0 \rightarrow \rho^0\gamma)$ and uses lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.071 \pm 0.009$.

² Also reports $|V_{td}/V_{ts}| = 0.233^{+0.025}_{-0.024}^{+0.022}_{-0.021}$.

³ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

⁴ Assumes $\Gamma(B \rightarrow \rho\gamma) = \Gamma(B^+ \rightarrow \rho^+\gamma) = 2 \Gamma(B^0 \rightarrow \rho^0\gamma)$ and uses lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.083 \pm 0.017$.

 $\Gamma(\rho/\omega\gamma)/\Gamma(K^*(892)\gamma)$ Γ_{85}/Γ_{65}

<u>VALUE</u> (units 10^{-2})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.84 \pm 0.50^{+0.27}_{-0.29}$	1	TANIGUCHI	08 BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.5	90	MOHAPATRA	05 BELL	Repl. by TANIGUCHI 08
------	----	-----------	---------	-----------------------

¹ Also reports $|V_{td}/V_{ts}| = 0.195^{+0.020}_{-0.019} \pm 0.015$.

 $\Gamma(\pi^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{86}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.585 \pm 0.025 \pm 0.070$	¹ ALBRECHT	93I ARG	$e^+ e^- \rightarrow \gamma(4S)$

¹ ALBRECHT 93 excludes π^\pm from K_S^0 and Λ decays. If included, they find $4.105 \pm 0.025 \pm 0.080$.

 $\Gamma(\pi^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{87}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.35 \pm 0.02 \pm 0.11$	¹ ABE	01J BELL	$e^+ e^- \rightarrow \gamma(4S)$

¹ From fully inclusive π^0 yield with no corrections from decays of K_S^0 or other particles.

 $\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$ Γ_{88}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.176 \pm 0.011 \pm 0.012$	KUBOTA	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(\rho^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{89}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.208 \pm 0.042 \pm 0.032$	ALBRECHT	94J ARG	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$	Γ_{90}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.81	90	ALBRECHT	94J	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$	Γ_{91}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0343 ± 0.0012 OUR AVERAGE			
0.0353 $\pm 0.0005 \pm 0.0030$	HUANG 07	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.0341 $\pm 0.0006 \pm 0.0012$	AUBERT 04S	BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.0390 $\pm 0.0030 \pm 0.0035$	ALBRECHT 94J	ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.023 $\pm 0.006 \pm 0.005$	BORTOLETTO86	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$	Γ_{92}/Γ		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<2.2 \times 10^{-5}$	90	¹ BERGFELD 98	CLE2

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\pi^+ \text{ gluon (charmless)})/\Gamma_{\text{total}}$	Γ_{94}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.72^{+0.50}_{-0.47} \pm 0.59$	¹ DEL-AMO-SA..11	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ $B \rightarrow \pi^+ X$ with $m_X < 1.71 \text{ GeV}/c^2$.

$\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})/\Gamma_{\text{total}}$	Γ_{95}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.045 \pm 0.003 \pm 0.012$		¹ AUBERT 07C	BABR	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.064 $\pm 0.008 \pm 0.008$		² CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.14 ± 0.09		³ ALBRECHT 88E	ARG	$e^+ e^- \rightarrow \gamma(4S)$
<0.112	90	⁴ ALAM 87	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

¹ AUBERT 07C reports $0.045 \pm 0.003 \pm 0.012$ from a measurement of $[\Gamma(B \rightarrow \Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$.

² CRAWFORD 92 result derived from lepton baryon correlations. Assumes all charmed baryons in B^0 and B^\pm decay are Λ_c .

³ ALBRECHT 88E measured $B(B \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (0.30 \pm 0.12 \pm 0.06)\%$ and used $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (2.2 \pm 1.0)\%$ from ABRAMS 80 to obtain above number.

⁴ Assuming all baryons result from charmed baryons, ALAM 86 conclude the branching fraction is $7.4 \pm 2.9\%$. The limit given above is model independent.

$\Gamma(\Lambda_c^+ \text{ anything})/\Gamma(\bar{\Lambda}_c^- \text{ anything})$	Γ_{96}/Γ_{97}		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.19 \pm 0.13 \pm 0.04$	¹ AMMAR 97	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ AMMAR 97 uses a high-momentum lepton tag ($P_\ell > 1.4 \text{ GeV}/c^2$).

$\Gamma(\bar{\Lambda}_c^- \mu^+ \text{anything})/\Gamma(\bar{\Lambda}_c^- \text{anything})$ Γ_{100}/Γ_{97}

<u>VALUE</u> (units 10^{-2})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-2.0 \pm 2.0 \pm 1.9$	LEES	12	BABR $e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(\bar{\Lambda}_c^- \ell^+ \text{anything})/\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{anything})$ Γ_{98}/Γ_{95}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.5 \times 10^{-2}$	90	¹ LEES	12	BABR $e^+ e^- \rightarrow \gamma(4S)$

¹ LEES 12 quotes also the measurement $\Gamma(B \rightarrow \bar{\Lambda}_c^- \ell^+ \text{anything})/\Gamma(B \rightarrow \Lambda_c^+ / \bar{\Lambda}_c^- \text{anything}) = (1.2 \pm 0.7 \pm 0.4) \times 10^{-2}$.

 $\Gamma(\bar{\Lambda}_c^- e^+ \text{anything})/\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{anything})$ Γ_{99}/Γ_{95}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.05	90	¹ BONVICINI	98	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ BONVICINI 98 uses the electron with momentum above 0.6 GeV/c.

 $\Gamma(\bar{\Lambda}_c^- e^+ \text{anything})/\Gamma(\bar{\Lambda}_c^- \text{anything})$ Γ_{99}/Γ_{97}

<u>VALUE</u> (units 10^{-2})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.5 \pm 1.1 \pm 0.6$	¹ LEES	12	BABR $e^+ e^- \rightarrow \gamma(4S)$

¹ Uses the full reconstruction of the recoiling B in a hadronic decay as a tag.

 $\Gamma(\bar{\Lambda}_c^- \ell^+ \text{anything})/\Gamma(\bar{\Lambda}_c^- \text{anything})$ Γ_{98}/Γ_{97}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.5 \times 10^{-2}$	90	¹ LEES	12	BABR $e^+ e^- \rightarrow \gamma(4S)$

¹ LEES 12 quotes also the measurement $\Gamma(B \rightarrow \bar{\Lambda}_c^- \ell^+ \text{anything})/\Gamma(B \rightarrow \bar{\Lambda}_c^- \text{anything}) = (1.7 \pm 1.0 \pm 0.6) \times 10^{-2}$.

 $\Gamma(\bar{\Lambda}_c^- p \text{anything})/\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{anything})$ Γ_{101}/Γ_{95}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.57 \pm 0.05 \pm 0.05$	BONVICINI	98	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(\bar{\Lambda}_c^- p e^+ \nu_e)/\Gamma(\bar{\Lambda}_c^- p \text{anything})$ $\Gamma_{102}/\Gamma_{101}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.04	90	¹ BONVICINI	98	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ BONVICINI 98 uses the electron with momentum above 0.6 GeV/c.

 $\Gamma(\bar{\Sigma}_c^{--} \text{anything})/\Gamma_{\text{total}}$ Γ_{103}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.0042 \pm 0.0021 \pm 0.0011$	77	¹ PROCARIO	94	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ PROCARIO 94 reports $[\Gamma(B \rightarrow \bar{\Sigma}_c^{--} \text{anything})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = 0.00021 \pm 0.00008 \pm 0.00007$ which we divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{\Sigma}_c^- \text{ anything})/\Gamma_{\text{total}}$		Γ_{104}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.010	90	1 PROCARIO	94	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ PROCARIO 94 reports $[\Gamma(B \rightarrow \bar{\Sigma}_c^- \text{ anything})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)] < 0.00048$ which we divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 5.0 \times 10^{-2}$.

$\Gamma(\bar{\Sigma}_c^0 \text{ anything})/\Gamma_{\text{total}}$		Γ_{105}/Γ		
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0046 ± 0.0021 ± 0.0012	76	1 PROCARIO	94	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ PROCARIO 94 reports $[\Gamma(B \rightarrow \bar{\Sigma}_c^0 \text{ anything})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = 0.00023 \pm 0.00008 \pm 0.00007$ which we divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{\Sigma}_c^0 N(N = p \text{ or } n))/\Gamma_{\text{total}}$		Γ_{106}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.5 × 10⁻³	90	1 PROCARIO	94	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ PROCARIO 94 reports < 0.0017 from a measurement of $[\Gamma(B \rightarrow \bar{\Sigma}_c^0 N(N = p \text{ or } n))/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.043$, which we rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 5.0 \times 10^{-2}$.

$\Gamma(\Xi_c^0 \text{ anything} \times B(\Xi_c^0 \rightarrow \Xi^- \pi^+))/\Gamma_{\text{total}}$		Γ_{107}/Γ	
VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
0.193 ± 0.030 OUR AVERAGE			Error includes scale factor of 1.1.
0.211 ± 0.019 ± 0.025	1 AUBERT,B 05M BABR	e ⁺ e ⁻ → γ(4S)	
0.144 ± 0.048 ± 0.021	2 BARISH 97 CLE2	e ⁺ e ⁻ → γ(4S)	

¹ The yield is obtained by requiring the momentum $P < 2.15 \text{ GeV}/c$.

² BARISH 97 find $79 \pm 27 \Xi_c^0$ events.

$\Gamma(\Xi_c^+ \text{ anything} \times B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+))/\Gamma_{\text{total}}$		Γ_{108}/Γ	
VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
0.453 ± 0.096 + 0.085 - 0.065	1 BARISH 97 CLE2	e ⁺ e ⁻ → γ(4S)	

¹ BARISH 97 find $125 \pm 28 \Xi_c^+$ events.

$\Gamma(p/\bar{p} \text{ anything})/\Gamma_{\text{total}}$		Γ_{109}/Γ	
INCLUDES p AND \bar{p} FROM Λ AND $\bar{\Lambda}$ DECAY.	DOCUMENT ID	TECN	COMMENT
0.080 ± 0.004 OUR AVERAGE			
0.080 ± 0.005 ± 0.005	ALBRECHT 93I ARG	e ⁺ e ⁻ → γ(4S)	
0.080 ± 0.005 ± 0.003	CRAWFORD 92 CLEO	e ⁺ e ⁻ → γ(4S)	
0.082 ± 0.005 + 0.013 - 0.010 2163	1 ALBRECHT 89K ARG	e ⁺ e ⁻ → γ(4S)	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
>0.021	2 ALAM 83B CLEO	e ⁺ e ⁻ → γ(4S)	

¹ ALBRECHT 89K include direct and nondirect protons.

² ALAM 83B reported their result as $> 0.036 \pm 0.006 \pm 0.009$. Data are consistent with equal yields of p and \bar{p} . Using assumed yields below cut, $B(B \rightarrow p + X) = 0.03$ not including protons from Λ decays.

$\Gamma(p/\bar{p} \text{ (direct) anything})/\Gamma_{\text{total}}$

Γ_{110}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.055±0.005 OUR AVERAGE				
0.055±0.005±0.0035		ALBRECHT 93I	ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.056±0.006±0.005		CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.055±0.016	1220	¹ ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \gamma(4S)$

¹ ALBRECHT 89K subtract contribution of Λ decay from the inclusive proton yield.

$\Gamma(\Lambda/\bar{\Lambda} \text{ anything})/\Gamma_{\text{total}}$

Γ_{111}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.040±0.005 OUR AVERAGE				
0.038±0.004±0.006	2998	CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.042±0.005±0.006	943	ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.022±0.003±0.0022 ¹ ACKERSTAFF 97N OPAL $e^+ e^- \rightarrow Z$

>0.011 ² ALAM 83B CLEO $e^+ e^- \rightarrow \gamma(4S)$

¹ ACKERSTAFF 97N assumes $B(b \rightarrow B) = 0.868 \pm 0.041$, i.e., an admixture of B^0 , B^\pm , and B_S .

² ALAM 83B reported their result as $> 0.022 \pm 0.007 \pm 0.004$. Values are for $(B(\Lambda X) + B(\bar{\Lambda} X))/2$. Data are consistent with equal yields of p and \bar{p} . Using assumed yields below cut, $B(B \rightarrow \Lambda X) = 0.03$.

$\Gamma(\Lambda \text{ anything})/\Gamma(\bar{\Lambda} \text{ anything})$

$\Gamma_{112}/\Gamma_{113}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.43±0.09±0.07				

¹ AMMAR 97 uses a high-momentum lepton tag ($P_\ell > 1.4 \text{ GeV}/c^2$).

$\Gamma(\Xi^-/\bar{\Xi}^+ \text{ anything})/\Gamma_{\text{total}}$

Γ_{114}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0027±0.0006 OUR AVERAGE				
0.0027±0.0005±0.0004	147	CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.0028±0.0014	54	ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\text{baryons anything})/\Gamma_{\text{total}}$

Γ_{115}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.068±0.005±0.003				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.076±0.014 ² ALBRECHT 89K ARG $e^+ e^- \rightarrow \gamma(4S)$

¹ ALBRECHT 920 result is from simultaneous analysis of p and Λ yields, $p\bar{p}$ and $\Lambda\bar{p}$ correlations, and various lepton-baryon and lepton-baryon-antibaryon correlations. Supersedes ALBRECHT 89K.

² ALBRECHT 89K obtain this result by adding their measurements ($5.5 \pm 1.6\%$) for direct protons and ($4.2 \pm 0.5 \pm 0.6\%$) for inclusive Λ production. They then assume ($5.5 \pm 1.6\%$) for neutron production and add it in also. Since each B decay has two baryons, they divide by 2 to obtain ($7.6 \pm 1.4\%$).

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.0 ± 2.3	$\begin{array}{c} +1.3 \\ -1.1 \end{array}$	² KANEKO	03	BELL	Repl. by IWASAKI 05
< 57	90	GLENN	98	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
<50000	90	BEBEK	81	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

¹ Requires $M_{\ell^+ \ell^-} > 0.2 \text{ GeV}/c^2$.

² Requires $M_{e^+ e^-} > 0.2 \text{ GeV}/c^2$.

$\Gamma(s\mu^+\mu^-)/\Gamma_{\text{total}}$

Γ_{120}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
4.3 ±1.2 OUR AVERAGE				

4.13 ± 1.05

¹ IWASAKI 05 BELL $e^+ e^- \rightarrow \gamma(4S)$

$5.0 \pm 2.8 \pm 1.2$

AUBERT,B 04I BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.9 ± 2.1

KANEKO 03 BELL Repl. by IWASAKI 05

< 58 90 GLENN 98 CLEO $e^+ e^- \rightarrow \gamma(4S)$

<17000 90 CHADWICK 81 CLEO $e^+ e^- \rightarrow \gamma(4S)$

¹ Requires $M_{\ell^+ \ell^-} > 0.2 \text{ GeV}/c^2$.

$[\Gamma(s\mu^+\mu^-) + \Gamma(s\mu^+\mu^-)]/\Gamma_{\text{total}}$

$(\Gamma_{119} + \Gamma_{120})/\Gamma$

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.2 × 10⁻⁵				

90 GLENN 98 CLEO $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0024 90 ¹ BEAN 87 CLEO Repl. by GLENN 98

<0.0062 90 ² AVERY 84 CLEO Repl. by BEAN 87

¹ BEAN 87 reports $[(\mu^+\mu^-) + (e^+e^-)]/2$ and we converted it.

² Determine ratio of B^+ to B^0 semileptonic decays to be in the range 0.25–2.9.

$\Gamma(s\ell^+\ell^-)/\Gamma_{\text{total}}$

Γ_{121}/Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
4.5 ±1.0 OUR AVERAGE			

4.11 ± 0.83

¹ IWASAKI 05 BELL $e^+ e^- \rightarrow \gamma(4S)$

$5.6 \pm 1.5 \pm 1.3$

² AUBERT,B 04I BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.1 ± 1.4

² KANEKO 03 BELL Repl. by IWASAKI 05

¹ Requires $M_{\ell^+ \ell^-} > 0.2 \text{ GeV}/c^2$.

² Requires $M_{e^+ e^-} > 0.2 \text{ GeV}/c^2$.

$\Gamma(\pi\ell^+\ell^-)/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.9 \times 10^{-8}$	90	1 LEES	13M BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<6.2 \times 10^{-8}$	90	1 WEI	08A BELL	$e^+ e^- \rightarrow \gamma(4S)$	
$<9.1 \times 10^{-8}$	90	1 AUBERT	07AG BABR	$e^+ e^- \rightarrow \gamma(4S)$	

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\pi e^+ e^-)/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<11.0 \times 10^{-8}$	90	1 LEES	13M BABR	$e^+ e^- \rightarrow \gamma(4S)$	

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\pi\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.0 \times 10^{-8}$	90	1 LEES	13M BABR	$e^+ e^- \rightarrow \gamma(4S)$	

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K e^+ e^-)/\Gamma_{\text{total}}$ Γ_{125}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT	
4.4±0.6 OUR AVERAGE					
$3.9^{+0.9}_{-0.8} \pm 0.2$		1 AUBERT	09T BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$4.8^{+0.8}_{-0.7} \pm 0.3$		1 WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$3.3^{+0.9}_{-0.8} \pm 0.2$		1 AUBERT,B	06J BABR	Repl. by AUBERT 09T	
$7.4^{+1.8}_{-1.6} \pm 0.5$		1 AUBERT	03U BABR	Repl. by AUBERT,B 06J	
$4.8^{+1.5}_{-1.3} \pm 0.3$		1,2 ISHIKAWA	03 BELL	Repl. by WEI 09A	
<13	90	ABE	02 BELL	Repl. by ISHIKAWA 03	

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.² The second error is a total of systematic uncertainties including model dependence. $\Gamma(K^*(892)e^+e^-)/\Gamma_{\text{total}}$ Γ_{126}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT	
11.9±2.0 OUR AVERAGE Error includes scale factor of 1.2.					
$9.9^{+2.3}_{-2.1} \pm 0.6$		1 AUBERT	09T BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$13.9^{+2.3}_{-2.0} \pm 1.2$		1 WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.9 \pm 0.7 \pm 0.2$	¹ AUBERT	09T	BABR	Repl. by LEES 12S
$3.4 \pm 0.7 \pm 0.2$	¹ AUBERT,B	06J	BABR	Repl. by AUBERT 09T
$6.5^{+1.4}_{-1.3} \pm 0.4$	² AUBERT	03U	BABR	Repl. by AUBERT,B 06J
$4.8^{+1.0}_{-0.9} \pm 0.3$	³ ISHIKAWA	03	BELL	Repl. by WEI 09A
$7.5^{+2.5}_{-2.1} \pm 0.6$	⁴ ABE	02	BELL	Repl. by ISHIKAWA 03
< 5.1	90	¹ AUBERT	02L	$e^+ e^- \rightarrow \gamma(4S)$
<17	90	⁵ ANDERSON	01B	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes all four $B \rightarrow K\ell^+\ell^-$ modes having equal partial widths in the fit.

³ Assumes equal production rate for charge and neutral B meson pairs, isospin invariance, lepton universality for $B \rightarrow K\ell^+\ell^-$, and $B(B \rightarrow K^*(892)\mu^+\mu^-) = 1.33$. The second error is total systematic uncertainties including model dependence.

⁴ Assumes lepton universality.

⁵ The result is for di-lepton masses above 0.5 GeV.

$\Gamma(K^*(892)\ell^+\ell^-)/\Gamma_{\text{total}}$

Γ_{130}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
10.5±1.0 OUR AVERAGE				
$10.2^{+1.4}_{-1.3} \pm 0.5$		LEES	12S	BABR $e^+ e^- \rightarrow \gamma(4S)$
$10.7^{+1.1}_{-1.0} \pm 0.9$		WEI	09A	BELL $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$11.1^{+1.9}_{-1.8} \pm 0.7$		¹ AUBERT	09T	BABR Repl. by LEES 12S
$7.8^{+1.9}_{-1.7} \pm 1.1$		¹ AUBERT,B	06J	BABR Repl. by AUBERT 09T
$8.8^{+3.3}_{-2.9} \pm 1.0$		² AUBERT	03U	BABR Repl. by AUBERT,B 06J
$11.5^{+2.6}_{-2.4} \pm 0.8$		³ ISHIKAWA	03	BELL Repl. by WEI 09A
<31	90	^{1,4} AUBERT	02L	BABR Repl. by AUBERT 03U
<33	90	⁵ ANDERSON	01B	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes the partial width ratio of electron and muon modes to be $\Gamma(B \rightarrow K^*(892)e^+e^-)/\Gamma(B \rightarrow K^*(892)\mu^+\mu^-) = 1.33$.

³ Assumes equal production rate for charge and neutral B meson pairs, isospin invariance, lepton universality for $B \rightarrow K\ell^+\ell^-$, and $B(B \rightarrow K^*(892)\mu^+\mu^-) = 1.33$. The second error is total systematic uncertainties including model dependence.

⁴ For averaging $K^*(892)\mu^+\mu^-$ and $K^*(892)e^+e^-$ modes, AUBERT 02L assumed $B(B \rightarrow K^*(892)e^+e^-)/B(B \rightarrow K^*(892)\mu^+\mu^-) = 1.2$.

⁵ The result is for di-lepton masses above 0.5 GeV.

$\Gamma(K^*(892)e^\pm\mu^\mp)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{137}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.1	90	¹ AUBERT,B	06J BABR	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<62	90	¹ EDWARDS	02B CLE2	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

CP VIOLATION A_{CP} is defined as

$$\frac{B(\bar{B} \rightarrow f) - B(B \rightarrow f)}{B(\bar{B} \rightarrow f) + B(B \rightarrow f)},$$

the CP -violation charge asymmetry of inclusive B^\pm and B^0 decay. $A_{CP}(B \rightarrow K^*(892)\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.003±0.017 OUR AVERAGE			
-0.003±0.017±0.007	¹ AUBERT	09AO BABR	$e^+e^- \rightarrow \gamma(4S)$
-0.015±0.044±0.012	² NAKAO	04 BELL	$e^+e^- \rightarrow \gamma(4S)$
+0.08 ±0.13 ±0.03	² COAN	00 CLE2	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.013±0.036±0.010	³ AUBERT,BE	04A BABR	Repl. by AUBERT 09AO
-0.044±0.076±0.012	⁴ AUBERT	02C BABR	Repl. by AUBERT,BE 04A

¹ Corresponds to a 90% CL interval $-0.033 < A_{CP} < 0.028$.² Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.³ Corresponds to a 90% CL allowed region, $-0.074 < A_{CP} < 0.049$.⁴ A 90% CL range is $-0.170 < A_{CP} < 0.082$. $A_{CP}(b \rightarrow s\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.008±0.029 OUR AVERAGE			
-0.011±0.030±0.014	¹ AUBERT	08BJ BABR	$e^+e^- \rightarrow \gamma(4S)$
0.002±0.050±0.030	² NISHIDA	04 BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.025±0.050±0.015	³ AUBERT,B	04E BABR	Repl. by AUBERT 08BJ
1	Uses a sum of exclusively reconstructed $B \rightarrow X_s$ decay modes, with X_s mass between 0.6 and 2.8 GeV/c ² .		
2	This measurement is performed inclusively for recoil mass X_s less than 2.1 GeV, which corresponds to $-0.093 < A_{CP} < 0.096$ at 90% CL.		
3	Corresponds to $-0.06 < A_{CP} < 0.11$ at 90% CL.		

$A_{CP}(b \rightarrow (s+d)\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.01 ± 0.05 OUR AVERAGE			
0.057 ± 0.060 ± 0.018	LEES 12V	BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.10 ± 0.18 ± 0.05	¹ AUBERT 080	BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.110 ± 0.115 ± 0.017	AUBERT,BE 06B	BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.079 ± 0.108 ± 0.022	² COAN 01	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side. Requires $E_\gamma > 2.2$ GeV.

² Corresponds to $-0.27 < A_{CP} < 0.10$ at 90% CL.

 $A_{CP}(B \rightarrow X_s \ell^+ \ell^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.22 ± 0.26 ± 0.02			

¹ The final state flavor is determined by the kaon and pion charges where modes with $X_s = K_S^0$, $K_S^0 \pi^0$ or $K_S^0 \pi^+ \pi^-$ are not used.

 $A_{CP}(B \rightarrow K^* e^+ e^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.18 ± 0.15 ± 0.01			

 $A_{CP}(B \rightarrow K^* \mu^+ \mu^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.03 ± 0.13 ± 0.02			

 $A_{CP}(B \rightarrow K^* \ell^+ \ell^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.04 ± 0.07 OUR AVERAGE			

0.03 ± 0.13 ± 0.01

+0.01^{+0.16}_{-0.15} ± 0.01

-0.10 ± 0.10 ± 0.01

¹ Measured in the union of $0.10 < q^2 < 8.12$ GeV^2/c^4 and $q^2 > 10.11$ GeV^2/c^4 .
LEES 12S reports also individual measurements $A_{CP}(B \rightarrow K^* \ell^+ \ell^-) = -0.13^{+0.18}_{-0.19} \pm 0.01$ for $0.10 < q^2 < 8.12$ GeV^2/c^4 and $A_{CP}(B \rightarrow K^* \ell^+ \ell^-) = 0.16^{+0.18}_{-0.19} \pm 0.01$ for $q^2 > 10.11$ GeV^2/c^4 .

 $A_{CP}(B \rightarrow \eta \text{anything})$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.13 ± 0.04 ± 0.02			

¹ Uses $B \rightarrow \eta X_s$ with $0.4 < m_{X_s} < 2.6$ GeV/c^2 .

$F_L(B \rightarrow K^* \ell^+ \ell^-)$ ($4.3 < q^2 < 8.6 \text{ GeV}^2/c^4$)


VALUE	DOCUMENT ID	TECN	COMMENT
0.64±0.06 OUR AVERAGE			
0.57±0.07±0.03	AAIJ	13Y LHCb	$p\bar{p}$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
0.81 ^{+0.13} _{-0.12} ±0.05	CHATRCHYAN 13BL	CMS	$p\bar{p}$ at 7 TeV
0.68 ^{+0.15} _{-0.17} ±0.09	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
0.64 ^{+0.23} _{-0.24} ±0.07	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.60 ^{+0.06} _{-0.07} ±0.01	AAIJ	12U LHCb	Repl. by AAIJ 13Y
0.82 ^{+0.19} _{-0.23} ±0.07	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $F_L(B \rightarrow K^* \ell^+ \ell^-)$ ($10.09 < q^2 < 12.86 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.43±0.06 OUR AVERAGE			
0.48 ^{+0.08} _{-0.09} ±0.03	AAIJ	13Y LHCb	$p\bar{p}$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
0.45 ^{+0.10} _{-0.11} ±0.04	CHATRCHYAN 13BL	CMS	$p\bar{p}$ at 7 TeV
0.47±0.14±0.03	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
0.17 ^{+0.17} _{-0.15} ±0.03	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.41±0.11±0.03	AAIJ	12U LHCb	Repl. by AAIJ 13Y
0.31 ^{+0.19} _{-0.18} ±0.02	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $F_L(B \rightarrow K^* \ell^+ \ell^-)$ ($14.18 < q^2 < 16.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.35±0.08 OUR AVERAGE			
Error includes scale factor of 1.4. See the ideogram below.			
0.33 ^{+0.08} _{-0.07} ^{+0.02} _{-0.03}	AAIJ	13Y LHCb	$p\bar{p}$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
0.53±0.12±0.03	CHATRCHYAN 13BL	CMS	$p\bar{p}$ at 7 TeV
0.29 ^{+0.14} _{-0.13} ±0.05	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
-0.15 ^{+0.27} _{-0.23} ±0.07	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.37±0.09±0.05	AAIJ	12U LHCb	Repl. by AAIJ 13Y
0.55 ^{+0.17} _{-0.18} ±0.02	AALTONEN	11L CDF	Repl. by AALTONEN 12I



$F_L(B \rightarrow K^* \ell^+ \ell^-) (14.18 < q^2 < 16.0 \text{ GeV}^2/c^4)$

$F_L(B \rightarrow K^* \ell^+ \ell^-) (16.0 < q^2 < 19.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.37±0.06 OUR AVERAGE	Error includes scale factor of 1.2.		
0.38 ^{+0.09} _{-0.07} ± 0.03	AAIJ	13Y LHCb	$p\bar{p}$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
0.44 ± 0.07 ± 0.03	CHATRCHYAN	13BL CMS	$p\bar{p}$ at 7 TeV
0.20 ^{+0.19} _{-0.17} ± 0.05	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
0.12 ^{+0.15} _{-0.13} ± 0.02	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.26 ^{+0.10} _{-0.08} ± 0.03	AAIJ	12U LHCb	Repl. by AAIJ 13Y
0.09 ^{+0.18} _{-0.14} ± 0.03	AALTONEN	11L CDF	Repl. by AALTONEN 12I

$F_L(B \rightarrow K^* \ell^+ \ell^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.66±0.06 OUR AVERAGE			
0.65 ^{+0.08} _{-0.07} ± 0.03	AAIJ	13Y LHCb	$p\bar{p}$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
0.68 ± 0.10 ± 0.02	CHATRCHYAN	13BL CMS	$p\bar{p}$ at 7 TeV
0.69 ^{+0.19} _{-0.21} ± 0.08	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
0.67 ± 0.23 ± 0.05	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.55 ± 0.10 ± 0.03	AAIJ	12U LHCb	Repl. by AAIJ 13Y
0.50 ^{+0.27} _{-0.30} ± 0.03	AALTONEN	11L CDF	Repl. by AALTONEN 12I

$F_L(B \rightarrow K^* \ell^+ \ell^-)$ ($0.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.33^{+0.14}_{-0.13} \pm 0.03$	AALTONEN	12I	CDF $p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.47^{+0.23}_{-0.24} \pm 0.03$	AALTONEN	11L	CDF Repl. by AALTONEN 12I

PARTIAL BRANCHING FRACTIONS IN $B \rightarrow K^{(*)} \ell^+ \ell^-$ **$B(B \rightarrow K^* \ell^+ \ell^-)$ ($q^2 < 2.0 \text{ GeV}^2/c^4$)**

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.68 ± 0.23 OUR AVERAGE			
$1.89^{+0.52}_{-0.46} \pm 0.06$	1 LEES	12S	BABR $e^+ e^- \rightarrow \gamma(4S)$
$1.73 \pm 0.33 \pm 0.10$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
$1.46^{+0.40}_{-0.35} \pm 0.11$	WEI	09A	BELL $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.98 \pm 0.40 \pm 0.09$	AALTONEN	11L	CDF Repl. by AALTONEN 11AI

¹ The value reported here from LEES 12S refers to $0.1 < q^2 < 2.0 \text{ GeV}^2/c^2$.

 $B(B \rightarrow K^* \ell^+ \ell^-)$ ($2.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.87 ± 0.17 OUR AVERAGE			
$0.95^{+0.35}_{-0.30} \pm 0.04$	LEES	12S	BABR $e^+ e^- \rightarrow \gamma(4S)$
$0.82 \pm 0.26 \pm 0.06$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
$0.86^{+0.31}_{-0.27} \pm 0.07$	WEI	09A	BELL $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.00 \pm 0.38 \pm 0.09$	AALTONEN	11L	CDF Repl. by AALTONEN 11AI

 $B(B \rightarrow K^* \ell^+ \ell^-)$ ($4.3 < q^2 < 8.68 \text{ GeV}^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.67 ± 0.29 OUR AVERAGE			
$1.82^{+0.56}_{-0.52} \pm 0.09$	1 LEES	12S	BABR $e^+ e^- \rightarrow \gamma(4S)$
$1.72 \pm 0.41 \pm 0.14$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
$1.37^{+0.47}_{-0.42} \pm 0.39$	WEI	09A	BELL $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.69 \pm 0.57 \pm 0.15$	AALTONEN	11L	CDF Repl. by AALTONEN 11AI

¹ The value reported here from LEES 12S refers to $4.3 < q^2 < 8.12 \text{ GeV}^2/c^2$.

$B(B \rightarrow K^* \ell^+ \ell^-)$ ($10.09 < q^2 < 12.86 \text{ GeV}^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.93±0.25 OUR AVERAGE			
$1.86^{+0.52}_{-0.48} \pm 0.10$	¹ LEES	12S BABR	$e^+ e^- \rightarrow \gamma(4S)$
$1.77 \pm 0.34 \pm 0.11$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$2.24^{+0.44}_{-0.40} \pm 0.19$	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.97 \pm 0.47 \pm 0.17$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

¹ The value reported here from LEES 12S refers to $10.11 < q^2 < 12.89 \text{ GeV}^2/c^2$.


$B(B \rightarrow K^* \ell^+ \ell^-)$ ($14.18 < q^2 < 16.0 \text{ GeV}^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.21±0.17 OUR AVERAGE			
$1.46^{+0.41}_{-0.36} \pm 0.06$	¹ LEES	12S BABR	$e^+ e^- \rightarrow \gamma(4S)$
$1.21 \pm 0.24 \pm 0.07$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$1.05^{+0.29}_{-0.26} \pm 0.08$	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.51 \pm 0.36 \pm 0.13$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

¹ The value reported here from LEES 12S refers to $14.21 < q^2 < 16.0 \text{ GeV}^2/c^2$.

$B(B \rightarrow K^* \ell^+ \ell^-)$ ($16.0 < q^2 \text{ GeV}^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.3 ±0.4 OUR AVERAGE		Error includes scale factor of 2.3. See the ideogram below.	
$1.02^{+0.47}_{-0.42} \pm 0.06$	LEES	12S BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.88 \pm 0.22 \pm 0.05$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$2.04^{+0.27}_{-0.24} \pm 0.16$	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.35 \pm 0.37 \pm 0.12$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI



$B(B \rightarrow K^* \ell^+ \ell^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.64±0.26 OUR AVERAGE			
2.05 ^{+0.53} _{-0.48} ± 0.07	LEES	12S BABR	$e^+ e^- \rightarrow \gamma(4S)$
1.48 ± 0.39 ± 0.12	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
1.49 ^{+0.45} _{-0.40} ± 0.12	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.60 ± 0.54 ± 0.14	AALTONEN	11L CDF	Repl. by AALTONEN 11AI


$B(B \rightarrow K^* \ell^+ \ell^-) (0.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
2.53±0.43±0.15			
AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.98 ± 0.55 ± 0.18	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

$B(B \rightarrow K \ell^+ \ell^-) (q^2 < 2.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.51±0.16 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.			
0.71 ^{+0.20} _{-0.18} ± 0.02	¹ LEES	12S BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.33 ± 0.10 ± 0.02	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
0.81 ^{+0.18} _{-0.16} ± 0.05	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.38 ± 0.16 ± 0.03	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

¹ The value reported here from LEES 12S refers to $0.1 < q^2 < 2.0 \text{ GeV}^2/c^2$.



$B(B \rightarrow K \ell^+ \ell^-) (2.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.57^{+0.10}_{-0.09} OUR AVERAGE Error includes scale factor of 1.2.

$0.49^{+0.15}_{-0.13} \pm 0.01$	LEES	12S	BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.77 \pm 0.14 \pm 0.05$	AALTONEN	11AI	CDF	$p\bar{p}$ at 1.96 TeV
$0.46^{+0.14}_{-0.12} \pm 0.03$	WEI	09A	BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.58 \pm 0.19 \pm 0.04$ AALTONEN 11L CDF Repl. by AALTONEN 11AI

$B(B \rightarrow K \ell^+ \ell^-) (4.3 < q^2 < 8.68 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.00^{+0.11}_{-0.11} OUR AVERAGE

$0.94^{+0.20}_{-0.19} \pm 0.02$	1 LEES	12S	BABR	$e^+ e^- \rightarrow \gamma(4S)$
$1.05 \pm 0.17 \pm 0.07$	AALTONEN	11AI	CDF	$p\bar{p}$ at 1.96 TeV
$1.00^{+0.19}_{-0.18} \pm 0.06$	WEI	09A	BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •


$0.93 \pm 0.25 \pm 0.06$ AALTONEN 11L CDF Repl. by AALTONEN 11AI

¹ The value reported here from LEES 12S refers to $4.3 < q^2 < 8.12 \text{ GeV}^2/c^2$.

$B(B \rightarrow K\ell^+\ell^-)$ ($10.09 < q^2 < 12.86 \text{ GeV}^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.57 ± 0.11 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.		
$0.90^{+0.20}_{-0.19} \pm 0.04$	¹ LEES	12S BABR $e^+e^- \rightarrow \gamma(4S)$	
$0.48 \pm 0.10 \pm 0.03$	AALTONEN	11AI CDF $p\bar{p}$ at 1.96 TeV	
$0.55^{+0.16}_{-0.14} \pm 0.03$	WEI	09A BELL $e^+e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.72 \pm 0.17 \pm 0.05$	AALTONEN	11L CDF Repl. by AALTONEN 11AI	

¹ The value reported here from LEES 12S refers to $10.11 < q^2 < 12.89 \text{ GeV}^2/c^2$.



$B(B \rightarrow K\ell^+\ell^-)$ ($14.18 < q^2 < 16.0 \text{ GeV}^2/c^4$)


VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.49 ± 0.07 OUR AVERAGE			
$0.49^{+0.15}_{-0.14} \pm 0.02$	¹ LEES	12S BABR $e^+e^- \rightarrow \gamma(4S)$	
$0.52 \pm 0.09 \pm 0.03$	AALTONEN	11AI CDF $p\bar{p}$ at 1.96 TeV	
$0.38^{+0.19}_{-0.12} \pm 0.02$	WEI	09A BELL $e^+e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.38 \pm 0.12 \pm 0.03$	AALTONEN	11L CDF Repl. by AALTONEN 11AI	

¹ The value reported here from LEES 12S refers to $14.21 < q^2 < 16.0 \text{ GeV}^2/c^2$.

$B(B \rightarrow K\ell^+\ell^-)$ ($16.0 < q^2$ GeV $^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.52±0.16 OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.		
$0.67^{+0.23}_{-0.21} \pm 0.05$	LEES	12S BABR	$e^+e^- \rightarrow \gamma(4S)$
$0.38 \pm 0.09 \pm 0.02$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$0.98^{+0.20}_{-0.18} \pm 0.06$	WEI	09A BELL	$e^+e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.35 \pm 0.13 \pm 0.02$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

WEIGHTED AVERAGE
 0.52 ± 0.16 (Error scaled by 2.1)



$$B(B \rightarrow K\ell^+\ell^-) \text{ (} 16.0 < q^2 \text{ GeV}^2/c^4 \text{)} \text{ (units } 10^{-7})$$

$B(B \rightarrow K\ell^+\ell^-)$ ($1.0 < q^2 < 6.0$ GeV $^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.33±0.13 OUR AVERAGE			
$1.36^{+0.27}_{-0.24} \pm 0.03$	LEES	12S BABR	$e^+e^- \rightarrow \gamma(4S)$
$1.29 \pm 0.18 \pm 0.08$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$1.36^{+0.23}_{-0.21} \pm 0.08$	WEI	09A BELL	$e^+e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$1.01 \pm 0.26 \pm 0.07$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

$B(B \rightarrow K\ell^+\ell^-) (0.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$1.07 \pm 0.17 \pm 0.07$	AALTONEN 11AI	CDF	$p\bar{p}$ at 1.96 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.96 \pm 0.25 \pm 0.06$	AALTONEN 11L	CDF	Repl. by AALTONEN 11AI

**LEPTON FORWARD-BACKWARD ASYMMETRY
IN $B \rightarrow K^{(*)}\ell^+\ell^-$ DECAY**

The forward-backward angular asymmetry of the lepton pair in $B \rightarrow K^{(*)}\ell^+\ell^-$ decay is defined as

$$A_{FB}(s) = \frac{N(\cos\theta > 0) - N(\cos\theta < 0)}{N(\cos\theta > 0) + N(\cos\theta < 0)},$$

where $s = q^2/m_B^2$, and θ is the angle of the lepton with respect to the flight direction of the B meson, measured in the dilepton rest frame. In addition, the fraction of longitudinal polarization F_L of the K^* and F_S , the relative contribution from scalar and pseudoscalar penguin amplitudes in $B \rightarrow K\ell^+\ell^-$, can be measured from the angular distribution of its decay products.

 $A_{FB}(B \rightarrow K^*\ell^+\ell^-) (q^2 > 0.1 \text{ GeV}^2/c^4)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$0.50 \pm 0.15 \pm 0.02$		¹ ISHIKAWA 06	BELL	$e^+e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
>0.55	95	² AUBERT,B 06J	BABR	$e^+e^- \rightarrow \gamma(4S)$


¹ Using an unbinned max. likelihood fits to the M_{bc} distribution in five q^2 bins for $\cos\theta > 0$ and $\cos\theta < 0$.

² Results with different q^2 cuts are also reported.

 $A_{FB}(B \rightarrow K^*\ell^+\ell^-) (0.1 < q^2 < 2.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.01 ± 0.14 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.		
$-0.02 \pm 0.12 \pm 0.01$	AAIJ 13Y	LHCb	$p\bar{p}$ at 7 TeV, $K^{*0}\mu^+\mu^-$
$-0.35^{+0.26}_{-0.23} \pm 0.10$	AALTONEN 12I	CDF	$p\bar{p}$ at 1.96 TeV
$0.47^{+0.26}_{-0.32} \pm 0.03$	WEI 09A	BELL	$e^+e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$-0.29^{+0.37}_{-0.00} \pm 0.18$	¹ CHATRCHYAN 13BL	CMS	$p\bar{p}$ at 7 TeV
$-0.15 \pm 0.20 \pm 0.06$	AAIJ 12U	LHCb	Repl. by AAIJ 13Y
$0.13^{+1.65}_{-0.75} \pm 0.25$	AALTONEN 11L	CDF	Repl. by AALTONEN 12I

¹ CHATRCHYAN 13BL uses, for this bin, $1.0 < q^2 < 2.0 \text{ GeV}^2/c^4$.



$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (m_{\ell\ell} < 2.5 \text{ GeV}/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.24^{+0.18}_{-0.23} \pm 0.05$	AUBERT	09N BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (m_{\ell\ell} > 3.2 \text{ GeV}/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.76^{+0.52}_{-0.32} \pm 0.07$	AUBERT	09N BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (2.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.15±0.07 OUR AVERAGE			
-0.20±0.08±0.01	AAIJ 13Y	LHCb	$p p$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
-0.07±0.20±0.02	CHATRCHYAN 13BL	CMS	$p p$ at 7 TeV
$0.29^{+0.32}_{-0.35} \pm 0.15$	AALTONEN 12I	CDF	$p \bar{p}$ at 1.96 TeV
$0.11^{+0.31}_{-0.36} \pm 0.07$	WEI 09A	BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.05^{+0.16}_{-0.20} \pm 0.04$	AAIJ 12U	LHCb	Repl. by AAIJ 13Y
$0.19^{+0.40}_{-0.41} \pm 0.14$	AALTONEN 11L	CDF	Repl. by AALTONEN 12I


$A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($0.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.08^{+0.21}_{-0.20} \pm 0.05$	AALTONEN	12I	CDF $p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.21^{+0.31}_{-0.33} \pm 0.05$	AALTONEN	11L	CDF Repl. by AALTONEN 12I

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
-0.12 ± 0.07 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
$-0.17 \pm 0.06 \pm 0.01$	AAIJ	13Y	LHCb $p\bar{p}$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
$-0.07 \pm 0.12 \pm 0.01$	CHATRCHYAN	13BL	CMS $p\bar{p}$ at 7 TeV
$0.29^{+0.20}_{-0.23} \pm 0.07$	AALTONEN	12I	CDF $p\bar{p}$ at 1.96 TeV
$0.26^{+0.27}_{-0.30} \pm 0.07$	WEI	09A	BELL $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$-0.06^{+0.13}_{-0.14} \pm 0.07$	AAIJ	12U	LHCb Repl. by AAIJ 13Y
$0.43^{+0.36}_{-0.37} \pm 0.06$	AALTONEN	11L	CDF Repl. by AALTONEN 12I

WEIGHTED AVERAGE
 -0.12 ± 0.07 (Error scaled by 1.3)



$A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$)

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($4.3 < q^2 < 8.6 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.13^{+0.06}_{-0.05}$ OUR AVERAGE	Error includes scale factor of 1.1.		
$0.16^{+0.06}_{-0.05} \pm 0.01$	AAIJ	13Y LHCb	$p\bar{p}$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
$-0.01 \pm 0.11 \pm 0.03$	CHATRCHYAN	13BL CMS	$p\bar{p}$ at 7 TeV
$0.01 \pm 0.20 \pm 0.09$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$0.45^{+0.15}_{-0.21} \pm 0.15$	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.27^{+0.06}_{-0.08} \pm 0.02$	AAIJ	12U LHCb	Repl. by AAIJ 13Y
$-0.06^{+0.30}_{-0.28} \pm 0.05$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($10.09 < q^2 < 12.86 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.34 ± 0.05 OUR AVERAGE			
$0.28^{+0.07}_{-0.06} \pm 0.02$	AAIJ	13Y LHCb	$p\bar{p}$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
$0.40 \pm 0.08 \pm 0.05$	CHATRCHYAN	13BL CMS	$p\bar{p}$ at 7 TeV
$0.38^{+0.16}_{-0.19} \pm 0.09$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$0.43^{+0.18}_{-0.20} \pm 0.03$	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.27^{+0.11}_{-0.13} \pm 0.02$	AAIJ	12U LHCb	Repl. by AAIJ 13Y
$0.66^{+0.23}_{-0.20} \pm 0.07$	AALTONEN	11L CDF	Repl. by AALTONEN 12I


 $A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($14.18 < q^2 < 16.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.47^{+0.07}_{-0.06}$ OUR AVERAGE	Error includes scale factor of 1.2.		
$0.51^{+0.07}_{-0.05} \pm 0.02$	AAIJ	13Y LHCb	$p\bar{p}$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
$0.29 \pm 0.09 \pm 0.05$	CHATRCHYAN	13BL CMS	$p\bar{p}$ at 7 TeV
$0.44^{+0.18}_{-0.21} \pm 0.10$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$0.70^{+0.16}_{-0.22} \pm 0.10$	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.47^{+0.06}_{-0.08} \pm 0.03$	AAIJ	12U LHCb	Repl. by AAIJ 13Y
$0.42 \pm 0.16 \pm 0.09$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($16.0 < q^2 < 19.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.40±0.06 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
$0.30 \pm 0.08^{+0.01}_{-0.02}$	AAIJ	13Y LHCb	$p\bar{p}$ at 7 TeV, $K^{*0} \mu^+ \mu^-$
$0.41 \pm 0.05 \pm 0.03$	CHATRCHYAN	13BL CMS	$p\bar{p}$ at 7 TeV
$0.65^{+0.17}_{-0.18} \pm 0.16$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$0.66^{+0.11}_{-0.16} \pm 0.04$	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.16^{+0.11}_{-0.13} \pm 0.06$	AAIJ	12U LHCb	Repl. by AAIJ 13Y
$0.70^{+0.16}_{-0.25} \pm 0.10$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

WEIGHTED AVERAGE
0.40±0.06 (Error scaled by 1.3)



$$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) \quad (16.0 < q^2 < 19.0 \text{ GeV}^2/c^4)$$

$A_{FB}(B \rightarrow K \ell^+ \ell^-)$ ($q^2 > 0.1 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.11±0.12 OUR AVERAGE			
$0.15^{+0.21}_{-0.23} \pm 0.08$	¹ AUBERT,B	06J BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.10 \pm 0.14 \pm 0.01$	² ISHIKAWA	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$

¹ Results with different q^2 cuts are also reported.

² Using an unbinned max. likelihood fits to the M_{bc} distribution in five q^2 bins for $\cos \theta > 0$ and $\cos \theta < 0$.

$A_{FB}(B \rightarrow K\ell^+\ell^-)$ ($q^2 < 2.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.00^{+0.06}_{-0.05}$ OUR AVERAGE			
$0.00^{+0.06}_{-0.05} \pm 0.01$	AAIJ	13H LHCb	$p\bar{p}$ at 7 TeV
$0.13^{+0.42}_{-0.43} \pm 0.07$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$0.06^{+0.32}_{-0.35} \pm 0.02$	WEI	09A BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$-0.15^{+0.46}_{-0.39} \pm 0.08$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K\ell^+\ell^-)$ ($2.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.09^{+0.10}_{-0.07}$ OUR AVERAGE Error includes scale factor of 1.4.			
$0.07^{+0.08}_{-0.05} \pm 0.02$	AAIJ	13H LHCb	$p\bar{p}$ at 7 TeV
$0.32^{+0.15}_{-0.16} \pm 0.05$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$-0.43^{+0.38}_{-0.40} \pm 0.09$	WEI	09A BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.72^{+0.40}_{-0.35} \pm 0.07$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K\ell^+\ell^-)$ ($0.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.31 \pm 0.16 \pm 0.04$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.36^{+0.24}_{-0.26} \pm 0.06$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K\ell^+\ell^-)$ ($1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.034^{+0.040}_{-0.029}$ OUR AVERAGE			
$0.02^{+0.05}_{-0.03} \pm 0.02$	AAIJ	13H LHCb	$p\bar{p}$ at 7 TeV
$0.13 \pm 0.09 \pm 0.02$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$-0.04^{+0.13}_{-0.16} \pm 0.05$	WEI	09A BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.08^{+0.27}_{-0.22} \pm 0.07$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

$A_{FB}(B \rightarrow K\ell^+\ell^-)$ ($4.3 < q^2 < 8.6 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
-0.04^{+0.04}_{-0.05} OUR AVERAGE			
-0.02 ^{+0.03} _{-0.05} ± 0.03	AAIJ	13H LHCb	$p p$ at 7 TeV
0.01 ^{+0.13} _{-0.10} ± 0.01	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
-0.20 ^{+0.12} _{-0.14} ± 0.03	WEI	09A BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.20 ^{+0.17} _{-0.28} ± 0.03	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K\ell^+\ell^-)$ ($10.09 < q^2 < 12.86 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
-0.05±0.06 OUR AVERAGE			
-0.03 ± 0.07 ± 0.01	AAIJ	13H LHCb	$p p$ at 7 TeV
-0.03 ^{+0.11} _{-0.10} ± 0.04	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
-0.21 ^{+0.17} _{-0.15} ± 0.06	WEI	09A BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.10 ^{+0.17} _{-0.15} ± 0.07	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K\ell^+\ell^-)$ ($14.18 < q^2 < 16.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
-0.02^{+0.07}_{-0.05} OUR AVERAGE			
-0.01 ^{+0.12} _{-0.06} ± 0.01	AAIJ	13H LHCb	$p p$ at 7 TeV
-0.05 ^{+0.09} _{-0.11} ± 0.03	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
0.04 ^{+0.32} _{-0.26} ± 0.05	WEI	09A BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.03 ^{+0.49} _{-0.16} ± 0.04	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K\ell^+\ell^-)$ ($16.0 < q^2 < 18.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
-0.09 ^{+0.07} _{-0.09} ^{+0.02} _{-0.01}	AAIJ	13H LHCb	$p p$ at 7 TeV

 $A_{FB}(B \rightarrow K\ell^+\ell^-)$ ($18.0 < q^2 < 22.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.02 ^{+0.11} _{-0.01} ± 0.01	AAIJ	13H LHCb	$p p$ at 7 TeV

$A_{FB}(B \rightarrow K\ell^+\ell^-)$ ($q^2 > 16.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.04$^{+0.09}_{-0.07}$ OUR AVERAGE			
$0.09^{+0.17}_{-0.13} \pm 0.03$	AALTONEN 12I	CDF	$p\bar{p}$ at 1.96 TeV
$0.02^{+0.11}_{-0.08} \pm 0.02$	WEI 09A	BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.07^{+0.30}_{-0.23} \pm 0.02$	AALTONEN 11L	CDF	Repl. by AALTONEN 12I

$F_S(B \rightarrow K\ell^+\ell^-)$ ($q^2 > 0.1 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.81^{+0.58}_{-0.61} \pm 0.46$	¹ AUBERT,B 06J	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Results with different q^2 cuts are also reported.

ISOSPIN ASYMMETRY

Δ_{0-} is defined as

$$\frac{\Gamma(B^0 \rightarrow f_d) - \Gamma(B^+ \rightarrow f_u)}{\Gamma(B^0 \rightarrow f) + \Gamma(B^+ \rightarrow f)},$$

the isospin asymmetry of inclusive neutral and charged B decay.

$\Delta_{0-}(B(B \rightarrow X_s \gamma))$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.01 ± 0.06 OUR AVERAGE			
-0.06 ± 0.15 ± 0.07	1,2 AUBERT 080	BABR	$e^+ e^- \rightarrow \gamma(4S)$
$-0.006 \pm 0.058 \pm 0.026$	AUBERT,B 05R	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ The result is for $E_\gamma > 2.2 \text{ GeV}$.

² Uses a fully reconstructed B meson as a tag on the recoil side.

$\Delta_{0+}(B \rightarrow K^*(892)\gamma)$

Δ_{0+} describes the isospin asymmetry between $\Gamma(B^0 \rightarrow K^*(892)^0 \gamma)$ and $\Gamma(B^+ \rightarrow K^*(892)^+ \gamma)$.

VALUE	DOCUMENT ID	TECN	COMMENT
0.052±0.026 OUR AVERAGE			
$0.066 \pm 0.021 \pm 0.022$	¹ AUBERT 09AO	BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.012 \pm 0.044 \pm 0.026$	NAKAO 04	BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.050 \pm 0.045 \pm 0.037$	² AUBERT,BE 04A	BABR	Repl. by AUBERT 09AO

¹ Uses the production ratio of charged and neutral B from $\gamma(4S)$ decays and the lifetime ratio $\tau_{B^+}/\tau_{B^0} = 1.071 \pm 0.009$. The 90% CL interval is $0.017 < \Delta_{0+} < 0.116$

² Uses the production ratio of charged and neutral B from $\gamma(4S)$ decays $R^+/0 = 1.006 \pm 0.048$ and the lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.083 \pm 0.017$. The 90% CL interval is $-0.046 < \Delta_{0+} < 0.146$.

$B \rightarrow X_c \ell \nu$ HADRONIC MASS MOMENTS **$\langle M_X^2 - \bar{M}_D^2 \rangle$ (First Moments)**

VALUE (GeV ²)	DOCUMENT ID	TECN	COMMENT
0.36 ±0.08 OUR AVERAGE			Error includes scale factor of 1.8.
0.467±0.038±0.068	¹ ACOSTA 05F	CDF	$p\bar{p}$ at 1.96 TeV
0.293±0.012±0.058	² CSORNA 04	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.251±0.023±0.062	³ CRONIN-HEN..01B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ Moments are measured with a minimum lepton momentum of 0.7 GeV/c in the B rest frame;
² Uses minimum lepton energy of 1.5 GeV and also reports moments with $E_\ell > 1.0$ GeV.
³ The leptons are required to have $P_\ell > 1.5$ GeV/c.

 $\langle M_X^2 \rangle$ (First Moments)

VALUE (GeV ²)	DOCUMENT ID	TECN	COMMENT
4.156±0.029 OUR AVERAGE			
4.144±0.028±0.022	¹ SCHWANDA 07	BELL	$e^+ e^- \rightarrow \gamma(4S)$
4.18 ± 0.04 ± 0.03	¹ AUBERT,B 04	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ The leptons are required to have $E_\ell > 1.5$ GeV/c.

 $\langle (M_X^2 - \bar{M}_D^2)^2 \rangle$ (Second Moments)

VALUE (GeV ⁴)	DOCUMENT ID	TECN	COMMENT
0.55 ±0.08 OUR AVERAGE			
0.515±0.061±0.064	¹ SCHWANDA 07	BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.629±0.031±0.143	² CSORNA 04	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.05 ± 0.26 ± 0.13	³ ACOSTA 05F	CDF	$p\bar{p}$ at 1.96 TeV
0.576±0.048±0.168	¹ CRONIN-HEN..01B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ The leptons are required to have $E_\ell > 1.5$ GeV/c.

² Uses minimum lepton energy of 1.5 GeV and also reports moments with $E_\ell > 1.0$ GeV.

³ Moments are measured with a minimum lepton momentum of 0.7 GeV/c in the B rest frame;

 $\langle (M_X^2 - \bar{M}_D^2)^2 \rangle$ (Second Moments)

VALUE (GeV ⁴)	DOCUMENT ID	TECN	COMMENT
0.639±0.056±0.178	¹ CRONIN-HEN..01B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ The leptons are required to have $E_\ell > 1.5$ GeV/c.

 $B \rightarrow X_c \ell \nu$ LEPTON MOMENTUM MOMENTS **$R_0 (\Gamma_{E_l>1.7\text{GeV}} / \Gamma_{E_l>1.5\text{GeV}})$**


VALUE	DOCUMENT ID	TECN	COMMENT
0.6187±0.0014±0.0016	¹ MAHMOOD 03	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ The leptons are required to have $E_l > 1.5$ GeV in the B rest frame.

$R_1 (\langle E_l \rangle_{E_l > 1.5 GeV})$

VALUE	DOCUMENT ID	TECN	COMMENT
1.7797 ± 0.0018 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.		
$1.7743 \pm 0.0019 \pm 0.0014$	¹ AUBERT,B 04A BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$1.7792 \pm 0.0021 \pm 0.0027$	² MAHMOOD 04 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
$1.7810 \pm 0.0007 \pm 0.0009$	³ MAHMOOD 03 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

- ¹ The leptons are required to have $E_l > 1.5$ GeV in the B rest frame. The result with $E_l > 0.6$ GeV is also given.
² Uses $E_e > 1.5$ GeV and also reports moments with other minimum minimum E_e conditions, as low as $E_e > 0.6$ GeV.
³ The leptons are required to have $E_l > 1.5$ GeV in the B rest frame.



$R_1 (\langle E_l \rangle_{E_l > 1.5 GeV})$

$R_2 (\langle E_l^2 - \bar{E}_l^2 \rangle_{E_l > 1.5 GeV})$

VALUE (10^{-3} GeV 2)	DOCUMENT ID	TECN	COMMENT
30.8 ± 0.8 OUR AVERAGE			
$30.3 \pm 0.9 \pm 0.5$	¹ AUBERT,B 04A BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$31.6 \pm 0.8 \pm 1.0$	² MAHMOOD 04 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	

- ¹ The leptons are required to have $E_l > 1.5$ GeV in the B rest frame. The result with $E_l > 0.6$ GeV is also given.
² Uses $E_e > 1.5$ GeV and also reports moments with other minimum minimum E_e conditions, as low as $E_e > 0.6$ GeV.

$R_3 (\langle E_l^3 - \bar{E}_l^3 \rangle_{E_l > 1.5 \text{GeV}})$

VALUE (10^{-3} GeV 3)	DOCUMENT ID	TECN	COMMENT
$2.12 \pm 0.47 \pm 0.20$	¹ AUBERT,B	04A BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ The leptons are required to have $E_l > 1.5$ GeV in the B rest frame. The result with $E_l > 0.6$ GeV is also given.

 $B \rightarrow X_s \gamma$ PHOTON ENERGY MOMENTS $\langle E_\gamma \rangle$

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
2.314 ± 0.011 OUR AVERAGE			
$2.346 \pm 0.018^{+0.027}_{-0.022}$	1,2 LEES	12U BABR	$e^+ e^- \rightarrow \gamma(4S)$
$2.304 \pm 0.014 \pm 0.017$	2,3 LEES	12V BABR	$e^+ e^- \rightarrow \gamma(4S)$
$2.311 \pm 0.009 \pm 0.015$	³ LIMOSANI	09 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$2.289 \pm 0.058 \pm 0.027$	3,4 AUBERT	080 BABR	$e^+ e^- \rightarrow \gamma(4S)$
$2.309 \pm 0.023 \pm 0.023$	2,3 SCHWANDA	08 BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.288 \pm 0.025 \pm 0.023$ ³ AUBERT,BE 06B BABR Repl. by LEES 12V

¹ LEES 12U uses $E_\gamma > 1.897$ GeV to calculate the moments; the moments are used to calculate the HQET parameters $m_b = 4.579^{+0.032}_{-0.029}$ GeV/c 2 and $\mu_\pi^2 = 0.257^{+0.034}_{-0.039}$ GeV 2 in the shape function model. The same HQET parameters are also determined in the kinetic model.

² Results for different E_γ threshold values are also measured.

³ The result is for $E_\gamma > 1.9$ GeV.

⁴ Uses a fully reconstructed B meson as a tag on the recoil side.

 $\langle E_\gamma^2 \rangle - \langle E_\gamma \rangle^2$

VALUE (10^{-2} GeV 2)	DOCUMENT ID	TECN	COMMENT
3.03 ± 0.25 OUR AVERAGE			
$2.11 \pm 0.57^{+0.55}_{-0.69}$	1,2 LEES	12U BABR	$e^+ e^- \rightarrow \gamma(4S)$
$3.62 \pm 0.33 \pm 0.33$	2,3 LEES	12V BABR	$e^+ e^- \rightarrow \gamma(4S)$
$3.02 \pm 0.19 \pm 0.30$	³ LIMOSANI	09 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$3.34 \pm 1.24 \pm 0.62$	3,4 AUBERT	080 BABR	$e^+ e^- \rightarrow \gamma(4S)$
$2.17 \pm 0.60 \pm 0.55$	2,3 SCHWANDA	08 BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.28 \pm 0.40 \pm 0.43$ ³ AUBERT,BE 06B BABR Repl. by LEES 12V

¹ LEES 12U uses $E_\gamma > 1.897$ GeV to calculate the moments; the moments are used to calculate the HQET parameters $m_b = 4.579^{+0.032}_{-0.029}$ GeV/c 2 and $\mu_\pi^2 = 0.257^{+0.034}_{-0.039}$ GeV 2 in the shape function model. The same HQET parameters are also determined in the kinetic model.

² Results for different E_γ threshold values are also measured.

³ The result is for $E_\gamma > 1.9$ GeV.

⁴ Uses a fully reconstructed B meson as a tag on the recoil side.

