



$$I(J^P) = 0(0^-)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

B_c^\pm MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
6.2756 ± 0.0011 OUR AVERAGE			
6.27628 ± 0.00144 ± 0.00036	¹ AAIJ	13AS LHCb	$p\bar{p}$ at 7, 8 TeV
6.2737 ± 0.0013 ± 0.0016	² AAIJ	12AV LHCb	$p\bar{p}$ at 7 TeV
6.2756 ± 0.0029 ± 0.0025	³ AALTONEN	08M CDF	$p\bar{p}$ at 1.96 TeV
6.300 ± 0.014 ± 0.005	³ ABAZOV	08T D0	$p\bar{p}$ at 1.96 TeV
6.4 ± 0.39 ± 0.13	⁴ ABE	98M CDF	$p\bar{p}$ at 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
6.2857 ± 0.0053 ± 0.0012	³ ABULENCIA	06C CDF	Repl. by AALTONEN 08M
6.32 ± 0.06	⁵ ACKERSTAFF	98O OPAL	$e^+e^- \rightarrow Z$

¹ AAIJ 13AS uses the $B_c^+ \rightarrow J/\psi D_s^+$.

² AAIJ 12AV uses the $B_c^+ \rightarrow J/\psi \pi^+$ mode and also measures the mass difference $M(B_c^+) - M(B^+) = 994.6 \pm 1.3 \pm 0.6 \text{ MeV}/c^2$.

³ Measured using a fully reconstructed decay mode of $B_c \rightarrow J/\psi \pi$.

⁴ ABE 98M observed $20.4_{-5.5}^{+6.2}$ events in the $B_c^+ \rightarrow J/\psi(1S) \ell \nu_\ell$ with a significance of > 4.8 standard deviations. The mass value is estimated from $m(J/\psi(1S) \ell)$.

⁵ ACKERSTAFF 98O observed 2 candidate events in the $B_c^+ \rightarrow J/\psi(1S) \pi^+$ channel with an estimated background of 0.63 ± 0.20 events.

B_c^\pm MEAN LIFE

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

VALUE (10^{-12} s)	DOCUMENT ID	TECN	COMMENT
0.452 ± 0.033 OUR EVALUATION			
0.500 ± 0.013 OUR AVERAGE			
0.509 ± 0.008 ± 0.012	⁶ AAIJ	14G LHCb	$p\bar{p}$ at 8 TeV
0.452 ± 0.048 ± 0.027	⁷ AALTONEN	13 CDF	$p\bar{p}$ at 1.96 TeV
0.448 $^{+0.038}_{-0.036}$ ± 0.032	⁸ ABAZOV	09H D0	$p\bar{p}$ at 1.96 TeV
0.463 $^{+0.073}_{-0.065}$ ± 0.036	⁸ ABULENCIA	06O CDF	$p\bar{p}$ at 1.96 TeV
0.46 $^{+0.18}_{-0.16}$ ± 0.03	⁸ ABE	98M CDF	$p\bar{p}$ 1.8 TeV

⁶ Measured using $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu X$ decays.

⁷ Uses fully reconstructed $B_c^+ \rightarrow J/\psi \pi^+$ decays.

⁸ The lifetime is measured from the $J/\psi e$ decay vertices.

B_c^+ DECAY MODES $\times B(\bar{b} \rightarrow B_c)$

B_c^- modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Confidence level
The following quantities are not pure branching ratios; rather the fraction $\Gamma_i/\Gamma \times B(\bar{b} \rightarrow B_c)$.		
Γ_1 $J/\psi(1S)\ell^+\nu_\ell$ anything	$(5.2^{+2.4}_{-2.1}) \times 10^{-5}$	
Γ_2 $J/\psi(1S)\pi^+$	seen	
Γ_3 $J/\psi(1S)K^+$	seen	
Γ_4 $J/\psi(1S)\pi^+\pi^+\pi^-$	seen	
Γ_5 $J/\psi(1S)a_1(1260)$	$< 1.2 \times 10^{-3}$	90%
Γ_6 $J/\psi(1S)K^+K^-\pi^+$	seen	
Γ_7 $\psi(2S)\pi^+$	seen	
Γ_8 $J/\psi(1S)D_s^+$	seen	
Γ_9 $J/\psi(1S)D_s^{*+}$	seen	
Γ_{10} $D^*(2010)^+\bar{D}^0$	$< 6.2 \times 10^{-3}$	90%
Γ_{11} D^+K^{*0}	$< 0.20 \times 10^{-6}$	90%
Γ_{12} $D^+\bar{K}^{*0}$	$< 0.16 \times 10^{-6}$	90%
Γ_{13} $D_s^+K^{*0}$	$< 0.28 \times 10^{-6}$	90%
Γ_{14} $D_s^+\bar{K}^{*0}$	$< 0.4 \times 10^{-6}$	90%
Γ_{15} $D_s^+\phi$	$< 0.32 \times 10^{-6}$	90%
Γ_{16} K^+K^0	$< 4.6 \times 10^{-7}$	90%
Γ_{17} $B_s^0\pi^+ / B(\bar{b} \rightarrow B_s)$	$(2.37^{+0.37}_{-0.35}) \times 10^{-3}$	

B_c^+ BRANCHING RATIOS

$\Gamma(J/\psi(1S)\ell^+\nu_\ell \text{ anything})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$	$\Gamma_1/\Gamma \times B$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT

$(5.2^{+2.4}_{-2.1}) \times 10^{-5}$ ⁹ ABE 98M CDF $p\bar{p}$ 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.6 \times 10^{-4}$	90	¹⁰ ACKERSTAFF	98O	OPAL	$e^+e^- \rightarrow Z$
$< 1.9 \times 10^{-4}$	90	¹¹ ABREU	97E	DLPH	$e^+e^- \rightarrow Z$
$< 1.2 \times 10^{-4}$	90	¹² BARATE	97H	ALEP	$e^+e^- \rightarrow Z$

⁹ ABE 98M result is derived from the measurement of $[\sigma(B_c) \times B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell)] / [\sigma(B^+) \times B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.132^{+0.041}_{-0.037}(\text{stat}) \pm 0.031(\text{sys})^{+0.032}_{-0.020}(\text{lifetime})$ by using PDG 98 values of $B(b \rightarrow B^+)$ and $B(B^+ \rightarrow J/\psi(1S)K^+)$.

¹⁰ ACKERSTAFF 98O reports $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell) < 6.95 \times 10^{-5}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

¹¹ ABREU 97E value listed is for an assumed $\tau_{B_c} = 0.4$ ps and improves to 1.6×10^{-4} for $\tau_{B_c} = 1.4$ ps.

¹² BARATE 97H reports $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \cdot B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell) < 5.2 \times 10^{-5}$ at 90%CL. We rescale to our PDG 96 values of $B(Z \rightarrow b\bar{b})$. A $B_c^+ \rightarrow J/\psi(1S)\mu^+\nu_\mu$ candidate event is found, compared to all the known background sources 2×10^{-3} , which gives $m_{B_c} = 5.96^{+0.25}_{-0.19}$ GeV and $\tau_{B_c} = 1.77 \pm 0.17$ ps.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_2/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
seen		AALTONEN 13	CDF	$p\bar{p}$ at 1.96 TeV
seen		¹³ AAIJ 12AV	LHCB	pp at 7 TeV
seen		AALTONEN 08M	CDF	$p\bar{p}$ at 1.96 TeV
seen		ABAZOV 08T	D0	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.4 \times 10^{-4}$	90	¹⁴ ACKERSTAFF 98O	OPAL	$e^+e^- \rightarrow Z$
$< 3.4 \times 10^{-4}$	90	¹⁵ ABREU 97E	DLPH	$e^+e^- \rightarrow Z$
$< 8.2 \times 10^{-5}$	90	¹⁶ BARATE 97H	ALEP	$e^+e^- \rightarrow Z$
$< 2.0 \times 10^{-5}$	95	¹⁷ ABE 96R	CDF	$p\bar{p}$ 1.8 TeV

¹³AAIJ 12AV reports a measurement of $B(B_c^+ \rightarrow J/\psi\pi^+)/B(B^+ \rightarrow J/\psi K^+) f_c/f_u = (0.68 \pm 0.10 \pm 0.03 \pm 0.05)\%$ at $p_T(B) > 4$ GeV and $2.5 < \eta(B) < 4.5$.

¹⁴ACKERSTAFF 98O reports $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)\pi^+) < 1.06 \times 10^{-4}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

¹⁵ABREU 97E value listed is for an assumed $\tau_{B_c} = 0.4$ ps and improves to 2.7×10^{-4} for $\tau_{B_c} = 1.4$ ps.

¹⁶BARATE 97H reports $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \cdot B(B_c \rightarrow J/\psi(1S)\pi) < 3.6 \times 10^{-5}$ at 90%CL. We rescale to our PDG 96 values of $B(Z \rightarrow b\bar{b})$.

¹⁷ABE 96R reports $B(b \rightarrow B_c X)/B(b \rightarrow B^+ X) \cdot B(B_c^+ \rightarrow J/\psi(1S)\pi^+)/B(B^+ \rightarrow J/\psi(1S)K^+) < 0.053$ at 95%CL for $\tau_{B_c} = 0.8$ ps. It changes from 0.15 to 0.04 for $0.17 \text{ ps} < \tau_{B_c} < 1.6$ ps. We rescale to our PDG 96 values of $B(b \rightarrow B^+) = 0.378 \pm 0.022$ and $B(B^+ \rightarrow J/\psi(1S)K^+) = 0.00101 \pm 0.00014$.

$\Gamma(J/\psi(1S)K^+)/\Gamma(J/\psi(1S)\pi^+)$ Γ_3/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.069 \pm 0.019 \pm 0.005$	50	AAIJ 13BY	LHCB	pp at 7 TeV

$\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_4/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
seen		AAIJ 12Y	LHCB	pp at 7 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 5.7 \times 10^{-4}$	90	¹⁸ ABREU 97E	DLPH	$e^+e^- \rightarrow Z$
------------------------	----	-------------------------	------	------------------------

¹⁸ABREU 97E value listed is independent of $0.4 \text{ ps} < \tau_{B_c} < 1.4$ ps.

$\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+)$ Γ_4/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
$2.41 \pm 0.30 \pm 0.33$	AAIJ 12Y	LHCB	pp at 7 TeV

$\Gamma(J/\psi(1S) a_1(1260))/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_5/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-3}$	90	19 ACKERSTAFF 98O	OPAL	$e^+e^- \rightarrow Z$

¹⁹ ACKERSTAFF 98O reports $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S) a_1(1260)) < 5.29 \times 10^{-4}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

$\Gamma(J/\psi(1S) K^+ K^- \pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_6/\Gamma \times B$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	20 AAIJ	13CA LHCB	pp at 7, 8 TeV

²⁰ A signal yield of 78 ± 14 decays is reported with a significance of 6.2 standard deviations using an integrated luminosity of 3 fb^{-1} data.

$\Gamma(J/\psi(1S) K^+ K^- \pi^+)/\Gamma(J/\psi(1S) \pi^+)$ Γ_6/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
$0.53 \pm 0.10 \pm 0.05$	21 AAIJ	13CA LHCB	pp at 7, 8 TeV

²¹ A signal yield of 78 ± 14 decays is reported with a significance of 6.2 standard deviations using an integrated luminosity of 3 fb^{-1} data.

$\Gamma(\psi(2S) \pi^+)/\Gamma(J/\psi(1S) \pi^+)$ Γ_7/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
$0.250 \pm 0.068 \pm 0.015$	22 AAIJ	13AMLHCB	pp at 7 TeV

²² The last uncertainty is due to the measurement of $B(\psi(2S) \rightarrow \mu^+ \mu^-)/B(J/\psi \rightarrow \mu^+ \mu^-)$.

$\Gamma(J/\psi(1S) D_s^+)/\Gamma(J/\psi(1S) \pi^+)$ Γ_8/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
$2.90 \pm 0.57 \pm 0.24$	AAIJ	13AS LHCB	pp at 7, 8 TeV

$\Gamma(J/\psi(1S) D_s^{*+})/\Gamma(J/\psi(1S) D_s^+)$ Γ_9/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
$2.37 \pm 0.56 \pm 0.10$	AAIJ	13AS LHCB	pp at 7, 8 TeV

$\Gamma(D^*(2010)^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{10}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.2 \times 10^{-3}$	90	23 BARATE 98Q	ALEP	$e^+e^- \rightarrow Z$

²³ BARATE 98Q reports $B(Z \rightarrow B_c X) \times B(B_c \rightarrow D^*(2010)^+ \bar{D}^0) < 1.9 \times 10^{-3}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

$\Gamma(D^+ K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{11}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.20	90	24 AAIJ	13R LHCB	pp at 7 TeV

²⁴ AAIJ 13R reports $[\Gamma(B_c^+ \rightarrow D^+ K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.5 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.2 \times 10^{-2}$.

$\Gamma(D^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{12}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.16	90	25 AAIJ	13R LHCB	<i>pp</i> at 7 TeV
²⁵ AAIJ 13R reports $[\Gamma(B_c^+ \rightarrow D^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.4 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.2 \times 10^{-2}$.				

$\Gamma(D_s^+ K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{13}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.28	90	26 AAIJ	13R LHCB	<i>pp</i> at 7 TeV
²⁶ AAIJ 13R reports $[\Gamma(B_c^+ \rightarrow D_s^+ K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.7 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.2 \times 10^{-2}$.				

$\Gamma(D_s^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{14}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	27 AAIJ	13R LHCB	<i>pp</i> at 7 TeV
²⁷ AAIJ 13R reports $[\Gamma(B_c^+ \rightarrow D_s^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 1.1 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.2 \times 10^{-2}$.				

$\Gamma(D_s^+ \phi)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{15}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.32	90	28 AAIJ	13R LHCB	<i>pp</i> at 7 TeV
²⁸ AAIJ 13R reports $[\Gamma(B_c^+ \rightarrow D_s^+ \phi)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.8 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.2 \times 10^{-2}$.				

$\Gamma(K^+ K^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{16}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.6 $\times 10^{-7}$	90	29 AAIJ	13BS LHCB	<i>pp</i> at 7 TeV
²⁹ Derived from $\Gamma(K^+ K^0)/\Gamma \times B(\bar{b} \rightarrow B_c) / (B(B^+ \rightarrow K^0 \pi^+) B(\bar{b} \rightarrow B^+)) < 5.8\%$ at 90% CL using normalization mode $B(B^+ \rightarrow K^0 \pi^+) = (23.97 \pm 0.53 \pm 0.71) \times 10^{-6}$ and assuming a <i>B</i> production ratio $f(\bar{b} \rightarrow B_u^+) = 0.33$.				

$\Gamma(B_s^0 \pi^+ / B(\bar{b} \rightarrow B_s)) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{17}/\Gamma \times B$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$2.37 \pm 0.31 \pm 0.11$ -0.13	30 AAIJ	13BU LHCB	<i>pp</i> at 7, 8 TeV
³⁰ The last uncertainty is due to the uncertainty of the B_c^+ lifetime measurement.			

POLARIZATION IN B_c^+ DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal (L) or both are transverse and parallel (\parallel) or perpendicular (\perp) to each other with the parameters Γ_L/Γ , Γ_\perp/Γ , and the relative phases ϕ_\parallel and ϕ_\perp . See the definitions in the note on “Polarization in B Decays” review in the B^0 Particle Listings.

Γ_L/Γ in $B_c^+ \rightarrow J/\psi D_s^{*+}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.48 ± 0.20	³¹ AAIJ	13AS LHCb	pp at 7, 8 TeV
³¹ AAIJ 13AS measures $1 - \Gamma_L/\Gamma = 0.52 \pm 0.20$.			

B_c^\pm REFERENCES

AAIJ	14G	EPJ C74 2839	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AM	PR D87 071103	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AS	PR D87 112012	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D89 019901 (errat.)	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BS	PL B726 646	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BU	PRL 111 181801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BY	JHEP 1309 075	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13CA	JHEP 1311 094	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13R	JHEP 1302 043	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	13	PR D87 011101	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	12AV	PRL 109 232001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12Y	PRL 108 251802	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	09H	PRL 102 092001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
AALTONEN	08M	PRL 100 182002	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	08T	PRL 101 012001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABULENCIA	06C	PRL 96 082002	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABULENCIA	06O	PRL 97 012002	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABE	98M	PRL 81 2432	F. Abe <i>et al.</i>	(CDF Collab.)
Also		PR D58 112004	F. Abe <i>et al.</i>	(CDF Collab.)
ACKERSTAFF	98O	PL B420 157	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
PDG	98	EPJ C3 1	C. Caso <i>et al.</i>	(PDG Collab.)
ABREU	97E	PL B398 207	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BARATE	97H	PL B402 213	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABE	96R	PRL 77 5176	F. Abe <i>et al.</i>	(CDF Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	(PDG Collab.)