

**$J/\psi(1S)$** 

$$J^G(J^{PC}) = 0^-(1^{--})$$

 **$J/\psi(1S)$  MASS**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>3096.916±0.011 OUR AVERAGE</b>				
3096.917±0.010±0.007		AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3096.89 ±0.09	502	<sup>1</sup> ARTAMONOV 00	OLYA	$e^+e^- \rightarrow \text{hadrons}$
3096.91 ±0.03 ±0.01		<sup>2</sup> ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
3096.95 ±0.1 ±0.3	193	BAGLIN 87	SPEC	$\bar{p}p \rightarrow e^+e^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3097.5 ±0.3		GRIBUSHIN 96	FMPS	515 $\pi^- \text{Be} \rightarrow 2\mu X$
3098.4 ±2.0	38k	LEMOIGNE 82	GOLI	185 $\pi^- \text{Be} \rightarrow \gamma\mu^+\mu^-A$
3096.93 ±0.09	502	<sup>3</sup> ZHOLENTZ 80	REDE	$e^+e^-$
3097.0 ±1		<sup>4</sup> BRANDELIK 79C	DASP	$e^+e^-$

<sup>1</sup> Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

<sup>2</sup> Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the  $\psi(2S)$  mass from AULCHENKO 03.

<sup>3</sup> Superseded by ARTAMONOV 00.

<sup>4</sup> From a simultaneous fit to  $e^+e^-$ ,  $\mu^+\mu^-$  and hadronic channels assuming  $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ .

 **$J/\psi(1S)$  WIDTH**

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>92.9± 2.8 OUR AVERAGE</b> Error includes scale factor of 1.1.				
96.1± 3.2	13k	<sup>1</sup> ADAMS 06A	CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
84.4± 8.9		BAI 95B	BES	$e^+e^-$
91 ±11 ±6		<sup>2</sup> ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
85.5 <sup>+</sup> <sub>-</sub> 6.1 5.8		<sup>3</sup> HSUEH 92	RVUE	See $\Upsilon$ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
94.1± 2.7		<sup>4</sup> ANASHIN 10	KEDR	3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
93.7± 3.5	7.8k	<sup>1</sup> AUBERT 04	BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

<sup>1</sup> Calculated by us from the reported values of  $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$  using  $B(e^+e^-) = (5.94 \pm 0.06)\%$  and  $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$ .

<sup>2</sup> The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

<sup>3</sup> Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.

<sup>4</sup> Assuming  $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$  and using  $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$ .

## $J/\psi(1S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ hadrons	(87.7 $\pm$ 0.5 ) %	
$\Gamma_2$ virtual $\gamma \rightarrow$ hadrons	(13.50 $\pm$ 0.30 ) %	
$\Gamma_3$ $g g g$	(64.1 $\pm$ 1.0 ) %	
$\Gamma_4$ $\gamma g g$	( 8.8 $\pm$ 1.1 ) %	
$\Gamma_5$ $e^+ e^-$	( 5.971 $\pm$ 0.032) %	
$\Gamma_6$ $e^+ e^- \gamma$	[a] ( 8.8 $\pm$ 1.4 ) $\times 10^{-3}$	
$\Gamma_7$ $\mu^+ \mu^-$	( 5.961 $\pm$ 0.033) %	

### Decays involving hadronic resonances

$\Gamma_8$ $\rho \pi$	( 1.69 $\pm$ 0.15 ) %	S=2.4
$\Gamma_9$ $\rho^0 \pi^0$	( 5.6 $\pm$ 0.7 ) $\times 10^{-3}$	
$\Gamma_{10}$ $a_2(1320) \rho$	( 1.09 $\pm$ 0.22 ) %	
$\Gamma_{11}$ $\omega \pi^+ \pi^+ \pi^- \pi^-$	( 8.5 $\pm$ 3.4 ) $\times 10^{-3}$	
$\Gamma_{12}$ $\omega \pi^+ \pi^- \pi^0$	( 4.0 $\pm$ 0.7 ) $\times 10^{-3}$	
$\Gamma_{13}$ $\omega \pi^+ \pi^-$	( 8.6 $\pm$ 0.7 ) $\times 10^{-3}$	S=1.1
$\Gamma_{14}$ $\omega f_2(1270)$	( 4.3 $\pm$ 0.6 ) $\times 10^{-3}$	
$\Gamma_{15}$ $K^*(892)^0 \bar{K}^*(892)^0$	( 2.3 $\pm$ 0.7 ) $\times 10^{-4}$	
$\Gamma_{16}$ $K^*(892)^\pm K^*(892)^\mp$	( 1.00 $^{+0.22}_{-0.40}$ ) $\times 10^{-3}$	
$\Gamma_{17}$ $K^*(892)^\pm K^*(800)^\mp$	( 1.1 $^{+1.0}_{-0.6}$ ) $\times 10^{-3}$	
$\Gamma_{18}$ $\eta K^*(892)^0 \bar{K}^*(892)^0$	( 1.15 $\pm$ 0.26 ) $\times 10^{-3}$	
$\Gamma_{19}$ $K^*(892)^0 \bar{K}_2^*(1430)^0 + c.c.$	( 6.0 $\pm$ 0.6 ) $\times 10^{-3}$	
$\Gamma_{20}$ $K^*(892)^0 \bar{K}_2^*(1770)^0 + c.c. \rightarrow$ $K^*(892)^0 K^- \pi^+ + c.c.$	( 6.9 $\pm$ 0.9 ) $\times 10^{-4}$	
$\Gamma_{21}$ $\omega K^*(892) \bar{K} + c.c.$	( 6.1 $\pm$ 0.9 ) $\times 10^{-3}$	
$\Gamma_{22}$ $K^+ K^*(892)^- + c.c.$	( 5.12 $\pm$ 0.30 ) $\times 10^{-3}$	
$\Gamma_{23}$ $K^+ K^*(892)^- + c.c. \rightarrow$ $K^+ K^- \pi^0$	( 1.97 $\pm$ 0.20 ) $\times 10^{-3}$	
$\Gamma_{24}$ $K^+ K^*(892)^- + c.c. \rightarrow$ $K^0 K^\pm \pi^\mp + c.c.$	( 3.0 $\pm$ 0.4 ) $\times 10^{-3}$	
$\Gamma_{25}$ $K^0 \bar{K}^*(892)^0 + c.c.$	( 4.39 $\pm$ 0.31 ) $\times 10^{-3}$	
$\Gamma_{26}$ $K^0 \bar{K}^*(892)^0 + c.c. \rightarrow$ $K^0 K^\pm \pi^\mp + c.c.$	( 3.2 $\pm$ 0.4 ) $\times 10^{-3}$	
$\Gamma_{27}$ $K_1(1400)^\pm K^\mp$	( 3.8 $\pm$ 1.4 ) $\times 10^{-3}$	
$\Gamma_{28}$ $\bar{K}^*(892)^0 K^+ \pi^- + c.c.$	seen	
$\Gamma_{29}$ $\omega \pi^0 \pi^0$	( 3.4 $\pm$ 0.8 ) $\times 10^{-3}$	
$\Gamma_{30}$ $b_1(1235)^\pm \pi^\mp$	[b] ( 3.0 $\pm$ 0.5 ) $\times 10^{-3}$	
$\Gamma_{31}$ $\omega K^\pm K_S^0 \pi^\mp$	[b] ( 3.4 $\pm$ 0.5 ) $\times 10^{-3}$	
$\Gamma_{32}$ $b_1(1235)^0 \pi^0$	( 2.3 $\pm$ 0.6 ) $\times 10^{-3}$	
$\Gamma_{33}$ $\eta K^\pm K_S^0 \pi^\mp$	[b] ( 2.2 $\pm$ 0.4 ) $\times 10^{-3}$	
$\Gamma_{34}$ $\phi K^*(892) \bar{K} + c.c.$	( 2.18 $\pm$ 0.23 ) $\times 10^{-3}$	

$\Gamma_{35}$	$\omega K\bar{K}$		$(1.70 \pm 0.32) \times 10^{-3}$	
$\Gamma_{36}$	$\omega f_0(1710) \rightarrow \omega K\bar{K}$		$(4.8 \pm 1.1) \times 10^{-4}$	
$\Gamma_{37}$	$\phi 2(\pi^+\pi^-)$		$(1.66 \pm 0.23) \times 10^{-3}$	
$\Gamma_{38}$	$\Delta(1232)^{++} \bar{p}\pi^-$		$(1.6 \pm 0.5) \times 10^{-3}$	
$\Gamma_{39}$	$\omega\eta$		$(1.74 \pm 0.20) \times 10^{-3}$	S=1.6
$\Gamma_{40}$	$\phi K\bar{K}$		$(1.83 \pm 0.24) \times 10^{-3}$	S=1.5
$\Gamma_{41}$	$\phi f_0(1710) \rightarrow \phi K\bar{K}$		$(3.6 \pm 0.6) \times 10^{-4}$	
$\Gamma_{42}$	$\phi f_2(1270)$		$(7.2 \pm 1.3) \times 10^{-4}$	
$\Gamma_{43}$	$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$		$(1.10 \pm 0.29) \times 10^{-3}$	
$\Gamma_{44}$	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ \text{ (or c.c.)}$	[b]	$(1.10 \pm 0.12) \times 10^{-3}$	
$\Gamma_{45}$	$\phi f_2'(1525)$		$(8 \pm 4) \times 10^{-4}$	S=2.7
$\Gamma_{46}$	$\phi\pi^+\pi^-$		$(9.4 \pm 0.9) \times 10^{-4}$	S=1.2
$\Gamma_{47}$	$\phi\pi^0\pi^0$		$(5.6 \pm 1.6) \times 10^{-4}$	
$\Gamma_{48}$	$\phi K^\pm K_S^0 \pi^\mp$	[b]	$(7.2 \pm 0.8) \times 10^{-4}$	
$\Gamma_{49}$	$\omega f_1(1420)$		$(6.8 \pm 2.4) \times 10^{-4}$	
$\Gamma_{50}$	$\phi\eta$		$(7.5 \pm 0.8) \times 10^{-4}$	S=1.5
$\Gamma_{51}$	$\Xi^0 \Xi^0$		$(1.20 \pm 0.24) \times 10^{-3}$	
$\Gamma_{52}$	$\Xi(1530)^- \Xi^+$		$(5.9 \pm 1.5) \times 10^{-4}$	
$\Gamma_{53}$	$\rho K^- \bar{\Sigma}(1385)^0$		$(5.1 \pm 3.2) \times 10^{-4}$	
$\Gamma_{54}$	$\omega\pi^0$		$(4.5 \pm 0.5) \times 10^{-4}$	S=1.4
$\Gamma_{55}$	$\phi\eta'(958)$		$(4.0 \pm 0.7) \times 10^{-4}$	S=2.1
$\Gamma_{56}$	$\phi f_0(980)$		$(3.2 \pm 0.9) \times 10^{-4}$	S=1.9
$\Gamma_{57}$	$\phi f_0(980) \rightarrow \phi\pi^+\pi^-$		$(1.8 \pm 0.4) \times 10^{-4}$	
$\Gamma_{58}$	$\phi f_0(980) \rightarrow \phi\pi^0\pi^0$		$(1.7 \pm 0.7) \times 10^{-4}$	
$\Gamma_{59}$	$\eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$		$(3.2 \pm 1.0) \times 10^{-4}$	
$\Gamma_{60}$	$\phi a_0(980)^0 \rightarrow \phi\eta\pi^0$		$(5 \pm 4) \times 10^{-6}$	
$\Gamma_{61}$	$\Xi(1530)^0 \Xi^0$		$(3.2 \pm 1.4) \times 10^{-4}$	
$\Gamma_{62}$	$\Sigma(1385)^- \bar{\Sigma}^+ \text{ (or c.c.)}$	[b]	$(3.1 \pm 0.5) \times 10^{-4}$	
$\Gamma_{63}$	$\phi f_1(1285)$		$(2.6 \pm 0.5) \times 10^{-4}$	S=1.1
$\Gamma_{64}$	$\eta\pi^+\pi^-$		$(4.0 \pm 1.7) \times 10^{-4}$	
$\Gamma_{65}$	$\rho\eta$		$(1.93 \pm 0.23) \times 10^{-4}$	
$\Gamma_{66}$	$\omega\eta'(958)$		$(1.82 \pm 0.21) \times 10^{-4}$	
$\Gamma_{67}$	$\omega f_0(980)$		$(1.4 \pm 0.5) \times 10^{-4}$	
$\Gamma_{68}$	$\rho\eta'(958)$		$(1.05 \pm 0.18) \times 10^{-4}$	
$\Gamma_{69}$	$a_2(1320)^\pm \pi^\mp$	[b]	$< 4.3 \times 10^{-3}$	CL=90%
$\Gamma_{70}$	$K\bar{K}_2^*(1430) + \text{c.c.}$		$< 4.0 \times 10^{-3}$	CL=90%
$\Gamma_{71}$	$K_1(1270)^\pm K^\mp$		$< 3.0 \times 10^{-3}$	CL=90%
$\Gamma_{72}$	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$		$< 2.9 \times 10^{-3}$	CL=90%
$\Gamma_{73}$	$\phi\pi^0$		$< 6.4 \times 10^{-6}$	CL=90%
$\Gamma_{74}$	$\phi\eta(1405) \rightarrow \phi\eta\pi\pi$		$< 2.5 \times 10^{-4}$	CL=90%
$\Gamma_{75}$	$\omega f_2'(1525)$		$< 2.2 \times 10^{-4}$	CL=90%
$\Gamma_{76}$	$\omega X(1835) \rightarrow \omega\rho\bar{p}$		$< 3.9 \times 10^{-6}$	CL=95%
$\Gamma_{77}$	$\eta\phi(2170) \rightarrow$ $\eta K^*(892)^0 \bar{K}^*(892)^0$		$< 2.52 \times 10^{-4}$	CL=90%

$\Gamma_{78}$	$\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}$	$< 8.2$	$\times 10^{-6}$	CL=90%
$\Gamma_{79}$	$\Delta(1232)^+ \bar{p}$	$< 1$	$\times 10^{-4}$	CL=90%
$\Gamma_{80}$	$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$	$< 4.1$	$\times 10^{-6}$	CL=90%
$\Gamma_{81}$	$\Theta(1540) \bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + \text{c.c.}$	$< 1.1$	$\times 10^{-5}$	CL=90%
$\Gamma_{82}$	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	$< 2.1$	$\times 10^{-5}$	CL=90%
$\Gamma_{83}$	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	$< 1.6$	$\times 10^{-5}$	CL=90%
$\Gamma_{84}$	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	$< 5.6$	$\times 10^{-5}$	CL=90%
$\Gamma_{85}$	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	$< 1.1$	$\times 10^{-5}$	CL=90%
$\Gamma_{86}$	$\Sigma^0 \bar{\Lambda}$	$< 9$	$\times 10^{-5}$	CL=90%

### Decays into stable hadrons

$\Gamma_{87}$	$2(\pi^+ \pi^-) \pi^0$	$( 4.1 \pm 0.5 ) \%$		S=2.4
$\Gamma_{88}$	$3(\pi^+ \pi^-) \pi^0$	$( 2.9 \pm 0.6 ) \%$		
$\Gamma_{89}$	$\pi^+ \pi^- \pi^0$	$( 2.11 \pm 0.07 ) \%$		S=1.5
$\Gamma_{90}$	$\pi^+ \pi^- \pi^0 K^+ K^-$	$( 1.79 \pm 0.29 ) \%$		S=2.2
$\Gamma_{91}$	$4(\pi^+ \pi^-) \pi^0$	$( 9.0 \pm 3.0 ) \times 10^{-3}$		
$\Gamma_{92}$	$\pi^+ \pi^- K^+ K^-$	$( 6.6 \pm 0.5 ) \times 10^{-3}$		
$\Gamma_{93}$	$\pi^+ \pi^- K^+ K^- \eta$	$( 1.84 \pm 0.28 ) \times 10^{-3}$		
$\Gamma_{94}$	$\pi^0 \pi^0 K^+ K^-$	$( 2.45 \pm 0.31 ) \times 10^{-3}$		
$\Gamma_{95}$	$K \bar{K} \pi$	$( 6.1 \pm 1.0 ) \times 10^{-3}$		
$\Gamma_{96}$	$2(\pi^+ \pi^-)$	$( 3.57 \pm 0.30 ) \times 10^{-3}$		
$\Gamma_{97}$	$3(\pi^+ \pi^-)$	$( 4.3 \pm 0.4 ) \times 10^{-3}$		
$\Gamma_{98}$	$2(\pi^+ \pi^- \pi^0)$	$( 1.62 \pm 0.21 ) \%$		
$\Gamma_{99}$	$2(\pi^+ \pi^-) \eta$	$( 2.29 \pm 0.24 ) \times 10^{-3}$		
$\Gamma_{100}$	$3(\pi^+ \pi^-) \eta$	$( 7.2 \pm 1.5 ) \times 10^{-4}$		
$\Gamma_{101}$	$\rho \bar{\rho}$	$( 2.120 \pm 0.029 ) \times 10^{-3}$		
$\Gamma_{102}$	$\rho \bar{\rho} \pi^0$	$( 1.19 \pm 0.08 ) \times 10^{-3}$		S=1.1
$\Gamma_{103}$	$\rho \bar{\rho} \pi^+ \pi^-$	$( 6.0 \pm 0.5 ) \times 10^{-3}$		S=1.3
$\Gamma_{104}$	$\rho \bar{\rho} \pi^+ \pi^- \pi^0$	[c] $( 2.3 \pm 0.9 ) \times 10^{-3}$		S=1.9
$\Gamma_{105}$	$\rho \bar{\rho} \eta$	$( 2.00 \pm 0.12 ) \times 10^{-3}$		
$\Gamma_{106}$	$\rho \bar{\rho} \rho$	$< 3.1$	$\times 10^{-4}$	CL=90%
$\Gamma_{107}$	$\rho \bar{\rho} \omega$	$( 9.8 \pm 1.0 ) \times 10^{-4}$		S=1.3
$\Gamma_{108}$	$\rho \bar{\rho} \eta'(958)$	$( 2.1 \pm 0.4 ) \times 10^{-4}$		
$\Gamma_{109}$	$\rho \bar{\rho} \phi$	$( 4.5 \pm 1.5 ) \times 10^{-5}$		
$\Gamma_{110}$	$n \bar{n}$	$( 2.09 \pm 0.16 ) \times 10^{-3}$		
$\Gamma_{111}$	$n \bar{n} \pi^+ \pi^-$	$( 4 \pm 4 ) \times 10^{-3}$		
$\Gamma_{112}$	$\Sigma^+ \bar{\Sigma}^-$	$( 1.50 \pm 0.24 ) \times 10^{-3}$		
$\Gamma_{113}$	$\Sigma^0 \bar{\Sigma}^0$	$( 1.29 \pm 0.09 ) \times 10^{-3}$		
$\Gamma_{114}$	$2(\pi^+ \pi^-) K^+ K^-$	$( 4.7 \pm 0.7 ) \times 10^{-3}$		S=1.3
$\Gamma_{115}$	$\rho \bar{n} \pi^-$	$( 2.12 \pm 0.09 ) \times 10^{-3}$		
$\Gamma_{116}$	$n N(1440)$	seen		
$\Gamma_{117}$	$n N(1520)$	seen		
$\Gamma_{118}$	$n N(1535)$	seen		

$\Gamma_{119}$	$\Xi^- \Xi^+$	( 8.6 ± 1.1 ) × 10 <sup>-4</sup>	S=1.2
$\Gamma_{120}$	$\Lambda \bar{\Lambda}$	( 1.61 ± 0.15 ) × 10 <sup>-3</sup>	S=1.9
$\Gamma_{121}$	$\Lambda \bar{\Sigma}^- \pi^+$ (or c.c.)	[b] ( 8.3 ± 0.7 ) × 10 <sup>-4</sup>	S=1.2
$\Gamma_{122}$	$\rho K^- \bar{\Lambda}$	( 8.9 ± 1.6 ) × 10 <sup>-4</sup>	
$\Gamma_{123}$	$2(K^+ K^-)$	( 7.6 ± 0.9 ) × 10 <sup>-4</sup>	
$\Gamma_{124}$	$\rho K^- \bar{\Sigma}^0$	( 2.9 ± 0.8 ) × 10 <sup>-4</sup>	
$\Gamma_{125}$	$K^+ K^-$	( 2.70 ± 0.17 ) × 10 <sup>-4</sup>	
$\Gamma_{126}$	$K_S^0 K_L^0$	( 2.1 ± 0.4 ) × 10 <sup>-4</sup>	S=3.2
$\Gamma_{127}$	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	( 4.3 ± 1.0 ) × 10 <sup>-3</sup>	
$\Gamma_{128}$	$\Lambda \bar{\Lambda} \eta$	( 1.62 ± 0.17 ) × 10 <sup>-4</sup>	
$\Gamma_{129}$	$\Lambda \bar{\Lambda} \pi^0$	( 3.8 ± 0.4 ) × 10 <sup>-5</sup>	
$\Gamma_{130}$	$\bar{\Lambda} n K_S^0$ + c.c.	( 6.5 ± 1.1 ) × 10 <sup>-4</sup>	
$\Gamma_{131}$	$\pi^+ \pi^-$	( 1.47 ± 0.14 ) × 10 <sup>-4</sup>	
$\Gamma_{132}$	$\Lambda \bar{\Sigma}^+ + c.c.$	( 2.83 ± 0.23 ) × 10 <sup>-5</sup>	
$\Gamma_{133}$	$K_S^0 K_S^0$	< 1 × 10 <sup>-6</sup>	CL=95%

### Radiative decays

$\Gamma_{134}$	$3\gamma$	( 1.16 ± 0.22 ) × 10 <sup>-5</sup>	
$\Gamma_{135}$	$4\gamma$	< 9 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{136}$	$5\gamma$	< 1.5 × 10 <sup>-5</sup>	CL=90%
$\Gamma_{137}$	$\gamma \eta_c(1S)$	( 1.7 ± 0.4 ) %	S=1.6
$\Gamma_{138}$	$\gamma \eta_c(1S) \rightarrow 3\gamma$	( 3.8 <sup>+1.3</sup> <sub>-1.0</sub> ) × 10 <sup>-6</sup>	S=1.1
$\Gamma_{139}$	$\gamma \pi^+ \pi^- 2\pi^0$	( 8.3 ± 3.1 ) × 10 <sup>-3</sup>	
$\Gamma_{140}$	$\gamma \eta \pi \pi$	( 6.1 ± 1.0 ) × 10 <sup>-3</sup>	
$\Gamma_{141}$	$\gamma \eta_2(1870) \rightarrow \gamma \eta \pi^+ \pi^-$	( 6.2 ± 2.4 ) × 10 <sup>-4</sup>	
$\Gamma_{142}$	$\gamma \eta(1405/1475) \rightarrow \gamma K \bar{K} \pi$	[d] ( 2.8 ± 0.6 ) × 10 <sup>-3</sup>	S=1.6
$\Gamma_{143}$	$\gamma \eta(1405/1475) \rightarrow \gamma \gamma \rho^0$	( 7.8 ± 2.0 ) × 10 <sup>-5</sup>	S=1.8
$\Gamma_{144}$	$\gamma \eta(1405/1475) \rightarrow \gamma \eta \pi^+ \pi^-$	( 3.0 ± 0.5 ) × 10 <sup>-4</sup>	
$\Gamma_{145}$	$\gamma \eta(1405/1475) \rightarrow \gamma \gamma \phi$	< 8.2 × 10 <sup>-5</sup>	CL=95%
$\Gamma_{146}$	$\gamma \rho \rho$	( 4.5 ± 0.8 ) × 10 <sup>-3</sup>	
$\Gamma_{147}$	$\gamma \rho \omega$	< 5.4 × 10 <sup>-4</sup>	CL=90%
$\Gamma_{148}$	$\gamma \rho \phi$	< 8.8 × 10 <sup>-5</sup>	CL=90%
$\Gamma_{149}$	$\gamma \eta'(958)$	( 5.15 ± 0.16 ) × 10 <sup>-3</sup>	S=1.2
$\Gamma_{150}$	$\gamma 2\pi^+ 2\pi^-$	( 2.8 ± 0.5 ) × 10 <sup>-3</sup>	S=1.9
$\Gamma_{151}$	$\gamma f_2(1270) f_2(1270)$	( 9.5 ± 1.7 ) × 10 <sup>-4</sup>	
$\Gamma_{152}$	$\gamma f_2(1270) f_2(1270)$ (non resonant)	( 8.2 ± 1.9 ) × 10 <sup>-4</sup>	
$\Gamma_{153}$	$\gamma K^+ K^- \pi^+ \pi^-$	( 2.1 ± 0.6 ) × 10 <sup>-3</sup>	
$\Gamma_{154}$	$\gamma f_4(2050)$	( 2.7 ± 0.7 ) × 10 <sup>-3</sup>	
$\Gamma_{155}$	$\gamma \omega \omega$	( 1.61 ± 0.33 ) × 10 <sup>-3</sup>	
$\Gamma_{156}$	$\gamma \eta(1405/1475) \rightarrow \gamma \rho^0 \rho^0$	( 1.7 ± 0.4 ) × 10 <sup>-3</sup>	S=1.3
$\Gamma_{157}$	$\gamma f_2(1270)$	( 1.43 ± 0.11 ) × 10 <sup>-3</sup>	
$\Gamma_{158}$	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	( 8.5 <sup>+1.2</sup> <sub>-0.9</sub> ) × 10 <sup>-4</sup>	S=1.2

$\Gamma_{159}$	$\gamma f_0(1710) \rightarrow \gamma \pi \pi$	$( 4.0 \pm 1.0 ) \times 10^{-4}$	
$\Gamma_{160}$	$\gamma f_0(1710) \rightarrow \gamma \omega \omega$	$( 3.1 \pm 1.0 ) \times 10^{-4}$	
$\Gamma_{161}$	$\gamma \eta$	$( 1.104 \pm 0.034 ) \times 10^{-3}$	
$\Gamma_{162}$	$\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi$	$( 7.9 \pm 1.3 ) \times 10^{-4}$	
$\Gamma_{163}$	$\gamma f_1(1285)$	$( 6.1 \pm 0.8 ) \times 10^{-4}$	
$\Gamma_{164}$	$\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-$	$( 4.5 \pm 1.2 ) \times 10^{-4}$	
$\Gamma_{165}$	$\gamma f_2'(1525)$	$( 4.5 \begin{smallmatrix} +0.7 \\ -0.4 \end{smallmatrix} ) \times 10^{-4}$	
$\Gamma_{166}$	$\gamma f_2(1640) \rightarrow \gamma \omega \omega$	$( 2.8 \pm 1.8 ) \times 10^{-4}$	
$\Gamma_{167}$	$\gamma f_2(1910) \rightarrow \gamma \omega \omega$	$( 2.0 \pm 1.4 ) \times 10^{-4}$	
$\Gamma_{168}$	$\gamma f_0(1800) \rightarrow \gamma \omega \phi$	$( 2.5 \pm 0.6 ) \times 10^{-4}$	
$\Gamma_{169}$	$\gamma f_2(1950) \rightarrow$ $\gamma K^*(892) \bar{K}^*(892)$	$( 7.0 \pm 2.2 ) \times 10^{-4}$	
$\Gamma_{170}$	$\gamma K^*(892) \bar{K}^*(892)$	$( 4.0 \pm 1.3 ) \times 10^{-3}$	
$\Gamma_{171}$	$\gamma \phi \phi$	$( 4.0 \pm 1.2 ) \times 10^{-4}$	S=2.1
$\Gamma_{172}$	$\gamma p \bar{p}$	$( 3.8 \pm 1.0 ) \times 10^{-4}$	
$\Gamma_{173}$	$\gamma \eta(2225)$	$( 3.3 \pm 0.5 ) \times 10^{-4}$	
$\Gamma_{174}$	$\gamma \eta(1760) \rightarrow \gamma \rho^0 \rho^0$	$( 1.3 \pm 0.9 ) \times 10^{-4}$	
$\Gamma_{175}$	$\gamma \eta(1760) \rightarrow \gamma \omega \omega$	$( 1.98 \pm 0.33 ) \times 10^{-3}$	
$\Gamma_{176}$	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	$( 2.6 \pm 0.4 ) \times 10^{-4}$	
$\Gamma_{177}$	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$( 7.7 \begin{smallmatrix} +1.5 \\ -0.9 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{178}$	$\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-)$	$( 2.4 \begin{smallmatrix} +0.7 \\ -0.8 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{179}$	$\gamma (K \bar{K} \pi) [J^{PC} = 0^{-+}]$	$( 7 \pm 4 ) \times 10^{-4}$	S=2.1
$\Gamma_{180}$	$\gamma \pi^0$	$( 3.49 \begin{smallmatrix} +0.33 \\ -0.30 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{181}$	$\gamma p \bar{p} \pi^+ \pi^-$	$< 7.9 \times 10^{-4}$	CL=90%
$\Gamma_{182}$	$\gamma \Lambda \bar{\Lambda}$	$< 1.3 \times 10^{-4}$	CL=90%
$\Gamma_{183}$	$\gamma f_0(2200)$		
$\Gamma_{184}$	$\gamma f_J(2220)$	$> 2.50 \times 10^{-3}$	CL=99.9%
$\Gamma_{185}$	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$( 8 \pm 4 ) \times 10^{-5}$	
$\Gamma_{186}$	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 3.6 \times 10^{-5}$	
$\Gamma_{187}$	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$( 1.5 \pm 0.8 ) \times 10^{-5}$	
$\Gamma_{188}$	$\gamma f_0(1500)$	$( 1.01 \pm 0.32 ) \times 10^{-4}$	
$\Gamma_{189}$	$\gamma A \rightarrow \gamma \text{invisible}$	$[e] < 6.3 \times 10^{-6}$	CL=90%
$\Gamma_{190}$	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	$[f] < 2.1 \times 10^{-5}$	CL=90%

### Weak decays

$\Gamma_{191}$	$D^- e^+ \nu_e + \text{c.c.}$	$< 1.2 \times 10^{-5}$	CL=90%
$\Gamma_{192}$	$\bar{D}^0 e^+ e^- + \text{c.c.}$	$< 1.1 \times 10^{-5}$	CL=90%
$\Gamma_{193}$	$D_s^- e^+ \nu_e + \text{c.c.}$	$< 3.6 \times 10^{-5}$	CL=90%
$\Gamma_{194}$	$D^- \pi^+ + \text{c.c.}$	$< 7.5 \times 10^{-5}$	CL=90%
$\Gamma_{195}$	$\bar{D}^0 K^0 + \text{c.c.}$	$< 1.7 \times 10^{-4}$	CL=90%
$\Gamma_{196}$	$D_s^- \pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%

**Charge conjugation (C), Parity (P),  
Lepton Family number (LF) violating modes**

$\Gamma_{197}$	$\gamma\gamma$	C	< 5	$\times 10^{-6}$	CL=90%
$\Gamma_{198}$	$e^\pm \mu^\mp$	LF	< 1.6	$\times 10^{-7}$	CL=90%
$\Gamma_{199}$	$e^\pm \tau^\mp$	LF	< 8.3	$\times 10^{-6}$	CL=90%
$\Gamma_{200}$	$\mu^\pm \tau^\mp$	LF	< 2.0	$\times 10^{-6}$	CL=90%

**Other decays**

$\Gamma_{201}$	invisible		< 7	$\times 10^{-4}$	CL=90%
----------------	-----------	--	-----	------------------	--------

[a] For  $E_\gamma > 100$  MeV.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

[c] Includes  $p\bar{p}\pi^+\pi^-\gamma$  and excludes  $p\bar{p}\eta$ ,  $p\bar{p}\omega$ ,  $p\bar{p}\eta'$ .

[d] See the "Note on the  $\eta(1405)$ " in the  $\eta(1405)$  Particle Listings.

[e] For a narrow state  $A$  with mass less than 960 MeV.

[f] For a narrow scalar or pseudoscalar  $A^0$  with mass 0.21–3.0 GeV.

**$J/\psi(1S)$  PARTIAL WIDTHS**

**$\Gamma(\text{hadrons})$**

**$\Gamma_1$**

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$74.1 \pm 8.1$	BAI	95B	BES $e^+e^-$
$59 \pm 24$	BALDINI-...	75	FRAG $e^+e^-$
$59 \pm 14$	BOYARSKI	75	MRK1 $e^+e^-$
$50 \pm 25$	ESPOSITO	75B	FRAM $e^+e^-$

**$\Gamma(e^+e^-)$**

**$\Gamma_5$**

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.55 ± 0.14 ± 0.02 OUR EVALUATION</b>				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$5.71 \pm 0.16$	13k	<sup>1</sup> ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
$5.57 \pm 0.19$	7.8k	<sup>1</sup> AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$
$5.14 \pm 0.39$		BAI	95B	BES $e^+e^-$
$5.36^{+0.29}_{-0.28}$		<sup>2</sup> HSUEH	92	RVUE See $\Upsilon$ mini-review
$4.72 \pm 0.35$		ALEXANDER	89	RVUE See $\Upsilon$ mini-review
$4.4 \pm 0.6$		<sup>2</sup> BRANDELIK	79C	DASP $e^+e^-$
$4.6 \pm 0.8$		<sup>3</sup> BALDINI-...	75	FRAG $e^+e^-$
$4.8 \pm 0.6$		BOYARSKI	75	MRK1 $e^+e^-$
$4.6 \pm 1.0$		ESPOSITO	75B	FRAM $e^+e^-$

<sup>1</sup> Calculated by us from the reported values of  $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$  using  $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$ .

<sup>2</sup> From a simultaneous fit to  $e^+e^-$ ,  $\mu^+\mu^-$ , and hadronic channels assuming  $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ .

<sup>3</sup> Assuming equal partial widths for  $e^+e^-$  and  $\mu^+\mu^-$ .

$\Gamma(\mu^+ \mu^-)$   $\Gamma_7$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.13 ± 0.52	BAI	95B	BES	$e^+ e^-$
4.8 ± 0.6	BOYARSKI	75	MRK1	$e^+ e^-$
5 ± 1	ESPOSITO	75B	FRAM	$e^+ e^-$

$\Gamma(\gamma\gamma)$   $\Gamma_{197}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<5.4	90	BRANDELIK	79C	DASP	$e^+ e^-$
------	----	-----------	-----	------	-----------

$J/\psi(1S) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into  $e^+ e^-$  and with the total width is obtained from the integrated cross section into channel  $i$  in the  $e^+ e^-$  annihilation.

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_1 \Gamma_5 / \Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4 ± 0.8	<sup>1</sup> BALDINI-...	75	FRAG	$e^+ e^-$
3.9 ± 0.8	<sup>1</sup> ESPOSITO	75B	FRAM	$e^+ e^-$

<sup>1</sup>Data redundant with branching ratios or partial widths above.

$\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_5 \Gamma_5 / \Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

<b>332.3 ± 6.4 ± 4.8</b>	ANASHIN	10	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-$
--------------------------	---------	----	------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

350 ± 20	BRANDELIK	79C	DASP	$e^+ e^-$
320 ± 70	<sup>1</sup> BALDINI-...	75	FRAG	$e^+ e^-$
340 ± 90	<sup>1</sup> ESPOSITO	75B	FRAM	$e^+ e^-$
360 ± 100	<sup>1</sup> FORD	75	SPEC	$e^+ e^-$

<sup>1</sup>Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_7 \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**334 ± 5 OUR AVERAGE**

331.8 ± 5.2 ± 6.3		ANASHIN	10	KEDR	3.097 $e^+ e^- \rightarrow \mu^+ \mu^-$
338.4 ± 5.8 ± 7.1	13k	ADAMS	06A	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
330.1 ± 7.7 ± 7.3	7.8k	AUBERT	04	BABR	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

510 ± 90		DASP	75	DASP	$e^+ e^-$
380 ± 50		<sup>1</sup> ESPOSITO	75B	FRAM	$e^+ e^-$

<sup>1</sup>Data redundant with branching ratios or partial widths above.



$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{12}\Gamma_5/\Gamma$

VALUE ( $10^{-2}$ keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.2±0.3±0.2</b>	170	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{13}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>53.6±5.0±0.4</b>	788	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$

<sup>1</sup> AUBERT 07AU reports [ $\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)$ ] =  $47.8 \pm 3.1 \pm 3.2$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{19}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>33±4±1</b>	$317 \pm 23$	<sup>1,2</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

<sup>1</sup> Dividing by 2/3 to take into account that  $B(K^{*0} \rightarrow K^+\pi^-) = 2/3$ .

<sup>2</sup> AUBERT 07AK reports [ $\Gamma(J/\psi(1S) \rightarrow K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(K_2^*(1430) \rightarrow K\pi)$ ] =  $16.4 \pm 1.1 \pm 1.4$  eV which we divide by our best value  $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0\bar{K}_2(1770)^0 + \text{c.c.} \rightarrow K^*(892)^0K^-\pi^+ + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{20}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8±0.4±0.3</b>	$110 \pm 14$	<sup>1</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

<sup>1</sup> Dividing by 2/3 to take into account that  $B(K^{*0} \rightarrow K^+\pi^-) = 2/3$ .

$\Gamma(K^+K^*(892)^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{22}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>29.0±1.7±1.3</b>	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^+K^*(892)^-\gamma$

$\Gamma(K^+K^*(892)^- + \text{c.c.} \rightarrow K^+K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{23}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.96±0.85±0.70</b>	155	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\gamma$

$\Gamma(K^+K^*(892)^- + \text{c.c.} \rightarrow K^0K^\pm\pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{24}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.76±1.70±1.00</b>	89	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K_S^0K^\pm\pi^\mp\gamma$

$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{25}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>26.6±2.5±1.5</b>	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^0\bar{K}^*(892)^0\gamma$

$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0K^\pm\pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{26}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>17.70±1.70±1.00</b>	94	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K_S^0K^\pm\pi^\mp\gamma$

**$\Gamma(\omega K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{35}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.70±1.98±0.03</b>	24	<sup>1</sup> AUBERT	07AU BABR	10.6 e <sup>+</sup> e <sup>-</sup> → ω K <sup>+</sup> K <sup>-</sup> γ

<sup>1</sup> AUBERT 07AU reports [ $\Gamma(J/\psi(1S) \rightarrow \omega K\bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B(ω(782) → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>)] = 3.3 ± 1.3 ± 1.2 eV which we divide by our best value B(ω(782) → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>) = (89.2 ± 0.7) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\phi 2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{37}\Gamma_5/\Gamma$**

VALUE (10 <sup>-2</sup> keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.96±0.19±0.01</b>	35	<sup>1</sup> AUBERT	06D BABR	10.6 e <sup>+</sup> e <sup>-</sup> → φ 2(π <sup>+</sup> π <sup>-</sup> ) γ

<sup>1</sup> AUBERT 06D reports [ $\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+\pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B(φ(1020) → K<sup>+</sup>K<sup>-</sup>)] = (0.47 ± 0.09 ± 0.03) × 10<sup>-2</sup> keV which we divide by our best value B(φ(1020) → K<sup>+</sup>K<sup>-</sup>) = (48.9 ± 0.5) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{46}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.8 ± 0.4 OUR AVERAGE</b>				

4.52±0.48±0.04	254 ± 23	<sup>1</sup> SHEN	09 BELL	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> γ
5.33±0.71±0.05	103	<sup>2</sup> AUBERT, BE	06D BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> γ

<sup>1</sup> SHEN 09 reports 4.50 ± 0.41 ± 0.26 eV from a measurement of [ $\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B(φ(1020) → K<sup>+</sup>K<sup>-</sup>)] assuming B(φ(1020) → K<sup>+</sup>K<sup>-</sup>) = (49.2 ± 0.6) × 10<sup>-2</sup>, which we rescale to our best value B(φ(1020) → K<sup>+</sup>K<sup>-</sup>) = (48.9 ± 0.5) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> AUBERT, BE 06D reports [ $\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B(φ(1020) → K<sup>+</sup>K<sup>-</sup>)] = 2.61 ± 0.30 ± 0.18 eV which we divide by our best value B(φ(1020) → K<sup>+</sup>K<sup>-</sup>) = (48.9 ± 0.5) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{47}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.15±0.88±0.03</b>	23	<sup>1</sup> AUBERT, BE	06D BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>0</sup> π <sup>0</sup> γ

<sup>1</sup> AUBERT, BE 06D reports [ $\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B(φ(1020) → K<sup>+</sup>K<sup>-</sup>)] = 1.54 ± 0.40 ± 0.16 eV which we divide by our best value B(φ(1020) → K<sup>+</sup>K<sup>-</sup>) = (48.9 ± 0.5) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{50}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.1±2.7±0.4</b>	6	<sup>1</sup> AUBERT	07AU BABR	10.6 e <sup>+</sup> e <sup>-</sup> → φ η γ

<sup>1</sup> AUBERT 07AU quotes  $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$  eV.

$\Gamma(\phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{57} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**1.21 ± 0.23 OUR AVERAGE** Error includes scale factor of 1.2.

1.48 ± 0.27 ± 0.09 60 ± 11 <sup>1</sup> SHEN 09 BELL 10.6  $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

1.02 ± 0.24 ± 0.01 20 ± 5 <sup>2</sup> AUBERT 07AK BABR 10.6  $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

<sup>1</sup> Multiplied by 2/3 to take into account the  $\phi \pi^+ \pi^-$  mode only. Using  $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$ .

<sup>2</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.50 \pm 0.11 \pm 0.04$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{58} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**0.96 ± 0.40 ± 0.01** 7.0 ± 2.8 <sup>1</sup> AUBERT 07AK BABR 10.6  $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

<sup>1</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.47 \pm 0.19 \pm 0.05$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{64} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**2.23 ± 0.97 ± 0.03** 9 <sup>1</sup> AUBERT 07AU BABR 10.6  $e^+ e^- \rightarrow \eta \pi^+ \pi^- \gamma$

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \eta \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+ \pi^- \pi^0)] = 0.51 \pm 0.22 \pm 0.03$  eV which we divide by our best value  $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (22.92 \pm 0.28) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{15} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**1.28 ± 0.40 ± 0.11** 25 ± 8 <sup>1</sup> AUBERT 07AK BABR 10.6  $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

<sup>1</sup> Dividing by  $(2/3)^2$  to take twice into account that  $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$ .

$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{42} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**4.0 ± 0.7 ± 0.1** 44 ± 7 <sup>1,2</sup> AUBERT 07AK BABR 10.6  $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

<sup>1</sup> Using  $B(\phi \rightarrow (K + K)^-) = (49.3 \pm 0.6)\%$ .

<sup>2</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = 3.41 \pm 0.55 \pm 0.28$  eV which we divide by our best value  $B(f_2(1270) \rightarrow \pi \pi) = (84.8^{+2.4}_{-1.2}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+ \pi^-) \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{87} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**303 ± 5 ± 18** 4990 AUBERT 07AU BABR 10.6  $e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0 \gamma$

$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{89}\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.122±0.005±0.008</b>	AUBERT,B	04N	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> γ

$\Gamma(\pi^+\pi^-\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{90}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>107.0±4.3±6.4</b>	768	AUBERT	07AU	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> γ

$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{92}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>36.3±1.3±2.1</b>	1586 ± 58	AUBERT	07AK	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

33.6±2.7±2.7	233	<sup>1</sup> AUBERT	05D	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> γ
--------------	-----	---------------------	-----	---

<sup>1</sup>Superseded by AUBERT 07AK.

$\Gamma(\pi^+\pi^-K^+K^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{93}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>25.9±3.9±0.1</b>	73	<sup>1</sup> AUBERT	07AU	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> ηγ

<sup>1</sup>AUBERT 07AU reports [ $\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-K^+K^-\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B(η → 2γ)] = 10.2 ± 1.3 ± 0.8 eV which we divide by our best value B(η → 2γ) = (39.41 ± 0.20) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{94}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>13.6±1.1±1.3</b>	203 ± 16	AUBERT	07AK	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> K <sup>+</sup> K <sup>-</sup> γ

$\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{96}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>20.4±0.9±0.4</b>		LEES	12E	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2π <sup>+</sup> 2π <sup>-</sup> γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

19.5±1.4±1.3	270	<sup>1</sup> AUBERT	05D	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )γ
--------------	-----	---------------------	-----	--

<sup>1</sup>Superseded by LEES 12E.

$\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{97}\Gamma_5/\Gamma$

VALUE (10 <sup>-2</sup> keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.37±0.16±0.14</b>	496	AUBERT	06D	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 3(π <sup>+</sup> π <sup>-</sup> )γ

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{98}\Gamma_5/\Gamma$

VALUE (10 <sup>-2</sup> keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.9±0.5±1.0</b>	761	AUBERT	06D	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> )γ

**$\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{99}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>13.1±2.4±0.1</b>	85	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 5.16 \pm 0.85 \pm 0.39$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.


**$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{101}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.9±0.5 OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.		

11.3±0.4±0.3	821	LEES	13O BABR	$e^+e^- \rightarrow p\bar{p}\gamma$
12.9±0.4±0.4	918	LEES	13Y BABR	$e^+e^- \rightarrow p\bar{p}\gamma$
12.0±0.6±0.5	438	AUBERT	06B	$e^+e^- \rightarrow p\bar{p}\gamma$
9.7±1.7		<sup>1</sup> ARMSTRONG	93B E760	$\bar{p}p \rightarrow e^+e^-$

<sup>1</sup> Using  $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$  MeV.

WEIGHTED AVERAGE  
11.9±0.5 (Error scaled by 1.4)



$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  (eV)

**$\Gamma(\Sigma^0\bar{\Sigma}^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{113}\Gamma_5/\Gamma$**

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>6.4±1.2±0.6</b>	AUBERT	07BD BABR	10.6 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$

**$\Gamma(2(\pi^+\pi^-)K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{114}\Gamma_5/\Gamma$**

VALUE ( $10^{-2}$ keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.75±0.23±0.17</b>	205	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow K^+K^-2(\pi^+\pi^-)\gamma$

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{120}\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT
<b>10.7±0.9±0.7</b>		AUBERT	07BD BABR	10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$

$\Gamma(2(K^+K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{123}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.11±0.39±0.30</b>	156 ± 15	AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow 2(K^+K^-)\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.0 ± 0.7 ± 0.6	38	<sup>1</sup> AUBERT	05D BABR	10.6 $e^+e^- \rightarrow 2(K^+K^-)\gamma$
<sup>1</sup> Superseded by AUBERT 07AK.				

$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{125}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.42±0.23±0.08</b>	51	LEES	13Q BABR	$e^+e^- \rightarrow K^+K^-\gamma$

### $J/\psi(1S)$ BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths)  $\times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  above.

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$				$\Gamma_1/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT
<b>0.877±0.005 OUR AVERAGE</b>				
0.878±0.005		BAI	95B BES	$e^+e^-$
0.86 ± 0.02		BOYARSKI	75 MRK1	$e^+e^-$

$\Gamma(\text{virtual}\gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$				$\Gamma_2/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT
<b>0.135±0.003</b>		<sup>1,2</sup> SETH	04 RVUE	$e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.17 ± 0.02		<sup>1</sup> BOYARSKI	75 MRK1	$e^+e^-$
<sup>1</sup> Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ .				
<sup>2</sup> Using $B(J/\psi \rightarrow \ell^+\ell^-) = (5.90 \pm 0.09)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C.				

$\Gamma(ggg)/\Gamma_{\text{total}}$				$\Gamma_3/\Gamma$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>64.1±1.0</b>	6 M	<sup>1</sup> BESSON	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- + \text{hadrons}$
<sup>1</sup> Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the PDG 08 values of $B(\ell^+\ell^-)$ , $B(\text{virtual}\gamma \rightarrow \text{hadrons})$ , and $B(\gamma\eta_c)$ . The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 08.				

**$\Gamma(\gamma g g)/\Gamma_{\text{total}}$**   **$\Gamma_4/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.79±1.05</b>	200 k	<sup>1</sup> BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- \gamma + \text{hadrons}$

<sup>1</sup> Calculated using the value  $\Gamma(\gamma g g)/\Gamma(g g g) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$  from BESSON 08 and the value of  $\Gamma(g g g)/\Gamma_{\text{total}}$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(g g g)/\Gamma_{\text{total}}$  measurement of BESSON 08.

**$\Gamma(\gamma g g)/\Gamma(g g g)$**   **$\Gamma_4/\Gamma_3$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>13.7±0.1±0.7</b>	6 M	BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

**$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$**   **$\Gamma_5/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.971±0.032 OUR AVERAGE</b>				
5.983±0.007±0.037	720k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.945±0.067±0.042	15k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 ±0.05 ±0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 ±0.33		BAI	95B	BES $e^+ e^-$
5.92 ±0.15 ±0.20		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ±0.9		BOYARSKI	75	MRK1 $e^+ e^-$

**$\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$**   **$\Gamma_6/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.8±1.3±0.4</b>	<sup>1</sup> ARMSTRONG	96	E760 $\bar{p} p \rightarrow e^+ e^- \gamma$

<sup>1</sup> For  $E_\gamma > 100$  MeV.

**$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$**   **$\Gamma_7/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.961±0.033 OUR AVERAGE</b>				
5.973±0.007±0.038	770k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.960±0.065±0.050	17k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 ±0.06 ±0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 ±0.33		BAI	95B	BES $e^+ e^-$
5.90 ±0.15 ±0.19		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ±0.9		BOYARSKI	75	MRK1 $e^+ e^-$

**$\Gamma(e^+ e^-)/\Gamma(\mu^+ \mu^-)$**   **$\Gamma_5/\Gamma_7$**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.0016±0.0031 OUR AVERAGE</b>			
1.0022±0.0044±0.0048	<sup>1</sup> AULCHENKO	14	KEDR $3.097 e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
1.0017±0.0017±0.0033	<sup>2</sup> ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.002 ±0.021 ±0.013	<sup>3</sup> ANASHIN	10	KEDR $3.097 e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
0.997 ±0.012 ±0.006	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.011 ±0.013 ±0.016	BAI	98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.00 ±0.07	BAI	95B	BES	$e^+ e^-$
1.00 ±0.05	BOYARSKI	75	MRK1	$e^+ e^-$
0.91 ±0.15	ESPOSITO	75B	FRAM	$e^+ e^-$
0.93 ±0.10	FORD	75	SPEC	$e^+ e^-$

<sup>1</sup> From 235.3k  $J/\psi \rightarrow e^+ e^-$  and 156.6k  $J/\psi \rightarrow \mu^+ \mu^-$  observed events.

<sup>2</sup> Not independent of the corresponding measurements of  $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$  and  $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ .

<sup>3</sup> Not independent of the corresponding measurements of  $\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$  and  $\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ .


## HADRONIC DECAYS

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$

$\Gamma_8/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.69 ±0.15</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 2.4. See the ideogram below.		
2.18 ±0.19		1,2 AUBERT,B	04N BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
2.184 ±0.005 ±0.201	220k	2,3 BAI	04H BES	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$
2.091 ±0.021 ±0.116		2,4 BAI	04H BES	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
1.21 ±0.20		BAI	96D BES	$e^+ e^- \rightarrow \rho\pi$
1.42 ±0.01 ±0.19		COFFMAN	88 MRK3	$e^+ e^-$
1.3 ±0.3	150	FRANKLIN	83 MRK2	$e^+ e^-$
1.6 ±0.4	183	ALEXANDER	78 PLUT	$e^+ e^-$
1.33 ±0.21		BRANDELIK	78B DASP	$e^+ e^-$
1.0 ±0.2	543	BARTEL	76 CNTR	$e^+ e^-$
1.3 ±0.3	153	JEAN-MARIE	76 MRK1	$e^+ e^-$

WEIGHTED AVERAGE  
1.69±0.15 (Error scaled by 2.4)





<sup>1</sup> From the ratio of  $\Gamma(e^+e^-)B(\pi^+\pi^-\pi^0)$  and  $\Gamma(e^+e^-)B(\mu^+\mu^-)$  (AUBERT 04).

<sup>2</sup> Not independent of their  $B(\pi^+\pi^-\pi^0)$ .

<sup>3</sup> From  $J/\psi \rightarrow \pi^+\pi^-\pi^0$  events directly.

<sup>4</sup> Obtained comparing the rates for  $\pi^+\pi^-\pi^0$  and  $\mu^+\mu^-$ , using  $J/\psi$  events produced via  $\psi(2S) \rightarrow \pi^+\pi^-J/\psi$  and with  $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$ .

### $\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$

$\Gamma_9/\Gamma_8$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.328±0.005±0.027</b>	COFFMAN 88	MRK3	$e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.35 ±0.08	ALEXANDER 78	PLUT	$e^+e^-$
0.32 ±0.08	BRANDELIK 78B	DASP	$e^+e^-$
0.39 ±0.11	BARTEL 76	CNTR	$e^+e^-$
0.37 ±0.09	JEAN-MARIE 76	MRK1	$e^+e^-$

### $\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$

$\Gamma_{10}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.9±2.2 OUR AVERAGE</b>				
11.7±0.7±2.5	7584	AUGUSTIN 89	DM2	$J/\psi \rightarrow \rho^0\rho^\pm\pi^\mp$
8.4±4.5	36	VANNUCCI 77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

### $\Gamma(\omega\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$

$\Gamma_{11}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>85±34</b>	140	VANNUCCI 77	MRK1	$e^+e^- \rightarrow 3(\pi^+\pi^-)\pi^0$

### $\Gamma(\omega\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

$\Gamma_{12}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.40±0.06±0.04</b>	170	<sup>1</sup> AUBERT 06D	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$

<sup>1</sup> Using  $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$  keV.

### $\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma_{13}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.6±0.7 OUR AVERAGE</b>				Error includes scale factor of 1.1.
9.7±0.6±0.6	788	<sup>1</sup> AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
7.0±1.6	18058	AUGUSTIN 89	DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.8±1.6	215	BURMESTER 77D	PLUT	$e^+e^-$
6.8±1.9	348	VANNUCCI 77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

<sup>1</sup> AUBERT 07AU quotes  $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 47.8 \pm 3.1 \pm 3.2$  eV.

### $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$

$\Gamma_{14}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.3±0.6 OUR AVERAGE</b>				
4.3±0.2±0.6	5860	AUGUSTIN 89	DM2	$e^+e^-$
4.0±1.6	70	BURMESTER 77D	PLUT	$e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.9±0.8	81	VANNUCCI 77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3 ± 0.7 ± 0.1</b>		25 ± 8	<sup>1</sup> AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5                      90                      VANNUCCI 77      MRK1     $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$   
<sup>1</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (1.28 \pm 0.40 \pm 0.11) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^\pm K^*(892)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.00 ± 0.19<sup>+0.11</sup><sub>-0.32</sub></b>	323	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$\Gamma(K^*(892)^\pm K^*(800)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.09 ± 0.18<sup>+0.94</sup><sub>-0.54</sub></b>	655	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.15 ± 0.13 ± 0.22</b>	209	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.0 ± 0.6 OUR AVERAGE</b>				
5.9 ± 0.6 ± 0.2	317 ± 23	<sup>1,2</sup> AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
6.7 ± 2.6	40	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

<sup>1</sup> Using  $B(K_2^*(1430)^0 \rightarrow K\pi) = (49.9 \pm 1.2)\%$ .

<sup>2</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (32.9 \pm 2.3 \pm 2.7) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>61 ± 9 OUR AVERAGE</b>				
62.0 ± 6.8 ± 10.6	899 ± 98	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$
65.3 ± 10.2 ± 13.5	176 ± 28	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
53 ± 14 ± 14	530 ± 140	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

**$\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$**   **$\Gamma_{22}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**5.12±0.30 OUR AVERAGE**

5.2 ±0.4 ±0.1		<sup>1</sup> AUBERT	08S	BABR 10.6 $e^+ e^- \rightarrow K^+ K^*(892)^- \gamma$
4.57±0.17±0.70	2285	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
5.26±0.13±0.53		COFFMAN	88	MRK3 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp, K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ±0.6	24	FRANKLIN	83	MRK2 $J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ±0.6	48	VANNUCCI	77	MRK1 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ±1.2	39	BRAUNSCH...	76	DASP $J/\psi \rightarrow K^\pm X$

<sup>1</sup> AUBERT 08S reports  $[\Gamma(J/\psi(1S) \rightarrow K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (29.0 \pm 1.7 \pm 1.3) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{23}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**1.97±0.20±0.05** 155 <sup>1</sup> AUBERT 08S BABR 10.6  $e^+ e^- \rightarrow K^+ K^- \pi^0 \gamma$

<sup>1</sup> AUBERT 08S reports  $[\Gamma(J/\psi(1S) \rightarrow K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (10.96 \pm 0.85 \pm 0.70) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.})/\Gamma_{\text{total}}$**   **$\Gamma_{24}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**3.0±0.4±0.1** 89 <sup>1</sup> AUBERT 08S BABR 10.6  $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

<sup>1</sup> AUBERT 08S reports  $[\Gamma(J/\psi(1S) \rightarrow K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (16.76 \pm 1.70 \pm 1.00) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$**   **$\Gamma_{25}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**4.39±0.31 OUR AVERAGE**

4.8 ±0.5 ±0.1		<sup>1</sup> AUBERT	08S	BABR 10.6 $e^+ e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$
3.96±0.15±0.60	1192	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
4.33±0.12±0.45		COFFMAN	88	MRK3 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ±0.6	45	VANNUCCI	77	MRK1 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
----------	----	----------	----	---

<sup>1</sup> AUBERT 08S reports  $[\Gamma(J/\psi(1S) \rightarrow K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (26.6 \pm 2.5 \pm 1.5) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) / \Gamma(K^+ K^*(892)^- + \text{c.c.})$   $\Gamma_{25} / \Gamma_{22}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.82 ± 0.05 ± 0.09</b>	COFFMAN	88	MRK3 $J/\psi \rightarrow K \bar{K}^*(892) + \text{c.c.}$

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) / \Gamma_{\text{total}}$   $\Gamma_{26} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.2 ± 0.4 ± 0.1</b>	94	<sup>1</sup> AUBERT	08S	BABR 10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

<sup>1</sup> AUBERT 08S reports  $[\Gamma(J/\psi(1S) \rightarrow K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) / \Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (17.70 \pm 1.70 \pm 1.00) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_1(1400)^\pm K^\mp) / \Gamma_{\text{total}}$   $\Gamma_{27} / \Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.8 ± 0.8 ± 1.2</b>	<sup>1</sup> BAI	99C	BES $e^+ e^-$

<sup>1</sup> Assuming  $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$   $\Gamma_{28} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>1</sup> ABLIKIM	06C	BES2 $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$

<sup>1</sup> A  $K_0^*(800)$  is observed by ABLIKIM 06C in the  $K^+ \pi^-$  mass spectrum of the  $\bar{K}^*(892)^0 K^+ \pi^-$  final state against the  $\bar{K}^*(892)$ . A corresponding branching fraction of the  $J/\psi(1S)$  is not presented.

$\Gamma(\omega \pi^0 \pi^0) / \Gamma_{\text{total}}$   $\Gamma_{29} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.4 ± 0.3 ± 0.7</b>	509	AUGUSTIN	89	DM2 $J/\psi \rightarrow \pi^+ \pi^- 3\pi^0$

$\Gamma(b_1(1235)^\pm \pi^\mp) / \Gamma_{\text{total}}$   $\Gamma_{30} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>30 ± 5 OUR AVERAGE</b>				
31 ± 6	4600	AUGUSTIN	89	DM2 $J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
29 ± 7	87	BURMESTER	77D	PLUT $e^+ e^-$

$\Gamma(\omega K^\pm K_S^0 \pi^\mp) / \Gamma_{\text{total}}$   $\Gamma_{31} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>34 ± 5 OUR AVERAGE</b>				
37.7 ± 0.8 ± 5.8	1972 ± 41	ABLIKIM	08E	BES2 $e^+ e^- \rightarrow J/\psi$
29.5 ± 1.4 ± 7.0	879 ± 41	BECKER	87	MRK3 $e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(b_1(1235)^0 \pi^0) / \Gamma_{\text{total}}$   $\Gamma_{32} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>23 ± 3 ± 5</b>	229	AUGUSTIN	89	DM2 $e^+ e^-$

$\Gamma(\eta K^\pm K_S^0 \pi^\mp) / \Gamma_{\text{total}}$   $\Gamma_{33} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>21.8 ± 2.2 ± 3.4</b>	232 ± 23	ABLIKIM	08E	BES2 $e^+ e^- \rightarrow J/\psi$

$\Gamma(\phi K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>21.8±2.3 OUR AVERAGE</b>				
20.8±2.7±3.9	195 ± 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
29.6±3.7±4.7	238 ± 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$
20.7±2.4±3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
20 ± 3 ± 3	155 ± 20	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\omega K\bar{K})/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>17.0± 3.2 OUR AVERAGE</b>				
13.6± 5.0±1.0	24	<sup>1</sup> AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \omega K^+ K^- \gamma$
19.8± 2.1±3.9		<sup>2</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN	77 MRK1	$e^+ e^-$

<sup>1</sup> AUBERT 07AU quotes  $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \omega K^+ K^-) \cdot B(\eta \rightarrow 3\pi) = 3.3 \pm 1.3 \pm 0.2 \text{ eV}$ .

<sup>2</sup> Addition of  $\omega K^+ K^-$  and  $\omega K^0 \bar{K}^0$  branching ratios.

$\Gamma(\omega f_0(1710) \rightarrow \omega K\bar{K})/\Gamma_{\text{total}}$   $\Gamma_{36}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.8±1.1±0.3</b>	<sup>1,2</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

<sup>1</sup> Includes unknown branching fraction  $f_0(1710) \rightarrow K\bar{K}$ .

<sup>2</sup> Addition of  $f_0(1710) \rightarrow K^+ K^-$  and  $f_0(1710) \rightarrow K^0 \bar{K}^0$  branching ratios.

$\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>16.6±2.3 OUR AVERAGE</b>				
17.3±3.3±1.2	35	<sup>1</sup> AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow \phi 2(\pi^+ \pi^-) \gamma$
16.0±1.0±3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

<sup>1</sup> Using  $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04 \text{ keV}$ .

$\Gamma(\Delta(1232)^{++} p \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma$


<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.58±0.23±0.40</b>	332	EATON	84 MRK2	$e^+ e^-$

$\Gamma(\omega \eta)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.74 ± 0.20 OUR AVERAGE</b>				
Error includes scale factor of 1.6. See the ideogram below.				
2.352±0.273	5k	<sup>1</sup> ABLIKIM	06F BES2	$J/\psi \rightarrow \omega \eta$
1.44 ± 0.40 ± 0.14	13	<sup>2</sup> AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow \omega \eta \gamma$
1.43 ± 0.10 ± 0.21	378	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.71 ± 0.08 ± 0.20		COFFMAN	88 MRK3	$e^+ e^- \rightarrow 3\pi \eta$

<sup>1</sup> Using  $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$ ,  $B(\eta \rightarrow \pi^+\pi^-\pi^0) = 22.6 \pm 0.4\%$ ,  $B(\eta \rightarrow \pi^+\pi^-\gamma) = 4.68 \pm 0.11\%$ , and  $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$ .

<sup>2</sup> Using  $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$  keV.



### $\Gamma(\phi K \bar{K})/\Gamma_{\text{total}}$

### $\Gamma_{40}/\Gamma$


VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>18.3± 2.4 OUR AVERAGE</b>		Error includes scale factor of 1.5. See the ideogram below.		
21.4± 0.4±2.2		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi\pi^+\pi^-$
48 $^{+20}_{-16}$ ±6	9.0 $^{+3.7}_{-3.0}$	1,2 HUANG	03	BELL $B^+ \rightarrow (\phi K^+ K^-) K^+$
14.6± 0.8±2.1		<sup>3</sup> FALVARD	88	DM2 $J/\psi \rightarrow$ hadrons
18 ± 8	14	FELDMAN	77	MRK1 $e^+e^-$

<sup>1</sup> We have multiplied  $K^+K^-$  measurement by 2 to obtain  $K\bar{K}$ .

<sup>2</sup> Using  $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$ .

<sup>3</sup> Addition of  $\phi K^+K^-$  and  $\phi K^0\bar{K}^0$  branching ratios.

WEIGHTED AVERAGE  
 $18.3 \pm 2.4$  (Error scaled by 1.5)



**$\Gamma(\phi f_0(1710) \rightarrow \phi K \bar{K})/\Gamma_{\text{total}}$**   **$\Gamma_{41}/\Gamma$**

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>3.6 \pm 0.2 \pm 0.6</math></b>	1,2 FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$

<sup>1</sup> Including interference with  $f'_2(1525)$ .  
<sup>2</sup> Includes unknown branching fraction  $f_0(1710) \rightarrow K \bar{K}$ .

**$\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$**   **$\Gamma_{42}/\Gamma$**

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.72 \pm 0.13 \pm 0.02</math></b>	$44 \pm 7$		1,2 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- < 0.45            90            FALVARD    88    DM2     $J/\psi \rightarrow \text{hadrons}$
- < 0.37            90            VANNUCCI   77    MRK1    $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

<sup>1</sup> Using  $B(f_2(1270) \rightarrow \pi\pi) = (84.8^{+2.4}_{-1.2})\%$   
<sup>2</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (4.02 \pm 0.65 \pm 0.33) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$**   **$\Gamma_{43}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.10 \pm 0.09 \pm 0.28</math></b>	233	EATON 84	MRK2	$e^+ e^-$

$\Gamma(\Sigma(1385)^-\bar{\Sigma}(1385)^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$   $\Gamma_{44}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.10±0.12 OUR AVERAGE</b>				
1.23±0.07±0.30	0.8k	ABLIKIM	12P BES2	$J/\psi \rightarrow \Sigma(1385)^-\bar{\Sigma}(1385)^+$
1.50±0.08±0.38	1k	ABLIKIM	12P BES2	$J/\psi \rightarrow \Sigma(1385)^+\bar{\Sigma}(1385)^-$
1.00±0.04±0.21	0.6k	HENRARD	87 DM2	$e^+e^- \rightarrow \Sigma^{*-}$
1.19±0.04±0.25	0.7k	HENRARD	87 DM2	$e^+e^- \rightarrow \Sigma^{*+}$
0.86±0.18±0.22	56	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^{*-}$
1.03±0.24±0.25	68	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^{*+}$

$\Gamma(\phi f_2'(1525))/\Gamma_{\text{total}}$   $\Gamma_{45}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8 ±4 OUR AVERAGE</b> Error includes scale factor of 2.7.				
12.3±0.6±2.0		<sup>1,2</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
4.8±1.8	46	<sup>1</sup> GIDAL	81 MRK2	$J/\psi \rightarrow K^+K^-K^+K^-$

<sup>1</sup> Re-evaluated using  $B(f_2'(1525) \rightarrow K\bar{K}) = 0.713$ .

<sup>2</sup> Including interference with  $f_0(1710)$ .

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{46}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.94±0.09 OUR AVERAGE</b> Error includes scale factor of 1.2.				
0.96±0.13	103	<sup>1</sup> AUBERT, BE 06D	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
1.09±0.02±0.13		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
0.78±0.03±0.12		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.1 ±0.9	23	FELDMAN	77 MRK1	$e^+e^-$

<sup>1</sup> Derived by us. AUBERT, BE 06D measures  $\Gamma(J/\psi \rightarrow e^+e^-) \times B(J/\psi \rightarrow \phi\pi^+\pi^-) \times B(\phi \rightarrow K^+K^-) = (2.61 \pm 0.30 \pm 0.18) \text{ eV}$

$\Gamma(\phi\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.56±0.16</b>	23	<sup>1</sup> AUBERT, BE 06D	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$

<sup>1</sup> Derived by us. AUBERT, BE 06D measures  $\Gamma(J/\psi \rightarrow e^+e^-) \times B(J/\psi \rightarrow \phi\pi^0\pi^0) \times B(\phi \rightarrow K^+K^-) = (1.54 \pm 0.40 \pm 0.16) \text{ eV}$

$\Gamma(\phi K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{48}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.2±0.8 OUR AVERAGE</b>				
7.4±0.6±1.4	227 ± 19	ABLIKIM	08E BES2	$e^+e^- \rightarrow J/\psi$
7.4±0.9±1.1		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
7 ±0.6±1.0	163 ± 15	BECKER	87 MRK3	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$   $\Gamma_{49}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.8<sup>+1.9</sup><sub>-1.6</sub>±1.7</b>	111 <sup>+31</sup> <sub>-26</sub>	BECKER	87 MRK3	$e^+e^- \rightarrow \text{hadrons}$




**$\Gamma(\phi\eta)/\Gamma_{\text{total}}$**   **$\Gamma_{50}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.75 \pm 0.08</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.5. See the ideogram below.		
$1.4 \pm 0.6 \pm 0.1$	6	<sup>1</sup> AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \phi\eta\gamma$
$0.898 \pm 0.024 \pm 0.089$		ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadr}$
$0.64 \pm 0.04 \pm 0.11$	346	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$0.661 \pm 0.045 \pm 0.078$		COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \eta$

<sup>1</sup> AUBERT 07AU quotes  $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+ K^-) \cdot B(\eta \rightarrow \gamma\gamma) = 0.84 \pm 0.37 \pm 0.05 \text{ eV}$ .

WEIGHTED AVERAGE  
 $0.75 \pm 0.08$  (Error scaled by 1.5)



**$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{51}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.20 \pm 0.12 \pm 0.21</math></b>	206	ABLIKIM	080 BES2	$e^+ e^- \rightarrow J/\psi$

**$\Gamma(\Xi(1530)^- \Xi^+)/\Gamma_{\text{total}}$**   **$\Gamma_{52}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.59 \pm 0.09 \pm 0.12</math></b>	$75 \pm 11$	HENRARD	87 DM2	$e^+ e^-$

**$\Gamma(p K^- \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$**   **$\Gamma_{53}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.51 \pm 0.26 \pm 0.18</math></b>	89	EATON	84 MRK2	$e^+ e^-$


### $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

$\Gamma_{54}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.45 ± 0.05 OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.		
0.538 ± 0.012 ± 0.065	2090	<sup>1</sup> ABLIKIM	06F	BES2 $J/\psi \rightarrow \omega\pi^0$
0.360 ± 0.028 ± 0.054	222	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
0.482 ± 0.019 ± 0.064		COFFMAN	88	MRK3 $e^+e^- \rightarrow \pi^0\pi^+\pi^-\pi^0$

<sup>1</sup> Using  $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$ .

WEIGHTED AVERAGE  
0.45 ± 0.05 (Error scaled by 1.4)




### $\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$

$\Gamma_{55}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.40 ± 0.07 OUR AVERAGE</b>			Error includes scale factor of 2.1. See the ideogram below.		
0.546 ± 0.031 ± 0.056			ABLIKIM	05B	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \text{hadr}$
0.41 ± 0.03 ± 0.08		167	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
0.308 ± 0.034 ± 0.036			COFFMAN	88	MRK3 $e^+e^- \rightarrow K^+K^-\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.3	90	VANNUCCI	77	MRK1	$e^+e^-$
-------	----	----------	----	------	----------



**$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$**   **$\Gamma_{56}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.2 \pm 0.9</math> OUR AVERAGE</b>		Error includes scale factor of 1.9.		
$4.6 \pm 0.4 \pm 0.8$		<sup>1</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
$2.6 \pm 0.6$	50	<sup>1</sup> GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

<sup>1</sup> Assuming  $B(f_0(980) \rightarrow \pi\pi) = 0.78$ .

**$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-)/\Gamma_{\text{total}}$**   **$\Gamma_{57}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.182 \pm 0.042 \pm 0.005</math></b>	$19.5 \pm 4.5$	<sup>1,2</sup> AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^- K^+ K^- \gamma$

<sup>1</sup> Using  $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$ .

<sup>2</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (1.01 \pm 0.22 \pm 0.08) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\phi f_0(980) \rightarrow \phi\pi^0\pi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{58}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.171 \pm 0.073 \pm 0.004</math></b>	$7.0 \pm 2.8$	<sup>1,2</sup> AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^0\pi^0 K^+ K^- \gamma$

<sup>1</sup> Using  $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$ .

<sup>2</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (0.95 \pm 0.39 \pm 0.10) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{59}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.23±0.75±0.73</b>	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$

$\Gamma(\phi a_0(980)^0 \rightarrow \phi\eta\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{60}/\Gamma$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
<b>5.0±2.7±2.5</b>	<sup>1</sup> ABLIKIM	11D BES3	$J/\psi \rightarrow \phi\eta\pi^0$

<sup>1</sup> Assuming  $a_0(980) - f_0(980)$  mixing and isospin breaking via  $\gamma^*$  and  $K^* K$  loops.

$\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$   $\Gamma_{61}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.32±0.12±0.07</b>	24 ± 9	HENRARD	87 DM2	$e^+e^-$

$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.31±0.05 OUR AVERAGE</b>				
0.30±0.03±0.07	74 ± 8	HENRARD	87 DM2	$e^+e^- \rightarrow \Sigma^{*-}$
0.34±0.04±0.07	77 ± 9	HENRARD	87 DM2	$e^+e^- \rightarrow \Sigma^{*+}$
0.29±0.11±0.10	26	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^{*-}$
0.31±0.11±0.11	28	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^{*+}$

$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.6±0.5 OUR AVERAGE</b>	Error includes scale factor of 1.1.			
3.2±0.6±0.4		JOUSSET	90 DM2	$J/\psi \rightarrow \phi 2(\pi^+\pi^-)$
2.1±0.5±0.4	25	<sup>1</sup> JOUSSET	90 DM2	$J/\psi \rightarrow \phi\eta\pi^+\pi^-$
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
0.6±0.2±0.1	16 ± 6	BECKER	87 MRK3	$J/\psi \rightarrow \phi K \bar{K} \pi$

<sup>1</sup> We attribute to the  $f_1(1285)$  the signal observed in the  $\pi^+\pi^-\eta$  invariant mass distribution at 1297 MeV.

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.40±0.17±0.03</b>	9	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$

<sup>1</sup> AUBERT 07AU quotes  $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \eta\pi^+\pi^-) \cdot B(\eta \rightarrow 3\pi) = 0.51 \pm 0.22 \pm 0.03$  eV.

$\Gamma(\rho\eta)/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.193±0.023 OUR AVERAGE</b>				
0.194±0.017±0.029	299	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.193±0.013±0.029		COFFMAN	88 MRK3	$e^+e^- \rightarrow \pi^+\pi^-\eta$

$\Gamma(\omega\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.182±0.021 OUR AVERAGE</b>				
0.226±0.043	218	<sup>1</sup> ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta'$
0.18 $^{+0.10}_{-0.08} \pm 0.03$	6	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.166±0.017±0.019		COFFMAN	88 MRK3	$e^+e^- \rightarrow 3\pi\eta'$

<sup>1</sup> Using  $B(\eta' \rightarrow \pi^+\pi^-\eta) = (44.3 \pm 1.5)\%$ ,  $B(\eta' \rightarrow \pi^+\pi^-\gamma) = 29.5 \pm 1.0\%$ ,  $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$ , and  $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$ .

$\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.41±0.27±0.47</b>			
	<sup>1</sup> AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

<sup>1</sup> Assuming  $B(f_0(980) \rightarrow \pi\pi) = 0.78$ .

$\Gamma(\rho\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{68}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.105±0.018 OUR AVERAGE</b>				
0.083±0.030±0.012	19	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.114±0.014±0.016		COFFMAN	88 MRK3	$J/\psi \rightarrow \pi^+\pi^-\eta'$

$\Gamma(a_2(1320)^\pm\pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{69}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;43</b>	90	BRAUNSCH...	76 DASP	$e^+e^-$

$\Gamma(K\bar{K}_2^*(1430)+\text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;40</b>	90	VANNUCCI	77 MRK1	$e^+e^- \rightarrow K^0\bar{K}_2^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<66	90	BRAUNSCH...	76 DASP	$e^+e^- \rightarrow K^\pm\bar{K}_2^{*\mp}$
-----	----	-------------	---------	--

$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{71}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;3.0</b>	90	<sup>1</sup> BAI	99C BES	$e^+e^-$

<sup>1</sup> Assuming  $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

$\Gamma(K_2^*(1430)^0\bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$   $\Gamma_{72}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;29</b>	90	VANNUCCI	77 MRK1	$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{73}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;6.4</b>	90	ABLIKIM	05B BES2	$e^+e^- \rightarrow J/\psi \rightarrow \phi\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.8	90	COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+K^-\pi^0$
------	----	---------	---------	----------------------------------

$\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_{74}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.5	90	<sup>1</sup> FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$

<sup>1</sup> Includes unknown branching fraction  $\eta(1405) \rightarrow \eta\pi\pi$ .

$\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$   $\Gamma_{75}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	<sup>1</sup> VANNUCCI	77	MRK1 $e^+e^- \rightarrow \pi^+\pi^-\pi^0 K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.8	90	<sup>1</sup> FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$
------	----	----------------------	----	---

<sup>1</sup> Re-evaluated assuming  $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$ .

$\Gamma(\omega X(1835) \rightarrow \omega p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{76}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3.9	95	ABLIKIM	13P	BES3 $J/\psi \rightarrow \gamma\pi^0 p\bar{p}$

$\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{77}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.52	90	ABLIKIM	10C	BES2 $J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

$\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{78}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 0.82	90	ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<20	90	HENRARD	87	DM2 $e^+e^-$
-----	----	---------	----	--------------

$\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}}$   $\Gamma_{79}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	90	HENRARD	87	DM2 $e^+e^-$

$\Gamma(\Lambda(1520)\bar{\Lambda} + \text{c.c.} \rightarrow \gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$   $\Gamma_{80}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4.1	90	ABLIKIM	12B	BES3 $J/\psi \rightarrow \Lambda\bar{\Lambda}\gamma$

$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{81}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	BAI	04G	BES2 $e^+e^-$

$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$   $\Gamma_{82}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	BAI	04G	BES2 $e^+e^-$

$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$   $\Gamma_{83}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	BAI	04G	BES2 $e^+e^-$

$\Gamma(\bar{\Theta}(1540)K^+n \rightarrow K_S^0 \bar{p}K^+n)/\Gamma_{\text{total}}$   $\Gamma_{84}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5.6</b>	90	BAI	04G	BES2 $e^+e^-$

$\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$   $\Gamma_{85}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.1</b>	90	BAI	04G	BES2 $e^+e^-$

$\Gamma(\Sigma^0 \bar{\Lambda})/\Gamma_{\text{total}}$   $\Gamma_{86}/\Gamma$


VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.9</b>	90	HENRARD	87	DM2 $e^+e^-$

————— STABLE HADRONS —————

$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{87}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4.1 \pm 0.5</math> OUR AVERAGE</b>		Error includes scale factor of 2.4. See the ideogram below.		
$5.46 \pm 0.34 \pm 0.14$	4990	<sup>1</sup> AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow 2(\pi^+ \pi^-)\pi^0 \gamma$
$3.25 \pm 0.49$	46055	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
$3.17 \pm 0.42$	147	FRANKLIN	83 MRK2	$e^+e^- \rightarrow \text{hadrons}$
$3.64 \pm 0.52$	1500	BURMESTER	77D PLUT	$e^+e^-$
$4 \pm 1$	675	JEAN-MARIE	76 MRK1	$e^+e^-$

WEIGHTED AVERAGE  
 $4.1 \pm 0.5$  (Error scaled by 2.4)



<sup>1</sup>AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}})] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = 0.303 \pm 0.005 \pm 0.018$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\omega\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-\pi^0))$

$\Gamma_{13}/\Gamma_{87}$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.3	<sup>1</sup> JEAN-MARIE 76	MRK1	$e^+e^-$
<sup>1</sup> Final state $(\pi^+\pi^-\pi^0)$ under the assumption that $\pi\pi$ is isospin 0.			

### $\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$

$\Gamma_{88}/\Gamma$


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.029±0.006 OUR AVERAGE</b>				
0.028±0.009	11	FRANKLIN 83	MRK2	$e^+e^- \rightarrow$ hadrons
0.029±0.007	181	JEAN-MARIE 76	MRK1	$e^+e^-$

### $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

$\Gamma_{89}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>21.1 ±0.7 OUR AVERAGE</b>		Error includes scale factor of 1.5. See the ideogram below.		
21.37±0.04 <sup>+0.64</sup> <sub>-0.62</sub>	1.8M	<sup>1,2</sup> ABLIKIM 12H	BES3	$e^+e^- \rightarrow J/\psi$
22.9 ±2.0 ±0.4	256	<sup>3</sup> AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
21.8 ±1.9		<sup>4,5</sup> AUBERT,B 04N	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
21.84±0.05±2.01	220k	<sup>1,5</sup> BAI 04H	BES	$e^+e^-$
20.91±0.21±1.16		<sup>5,6</sup> BAI 04H	BES	$e^+e^-$
15 ±2	168	FRANKLIN 83	MRK2	$e^+e^-$

WEIGHTED AVERAGE  
21.1±0.7 (Error scaled by 1.5)





<sup>1</sup> From  $J/\psi \rightarrow \pi^+ \pi^- \pi^0$  events directly.

<sup>2</sup> The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of  $J/\psi$  events.

<sup>3</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}} = 0.813 \pm 0.013$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> From the ratio of  $\Gamma(e^+ e^-) B(\pi^+ \pi^- \pi^0)$  and  $\Gamma(e^+ e^-) B(\mu^+ \mu^-)$  (AUBERT 04).

<sup>5</sup> Mostly  $\rho\pi$ , see also  $\rho\pi$  subsection.

<sup>6</sup> Obtained comparing the rates for  $\pi^+ \pi^- \pi^0$  and  $\mu^+ \mu^-$ , using  $J/\psi$  events produced via  $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$  and with  $B(J/\psi \rightarrow \mu^+ \mu^-) = 5.88 \pm 0.10\%$ .

**$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{90}/\Gamma$**

VALUE (units  $10^{-2}$ )      EVTS      DOCUMENT ID      TECN      COMMENT

**1.79±0.29 OUR AVERAGE** Error includes scale factor of 2.2.

1.93±0.14±0.05      768      <sup>1</sup> AUBERT      07AU BABR      10.6  $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$

1.2 ± 0.3      309      VANNUCCI      77      MRK1       $e^+ e^-$

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- \pi^0 K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = 0.1070 \pm 0.0043 \pm 0.0064$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(4(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{91}/\Gamma$**

VALUE (units  $10^{-4}$ )      EVTS      DOCUMENT ID      TECN      COMMENT

**90±30**      13      JEAN-MARIE      76      MRK1       $e^+ e^-$

**$\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{92}/\Gamma$**

VALUE (units  $10^{-3}$ )      EVTS      DOCUMENT ID      TECN      COMMENT

**6.6±0.5 OUR AVERAGE**

6.5±0.4±0.2      1.6k      <sup>1</sup> AUBERT      07AK BABR      10.6  $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

7.2±2.3      205      VANNUCCI      77      MRK1       $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.1±0.7±0.2      233      <sup>2</sup> AUBERT      05D BABR      10.6  $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

<sup>1</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (36.3 \pm 1.3 \pm 2.1) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Superseded by AUBERT 07AK. AUBERT 05D reports  $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (33.6 \pm 2.7 \pm 2.7) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+\pi^-K^+K^-\eta)/\Gamma_{\text{total}}$   $\Gamma_{93}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.84 ± 0.28 ± 0.05</b>	73	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\eta\gamma$

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-K^+K^-\eta)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (10.2 \pm 1.3 \pm 0.8) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0\pi^0K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{94}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.45 ± 0.31 ± 0.06</b>	203 ± 16	<sup>1</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$

<sup>1</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \pi^0\pi^0K^+K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (13.6 \pm 1.1 \pm 1.3) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$   $\Gamma_{95}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>61 ± 10 OUR AVERAGE</b>				
55.2 ± 12.0	25	FRANKLIN	83	MRK2 $e^+e^- \rightarrow K^+K^-\pi^0$
78.0 ± 21.0	126	VANNUCCI	77	MRK1 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$   $\Gamma_{96}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.57 ± 0.30 OUR AVERAGE</b>				
3.53 ± 0.12 ± 0.29	1107	<sup>1</sup> ABLIKIM	05H BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\pi^+\pi^-, J/\psi \rightarrow 2(\pi^+\pi^-)$
4.0 ± 1.0	76	JEAN-MARIE	76	MRK1 $e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.51 ± 0.34 ± 0.09	270	<sup>2</sup> AUBERT	05D BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\gamma$
--------------------	-----	---------------------	----------	---

<sup>1</sup> Computed using  $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ .

<sup>2</sup> AUBERT 05D reports  $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (19.5 \pm 1.4 \pm 1.3) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$   $\Gamma_{97}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>43 ± 4 OUR AVERAGE</b>				
43.0 ± 2.9 ± 2.8	496	<sup>1</sup> AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 3(\pi^+\pi^-)\gamma$
40 ± 20	32	JEAN-MARIE	76	MRK1 $e^+e^-$

<sup>1</sup> Using  $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$  keV.

$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$   $\Gamma_{98}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.62±0.09±0.19</b>	761	<sup>1</sup> AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

<sup>1</sup> Using  $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$  keV.

$\Gamma(2(\pi^+\pi^-\eta))/\Gamma_{\text{total}}$   $\Gamma_{99}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.29±0.24 OUR AVERAGE</b>				
2.35±0.39±0.20	85	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\eta)\gamma$
2.26±0.08±0.27	4839	ABLIKIM	05C BES2	$e^+e^- \rightarrow 2(\pi^+\pi^-\eta)$

<sup>1</sup> AUBERT 07AU quotes  $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow 2(\pi^+\pi^-\eta)) \cdot B(\eta \rightarrow \gamma\gamma) = 5.16 \pm 0.85 \pm 0.39$  eV.

$\Gamma(3(\pi^+\pi^-\eta))/\Gamma_{\text{total}}$   $\Gamma_{100}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.24±0.96±1.11</b>	616	ABLIKIM	05C BES2	$e^+e^- \rightarrow 3(\pi^+\pi^-\eta)$

$\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{101}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.120±0.029 OUR AVERAGE</b>				
2.112±0.004±0.031	314k	ABLIKIM	12C BES3	$e^+e^-$
2.15 ±0.16 ±0.06	317	<sup>1</sup> WU	06 BELL	$B^+ \rightarrow \rho\bar{\rho}K^+$
2.26 ±0.01 ±0.14	63316	BAI	04E BES2	$e^+e^- \rightarrow J/\psi$
1.97 ±0.22	99	BALDINI	98 FENI	$e^+e^-$
1.91 ±0.04 ±0.30		PALLIN	87 DM2	$e^+e^-$
2.16 ±0.07 ±0.15	1420	EATON	84 MRK2	$e^+e^-$
2.5 ±0.4	133	BRANDELIK	79C DASP	$e^+e^-$
2.0 ±0.5		BESCH	78 BONA	$e^+e^-$
2.2 ±0.2	331	<sup>2</sup> PERUZZI	78 MRK1	$e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ±0.3	48	ANTONELLI	93 SPEC	$e^+e^-$
----------	----	-----------	---------	----------

<sup>1</sup> WU 06 reports  $[\Gamma(J/\psi(1S) \rightarrow \rho\bar{\rho})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.027 \pm 0.031) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Assuming angular distribution  $(1+\cos^2\theta)$ .

$\Gamma(\rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{102}/\Gamma$


VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.19±0.08 OUR AVERAGE</b>				Error includes scale factor of 1.1.
1.33±0.02±0.11	11k	ABLIKIM	09B BES2	$e^+e^-$
1.13±0.09±0.09	685	EATON	84 MRK2	$e^+e^-$
1.4 ±0.4		BRANDELIK	79C DASP	$e^+e^-$
1.00±0.15	109	PERUZZI	78 MRK1	$e^+e^-$

$\Gamma(\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma_{103}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.0 ± 0.5 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
6.46 ± 0.17 ± 0.43	1435	EATON	84	MRK2 $e^+e^-$
3.8 ± 1.6	48	BESCH	81	BONA $e^+e^-$
5.5 ± 0.6	533	PERUZZI	78	MRK1 $e^+e^-$

WEIGHTED AVERAGE  
6.0 ± 0.5 (Error scaled by 1.3)



$\Gamma(\rho\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

$\Gamma_{104}/\Gamma$

Including  $\rho\bar{p}\pi^+\pi^-\gamma$  and excluding  $\omega, \eta, \eta'$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3 ± 0.9 OUR AVERAGE</b>				Error includes scale factor of 1.9.
3.36 ± 0.65 ± 0.28	364	EATON	84	MRK2 $e^+e^-$
1.6 ± 0.6	39	PERUZZI	78	MRK1 $e^+e^-$

$\Gamma(\rho\bar{p}\eta)/\Gamma_{\text{total}}$

$\Gamma_{105}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.00 ± 0.12 OUR AVERAGE</b>				
1.91 ± 0.02 ± 0.17	13k	<sup>1</sup> ABLIKIM	09	BES2 $e^+e^-$
2.03 ± 0.13 ± 0.15	826	EATON	84	MRK2 $e^+e^-$
2.5 ± 1.2		BRANDELIK	79C	DASP $e^+e^-$
2.3 ± 0.4	197	PERUZZI	78	MRK1 $e^+e^-$


<sup>1</sup> From the combination of  $\rho\bar{p}\eta \rightarrow \rho\bar{p}\gamma\gamma$  and  $\rho\bar{p}\eta \rightarrow \rho\bar{p}\pi^+\pi^-\pi^0$  channels.

$\Gamma(p\bar{p}\rho)/\Gamma_{\text{total}}$   $\Gamma_{106}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.31</b>	90	EATON	84	MRK2 $e^+e^- \rightarrow \text{hadrons}\gamma$

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$   $\Gamma_{107}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.98±0.10 OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.		
0.90±0.02±0.09	2670	ABLIKIM	13P	BES3 $e^+e^-$
0.98±0.03±0.14	2449	ABLIKIM	08	BES2 $e^+e^-$
1.10±0.17±0.18	486	EATON	84	MRK2 $e^+e^-$
1.6 ±0.3	77	PERUZZI	78	MRK1 $e^+e^-$



$\Gamma(p\bar{p}\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{108}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.21 ±0.04 OUR AVERAGE</b>				
0.200±0.023±0.028	265 ± 31	<sup>1</sup> ABLIKIM	09	BES2 $e^+e^-$
0.68 ±0.23 ±0.17	19	EATON	84	MRK2 $e^+e^-$
1.8 ±0.6	19	PERUZZI	78	MRK1 $e^+e^-$

<sup>1</sup> From the combination of  $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$  and  $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$  channels.

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$   $\Gamma_{109}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.45±0.13±0.07</b>	FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$

$\Gamma(n\bar{n})/\Gamma_{\text{total}}$   $\Gamma_{110}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**2.09±0.16 OUR AVERAGE**

2.07±0.01±0.17	36k	ABLIKIM	12C	BES3 $e^+e^-$
2.31±0.49	79	BALDINI	98	FENI $e^+e^-$
1.8 ±0.9		BESCH	78	BONA $e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.90±0.55	40	ANTONELLI	93	SPEC $e^+e^-$
-----------	----	-----------	----	---------------

$\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{111}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**3.8±3.6**

	5	BESCH	81	BONA $e^+e^-$
--	---	-------	----	---------------

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$   $\Gamma_{112}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**1.50±0.10±0.22**

	399	ABLIKIM	080	BES2 $e^+e^- \rightarrow J/\psi$
--	-----	---------	-----	----------------------------------

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$   $\Gamma_{113}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**1.29±0.09 OUR AVERAGE**

1.15±0.24±0.03		<sup>1</sup> AUBERT	07BD	BABR 10.6 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$
1.33±0.04±0.11	1779	ABLIKIM	06	BES2 $J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.06±0.04±0.23	884 ± 30	PALLIN	87	DM2 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.58±0.16±0.25	90	EATON	84	MRK2 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.3 ±0.4	52	PERUZZI	78	MRK1 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.4 ±2.6	3	BESCH	81	BONA $e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$
----------	---	-------	----	--

<sup>1</sup>AUBERT 07BD reports  $[\Gamma(J/\psi(1S) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (6.4 \pm 1.2 \pm 0.6) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+\pi^-)K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{114}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**47 ± 7 OUR AVERAGE** Error includes scale factor of 1.3.

49.8± 4.2±3.4	205	<sup>1</sup> AUBERT	06D	BABR 10.6 $e^+e^- \rightarrow \omega K^+K^- 2(\pi^+\pi^-)\gamma$
---------------	-----	---------------------	-----	--

31 ±13	30	VANNUCCI	77	MRK1 $e^+e^-$
--------	----	----------	----	---------------

<sup>1</sup>Using  $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$  keV.

$\Gamma(\rho\bar{n}\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{115}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**2.12±0.09 OUR AVERAGE**

2.36±0.02±0.21	59k	ABLIKIM	06K	BES2 $J/\psi \rightarrow \rho\pi^-\bar{n}$
2.47±0.02±0.24	55k	ABLIKIM	06K	BES2 $J/\psi \rightarrow \bar{\rho}\pi^+n$
2.02±0.07±0.16	1288	EATON	84	MRK2 $e^+e^- \rightarrow \rho\pi^-$
1.93±0.07±0.16	1191	EATON	84	MRK2 $e^+e^- \rightarrow \bar{\rho}\pi^+$

1.7 ± 0.7	32	BESCH	81	BONA	$e^+e^- \rightarrow \rho\pi^-$
1.6 ± 1.2	5	BESCH	81	BONA	$e^+e^- \rightarrow \bar{\rho}\pi^+$
2.16 ± 0.29	194	PERUZZI	78	MRK1	$e^+e^- \rightarrow \rho\pi^-$
2.04 ± 0.27	204	PERUZZI	78	MRK1	$e^+e^- \rightarrow \bar{\rho}\pi^+$

$\Gamma(\Xi^- \Xi^+)/\Gamma_{total}$

$\Gamma_{119}/\Gamma$


VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.86 ± 0.11 OUR AVERAGE</b>		Error includes scale factor of 1.2.		
0.90 ± 0.03 ± 0.18	961 ± 35	ABLIKIM	12P	BES2 $J/\psi \rightarrow \Xi^- \Xi^+$
0.70 ± 0.06 ± 0.12	132 ± 11	HENRARD	87	DM2 $e^+e^- \rightarrow \Xi^- \Xi^+$
1.14 ± 0.08 ± 0.20	194	EATON	84	MRK2 $e^+e^- \rightarrow \Xi^- \Xi^+$
1.4 ± 0.5	51	PERUZZI	78	MRK1 $e^+e^- \rightarrow \Xi^- \Xi^+$

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{total}$

$\Gamma_{120}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.61 ± 0.15 OUR AVERAGE</b>		Error includes scale factor of 1.9. See the ideogram below.		
1.93 ± 0.21 ± 0.05		<sup>1</sup> AUBERT	07BD	BABR $10.6 e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
2.03 ± 0.03 ± 0.15	8887	ABLIKIM	06	BES2 $J/\psi \rightarrow \Lambda\bar{\Lambda}$
1.9 <sup>+0.5</sup> <sub>-0.4</sub> ± 0.1	46	<sup>2</sup> WU	06	BELL $B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
1.08 ± 0.06 ± 0.24	631	BAI	98G	BES $e^+e^-$
1.38 ± 0.05 ± 0.20	1847	PALLIN	87	DM2 $e^+e^-$
1.58 ± 0.08 ± 0.19	365	EATON	84	MRK2 $e^+e^-$
2.6 ± 1.6	5	BESCH	81	BONA $e^+e^-$
1.1 ± 0.2	196	PERUZZI	78	MRK1 $e^+e^-$

WEIGHTED AVERAGE  
1.61 ± 0.15 (Error scaled by 1.9)



<sup>1</sup> AUBERT 07BD reports  $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (10.7 \pm 0.9 \pm 0.7) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> WU 06 reports  $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.027 \pm 0.031) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda\bar{\Lambda})/\Gamma(p\bar{p})$	$\Gamma_{120}/\Gamma_{101}$			
VALUE	DOCUMENT ID	TECN	COMMENT	
<b><math>0.90^{+0.15}_{-0.14} \pm 0.10</math></b>	<sup>1</sup> WU	06	BELL	$B^+ \rightarrow p\bar{p}K^+, \Lambda\bar{\Lambda}K^+$

<sup>1</sup> Not independent of other  $J/\psi \rightarrow \Lambda\bar{\Lambda}, p\bar{p}$  branching ratios reported by WU 06.

$\Gamma(\Lambda\bar{\Sigma}^- \pi^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$	$\Gamma_{121}/\Gamma$			
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.83 \pm 0.07</math> OUR AVERAGE</b>	Error includes scale factor of 1.2.			
$0.770 \pm 0.051 \pm 0.083$	335	<sup>1</sup> ABLIKIM	07H BES2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+ \pi^-$
$0.747 \pm 0.056 \pm 0.076$	254	<sup>1</sup> ABLIKIM	07H BES2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$
$0.90 \pm 0.06 \pm 0.16$	$225 \pm 15$	HENRARD	87 DM2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+ \pi^-$
$1.11 \pm 0.06 \pm 0.20$	$342 \pm 18$	HENRARD	87 DM2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$
$1.53 \pm 0.17 \pm 0.38$	135	EATON	84 MRK2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+ \pi^-$
$1.38 \pm 0.21 \pm 0.35$	118	EATON	84 MRK2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$ .

$\Gamma(pK^-\bar{\Lambda})/\Gamma_{\text{total}}$	$\Gamma_{122}/\Gamma$			
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.89 \pm 0.07 \pm 0.14</math></b>	307	EATON	84 MRK2	$e^+e^-$

$\Gamma(2(K^+K^-))/\Gamma_{\text{total}}$	$\Gamma_{123}/\Gamma$			
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.76 \pm 0.09</math> OUR AVERAGE</b>				
$0.74 \pm 0.09 \pm 0.02$	$156 \pm 15$	<sup>1</sup> AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow 2(K^+K^-)\gamma$
$1.4^{+0.5}_{-0.4} \pm 0.2$	$11.0^{+4.3}_{-3.5}$	<sup>2</sup> HUANG	03 BELL	$B^+ \rightarrow 2(K^+K^-)K^+$
$0.7 \pm 0.3$		VANNUCCI	77 MRK1	$e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.72 \pm 0.17 \pm 0.02$  38 <sup>3</sup> AUBERT 05D BABR  $10.6 e^+e^- \rightarrow 2(K^+K^-)\gamma$

<sup>1</sup> AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow 2(K^+K^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (4.11 \pm 0.39 \pm 0.30) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using  $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$ .

<sup>3</sup> Superseded by AUBERT 07AK. AUBERT 05D reports  $[\Gamma(J/\psi(1S) \rightarrow 2(K^+K^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (4.0 \pm 0.7 \pm 0.6) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(pK^-\bar{\Sigma}^0)/\Gamma_{\text{total}}$   $\Gamma_{124}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.29±0.06±0.05</b>	90	EATON	84	MRK2 $e^+e^-$

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{125}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.70±0.17 OUR AVERAGE</b>				
2.86±0.09±0.19	1k	<sup>1</sup> METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^-K^+K^-$
2.39±0.24±0.22	107	BALTRUSAIT..85D	MRK3	$e^+e^-$
2.2 ±0.9	6	BRANDELIK	79C DASP	$e^+e^-$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(K_S^0K_L^0)/\Gamma_{\text{total}}$   $\Gamma_{126}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.1 ±0.4 OUR AVERAGE</b>				Error includes scale factor of 3.2.
2.62±0.15±0.14	0.3k	<sup>1</sup> METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^-K_S^0K_L^0$
1.82±0.04±0.13	2.1k	<sup>2</sup> BAI	04A BES2	$J/\psi \rightarrow K_S^0K_L^0 \rightarrow \pi^+\pi^-X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.18±0.12±0.18		JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
1.01±0.16±0.09	74	BALTRUSAIT..85D	MRK3	$e^+e^-$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Using  $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6868 \pm 0.0027$ .

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{127}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.30±0.13±0.99</b>	2.4k	ABLIKIM	12P BES2	$J/\psi$

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$   $\Gamma_{128}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.2±1.7 OUR AVERAGE</b>				
15.7±0.80±1.54	454	<sup>1</sup> ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
26.2±6.0 ±4.4	44	<sup>2</sup> ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^-p) = 63.9\%$  and  $B(\eta \rightarrow \gamma\gamma) = 39.31\%$ .

<sup>2</sup> Using  $B(\Lambda \rightarrow \pi^-p) = 63.9\%$  and  $B(\eta \rightarrow \gamma\gamma) = 39.4\%$ .

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{129}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.78±0.27±0.30</b>		323	<sup>1</sup> ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.4	90	<sup>2</sup> ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$
23 ±7 ±8	11	BAI	98G BES	$e^+e^-$
22 ±5 ±5	19	HENRARD	87 DM2	$e^+e^-$

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^-p) = 63.9\%$  and  $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$ .

<sup>2</sup> Using  $B(\Lambda \rightarrow \pi^-p) = 63.9\%$ .

$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{130}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.46 ± 0.20 ± 1.07</b>	1058	<sup>1</sup> ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$

<sup>1</sup> Using  $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$  and  $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$ .

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{131}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.47 ± 0.14 OUR AVERAGE</b>				
1.47 ± 0.13 ± 0.13	140	<sup>1</sup> METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+\pi^-)$
1.58 ± 0.20 ± 0.15	84	BALTRUSAIT..85D	MRK3	$e^+e^-$
1.0 ± 0.5	5	BRANDELIK	78B DASP	$e^+e^-$
1.6 ± 1.6	1	VANNUCCI	77 MRK1	$e^+e^-$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{132}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.83 ± 0.23 OUR AVERAGE</b>					
2.74 ± 0.24 ± 0.22	234 ± 21	<sup>1</sup> ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda\bar{\Sigma}^0$	
2.92 ± 0.22 ± 0.24	308 ± 24	<sup>2</sup> ABLIKIM	12B BES3	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<15                      90                      PERUZZI                      78                      MRK1                       $e^+e^- \rightarrow \Lambda X$

<sup>1</sup> ABLIKIM 12B quotes  $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$  which we multiply by 2.

<sup>2</sup> ABLIKIM 12B quotes  $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$  which we multiply by 2.

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{133}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.01</b>	95	<sup>1</sup> BAI	04D BES	$e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.052                      90                      <sup>1</sup> BALTRUSAIT..85C                      MRK3                       $e^+e^-$

<sup>1</sup> Forbidden by *CP*.

————— RADIATIVE DECAYS —————

$\Gamma(3\gamma)/\Gamma_{\text{total}}$   $\Gamma_{134}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.6 ± 2.2 OUR AVERAGE</b>					
11.3 ± 1.8 ± 2.0	113 ± 18	ABLIKIM	13I BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$	
12 ± 3 ± 2	24.2 <sup>+7.2</sup> <sub>-6.0</sub>	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55                      90                      PARTRIDGE                      80                      CBAL                       $e^+e^-$

$\Gamma(4\gamma)/\Gamma_{\text{total}}$   $\Gamma_{135}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;9</b>	90	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(5\gamma)/\Gamma_{\text{total}}$					$\Gamma_{136}/\Gamma$
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;15</b>	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$					$\Gamma_{137}/\Gamma$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>1.7 \pm 0.4</math> OUR AVERAGE</b>				Error includes scale factor of 1.6.	
$2.01 \pm 0.32 \pm 0.02$		<sup>1</sup> MITCHELL	09	CLEO $e^+ e^- \rightarrow \gamma X$	
$1.27 \pm 0.36$		GAISER	86	CBAL $J/\psi \rightarrow \gamma X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.79 \pm 0.20$	$273 \pm 43$	<sup>2</sup> AUBERT	06E	BABR $B^\pm \rightarrow K^\pm X_{c\bar{c}}$	
seen	16	BALTRUSAITIS	84	MRK3 $J/\psi \rightarrow 2\phi\gamma$	

<sup>1</sup> MITCHELL 09 reports  $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$  assuming  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.45 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Calculated by the authors using an average of  $B(J/\psi \rightarrow \gamma\eta_c) \times B(\eta_c \rightarrow K\bar{K}\pi)$  from BALTRUSAITIS 86, BISELLO 91, BAI 04 and  $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$  from AUBERT 06E.

$\Gamma(\gamma\eta_c(1S) \rightarrow 3\gamma)/\Gamma_{\text{total}}$					$\Gamma_{138}/\Gamma$
VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>3.8^{+1.3}_{-1.0}</math> OUR AVERAGE</b>				Error includes scale factor of 1.1.	
$4.5 \pm 1.2 \pm 0.6$	$33 \pm 9$	ABLIKIM	13i	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	
$1.2^{+2.7}_{-1.1} \pm 0.3$	$1.2^{+2.8}_{-1.1}$	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	

$\Gamma(\gamma\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{139}/\Gamma$
VALUE (units $10^{-3}$ )		DOCUMENT ID	TECN	COMMENT	
<b><math>8.3 \pm 0.2 \pm 3.1</math></b>		<sup>1</sup> BALTRUSAITIS	86B	MRK3 $J/\psi \rightarrow 4\pi\gamma$	
<sup>1</sup> $4\pi$ mass less than 2.0 GeV.					


$\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$					$\Gamma_{140}/\Gamma$
VALUE (units $10^{-3}$ )		DOCUMENT ID	TECN	COMMENT	
<b><math>6.1 \pm 1.0</math> OUR AVERAGE</b>					
$5.85 \pm 0.3 \pm 1.05$		<sup>1</sup> EDWARDS	83B	CBAL $J/\psi \rightarrow \eta\pi^+\pi^-$	
$7.8 \pm 1.2 \pm 2.4$		<sup>1</sup> EDWARDS	83B	CBAL $J/\psi \rightarrow \eta 2\pi^0$	
<sup>1</sup> Broad enhancement at 1700 MeV.					

$\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{141}/\Gamma$
VALUE (units $10^{-4}$ )		DOCUMENT ID	TECN	COMMENT	
<b><math>6.2 \pm 2.2 \pm 0.9</math></b>		BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$   $\Gamma_{142} / \Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.8 ± 0.6 OUR AVERAGE</b>	Error includes scale factor of 1.6. See the ideogram below.		
1.66 ± 0.1 ± 0.58	<sup>1,2</sup> BAI	00D BES	$J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$
3.8 ± 0.3 ± 0.6	<sup>3</sup> AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
4.0 ± 0.7 ± 1.0	<sup>3</sup> EDWARDS	82E CBAL	$J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
4.3 ± 1.7	<sup>3,4</sup> SCHARRE	80 MRK2	$e^+ e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.78 ± 0.21 ± 0.33	<sup>3,5,6</sup> AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
0.83 ± 0.13 ± 0.18	<sup>3,7,8</sup> AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
0.66 <sup>+0.17+0.24</sup> <sub>-0.16-0.15</sub>	<sup>3,6,9</sup> BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$
1.03 <sup>+0.21+0.26</sup> <sub>-0.18-0.19</sub>	<sup>3,8,10</sup> BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$

- <sup>1</sup> Interference with the  $J/\psi(1S)$  radiative transition to the broad  $K \bar{K} \pi$  pseudoscalar state around 1800 is  $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$ .
- <sup>2</sup> Interference with  $J/\psi \rightarrow \gamma f_1(1420)$  is  $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$ .
- <sup>3</sup> Includes unknown branching fraction  $\eta(1405) \rightarrow K \bar{K} \pi$ .
- <sup>4</sup> Corrected for spin-zero hypothesis for  $\eta(1405)$ .
- <sup>5</sup> From fit to the  $a_0(980) \pi 0^{-+}$  partial wave.
- <sup>6</sup>  $a_0(980) \pi$  mode.
- <sup>7</sup> From fit to the  $K^*(892) K 0^{-+}$  partial wave.
- <sup>8</sup>  $K^* K$  mode.
- <sup>9</sup> From  $a_0(980) \pi$  final state.
- <sup>10</sup> From  $K^*(890) K$  final state.



$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0)/\Gamma_{\text{total}}$   $\Gamma_{143}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.78 ± 0.20 OUR AVERAGE</b>	Error includes scale factor of 1.8.		
1.07 ± 0.17 ± 0.11	<sup>1</sup> BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
0.64 ± 0.12 ± 0.07	<sup>1</sup> COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

<sup>1</sup>Includes unknown branching fraction  $\eta(1405) \rightarrow \gamma\rho^0$ .

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{144}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.0 ± 0.5 OUR AVERAGE</b>				
2.6 ± 0.7 ± 0.4		BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
3.38 ± 0.33 ± 0.64		<sup>1</sup> BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.0 ± 0.6 ± 1.1	261	<sup>2</sup> AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-----------------	-----	-----------------------	----	---

<sup>1</sup>Via  $a_0(980)\pi$ .

<sup>2</sup>Includes unknown branching fraction to  $\eta\pi^+\pi^-$ .

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$   $\Gamma_{145}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.82</b>	95	BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma K^+ K^-$

$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$   $\Gamma_{146}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>4.5 ± 0.8 OUR AVERAGE</b>				
4.7 ± 0.3 ± 0.9		<sup>1</sup> BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
3.75 ± 1.05 ± 1.20		<sup>2</sup> BURKE	82	MRK2 $J/\psi \rightarrow 4\pi\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.09	90	<sup>3</sup> BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
-------	----	----------------------	-----	---------------------------------

<sup>1</sup> $4\pi$  mass less than 2.0 GeV.

<sup>2</sup> $4\pi$  mass less than 2.0 GeV. We have multiplied  $2\rho^0$  measurement by 3 to obtain  $2\rho$ .

<sup>3</sup> $4\pi$  mass in the range 2.0–25 GeV.

$\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$   $\Gamma_{147}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5.4</b>	90	ABLIKIM	08A	BES2 $e^+e^- \rightarrow J/\psi$

$\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$   $\Gamma_{148}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;8.8</b>	90	ABLIKIM	08A	BES2 $e^+e^- \rightarrow J/\psi$

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{149}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.15 ± 0.16 OUR AVERAGE</b>				Error includes scale factor of 1.2.
4.82 ± 0.23 ± 0.08		<sup>1</sup> ABLIKIM	11	BES3 $J/\psi \rightarrow \eta'\gamma$
5.24 ± 0.12 ± 0.11		PEDLAR	09	CLE3 $J/\psi \rightarrow \eta'\gamma$
5.55 ± 0.44	35k	ABLIKIM	06E	BES2 $J/\psi \rightarrow \eta'\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

4.50 ± 0.14 ± 0.53		BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta, \eta \rightarrow \gamma \gamma$
4.30 ± 0.31 ± 0.71		BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta, \eta \rightarrow \pi^+ \pi^- \pi^0$
4.04 ± 0.16 ± 0.85	622	AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
4.39 ± 0.09 ± 0.66	2420	AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$
4.1 ± 0.3 ± 0.6		BLOOM	83	CBAL	$e^+ e^- \rightarrow 3\gamma + \text{hadrons}$
2.9 ± 1.1	6	BRANDELIK	79C	DASP	$e^+ e^- \rightarrow 3\gamma$
2.4 ± 0.7	57	BARTEL	76	CNTR	$e^+ e^- \rightarrow 2\gamma \rho$

<sup>1</sup> ABLIKIM 11 reports  $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+ \pi^- \eta)] / [B(\eta \rightarrow 2\gamma)]$  assuming  $B(\eta'(958) \rightarrow \pi^+ \pi^- \eta) = (43.2 \pm 0.7) \times 10^{-2}$ ,  $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$ , which we rescale to our best values  $B(\eta'(958) \rightarrow \pi^+ \pi^- \eta) = (42.9 \pm 0.7) \times 10^{-2}$ ,  $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.


$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

$\Gamma_{150}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.8 ± 0.5 OUR AVERAGE</b>	Error includes scale factor of 1.9. See the ideogram below.		
4.32 ± 0.14 ± 0.73	<sup>1</sup> BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
2.08 ± 0.13 ± 0.35	<sup>2</sup> BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
3.05 ± 0.08 ± 0.45	<sup>2</sup> BALTRUSAIT..	86B	MRK3 $J/\psi \rightarrow 4\pi\gamma$
4.85 ± 0.45 ± 1.20	<sup>3</sup> BURKE	82	MRK2 $e^+ e^-$

- <sup>1</sup>  $4\pi$  mass less than 3.0 GeV.
- <sup>2</sup>  $4\pi$  mass less than 2.0 GeV.
- <sup>3</sup>  $4\pi$  mass less than 2.5 GeV.

WEIGHTED AVERAGE  
2.8 ± 0.5 (Error scaled by 1.9)



$\Gamma(\gamma f_2(1270) f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{151}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.5±0.7±1.6</b>	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

$\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant}))/\Gamma_{\text{total}}$   $\Gamma_{152}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.2±0.8±1.7</b>	<sup>1</sup> ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

<sup>1</sup> Subtracting contribution from intermediate  $\eta_c(1S)$  decays.

$\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{153}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.1±0.1±0.6</b>	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

$\Gamma(\gamma f_4(2050))/\Gamma_{\text{total}}$   $\Gamma_{154}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.7±0.5±0.5</b>	<sup>1</sup> BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

<sup>1</sup> Assuming branching fraction  $f_4(2050) \rightarrow \pi\pi/\text{total} = 0.167$ .

$\Gamma(\gamma\omega\omega)/\Gamma_{\text{total}}$   $\Gamma_{155}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.61±0.33 OUR AVERAGE</b>				
6.0 ± 4.8 ± 1.8		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma\omega\pi^+\pi^-$
1.41±0.2 ± 0.42	120 ± 17	BISELLO	87 SPEC	$e^+e^-$ , hadrons $\gamma$
1.76±0.09±0.45		BALTRUSAIT..85C	MRK3	$e^+e^- \rightarrow \text{hadrons}\gamma$

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$   $\Gamma_{156}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.7 ± 0.4 OUR AVERAGE</b>	Error includes scale factor of 1.3.		
2.1 ± 0.4	BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1.36±0.38	<sup>1,2</sup> BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

<sup>1</sup> Estimated by us from various fits.

<sup>2</sup> Includes unknown branching fraction to  $\rho^0\rho^0$ .

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{157}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.43±0.11 OUR AVERAGE</b>				
1.62±0.26 $^{+0.02}_{-0.05}$		<sup>1</sup> ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1.42±0.21 $^{+0.02}_{-0.04}$		<sup>2</sup> ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
1.33±0.05±0.20		<sup>3</sup> AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1.36±0.09±0.23		<sup>3</sup> BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1.48±0.25±0.30	178	EDWARDS	82B CBAL	$e^+e^- \rightarrow 2\pi^0\gamma$
2.0 ± 0.7	35	ALEXANDER	78 PLUT	$e^+e^-$
1.2 ± 0.6	30	<sup>4</sup> BRANDELIK	78B DASP	$e^+e^- \rightarrow \pi^+\pi^-\gamma$

<sup>1</sup> ABLIKIM 06V reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$  which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.8^{+2.4}_{-1.2}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ABLIKIM 06V reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$  which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.8^{+2.4}_{-1.2}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Estimated using  $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$ . The errors do not contain the uncertainty in the  $f_2(1270)$  decay.

<sup>4</sup> Restated by us to take account of spread of E1, M2, E3 transitions.

### $\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ $\Gamma_{158}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

**8.5  $\pm$  1.2** OUR AVERAGE Error includes scale factor of 1.2.

9.62  $\pm$  0.29  $^{+3.51}_{-1.86}$

<sup>1</sup> BAI 03G BES  $J/\psi \rightarrow \gamma K \bar{K}$

5.0  $\pm$  0.8  $^{+1.8}_{-0.4}$

<sup>2,3</sup> BAI 96C BES  $J/\psi \rightarrow \gamma K^+ K^-$

9.2  $\pm$  1.4  $\pm$  1.4

<sup>3</sup> AUGUSTIN 88 DM2  $J/\psi \rightarrow \gamma K^+ K^-$

10.4  $\pm$  1.2  $\pm$  1.6

<sup>3</sup> AUGUSTIN 88 DM2  $J/\psi \rightarrow \gamma K_S^0 K_S^0$

9.6  $\pm$  1.2  $\pm$  1.8

<sup>3</sup> BALTRUSAIT..87 MRK3  $J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.6  $\pm$  0.2  $^{+0.6}_{-0.2}$

<sup>3,4</sup> BAI 96C BES  $J/\psi \rightarrow \gamma K^+ K^-$

< 0.8

90

<sup>5</sup> BISELLO 89B  $J/\psi \rightarrow 4\pi\gamma$

1.6  $\pm$  0.4  $\pm$  0.3

<sup>6</sup> BALTRUSAIT..87 MRK3  $J/\psi \rightarrow \gamma \pi^+ \pi^-$

3.8  $\pm$  1.6

<sup>7</sup> EDWARDS 82D CBAL  $e^+ e^- \rightarrow \eta\eta\gamma$

<sup>1</sup> Includes unknown branching ratio to  $K^+ K^-$  or  $K_S^0 K_S^0$ .

<sup>2</sup> Assuming  $J^P = 2^+$  for  $f_0(1710)$ .

<sup>3</sup> Includes unknown branching fraction to  $K^+ K^-$  or  $K_S^0 K_S^0$ . We have multiplied  $K^+ K^-$  measurement by 2, and  $K_S^0 K_S^0$  by 4 to obtain  $K \bar{K}$  result.

<sup>4</sup> Assuming  $J^P = 0^+$  for  $f_0(1710)$ .

<sup>5</sup> Includes unknown branching fraction to  $\rho^0 \rho^0$ .

<sup>6</sup> Includes unknown branching fraction to  $\pi^+ \pi^-$ .

<sup>7</sup> Includes unknown branching fraction to  $\eta\eta$ .

### $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ $\Gamma_{159}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**4.0  $\pm$  1.0** OUR AVERAGE

3.96  $\pm$  0.06  $\pm$  1.12

<sup>1</sup> ABLIKIM 06V BES2  $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$

3.99  $\pm$  0.15  $\pm$  2.64

<sup>1</sup> ABLIKIM 06V BES2  $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.5  $\pm$  1.6  $\pm$  0.8

BAI 98H BES  $J/\psi \rightarrow \gamma \pi^0 \pi^0$

<sup>1</sup> Including unknown branching fraction to  $\pi\pi$ .



$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$   $\Gamma_{160} / \Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.31 \pm 0.06 \pm 0.08</math></b>	180	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma \omega \omega$

$\Gamma(\gamma \eta) / \Gamma_{\text{total}}$   $\Gamma_{161} / \Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.104 \pm 0.034</math> OUR AVERAGE</b>				
$1.101 \pm 0.029 \pm 0.022$		PEDLAR	09	CLE3 $J/\psi \rightarrow \eta \gamma$
$1.123 \pm 0.089$	11k	ABLIKIM	06E	BES2 $J/\psi \rightarrow \eta \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.88 \pm 0.08 \pm 0.11$		BLOOM	83	CBAL $e^+ e^-$
$0.82 \pm 0.10$		BRANDELIK	79C	DASP $e^+ e^-$
$1.3 \pm 0.4$	21	BARTEL	77	CNTR $e^+ e^-$

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$   $\Gamma_{162} / \Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.79 \pm 0.13</math> OUR AVERAGE</b>			
$0.68 \pm 0.04 \pm 0.24$	BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$0.76 \pm 0.15 \pm 0.21$	<sup>1,2</sup> AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.87 \pm 0.14^{+0.14}_{-0.11}$	<sup>1</sup> BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

<sup>1</sup> Included unknown branching fraction  $f_1(1420) \rightarrow K \bar{K} \pi$ .

<sup>2</sup> From fit to the  $K^*(892) K 1^{++}$  partial wave.

$\Gamma(\gamma f_1(1285)) / \Gamma_{\text{total}}$   $\Gamma_{163} / \Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.61 \pm 0.08</math> OUR AVERAGE</b>			
$0.69 \pm 0.16 \pm 0.20$	<sup>1</sup> BAI	04J	BES2 $J/\psi \rightarrow \gamma \gamma \rho^0$
$0.61 \pm 0.04 \pm 0.21$	<sup>2</sup> BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$0.45 \pm 0.09 \pm 0.17$	<sup>3</sup> BAI	99	BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$0.625 \pm 0.063 \pm 0.103$	<sup>4</sup> BOLTON	92	MRK3 $J/\psi \rightarrow \gamma f_1(1285)$
$0.70 \pm 0.08 \pm 0.16$	<sup>5</sup> BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

<sup>1</sup> Assuming  $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$ .

<sup>2</sup> Assuming  $\Gamma(f_1(1285) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}} = 0.090 \pm 0.004$ .

<sup>3</sup> Assuming  $\Gamma(f_1(1285) \rightarrow \eta \pi \pi) / \Gamma_{\text{total}} = 0.5 \pm 0.18$ .

<sup>4</sup> Obtained summing the sequential decay channels

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi \pi \pi \pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow \eta \pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow K \bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma \rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}.$$

<sup>5</sup> Using  $B(f_1(1285) \rightarrow a_0(980) \pi) = 0.37$ , and including unknown branching ratio for  $a_0(980) \rightarrow \eta \pi$ .

$\Gamma(\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{164} / \Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>4.5 \pm 1.0 \pm 0.7</math></b>	BAI	99	BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$   $\Gamma_{165}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**4.5  $^{+0.7}_{-0.4}$  OUR AVERAGE**

$3.85 \pm 0.17$	$^{+1.91}_{-0.73}$		<sup>1</sup> BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
$3.6 \pm 0.4$	$^{+1.4}_{-0.4}$		<sup>1</sup> BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
$5.6 \pm 1.4$	$\pm 0.9$		<sup>1</sup> AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$4.5 \pm 0.4$	$\pm 0.9$		<sup>1</sup> AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.8 \pm 1.6$	$\pm 1.4$		<sup>1</sup> BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.4	90	4	<sup>2</sup> BRANDELIK	79C DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<2.3	90	3	ALEXANDER	78 PLUT	$e^+ e^- \rightarrow K^+ K^- \gamma$

<sup>1</sup> Using  $B(f'_2(1525) \rightarrow K \bar{K}) = 0.888$ .

<sup>2</sup> Assuming isotropic production and decay of the  $f'_2(1525)$  and isospin.

$\Gamma(\gamma f_2(1640) \rightarrow \gamma \omega \omega)/\Gamma_{\text{total}}$   $\Gamma_{166}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>0.28 <math>\pm 0.05 \pm 0.17</math></b>	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$
--	-----	---------	---------	---

$\Gamma(\gamma f_2(1910) \rightarrow \gamma \omega \omega)/\Gamma_{\text{total}}$   $\Gamma_{167}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>0.20 <math>\pm 0.04 \pm 0.13</math></b>	151	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$
--	-----	---------	---------	---

$\Gamma(\gamma f_0(1800) \rightarrow \gamma \omega \phi)/\Gamma_{\text{total}}$   $\Gamma_{168}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**2.5  $\pm 0.6$  OUR AVERAGE**

$2.00 \pm 0.08$	$^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma \omega \phi$
$2.61 \pm 0.27$	$\pm 0.65$	95	ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma \omega \phi$

$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$   $\Gamma_{169}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

<b>0.7 <math>\pm 0.1 \pm 0.2</math></b>	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$
---	-----	---------	---

$\Gamma(\gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$   $\Gamma_{170}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>4.0 <math>\pm 0.3 \pm 1.3</math></b>	320	<sup>1</sup> BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$
---	-----	------------------	---------	---

<sup>1</sup> Summed over all charges.


$\Gamma(\gamma \phi \phi)/\Gamma_{\text{total}}$   $\Gamma_{171}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**4.0  $\pm 1.2$  OUR AVERAGE** Error includes scale factor of 2.1. See the ideogram below.

$7.5 \pm 0.6$	$\pm 1.2$	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
$3.4 \pm 0.8$	$\pm 0.6$	$33 \pm 7$	<sup>1</sup> BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$3.1 \pm 0.7$	$\pm 0.4$		<sup>1</sup> BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

<sup>1</sup>  $\phi\phi$  mass less than 2.9 GeV,  $\eta_c$  excluded.



**$\Gamma(\gamma\rho\bar{\rho})/\Gamma_{\text{total}}$**   **$\Gamma_{172}/\Gamma$**

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.38±0.07±0.07</b>		49	EATON	84	MRK2 $e^+e^-$
•••					We do not use the following data for averages, fits, limits, etc. •••
<0.11	90		PERUZZI	78	MRK1 $e^+e^-$

**$\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$**   **$\Gamma_{173}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.33±0.05 OUR AVERAGE</b>				
0.44±0.04±0.08	196 ± 19	<sup>1</sup> ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
0.33±0.08±0.05		<sup>1</sup> BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
0.27±0.06±0.06		<sup>1</sup> BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
0.24 <sup>+0.15</sup> <sub>-0.10</sub>		<sup>2,3</sup> BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

- <sup>1</sup> Includes unknown branching fraction to  $\phi\phi$ .
- <sup>2</sup> Estimated by us from various fits.
- <sup>3</sup> Includes unknown branching fraction to  $\rho^0\rho^0$ .

**$\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$**   **$\Gamma_{174}/\Gamma$**

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.13±0.09</b>	<sup>1,2</sup> BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

- <sup>1</sup> Estimated by us from various fits.
- <sup>2</sup> Includes unknown branching fraction to  $\rho^0\rho^0$ .

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$   $\Gamma_{175}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.98 ± 0.08 ± 0.32</b>	1045	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}}$   $\Gamma_{176}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.6 ± 0.4 OUR AVERAGE</b>				
2.87 ± 0.09 <sup>+0.49</sup> <sub>-0.52</sub>	4265	<sup>1</sup> ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
2.2 ± 0.4 ± 0.4	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

<sup>1</sup> From a fit of the  $\pi^+\pi^-\eta'$  mass distribution to a combination of  $\gamma f_1(1510)$ ,  $\gamma X(1835)$ , and two unconfirmed states  $\gamma X(2120)$ , and  $\gamma X(2370)$ , for  $M(p\bar{p}) < 2.8$  GeV, and accounting for backgrounds from non- $\eta'$  events and  $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$ .

$\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{177}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.77<sup>+0.15</sup><sub>-0.09</sub> OUR AVERAGE</b>				
0.90 <sup>+0.04+0.27</sup> <sub>-0.11-0.55</sub>		<sup>1</sup> ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
1.14 <sup>+0.43+0.42</sup> <sub>-0.30-0.26</sub>	231	<sup>2</sup> ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
0.70 ± 0.04 <sup>+0.19</sup> <sub>-0.08</sub>		BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

<sup>1</sup> From the fit including final state interaction effects in isospin 0 S-wave according to SIBIRTSEV 05A.

<sup>2</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma X(1835)$ ,  $\gamma R$  with  $M(R) = 2100$  MeV and  $\Gamma(R) = 160$  MeV, and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV.

$\Gamma(\gamma X(1840) \rightarrow \gamma 3(\pi^+\pi^-))/\Gamma_{\text{total}}$   $\Gamma_{178}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.44 ± 0.36<sup>+0.60</sup><sub>-0.74</sub></b>	0.6k	ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$

$\Gamma(\gamma(K\bar{K}\pi)[J^{PC} = 0^{-+}])/ \Gamma_{\text{total}}$   $\Gamma_{179}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.7 ± 0.4 OUR AVERAGE</b>	Error includes scale factor of 2.1.		
0.58 ± 0.03 ± 0.20	<sup>1</sup> BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
2.1 ± 0.1 ± 0.7	<sup>2</sup> BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

<sup>1</sup> For a broad structure around 1800 MeV.

<sup>2</sup> For a broad structure around 2040 MeV.

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{180}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.49<sup>+0.33</sup><sub>-0.30</sub> OUR AVERAGE</b>				
3.63 ± 0.36 ± 0.13		PEDLAR	09 CLE3	$J/\psi \rightarrow \pi^0\gamma$
3.13 <sup>+0.65</sup> <sub>-0.47</sub>	586	ABLIKIM	06E BES2	$J/\psi \rightarrow \pi^0\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3.6 ± 1.1 ± 0.7		BLOOM	83 CBAL	$e^+e^-$
7.3 ± 4.7	10	BRANDELIK	79C DASP	$e^+e^-$

$\Gamma(\gamma\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{181}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.79</b>	90	EATON	84	MRK2 $e^+e^-$

$\Gamma(\gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$   $\Gamma_{182}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.13</b>	90	HENRARD	87	DM2 $e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.16	90	BAI	98G	BES $e^+e^-$
-------	----	-----	-----	--------------

$\Gamma(\gamma f_0(2200))/\Gamma_{\text{total}}$   $\Gamma_{183}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5	<sup>1</sup> AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
-----	-----------------------	----	---

<sup>1</sup> Includes unknown branching fraction to  $K_S^0 K_S^0$ .

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$   $\Gamma_{184}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<b>&gt;250</b>	99.9		<sup>1</sup> HASAN	96	SPEC $\bar{p}p \rightarrow \pi^+\pi^-$
----------------	------	--	--------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

>300			<sup>2</sup> BAI	96B	BES $e^+e^- \rightarrow \gamma\bar{p}p, K\bar{K}$
------	--	--	------------------	-----	---

< 2.3	95		<sup>3</sup> AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K^+ K^-$
-------	----	--	-----------------------	----	---

< 1.6	95		<sup>3</sup> AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
-------	----	--	-----------------------	----	---

$12.4^{+6.4}_{-5.2} \pm 2.8$		23	<sup>3</sup> BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
------------------------------	--	----	------------------------------	------	---

$8.4^{+3.4}_{-2.8} \pm 1.6$		93	<sup>3</sup> BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
-----------------------------	--	----	------------------------------	------	-------------------------------------

<sup>1</sup> Using BAI 96B.

<sup>2</sup> Using BARNES 93.

<sup>3</sup> Includes unknown branching fraction to  $K^+ K^-$  or  $K_S^0 K_S^0$ .

$\Gamma(\gamma f_J(2220) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_{185}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

<b><math>0.84 \pm 0.26 \pm 0.30</math></b>	BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
--	-----	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.4 \pm 0.8 \pm 0.4$	BAI	98H	BES $J/\psi \rightarrow \gamma\pi^0\pi^0$
-----------------------	-----	-----	---

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$   $\Gamma_{186}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

<b>&lt; 3.6</b>	<sup>1</sup> DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
-----------------	------------------------------	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.9	<sup>1</sup> DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
-------	------------------------------	------	--

$6.6 \pm 2.9 \pm 2.4$	BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
-----------------------	-----	-----	--

$10.8 \pm 4.0 \pm 3.2$	BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
------------------------	-----	-----	--

<sup>1</sup> For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$   $\Gamma_{187}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>1.5 \pm 0.6 \pm 0.5</math></b>	BAI	96B	BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma p \bar{p}$

$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$   $\Gamma_{188}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>1.01 \pm 0.32</math> OUR AVERAGE</b>			
$1.00 \pm 0.03 \pm 0.45$	<sup>1</sup> ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
$1.02 \pm 0.09 \pm 0.45$	<sup>1</sup> ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$>5.7 \pm 0.8$  <sup>2,3</sup> BUGG 95 MRK3  $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$

<sup>1</sup> Including unknown branching fraction to  $\pi\pi$ .

<sup>2</sup> Including unknown branching ratio for  $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ .

<sup>3</sup> Assuming that  $f_0(1500)$  decays only to two S-wave dipions.

$\Gamma(\gamma A \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$   $\Gamma_{189}/\Gamma$   
**(narrow state A with  $m_A < 960$  MeV)**

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;6.3</math></b>	90	<sup>1</sup> INSLER	10	CLEO $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

<sup>1</sup> The limit varies with mass  $m_A$  of a narrow state A and is  $4.3 \times 10^{-6}$  for  $m_A = 0$  MeV, reaches its largest value of  $6.3 \times 10^{-6}$  at  $m_A = 500$  MeV, and is  $3.6 \times 10^{-6}$  at  $m_A = 960$  MeV.

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{190}/\Gamma$   
**(narrow state  $A^0$  with  $0.2 \text{ GeV} < m_{A^0} < 3 \text{ GeV}$ )**

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;2.1</math></b>	90	<sup>1</sup> ABLIKIM	12	BES3 $J/\psi \rightarrow \gamma \mu^+ \mu^-$

<sup>1</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of  $m_{A^0}$  ranges from  $4 \times 10^{-7}$  to  $2.1 \times 10^{-5}$ .

———— WEAK DECAYS ————

$\Gamma(D^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{191}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;1.2</math></b>	90	ABLIKIM	06M	BES2 $e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{192}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;1.1</math></b>	90	ABLIKIM	06M	BES2 $e^+ e^- \rightarrow J/\psi$

$\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{193}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;3.6</math></b>	90	<sup>1</sup> ABLIKIM	06M	BES2 $e^+ e^- \rightarrow J/\psi$

<sup>1</sup> Using  $B(D_s^- \rightarrow \phi \pi^-) = 4.4 \pm 0.5 \%$ .

$\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{194}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.5 \times 10^{-5}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{195}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{196}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-4}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{197}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.5$	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<16$	90	<sup>1</sup> WICHT 08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
$< 2.2$	90	ABLIKIM 07J	BES2	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
$<50$	90	BARTEL 77	CNTR	$e^+ e^-$

<sup>1</sup> WICHT 08 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] < 0.16 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow J/\psi(1S)K^+) = 1.027 \times 10^{-3}$ .

———— LEPTON FAMILY NUMBER (LF) VIOLATING MODES ————

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$   $\Gamma_{198}/\Gamma$

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6$	90	ABLIKIM 13L	BES3	$e^+ e^- \rightarrow J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<11$	90	BAI 03D	BES	$e^+ e^- \rightarrow J/\psi$

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{199}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<8.3$	90	ABLIKIM 04	BES	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{200}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0$	90	ABLIKIM 04	BES	$e^+ e^- \rightarrow J/\psi$

———— OTHER DECAYS ————

$\Gamma(\text{invisible})/\Gamma(e^+ e^-)$   $\Gamma_{201}/\Gamma_5$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.6 \times 10^{-2}$	90	LEES 13l	BABR	$B \rightarrow K^{(*)} J/\psi$

$\Gamma(\text{invisible})/\Gamma(\mu^+ \mu^-)$   $\Gamma_{201}/\Gamma_7$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-2}$	90	ABLIKIM 08G	BES2	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

**$J/\psi(1S)$  REFERENCES**

AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13L	PR D87 112007	Ablikim M. <i>et al.</i>	(BES III Collab.)
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DEL-AMO-SA...	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)



AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 Collab., E706 Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
HSUEH	92	PR D45 R2181	S. Hsueh, S. Palestini	(FNAL, TORI)
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)

BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
BALTRUSAIT...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BALTRUSAIT...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
BALTRUSAIT...	84	PRL 52 2126	Translated from YAF 41 733.	R.M. Baltrusaitis <i>et al.</i>
EATON	84	PR D29 804	M.W. Eaton <i>et al.</i>	(CIT, UCSC+)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(LBL, SLAC)
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(SLAC, CIT)
FRANKLIN	83	PRL 51 963	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
BURKE	82	PRL 49 632	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
EDWARDS	82B	PR D25 3065	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
Also		ARNS 33 143	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS	82E	PRL 49 259	E.D. Bloom, C. Peck	(SLAC, CIT)
LEMOIGNE	82	PL 113B 509	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
BESCH	81	ZPHY C8 1	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
GIDAL	81	PL 107B 153	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)
PARTRIDGE	80	PRL 44 712	G. Gidal <i>et al.</i>	(SLAC, LBL)
SCHARRE	80	PL 97B 329	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)
ZHOLENTZ	80	PL 96B 214	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
BRANDELIK	79C	ZPHY C1 233	Translated from YAF 34 1471.	A.A. Zholents <i>et al.</i>
ALEXANDER	78	PL 72B 493	R. Brandelik <i>et al.</i>	(DASP Collab.)
BESCH	78	PL 78B 347	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)
BRANDELIK	78B	PL 74B 292	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)
PERUZZI	78	PR D17 2901	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	77	PL 66B 489	I. Peruzzi <i>et al.</i>	(SLAC, LBL)
BURMESTER	77D	PL 72B 135	W. Bartel <i>et al.</i>	(DESY, HEIDP)
FELDMAN	77	PRPL 33C 285	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
VANNUCCI	77	PR D15 1814	G.J. Feldman, M.L. Perl	(LBL, SLAC)
BARTEL	76	PL 64B 483	F. Vannucci <i>et al.</i>	(SLAC, LBL)
BRAUNSCH...	76	PL 63B 487	W. Bartel <i>et al.</i>	(DESY, HEIDP)
JEAN-MARIE	76	PRL 36 291	W. Braunschweig <i>et al.</i>	(DASP Collab.)
BALDINI-...	75	PL 58B 471	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG
BOYARSKI	75	PRL 34 1357	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)
DASP	75	PL 56B 491	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC
ESPOSITO	75B	LNC 14 73	W. Braunschweig <i>et al.</i>	(DASP Collab.)
FORD	75	PRL 34 604	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
			R.L. Ford <i>et al.</i>	(SLAC, PENN)