

K₁(1650)*I(J^P) = ½(1⁺)*

OMMITTED FROM SUMMARY TABLE

This entry contains various peaks in strange meson systems ($K^+\phi$, $K\pi\pi$) reported in partial-wave analysis in the 1600–1900 mass region.

K₁(1650) MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
1650±50	FRAME 86	OMEG +	13	$K^+ p \rightarrow \phi K^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~1840	ARMSTRONG 83	OMEG –	18.5	$K^- p \rightarrow 3K p$
~1800	DAUM 81C	CNTR –	63	$K^- p \rightarrow K^- 2\pi p$

K₁(1650) WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
150±50	FRAME 86	OMEG +	13	$K^+ p \rightarrow \phi K^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~250	DAUM 81C	CNTR –	63	$K^- p \rightarrow K^- 2\pi p$

K₁(1650) DECAY MODES

Mode
$\Gamma_1 \quad K\pi\pi$
$\Gamma_2 \quad K\phi$

K₁(1650) REFERENCES

FRAME 86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)
ARMSTRONG 83	NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
DAUM 81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)