

**X(10650) $^\pm$**  $I^G(J^P) = ?^+(1^+)$ 

## OMITTED FROM SUMMARY TABLE

Observed by BONDAR 12 in  $\gamma(5S)$  decays to  $\gamma(nS)\pi^+\pi^-$  ( $n = 1, 2, 3$ ) and  $h_b(mP)\pi^+\pi^-$  ( $m = 1, 2$ ).  $J^P = 1^+$  is favored from angular analyses.

**X(10650) $^\pm$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10652.2<math>\pm</math>1.5</b>	<sup>1</sup> BONDAR	12	BELL $e^+e^- \rightarrow$ hadrons
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
10657 $\pm 6$ $\pm 3$	<sup>2</sup> BONDAR	12	BELL $e^+e^- \rightarrow \gamma(1S)\pi^+\pi^-$
10651 $\pm 2$ $\pm 3$	<sup>2</sup> BONDAR	12	BELL $e^+e^- \rightarrow \gamma(2S)\pi^+\pi^-$
10652 $\pm 1$ $\pm 2$	<sup>2</sup> BONDAR	12	BELL $e^+e^- \rightarrow \gamma(3S)\pi^+\pi^-$
10654 $\pm 3$ $^{+1}_{-2}$	<sup>2</sup> BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
10651 $^{+2}_{-3}$ $^{+3}_{-2}$	<sup>2</sup> BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

<sup>1</sup> Average of the BONDAR 12 measurements in separate channels.

<sup>2</sup> Superseded by the average measurement of BONDAR 12.

**X(10650) $^\pm$  WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>11.5<math>\pm</math>2.2</b>	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow$ hadrons
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
$16.3 \pm 9.8$ $^{+6.0}_{-2.0}$	<sup>4</sup> BONDAR	12	BELL $e^+e^- \rightarrow \gamma(1S)\pi^+\pi^-$
$13.3 \pm 3.3$ $^{+4.0}_{-3.0}$	<sup>4</sup> BONDAR	12	BELL $e^+e^- \rightarrow \gamma(2S)\pi^+\pi^-$
$8.4 \pm 2.0$ $\pm 2.0$	<sup>4</sup> BONDAR	12	BELL $e^+e^- \rightarrow \gamma(3S)\pi^+\pi^-$
$20.9$ $^{+5.4}_{-4.7}$ $^{+2.1}_{-5.7}$	<sup>4</sup> BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
$19$ $\pm 7$ $^{+11}_{-7}$	<sup>4</sup> BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

<sup>3</sup> Average of the BONDAR 12 measurements in separate channels.

<sup>4</sup> Superseded by the average measurement of BONDAR 12.

**$X(10650)^+$  DECAY MODES** $X(10650)^-$  decay modes are charge conjugates of the modes below.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \gamma(1S)\pi^+$	seen
$\Gamma_2 \gamma(2S)\pi^+$	seen
$\Gamma_3 \gamma(3S)\pi^+$	seen
$\Gamma_4 h_b(1P)\pi^+$	seen
$\Gamma_5 h_b(2P)\pi^+$	seen

 **$X(10650)^{\pm}$  BRANCHING RATIOS**

$\Gamma(\gamma(1S)\pi^+)/\Gamma_{\text{total}}$

VALUE
seen

DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
BONDAR 12	BELL	$e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$	

$\Gamma(\gamma(2S)\pi^+)/\Gamma_{\text{total}}$

VALUE
seen

DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
BONDAR 12	BELL	$e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$	

$\Gamma(\gamma(3S)\pi^+)/\Gamma_{\text{total}}$

VALUE
seen

DOCUMENT ID	TECN	COMMENT	$\Gamma_3/\Gamma$
BONDAR 12	BELL	$e^+ e^- \rightarrow \gamma(3S)\pi^+\pi^-$	

$\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$

VALUE
seen

DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma$
BONDAR 12	BELL	$e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$	

$\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$

VALUE
seen

DOCUMENT ID	TECN	COMMENT	$\Gamma_5/\Gamma$
BONDAR 12	BELL	$e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$	

 **$X(10650)^{\pm}$  REFERENCES**

BONDAR 12 PRL 108 122001

A. Bondar *et al.*

(BELLE Collab.)