

**$\eta_c(2S)$**  $I^G(J^{PC}) = 0^+(0^{-+})$ 

Quantum numbers are quark model predictions.

 **$\eta_c(2S)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3639.4 \pm 1.3</math> OUR AVERAGE</b>				Error includes scale factor of 1.2.
$3646.9 \pm 1.6 \pm 3.6$	$57 \pm 17$	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
$3637.6 \pm 2.9 \pm 1.6$	$127 \pm 18$	<sup>1</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi, K\bar{K}\pi^0$
$3638.5 \pm 1.5 \pm 0.8$	624	<sup>2</sup> DEL-AMO-SA..11M BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	
$3640.5 \pm 3.2 \pm 2.5$	1201	<sup>2</sup> DEL-AMO-SA..11M BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
$3636.1^{+3.9}_{-4.2}{}^{+0.7}_{-2.0}$	128	<sup>3</sup> VINOKUROVA 11 BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$	
$3626 \pm 5 \pm 6$	311	<sup>4</sup> ABE	07 BELL	$e^+ e^- \rightarrow J/\psi(c\bar{c})$
$3645.0 \pm 5.5^{+4.9}_{-7.8}$	$121 \pm 27$	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
$3642.9 \pm 3.1 \pm 1.5$	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3639 $\pm 7$	$98 \pm 52$	<sup>5</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
$3630.8 \pm 3.4 \pm 1.0$	$112 \pm 24$	<sup>6</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
$3654 \pm 6 \pm 8$	$39 \pm 11$	<sup>7</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 $\pm 5$		<sup>8</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>1</sup> From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.<sup>2</sup> Ignoring possible interference with continuum.<sup>3</sup> Accounts for interference with non-resonant continuum.<sup>4</sup> From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE, K 02 and ABE 04G.<sup>5</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.<sup>6</sup> Superseded by DEL-AMO-SANCHEZ 11M.<sup>7</sup> Superseded by VINOKUROVA 11.<sup>8</sup> Assuming mass of  $\psi(2S) = 3686$  MeV. **$\eta_c(2S)$  WIDTH**

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>11.3^{+3.2}_{-2.9}</math> OUR AVERAGE</b>					
$9.9 \pm 4.8 \pm 2.9$	$57 \pm 17$	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	
$16.9 \pm 6.4 \pm 4.8$	$127 \pm 18$	<sup>9</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi, K\bar{K}\pi^0$	
$13.4 \pm 4.6 \pm 3.2$	624	<sup>10</sup> DEL-AMO-SA..11M BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$		
$6.6^{+8.4}_{-5.1}{}^{+2.6}_{-0.9}$	128	<sup>11</sup> VINOKUROVA 11 BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$		
$6.3 \pm 12.4 \pm 4.0$	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 23	90	$98 \pm 52$	<sup>12</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
22 ± 14		$121 \pm 27$	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
$17.0 \pm 8.3 \pm 2.5$		$112 \pm 24$	<sup>13</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
< 55	90	$39 \pm 11$	<sup>14</sup> CHOI	02 BELL	$B \rightarrow KK_SK^- \pi^+$
< 8.0	95		<sup>15</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>9</sup> From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.

<sup>10</sup> Ignoring possible interference with continuum.

<sup>11</sup> Accounts for interference with non-resonant continuum.

<sup>12</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>13</sup> Superseded by DEL-AMO-SANCHEZ 11M.

<sup>14</sup> For a mass value of  $3654 \pm 6$  MeV. Superseded by VINOKUROVA 11.

<sup>15</sup> For a mass value of  $3594 \pm 5$  MeV

## $\eta_c(2S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ hadrons	not seen	
$\Gamma_2$ $K\bar{K}\pi$	( $1.9 \pm 1.2$ ) %	
$\Gamma_3$ $2\pi^+ 2\pi^-$	not seen	
$\Gamma_4$ $\rho^0 \rho^0$	not seen	
$\Gamma_5$ $3\pi^+ 3\pi^-$	not seen	
$\Gamma_6$ $K^+ K^- \pi^+ \pi^-$	not seen	
$\Gamma_7$ $K^{*0} \bar{K}^{*0}$	not seen	
$\Gamma_8$ $K^+ K^- \pi^+ \pi^- \pi^0$	( $1.4 \pm 1.0$ ) %	
$\Gamma_9$ $K^+ K^- 2\pi^+ 2\pi^-$	not seen	
$\Gamma_{10}$ $K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$	seen	
$\Gamma_{11}$ $2K^+ 2K^-$	not seen	
$\Gamma_{12}$ $\phi\phi$	not seen	
$\Gamma_{13}$ $p\bar{p}$	$< 2.0 \times 10^{-3}$	90%
$\Gamma_{14}$ $\gamma\gamma$	( $1.9 \pm 1.3$ ) $\times 10^{-4}$	
$\Gamma_{15}$ $\pi^+ \pi^- \eta$	not seen	
$\Gamma_{16}$ $\pi^+ \pi^- \eta'$	not seen	
$\Gamma_{17}$ $K^+ K^- \eta$	not seen	
$\Gamma_{18}$ $\pi^+ \pi^- \eta_c(1S)$	$< 25$ %	90%

## $\eta_c(2S)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	$\Gamma_{14}$
<u>VALUE (keV)</u>	
<u>DOCUMENT ID</u>	
<u>TECN</u>	
<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
1.3 ± 0.6 <sup>16</sup> ASNER      04 CLEO $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$	

<sup>16</sup> They measure  $\Gamma(\eta_c(2S)\gamma\gamma)$   $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma)$   $B(\eta_c(1S) \rightarrow K\bar{K}\pi)$ . The value for  $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$  is derived assuming that the branching fractions for  $\eta_c(2S)$  and  $\eta_c(1S)$  decays to  $K_S K\pi$  are equal and using  $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$  keV.

### $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(2\pi^+ 2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		$\Gamma_3 \Gamma_{14}/\Gamma$		
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<6.5	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+ \pi^-)$

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		$\Gamma_2 \Gamma_{14}/\Gamma$		
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>41±4±6</b>	624	17	DEL-AMO-SA..11M	BABR $\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>17</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		$\Gamma_6 \Gamma_{14}/\Gamma$		
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		$\Gamma_8 \Gamma_{14}/\Gamma$		
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>30±6±5</b>	1201	18	DEL-AMO-SA..11M	BABR $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

<sup>18</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(2K^+ 2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		$\Gamma_{11} \Gamma_{14}/\Gamma$		
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+ K^-)$

$\Gamma(\pi^+ \pi^- \eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		$\Gamma_{18} \Gamma_{14}/\Gamma$		
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<133	90	LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

### $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		$\Gamma_{13}/\Gamma \times \Gamma_{14}/\Gamma$		
VALUE (units $10^{-8}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 5.6	90	19,20,21 AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
< 8.0	90	19,20,22 AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
<12.0	90	20,22 AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$

<sup>19</sup> Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

<sup>20</sup> For a total width  $\Gamma=5$  MeV.

<sup>21</sup> For the resonance mass region  $3589\text{--}3599$   $\text{MeV}/c^2$ .

<sup>22</sup> For the resonance mass region  $3575\text{--}3660$   $\text{MeV}/c^2$ .

**$\eta_c(2S)$  BRANCHING RATIOS** **$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$** 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
<b>not seen</b>	ABREU	980	DLPH $e^+ e^- \rightarrow e^+ e^- + \text{hadrons}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				

seen 23 EDWARDS 82C CBAL  $e^+ e^- \rightarrow \gamma X$ 23 For a mass value of  $3594 \pm 5$  MeV **$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
<b>1.9±0.4±1.1</b>	$59 \pm 12$	24 AUBERT	08AB	BABR $B \rightarrow \eta_c(2S) K \rightarrow K\bar{K}\pi K$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	$127 \pm 18$	ABLIKIM	13K	BES3 $\psi(2S) \rightarrow \gamma K\bar{K}\pi$	
seen	$39 \pm 11$	25 CHOI	02	BELL $B \rightarrow K K_S K^- \pi^+$	

24 Derived from a measurement of  $[B(B^+ \rightarrow \eta_c(2S) K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$  and using  $B(B^+ \rightarrow \eta_c(2S) K^+) = (3.4 \pm 1.8) \times 10^{-4}$ , and  $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$ .

25 For a mass value of  $3654 \pm 6$  MeV **$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$** 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_3/\Gamma$
<b>not seen</b>	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$	

 **$\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$** 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma$
<b>not seen</b>	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$	

 **$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$** 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_6/\Gamma$
<b>not seen</b>	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$	

 **$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_8/\Gamma_2$
<b>0.73±0.17±0.17</b>	1201	26 DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

26 We have multiplied the value of  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$  reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$ . Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

 **$\Gamma(K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$** 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_7/\Gamma$
<b>not seen</b>	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$	

 **$\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{10}/\Gamma$
<b>seen</b>	$57 \pm 17$	ABLIKIM	13K	BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	

### $\Gamma(2K^+ 2K^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{11}/\Gamma$
<b>not seen</b>	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$	

### $\Gamma(\phi\phi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{12}/\Gamma$
<b>not seen</b>	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$	

### $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{14}/\Gamma$
-------	-----	-------------	------	---------	----------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<5 \times 10^{-4}$  90 27 WICHT 08 BELL  $B^\pm \rightarrow K^\pm \gamma\gamma$

not seen AMBROGIANI 01 E835  $\bar{p}p \rightarrow \gamma\gamma$

$<0.01$  90 LEE 85 CBAL  $\psi' \rightarrow \text{photons}$

27 WICHT 08 reports  $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S) K^+)] < 0.18 \times 10^{-6}$   
which we divide by our best value  $B(B^+ \rightarrow \eta_c(2S) K^+) = 3.4 \times 10^{-4}$ .

### $\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma(K\bar{K}\pi)$

### $\Gamma_{18}/\Gamma_2$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{18}/\Gamma_2$
<b>&lt;3.33</b>	90	28 LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$	

28 We divided the reported limit by 3 to take into account isospin relations.

## $\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS

### $\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$

### $\Gamma_3/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_3/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}$
<b>&lt;14.6 \times 10^{-6}</b>	90	29 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$	

29 Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

### $\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$

### $\Gamma_4/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}$
<b>&lt;12.7 \times 10^{-7}</b>	90	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$	

### $\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$

### $\Gamma_5/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_5/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}$
<b>&lt;13.2 \times 10^{-6}</b>	90	30 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$	

30 Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_6/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.6 \times 10^{-6}$	90	31 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

<sup>31</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_7/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<19.6 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_8/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43.0 \times 10^{-6}$	90	32 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

<sup>32</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_9/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	33 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

<sup>33</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_{10}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.03 ± 2.10 ± 0.7</b>	60	ABLIKIM	13K BES3	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 15.2$	90	34 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$
----------	----	-------------------	------	--

<sup>34</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow \phi \phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_{12}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.8 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{15} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.3 \times 10^{-6}$	90	35 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta$

35 Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta') / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{16} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<14.2 \times 10^{-6}$	90	36 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta'$

36 Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{17} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.9 \times 10^{-6}$	90	37 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \eta$

37 Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{18} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	38 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta_c(1S)$

38 Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{13} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-6}$	90	ABLIKIM	13V	BES3 $\psi(2S) \rightarrow \gamma p\bar{p}$

## $\eta_c(2S)$ REFERENCES

ABLIKIM	13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BES III Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)

AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)