




$$I(J^P) = 0(0^-)$$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

B_s^0 MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|----------|---------------------------------|
| 5366.79 ± 0.23 | | | | OUR FIT |
| 5366.7 ± 0.4 | | | | OUR AVERAGE |
| Error includes scale factor of 1.3. See the ideogram below. | | | | |
| 5366.90 ± 0.28 ± 0.23 | | 1 AAIJ | 12E LHCb | $p\bar{p}$ at 7 TeV |
| 5364.4 ± 1.3 ± 0.7 | | LOUVOT | 09 BELL | $e^+e^- \rightarrow \gamma(5S)$ |
| 5366.01 ± 0.73 ± 0.33 | | 2 ACOSTA | 06 CDF | $p\bar{p}$ at 1.96 TeV |
| 5369.9 ± 2.3 ± 1.3 | 32 | 3 ABE | 96B CDF | $p\bar{p}$ at 1.8 TeV |
| 5374 ± 16 ± 2 | 3 | ABREU | 94D DLPH | $e^+e^- \rightarrow Z$ |
| 5359 ± 19 ± 7 | 1 | 3 AKERS | 94J OPAL | $e^+e^- \rightarrow Z$ |
| 5368.6 ± 5.6 ± 1.5 | 2 | BUSKULIC | 93G ALEP | $e^+e^- \rightarrow Z$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 5370 ± 1 ± 3 | | DRUTSKOY | 07A BELL | Repl. by LOUVOT 09 |
| 5370 ± 40 | 6 | 4 AKERS | 94J OPAL | $e^+e^- \rightarrow Z$ |
| 5383.3 ± 4.5 ± 5.0 | 14 | ABE | 93F CDF | Repl. by ABE 96B |

WEIGHTED AVERAGE
5366.7 ± 0.4 (Error scaled by 1.3)



¹ Uses $B_s^0 \rightarrow J/\psi\phi$ fully reconstructed decays.

² Uses exclusively reconstructed final states containing a $J/\psi \rightarrow \mu^+\mu^-$ decays.

³ From the decay $B_s \rightarrow J/\psi(1S)\phi$.

⁴ From the decay $B_s \rightarrow D_s^- \pi^+$.

$m_{B_s^0} - m_B$

m_B is the average of our B masses $(m_{B^\pm} + m_{B^0})/2$.

| VALUE (MeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------|-----|-------------|------|---------|
|-------------|-----|-------------|------|---------|

87.33±0.23 OUR FIT

87.34±0.29 OUR AVERAGE

87.42±0.30±0.09

¹ AAIJ 12E LHCB pp at 7 TeV

86.64±0.80±0.08

² ACOSTA 06 CDF $p\bar{p}$ at 1.96 TeV

• • • We use the following data for averages but not for fits. • • •

89.7 ±2.7 ±1.2

ABE 96B CDF $p\bar{p}$ at 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

80 to 130

68

LEE-FRANZINI 90 CSB2 $e^+e^- \rightarrow \Upsilon(5S)$

¹ The reported result is $m_{B_s^0} - m_{B^+} = 87.52 \pm 0.30 \pm 0.12$ MeV. We convert it to the mass difference with respect to the average of $(m_{B^\pm} + m_{B^0})/2$.

² The reported result is $m_{B_s^0} - m_{B^0} = 86.38 \pm 0.90 \pm 0.06$ MeV. We convert it to the mass difference with respect to the average of $(m_{B^\pm} + m_{B^0})/2$.

$m_{B_s^0} - m_{B_s^0}$

See the $B_s^0-\bar{B}_s^0$ MIXING section near the end of these B_s^0 Listings.

B_s^0 MEAN LIFE

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements and asymmetric lifetime errors.

“OUR EVALUATION” is an average of $1 / [0.5 (\Gamma_{B_{sL}^0} + \Gamma_{B_{sH}^0})]$.

| VALUE (10^{-12} s) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|------|---------|
|-----------------------|------|-------------|------|---------|

1.510±0.005 OUR EVALUATION

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--|-----|-------------------------|----------|-------------------------|
| $1.518 \pm 0.041 \pm 0.027$ | | ¹ AALTONEN | 11AP CDF | $p\bar{p}$ at 1.96 TeV |
| $1.398 \pm 0.044^{+0.028}_{-0.025}$ | | ² ABAZOV | 06V D0 | $p\bar{p}$ at 1.96 TeV |
| $1.42^{+0.14}_{-0.13} \pm 0.03$ | | ³ ABREU | 00Y DLPH | $e^+e^- \rightarrow Z$ |
| $1.53^{+0.16}_{-0.15} \pm 0.07$ | | ⁴ ABREU,P | 00G DLPH | $e^+e^- \rightarrow Z$ |
| $1.36 \pm 0.09^{+0.06}_{-0.05}$ | | ⁵ ABE | 99D CDF | $p\bar{p}$ at 1.8 TeV |
| $1.72^{+0.20}_{-0.19}^{+0.18}_{-0.17}$ | | ⁶ ACKERSTAFF | 98F OPAL | $e^+e^- \rightarrow Z$ |
| $1.50^{+0.16}_{-0.15} \pm 0.04$ | | ⁵ ACKERSTAFF | 98G OPAL | $e^+e^- \rightarrow Z$ |
| $1.47 \pm 0.14 \pm 0.08$ | | ⁴ BARATE | 98C ALEP | $e^+e^- \rightarrow Z$ |
| 1.51 ± 0.11 | | ⁷ BARATE | 98C ALEP | $e^+e^- \rightarrow Z$ |
| $1.56^{+0.29}_{-0.26}^{+0.08}_{-0.07}$ | | ⁵ ABREU | 96F DLPH | Repl. by ABREU 00Y |
| $1.65^{+0.34}_{-0.31} \pm 0.12$ | | ⁴ ABREU | 96F DLPH | Repl. by ABREU 00Y |
| $1.76 \pm 0.20^{+0.15}_{-0.10}$ | | ⁸ ABREU | 96F DLPH | Repl. by ABREU 00Y |
| $1.60 \pm 0.26^{+0.13}_{-0.15}$ | | ⁹ ABREU | 96F DLPH | Repl. by ABREU,P 00G |
| 1.67 ± 0.14 | | ¹⁰ ABREU | 96F DLPH | $e^+e^- \rightarrow Z$ |
| $1.61^{+0.30}_{-0.29}^{+0.18}_{-0.16}$ | 90 | ⁴ BUSKULIC | 96E ALEP | Repl. by BARATE 98C |
| $1.54^{+0.14}_{-0.13} \pm 0.04$ | | ⁵ BUSKULIC | 96M ALEP | $e^+e^- \rightarrow Z$ |
| $1.42^{+0.27}_{-0.23} \pm 0.11$ | 76 | ⁵ ABE | 95R CDF | Repl. by ABE 99D |
| $1.74^{+1.08}_{-0.69} \pm 0.07$ | 8 | ¹¹ ABE | 95R CDF | Sup. by ABE 96N |
| $1.54^{+0.25}_{-0.21} \pm 0.06$ | 79 | ⁵ AKERS | 95G OPAL | Repl. by ACKERSTAFF 98G |
| $1.59^{+0.17}_{-0.15} \pm 0.03$ | 134 | ⁵ BUSKULIC | 95O ALEP | Sup. by BUSKULIC 96M |
| 0.96 ± 0.37 | 41 | ¹² ABREU | 94E DLPH | Sup. by ABREU 96F |
| $1.92^{+0.45}_{-0.35} \pm 0.04$ | 31 | ⁵ BUSKULIC | 94C ALEP | Sup. by BUSKULIC 95O |
| $1.13^{+0.35}_{-0.26} \pm 0.09$ | 22 | ⁵ ACTON | 93H OPAL | Sup. by AKERS 95G |

¹ AALTONEN 11AP combines the fully reconstructed $B_s^0 \rightarrow D_s^- \pi^+$ decays and partially reconstructed $B_s^0 \rightarrow D_s X$ decays.

² Measured using $D_s \mu^+$ vertices.

³ Uses $D_s^- \ell^+$, and $\phi \ell^+$ vertices.

⁴ Measured using D_s hadron vertices.

⁵ Measured using $D_s^- \ell^+$ vertices.

⁶ ACKERSTAFF 98F use fully reconstructed $D_s^- \rightarrow \phi \pi^-$ and $D_s^- \rightarrow K^{*0} K^-$ in the inclusive B_s^0 decay.

⁷ Combined results from $D_s^- \ell^+$ and D_s hadron.

⁸ Measured using $\phi \ell$ vertices.

⁹ Measured using inclusive D_s vertices.


¹⁰ Combined result for the four ABREU 96F methods.

¹¹ Exclusive reconstruction of $B_s \rightarrow \psi \phi$.

¹² ABREU 94E uses the flight-distance distribution of D_s vertices, ϕ -lepton vertices, and $D_s \mu$ vertices.

B_s^0 MEAN LIFE (Flavor specific)

| VALUE (10^{-12} s) | DOCUMENT ID | TECN | COMMENT |
|---|--------------|-----------|---|
| 1.511 ± 0.014 OUR EVALUATION | | | |
| 1.508 ± 0.019 OUR AVERAGE | | | Error includes scale factor of 1.3. See the ideogram below. |
| 1.479 ± 0.010 ± 0.021 | 1 ABAZOV | 15A D0 | $\rho \bar{p}$ at 1.96 TeV |
| 1.535 ± 0.015 ± 0.014 | 2 AAIJ | 14AX LHCb | pp at 7 TeV |
| 1.52 ± 0.15 ± 0.01 | 3 AAIJ | 14F LHCb | pp at 7, 8 TeV |
| 1.518 ± 0.041 ± 0.027 | 4 AALTONEN | 11AP CDF | $\rho \bar{p}$ at 1.96 TeV |
| 1.42 $^{+0.14}_{-0.13}$ ± 0.03 | 5 ABREU | 00Y DLPH | $e^+ e^- \rightarrow Z$ |
| 1.36 ± 0.09 $^{+0.06}_{-0.05}$ | 6 ABE | 99D CDF | $\rho \bar{p}$ at 1.8 TeV |
| 1.50 $^{+0.16}_{-0.15}$ ± 0.04 | 6 ACKERSTAFF | 98G OPAL | $e^+ e^- \rightarrow Z$ |
| 1.54 $^{+0.14}_{-0.13}$ ± 0.04 | 6 BUSKULIC | 96M ALEP | $e^+ e^- \rightarrow Z$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1.60 ± 0.06 ± 0.01 | 7 AAIJ | 14R LHCb | pp at 7 TeV |
| 1.398 ± 0.044 $^{+0.028}_{-0.025}$ | 8 ABAZOV | 06V D0 | Repl. by ABAZOV 15A |



- ¹ Measured using $B_S^0 \rightarrow D_S^- \mu^+ \nu X$ decays.
- ² Measured using the $B_S^0 \rightarrow D_S^- \pi^+$ decays.
- ³ Measured using $B_S^0 \rightarrow D^+ D_S^-$.
- ⁴ AALTONEN 11AP combines the fully reconstructed $B_S^0 \rightarrow D_S^- \pi^+$ decays and partially reconstructed $B_S^0 \rightarrow D_S X$ decays.
- ⁵ Uses $D_S^- \ell^+$, and $\phi \ell^+$ vertices.
- ⁶ Measured using $D_S^- \ell^+$ vertices.
- ⁷ Measured using $B_S^0 \rightarrow \pi^+ K^-$ decays. May not be flavor specific.
- ⁸ Measured using $D_S^- \mu^+$ vertices.

B_S^0 MEAN LIFE ($B_S \rightarrow J/\psi\phi$)

| VALUE (10^{-12} s) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|----------|------------------------|
| 1.479 ± 0.012 OUR EVALUATION | | | |
| 1.479 ± 0.012 OUR AVERAGE | | | |
| 1.480 ± 0.011 ± 0.005 | ¹ AAIJ | 14E LHCB | pp at 7 TeV |
| 1.444 ^{+0.098} _{-0.090} ± 0.020 | ¹ ABAZOV | 05B D0 | $p\bar{p}$ at 1.96 TeV |
| 1.34 ^{+0.23} _{-0.19} ± 0.05 | ² ABE | 98B CDF | $p\bar{p}$ at 1.8 TeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 1.529 ± 0.025 ± 0.012 | ² AALTONEN | 12D CDF | $p\bar{p}$ at 1.96 TeV |
| 1.39 ^{+0.13} _{-0.16} ^{+0.01} _{-0.02} | ² ABAZOV | 05W D0 | $p\bar{p}$ at 1.96 TeV |
| 1.40 ^{+0.15} _{-0.13} ± 0.02 | ² ACOSTA | 05 CDF | $p\bar{p}$ at 1.96 TeV |
| 1.34 ^{+0.23} _{-0.19} ± 0.05 | ³ ABE | 96N CDF | Repl. by ABE 98B |

- ¹ Measured using fully reconstructed $B_S \rightarrow J/\psi\phi$ decays.
- ² Measured using the time-dependent angular analysis of $B_S^0 \rightarrow J/\psi\phi$ decays.
- ³ ABE 96N uses 58 ± 12 exclusive $B_S \rightarrow J/\psi\phi$ events.

$\tau_{B_S^0}/\tau_{B^0}$ MEAN LIFE RATIO

$\tau_{B_S^0}$ is an average of $1 / [0.5 (\Gamma_{B_{sL}^0} + \Gamma_{B_{sH}^0})]$.

$\tau_{B_S^0}/\tau_{B^0}$ (direct measurements)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|-----------|------------------------|
| 0.993 ± 0.004 OUR EVALUATION | | | |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 0.964 ± 0.013 ± 0.007 | ¹ ABAZOV | 15A D0 | $p\bar{p}$ at 1.96 TeV |
| 1.010 ± 0.010 ± 0.008 | ² AAIJ | 14AX LHCB | pp at 7 TeV |
| 0.971 ± 0.009 ± 0.004 | ³ AAIJ | 14E LHCB | pp at 7 TeV |
| 1.052 ± 0.061 ± 0.015 | ⁴ ABAZOV | 09E D0 | $p\bar{p}$ at 1.96 TeV |
| 0.980 ^{+0.076} _{-0.071} ± 0.003 | ⁵ ABAZOV | 05B D0 | Repl. by ABAZOV 05W |
| 0.91 ± 0.09 ± 0.003 | ⁶ ABAZOV | 05W D0 | Repl. by ABAZOV 09E |

- ¹ Measured using $B_s^0 \rightarrow D_s^- \mu^+ \nu X$ and $B^0 \rightarrow D^- \mu^+ \nu X$ decays.
- ² Measured using the $B_s^0 \rightarrow D_s^- \pi^+$ decays.
- ³ Measured using $B_s^0 \rightarrow J/\psi \phi$ and $B^0 \rightarrow J/\psi K^{*0}$ decays.
- ⁴ Measured the angular and lifetime parameters for the time-dependent angular untagged decays $B_d^0 \rightarrow J/\psi K^{*0}$ and $B_s^0 \rightarrow J/\psi \phi$.
- ⁵ Measured mean life ratio using fully reconstructed decays.
- ⁶ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.

B_{sH}^0 MEAN LIFE

B_{sH}^0 is the heavy mass state of two B_s^0 CP eigenstates.

“OUR EVALUATION” has been obtained by the Heavy Flavor Averaging Group (HFAG) using the constraint of the flavor-specific lifetime average in a way similar to $\Delta\Gamma_{B_s^0}/\Gamma_{B_s^0}$.

| VALUE (10^{-12} s) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------|-----------|------------------------|
| 1.610 ± 0.012 OUR EVALUATION | | | |
| 1.70 ± 0.04 OUR AVERAGE | | | |
| 1.75 ± 0.12 ± 0.07 | ¹ AAIJ | 13AB LHCb | pp at 7 TeV |
| 1.700 ± 0.040 ± 0.026 | ² AAIJ | 12AN LHCb | pp at 7 TeV |
| 1.70 $^{+0.12}_{-0.11}$ ± 0.03 | ² AALTONEN | 11AB CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| | ³ AALTONEN | 12D CDF | $p\bar{p}$ at 1.96 TeV |
| 1.613 $^{+0.123}_{-0.113}$ | ^{4,5} AALTONEN | 08J CDF | Repl. by AALTONEN 12D |
| 1.58 $^{+0.39}_{-0.42}$ $^{+0.01}_{-0.02}$ | ⁵ ABAZOV | 05W D0 | Repl. by ABAZOV 08AM |
| 2.07 $^{+0.58}_{-0.46}$ ± 0.03 | ⁵ ACOSTA | 05 CDF | Repl. by AALTONEN 08J |

- ¹ Measured using a pure CP -odd final state $J/\psi K_S^0$ with the assumption that contributions from penguin diagrams are small.
- ² Measured using a pure CP -odd final state $J/\psi f_0(980)$.
- ³ Uses the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays assuming CP -violating angle $\beta_s(B^0 \rightarrow J/\psi \phi) = 0.02$.
- ⁴ Obtained from $\Delta\Gamma_s$ and Γ_s fit with a correlation of 0.6.
- ⁵ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.

B_{sL}^0 MEAN LIFE

B_{sL}^0 is the light mass state of two B_s^0 CP eigenstates.

“OUR EVALUATION” has been obtained by the Heavy Flavor Averaging Group (HFAG) using the constraint of the flavor-specific lifetime average in a way similar to $\Delta\Gamma_{B_s^0}/\Gamma_{B_s^0}$.

| VALUE (10^{-12} s) | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------|-------------------|----------|------------------|
| 1.422 ± 0.008 OUR EVALUATION | | | |
| 1.379 ± 0.026 ± 0.017 | ¹ AAIJ | 14F LHCb | pp at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--|-------------------------|-----|------|---------------------------|
| 1.407 ± 0.016 ± 0.007 | ² AAIJ | 14R | LHCB | $p\bar{p}$ at 7 TeV |
| 1.440 ± 0.096 ± 0.009 | ² AAIJ | 12 | LHCB | Repl. by AAIJ 14R |
| 1.455 ± 0.046 ± 0.006 | ² AAIJ | 12R | LHCB | Repl. by AAIJ 14R |
| | ³ AALTONEN | 12D | CDF | $\rho\bar{p}$ at 1.96 TeV |
| 1.437 ^{+0.054} _{-0.047} | ^{4,5} AALTONEN | 08J | CDF | Repl. by AALTONEN 12D |
| 1.24 ^{+0.14} _{-0.11} ± 0.01 -0.02 | ⁵ ABAZOV | 05W | D0 | Repl. by ABAZOV 08AM |
| 1.05 ^{+0.16} _{-0.13} ± 0.02 | ⁵ ACOSTA | 05 | CDF | Repl. by AALTONEN 08J |
| 1.27 ± 0.33 ± 0.08 | ⁶ BARATE | 00K | ALEP | $e^+e^- \rightarrow Z$ |

¹ Measured using $B_s^0 \rightarrow D_s^- D_s^+$. The effective lifetime is translated into a decay width of $\Gamma_L = 0.725 \pm 0.014 \pm 0.009 \text{ ps}^{-1}$.

² Measured using $B_s^0 \rightarrow K^+ K^-$ decays. There may still be CPV in the decay.

³ Uses the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays and assuming CP-violating angle $\beta_s(B^0 \rightarrow J/\psi\phi) = 0.02$.

⁴ Obtained from $\Delta\Gamma_s$ and Γ_s fit with a correlation of 0.6.

⁵ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

⁶ Uses $\phi\phi$ correlations from $B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}$.

$\Delta\Gamma_{B_s^0}/\Gamma_{B_s^0}$

$\Gamma_{B_s^0}$ and $\Delta\Gamma_{B_s^0}$ are the decay rate average and difference between two B_s^0 CP eigenstates (light – heavy).

“OUR EVALUATION” is an average of all available B_s flavor-specific lifetime measurements with the $\Delta\Gamma_{B_s^0}/\Gamma_s$ analyses performed by the Heavy

Flavor Averaging Group (HFAG) as described in our “Review on $B-\bar{B}$ Mixing” in the B^0 Section of these Listings.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|------------------------|-------------|--|
| 0.124 ± 0.011 OUR EVALUATION | | | | |
| | | ¹ AAIJ | 12D | LHCB $p\bar{p}$ at 7 TeV |
| | | ² ABAZOV | 12D | D0 $\rho\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.090 ± 0.009 ± 0.023 | | ³ ESEN | 13 | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |
| | | ⁴ AALTONEN | 12D | CDF $\rho\bar{p}$ at 1.96 TeV |
| 0.147 ^{+0.036} _{-0.030} ± 0.042 -0.041 | | ³ ESEN | 10 | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |
| 0.116 ^{+0.09} _{-0.10} ± 0.010 | | ⁵ AALTONEN | 08J | CDF Repl. by AALTONEN 12D |
| 0.24 ^{+0.28} _{-0.38} ± 0.03 -0.04 | | ^{5,6} ABAZOV | 05W | D0 Repl. by ABAZOV 08AM |
| 0.65 ^{+0.25} _{-0.33} ± 0.01 | | ⁵ ACOSTA | 05 | CDF Repl. by AALTONEN 08J |
| <0.46 | 95 | ⁷ ABREU | 00Y | DLPH $e^+e^- \rightarrow Z$ |
| <0.69 | 95 | ⁸ ABREU,P | 00G | DLPH $e^+e^- \rightarrow Z$ |
| <0.83 | 95 | ⁹ ABE | 99D | CDF $\rho\bar{p}$ at 1.8 TeV |
| <0.67 | 95 | ¹⁰ ACCIARRI | 98S | L3 $e^+e^- \rightarrow Z$ |

- ¹ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.
- ² Measured using fully reconstructed $B_s \rightarrow J/\psi \phi$ decays.
- ³ Assumes CP violation is negligible.
- ⁴ Uses the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays and assuming CP -violating angle $\beta_s(B^0 \rightarrow J/\psi \phi) = 0.02$.
- ⁵ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.
- ⁶ Uses $|A_0|^2 - |A_{||}|^2 = 0.355 \pm 0.066$ from ACOSTA 05.
- ⁷ Uses $D_s^- \ell^+$, and $\phi \ell^+$ vertices.
- ⁸ Measured using D_s hadron vertices.
- ⁹ ABE 99D assumes $\tau_{B_s^0} = 1.55 \pm 0.05$ ps.
- ¹⁰ ACCIARRI 98S assumes $\tau_{B_s^0} = 1.49 \pm 0.06$ ps and PDG 98 values of b production fraction.

$\Delta\Gamma_{B_s^0}$

"OUR EVALUATION" has been obtained by the Heavy Flavor Averaging Group (HFAG) using the constraint of the flavor-specific lifetime average in a way similar to $\Delta\Gamma_{B_s^0}/\Gamma_{B_s^0}$.

| <u>VALUE (10^{12} s^{-1})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------------------|-------------|------------------------|
| 0.082 ± 0.007 | OUR EVALUATION | | |
| 0.077 ± 0.008 | OUR AVERAGE | | |
| 0.0805 ± 0.0091 ± 0.0032 | ¹ AAIJ | 15I LHCb | $p\bar{p}$ at 7, 8 TeV |
| 0.053 ± 0.021 ± 0.010 | ² AAD | 14U ATLS | $p\bar{p}$ at 7 TeV |
| 0.068 ± 0.026 ± 0.007 | ³ AALTONEN | 12AJ CDF | $p\bar{p}$ at 1.96 TeV |
| 0.163 ^{+0.065} / _{-0.064} | ^{4,5} ABAZOV | 12D D0 | $p\bar{p}$ at 1.96 TeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 0.106 ± 0.011 ± 0.007 | ⁶ AAIJ | 13AR LHCb | Repl. by AAIJ 15I |
| 0.053 ± 0.021 ± 0.010 | ³ AAD | 12CV ATLS | Repl. by AAD 14U |
| 0.123 ± 0.029 ± 0.011 | ³ AAIJ | 12D LHCb | Repl. by AAIJ 13AR |
| 0.075 ± 0.035 ± 0.006 | ⁷ AALTONEN | 12D CDF | Repl. by AALTONEN 12AJ |
| 0.085 ^{+0.072} / _{-0.078} ± 0.001 | ⁸ ABAZOV | 09E D0 | Repl. by ABAZOV 08AM |
| 0.076 ^{+0.059} / _{-0.063} ± 0.006 | ⁹ AALTONEN | 08J CDF | Repl. by AALTONEN 12D |
| 0.19 ± 0.07 ^{+0.02} / _{-0.01} | ^{5,10} ABAZOV | 08AMD0 | Repl. by ABAZOV 12D |
| 0.12 ^{+0.08} / _{-0.10} ± 0.02 | ^{9,11} ABAZOV | 07 D0 | Repl. by ABAZOV 07N |
| 0.13 ± 0.09 | ¹² ABAZOV | 07N D0 | Repl. by ABAZOV 09E |
| 0.47 ^{+0.19} / _{-0.24} ± 0.01 | ⁹ ACOSTA | 05 CDF | Repl. by AALTONEN 08J |

- ¹ Measured using time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.
- ² Measured using the flavor tagged time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.
- ³ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.
- ⁴ The error includes both statistical and systematic uncertainties.
- ⁵ Measured using fully reconstructed $B_s \rightarrow J/\psi \phi$ decays.

⁶ AAIJ 13AR result comes from a combined fit to $B_s^0 \rightarrow J/\psi K^+ K^-$ and $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ data sets. Also reports $\Delta\Gamma_s = 0.100 \pm 0.016 \pm 0.003 \text{ ps}^{-1}$ from a fit to $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

⁷ Uses the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays and assuming CP -violating angle $\beta_s(B^0 \rightarrow J/\psi \phi) = 0.02$.

⁸ Measured the angular and lifetime parameters for the time-dependent angular untagged decays $B_d^0 \rightarrow J/\psi K^{*0}$ and $B_s^0 \rightarrow J/\psi \phi$.

⁹ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays and assuming CP -violating phase $\phi_s = 0$.

¹⁰ Obtains 90% CL interval $-0.06 < \Delta\Gamma_s < 0.30$.

¹¹ ABAZOV 07 reports $0.17 \pm 0.09 \pm 0.02$ with CP -violating phase ϕ_s as a free parameter.

¹² Combines D^0 measurements of time-dependent angular distributions in $B_s^0 \rightarrow J/\psi \phi$ and charge asymmetry in semileptonic decays. There is a 4-fold ambiguity in the solution.

$\Delta\Gamma_s^{CP} / \Gamma_s$

Γ_s and $\Delta\Gamma_s^{CP}$ are the decay rate average and difference between even, $\Gamma_s^{CP\text{-even}}$, and odd, $\Gamma_s^{CP\text{-odd}}$, CP eigenstates.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-----------------------|----------|---------------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| $0.072 \pm 0.021 \pm 0.022$ | | ¹ ABAZOV | 09I D0 | $\rho\bar{p}$ at 1.96 TeV |
| >0.012 | 95 | ¹ AALTONEN | 08F CDF | $\rho\bar{p}$ at 1.96 TeV |
| $0.079^{+0.038+0.031}_{-0.035-0.030}$ | | ¹ ABAZOV | 07Y D0 | Repl. by ABAZOV 09I |
| $0.25^{+0.21}_{-0.14}$ | | ² BARATE | 00K ALEP | $e^+e^- \rightarrow Z$ |

¹ Assumes $2 \text{B}(B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}) \simeq \Delta\Gamma_s^{CP} / \Gamma_s$.

² Uses $\phi\phi$ correlations from $B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}$.

$1 / \Gamma_{B_s^0}$

“OUR EVALUATION” has been obtained by the Heavy Flavor Averaging Group (HFAG) using the constraint of the flavor-specific lifetime average in a way similar to $\Delta\Gamma_{B_s^0} / \Gamma_{B_s^0}$.

| VALUE (10^{-12} s) | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------|--|----------|---------------------------|
| 1.510 ± 0.005 | OUR EVALUATION | | |
| 1.509 ± 0.010 | OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below. | | |
| $1.5145 \pm 0.0062 \pm 0.0034$ | ¹ AAIJ | 15I LHCB | pp at 7, 8 TeV |
| $1.477 \pm 0.015 \pm 0.009$ | ² AAD | 14U ATLS | pp at 7 TeV |
| $1.528 \pm 0.019 \pm 0.009$ | ³ AALTONEN | 12AJ CDF | $\rho\bar{p}$ at 1.96 TeV |
| $1.443^{+0.038}_{-0.035}$ | ^{3,4} ABAZOV | 12D D0 | $\rho\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----------------------|-----------------------|-----------|------------------------|
| 1.513 ± 0.009 ± 0.014 | ⁵ AAIJ | 13AR LHCb | Repl. by AAIJ 15I |
| 1.477 ± 0.015 ± 0.009 | ⁶ AAD | 12CV ATLS | Repl. by AAD 14U |
| 1.522 ± 0.021 ± 0.019 | ⁷ AAIJ | 12D LHCb | Repl. by AAIJ 13AR |
| 1.529 ± 0.025 ± 0.012 | ³ AALTONEN | 12D CDF | Repl. by AALTONEN 12AJ |
| 1.487 ± 0.060 ± 0.028 | ³ ABAZOV | 09E D0 | Repl. by ABAZOV 08AM |
| 1.52 ± 0.04 ± 0.02 | ³ AALTONEN | 08J CDF | Repl. by AALTONEN 12D |
| 1.52 ± 0.05 ± 0.01 | ³ ABAZOV | 08AMD0 | Repl. by ABAZOV 12D |

¹ AAIJ 15I reports $\Gamma_{B_s^0} = 0.6603 \pm 0.0027 \pm 0.0015 \text{ ps}^{-1}$ obtained from time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

² AAD 14U reports $\Gamma_{B_s^0} = 0.677 \pm 0.007 \pm 0.004 \text{ ps}^{-1}$ measured using a tagged, time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.


³ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.

⁴ The error includes both statistical and systematic uncertainties.

⁵ AAIJ 13AR reports $\Gamma_s = 0.661 \pm 0.004 \pm 0.006 \text{ ps}^{-1}$ obtained from combined fit to $B_s^0 \rightarrow J/\psi K^+ K^-$ and $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ data sets. Also reports a separate measurement of $\Gamma_s = 0.663 \pm 0.005 \pm 0.006 \text{ ps}^{-1}$ from $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

⁶ AAD 12CV reports $\Gamma_{B_s^0} = 0.677 \pm 0.007 \pm 0.004 \text{ ps}^{-1}$ measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.

⁷ AAIJ 12D reports average decay width of B_s^0 , $\Gamma_{B_s^0} = 0.657 \pm 0.009 \pm 0.008 \text{ ps}^{-1}$ that we converted to $1/\Gamma_{B_s^0}$.



B_s^0 DECAY MODES

These branching fractions all scale with $B(\bar{b} \rightarrow B_s^0)$.

The branching fraction $B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything})$ is not a pure measurement since the measured product branching fraction $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything})$ was used to determine $B(\bar{b} \rightarrow B_s^0)$, as described in the note on “ B^0 - \bar{B}^0 Mixing”

For inclusive branching fractions, *e.g.*, $B \rightarrow D^\pm \text{ anything}$, the values usually are multiplicities, not branching fractions. They can be greater than one.

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|--|------------------------------------|-----------------------------------|
| Γ_1 D_s^- anything | (93 ± 25) % | |
| Γ_2 $\ell \nu_\ell X$ | (9.6 ± 0.8) % | |
| Γ_3 $e^+ \nu X^-$ | (9.1 ± 0.8) % | |
| Γ_4 $\mu^+ \nu X^-$ | (10.2 ± 1.0) % | |
| Γ_5 $D_s^- \ell^+ \nu_\ell$ anything | [a] (7.9 ± 2.4) % | |
| Γ_6 $D_{s1}(2536)^- \mu^+ \nu_\mu$, $D_{s1}^- \rightarrow D^{*-} K_S^0$ | (2.5 ± 0.7) × 10 ⁻³ | |
| Γ_7 $D_{s1}(2536)^- X \mu^+ \nu$, $D_{s1}^- \rightarrow \bar{D}^0 K^+$ | (4.3 ± 1.7) × 10 ⁻³ | |
| Γ_8 $D_{s2}(2573)^- X \mu^+ \nu$, $D_{s2}^- \rightarrow \bar{D}^0 K^+$ | (2.6 ± 1.2) × 10 ⁻³ | |
| Γ_9 $D_s^- \pi^+$ | (3.04 ± 0.23) × 10 ⁻³ | |
| Γ_{10} $D_s^- \rho^+$ | (7.0 ± 1.5) × 10 ⁻³ | |
| Γ_{11} $D_s^- \pi^+ \pi^+ \pi^-$ | (6.3 ± 1.1) × 10 ⁻³ | |
| Γ_{12} $D_{s1}(2536)^- \pi^+$, $D_{s1}^- \rightarrow D_s^- \pi^+ \pi^-$ | (2.5 ± 0.8) × 10 ⁻⁵ | |
| Γ_{13} $D_s^\mp K^\pm$ | (2.03 ± 0.28) × 10 ⁻⁴ | S=1.3 |
| Γ_{14} $D_s^- K^+ \pi^+ \pi^-$ | (3.3 ± 0.7) × 10 ⁻⁴ | |
| Γ_{15} $D_s^+ D_s^-$ | (4.4 ± 0.5) × 10 ⁻³ | |
| Γ_{16} $D_s^- D^+$ | (2.8 ± 0.5) × 10 ⁻⁴ | |
| Γ_{17} $D^+ D^-$ | (2.2 ± 0.6) × 10 ⁻⁴ | |
| Γ_{18} $D^0 \bar{D}^0$ | (1.9 ± 0.5) × 10 ⁻⁴ | |
| Γ_{19} $D_s^{*-} \pi^+$ | (2.0 ± 0.5) × 10 ⁻³ | |
| Γ_{20} $D_s^{*-} \rho^+$ | (9.7 ± 2.2) × 10 ⁻³ | |
| Γ_{21} $D_s^{*+} D_s^- + D_s^{*-} D_s^+$ | (1.29 ± 0.22) % | S=1.1 |
| Γ_{22} $D_s^{*+} D_s^{*-}$ | (1.86 ± 0.30) % | |
| Γ_{23} $D_s^{(*)+} D_s^{(*)-}$ | (4.5 ± 1.4) % | |
| Γ_{24} $\bar{D}^0 K^- \pi^+$ | (9.9 ± 1.5) × 10 ⁻⁴ | |

| | | | |
|---------------|---|----------------------------------|--------|
| Γ_{25} | $\overline{D}^0 \overline{K}^*(892)^0$ | $(4.4 \pm 0.6) \times 10^{-4}$ | |
| Γ_{26} | $\overline{D}^0 \overline{K}^*(1410)$ | $(3.9 \pm 3.5) \times 10^{-4}$ | |
| Γ_{27} | $\overline{D}^0 \overline{K}_0^*(1430)$ | $(3.0 \pm 0.7) \times 10^{-4}$ | |
| Γ_{28} | $\overline{D}^0 \overline{K}_2^*(1430)$ | $(1.1 \pm 0.4) \times 10^{-4}$ | |
| Γ_{29} | $\overline{D}^0 \overline{K}^*(1680)$ | $< 7.8 \times 10^{-5}$ | CL=90% |
| Γ_{30} | $\overline{D}^0 \overline{K}_0^*(1950)$ | $< 1.1 \times 10^{-4}$ | CL=90% |
| Γ_{31} | $\overline{D}^0 \overline{K}_3^*(1780)$ | $< 2.6 \times 10^{-5}$ | CL=90% |
| Γ_{32} | $\overline{D}^0 \overline{K}_4^*(2045)$ | $< 3.1 \times 10^{-5}$ | CL=90% |
| Γ_{33} | $\overline{D}^0 K^- \pi^+$ (non-resonant) | $(2.1 \pm 0.8) \times 10^{-4}$ | |
| Γ_{34} | $D_{s2}^*(2573)^- \pi^+$, $D_{s2}^* \rightarrow \overline{D}^0 K^-$ | $(2.6 \pm 0.4) \times 10^{-4}$ | |
| Γ_{35} | $D_{s1}^*(2700)^- \pi^+$, $D_{s1}^* \rightarrow \overline{D}^0 K^-$ | $(1.6 \pm 0.8) \times 10^{-5}$ | |
| Γ_{36} | $D_{s1}^*(2860)^- \pi^+$, $D_{s1}^* \rightarrow \overline{D}^0 K^-$ | $(5 \pm 4) \times 10^{-5}$ | |
| Γ_{37} | $D_{s3}^*(2860)^- \pi^+$, $D_{s3}^* \rightarrow \overline{D}^0 K^-$ | $(2.2 \pm 0.6) \times 10^{-5}$ | |
| Γ_{38} | $\overline{D}^0 K^+ K^-$ | $(4.2 \pm 1.9) \times 10^{-5}$ | |
| Γ_{39} | $\overline{D}^0 \phi$ | $(3.0 \pm 0.8) \times 10^{-5}$ | |
| Γ_{40} | $D^{*\mp} \pi^\pm$ | $< 6.1 \times 10^{-6}$ | CL=90% |
| Γ_{41} | $J/\psi(1S) \phi$ | $(1.08 \pm 0.09) \times 10^{-3}$ | |
| Γ_{42} | $J/\psi(1S) \pi^0$ | $< 1.2 \times 10^{-3}$ | CL=90% |
| Γ_{43} | $J/\psi(1S) \eta$ | $(3.9 \pm 0.7) \times 10^{-4}$ | S=1.4 |
| Γ_{44} | $J/\psi(1S) K_S^0$ | $(1.87 \pm 0.17) \times 10^{-5}$ | |
| Γ_{45} | $J/\psi(1S) K^*(892)^0$ | $(4.4 \pm 0.9) \times 10^{-5}$ | |
| Γ_{46} | $J/\psi(1S) \eta'$ | $(3.3 \pm 0.4) \times 10^{-4}$ | |
| Γ_{47} | $J/\psi(1S) \pi^+ \pi^-$ | $(2.14 \pm 0.19) \times 10^{-4}$ | |
| Γ_{48} | $J/\psi(1S) f_0(500)$, $f_0 \rightarrow \pi^+ \pi^-$ | $< 1.7 \times 10^{-6}$ | CL=90% |
| Γ_{49} | $J/\psi(1S) \rho$, $\rho \rightarrow \pi^+ \pi^-$ | $< 1.2 \times 10^{-6}$ | CL=90% |
| Γ_{50} | $J/\psi(1S) f_0(980)$, $f_0 \rightarrow \pi^+ \pi^-$ | $(1.35 \pm 0.16) \times 10^{-4}$ | |
| Γ_{51} | $J/\psi(1S) f_0(980)_0$, $f_0 \rightarrow \pi^+ \pi^-$ | $(5.1 \pm 0.9) \times 10^{-5}$ | |
| Γ_{52} | $J/\psi(1S) f_2(1270)$, $f_2 \rightarrow \pi^+ \pi^-$ | | |
| Γ_{53} | $J/\psi(1S) f_2(1270)_0$, $f_2 \rightarrow \pi^+ \pi^-$ | $(2.6 \pm 0.7) \times 10^{-7}$ | |
| Γ_{54} | $J/\psi(1S) f_2(1270)_\parallel$, $f_2 \rightarrow \pi^+ \pi^-$ | $(3.8 \pm 1.3) \times 10^{-7}$ | |

| | | | |
|-----------------|--|---|--------|
| Γ ₅₅ | $J/\psi(1S) f_2(1270)_{\perp},$ $f_2 \rightarrow \pi^+ \pi^-$ | $(4.6 \pm 2.8) \times 10^{-7}$ | |
| Γ ₅₆ | $J/\psi(1S) f_0(1370),$ $f_0 \rightarrow \pi^+ \pi^-$ | | |
| Γ ₅₇ | $J/\psi(1S) f_0(1500),$ $f_0 \rightarrow \pi^+ \pi^-$ | $(7.4 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.6 \\ 1.4 \end{smallmatrix}) \times 10^{-6}$ | |
| Γ ₅₈ | $J/\psi(1S) f'_2(1525)_0,$ $f'_2 \rightarrow \pi^+ \pi^-$ | $(3.7 \pm 1.0) \times 10^{-7}$ | |
| Γ ₅₉ | $J/\psi(1S) f'_2(1525)_{\parallel},$ $f'_2 \rightarrow \pi^+ \pi^-$ | $(4.4 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 10.0 \\ 3.1 \end{smallmatrix}) \times 10^{-8}$ | |
| Γ ₆₀ | $J/\psi(1S) f'_2(1525)_{\perp},$ $f'_2 \rightarrow \pi^+ \pi^-$ | $(1.9 \pm 1.4) \times 10^{-7}$ | |
| Γ ₆₁ | $J/\psi(1S) f_0(1790),$ $f_0 \rightarrow \pi^+ \pi^-$ | $(1.7 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 4.0 \\ 0.4 \end{smallmatrix}) \times 10^{-6}$ | |
| Γ ₆₂ | $J/\psi(1S) \pi^+ \pi^-$ (nonres- onant) | | |
| Γ ₆₃ | $J/\psi(1S) \bar{K}^0 \pi^+ \pi^-$ | $< 4.4 \times 10^{-5}$ | CL=90% |
| Γ ₆₄ | $J/\psi(1S) K^+ K^-$ | $(7.9 \pm 0.7) \times 10^{-4}$ | |
| Γ ₆₅ | $J/\psi(1S) K^0 K^- \pi^+ + \text{c.c.}$ | $(9.3 \pm 1.3) \times 10^{-4}$ | |
| Γ ₆₆ | $J/\psi(1S) \bar{K}^0 K^+ K^-$ | $< 1.2 \times 10^{-5}$ | CL=90% |
| Γ ₆₇ | $J/\psi(1S) f'_2(1525)$ | $(2.6 \pm 0.6) \times 10^{-4}$ | |
| Γ ₆₈ | $J/\psi(1S) p \bar{p}$ | $< 4.8 \times 10^{-6}$ | CL=90% |
| Γ ₆₉ | $J/\psi(1S) \pi^+ \pi^- \pi^+ \pi^-$ | $(8.0 \pm 0.9) \times 10^{-5}$ | |
| Γ ₇₀ | $J/\psi(1S) f_1(1285)$ | $(7.1 \pm 1.4) \times 10^{-5}$ | |
| Γ ₇₁ | $\psi(2S) \eta$ | $(3.3 \pm 0.9) \times 10^{-4}$ | |
| Γ ₇₂ | $\psi(2S) \eta'$ | $(1.29 \pm 0.35) \times 10^{-4}$ | |
| Γ ₇₃ | $\psi(2S) \pi^+ \pi^-$ | $(7.3 \pm 1.3) \times 10^{-5}$ | |
| Γ ₇₄ | $\psi(2S) \phi$ | $(5.4 \pm 0.6) \times 10^{-4}$ | |
| Γ ₇₅ | $\chi_{c1} \phi$ | $(2.05 \pm 0.31) \times 10^{-4}$ | |
| Γ ₇₆ | $\pi^+ \pi^-$ | $(7.6 \pm 1.9) \times 10^{-7}$ | S=1.4 |
| Γ ₇₇ | $\pi^0 \pi^0$ | $< 2.1 \times 10^{-4}$ | CL=90% |
| Γ ₇₈ | $\eta \pi^0$ | $< 1.0 \times 10^{-3}$ | CL=90% |
| Γ ₇₉ | $\eta \eta$ | $< 1.5 \times 10^{-3}$ | CL=90% |
| Γ ₈₀ | $\rho^0 \rho^0$ | $< 3.20 \times 10^{-4}$ | CL=90% |
| Γ ₈₁ | $\phi \rho^0$ | $< 6.17 \times 10^{-4}$ | CL=90% |
| Γ ₈₂ | $\phi \phi$ | $(1.93 \pm 0.31) \times 10^{-5}$ | |
| Γ ₈₃ | $\pi^+ K^-$ | $(5.5 \pm 0.6) \times 10^{-6}$ | |
| Γ ₈₄ | $K^+ K^-$ | $(2.50 \pm 0.17) \times 10^{-5}$ | |
| Γ ₈₅ | $K^0 \bar{K}^0$ | $< 6.6 \times 10^{-5}$ | CL=90% |
| Γ ₈₆ | $K^0 \pi^+ \pi^-$ | $(1.5 \pm 0.4) \times 10^{-5}$ | |
| Γ ₈₇ | $K^0 K^{\pm} \pi^{\mp}$ | $(7.7 \pm 1.0) \times 10^{-5}$ | |
| Γ ₈₈ | $K^*(892)^- \pi^+$ | $(3.3 \pm 1.2) \times 10^{-6}$ | |
| Γ ₈₉ | $K^*(892)^{\pm} K^{\mp}$ | $(1.25 \pm 0.26) \times 10^{-5}$ | |

| | | | | | |
|---------------|-------------------------------|-----------|---------------------------|------------------|--------|
| Γ_{90} | $K^0 K^+ K^-$ | | < 3.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{91} | $\bar{K}^*(892)^0 \rho^0$ | | < 7.67 | $\times 10^{-4}$ | CL=90% |
| Γ_{92} | $\bar{K}^*(892)^0 K^*(892)^0$ | | (2.8 ± 0.7) | $\times 10^{-5}$ | |
| Γ_{93} | $\phi K^*(892)^0$ | | (1.13 ± 0.30) | $\times 10^{-6}$ | |
| Γ_{94} | $\rho \bar{\rho}$ | | $(2.8 \pm_{-1.7}^{+2.2})$ | $\times 10^{-8}$ | |
| Γ_{95} | $\Lambda_c^- \Lambda \pi^+$ | | (3.6 ± 1.6) | $\times 10^{-4}$ | |
| Γ_{96} | $\Lambda_c^- \Lambda_c^+$ | | < 8.0 | $\times 10^{-5}$ | CL=95% |
| Γ_{97} | $\gamma \gamma$ | <i>B1</i> | < 3.1 | $\times 10^{-6}$ | CL=90% |
| Γ_{98} | $\phi \gamma$ | | (3.52 ± 0.34) | $\times 10^{-5}$ | |

**Lepton Family number (LF) violating modes or
 $\Delta B = 1$ weak neutral current (B1) modes**

| | | | | | |
|----------------|---|-----------|----------------------|------------------|--------|
| Γ_{99} | $\mu^+ \mu^-$ | <i>B1</i> | (3.1 ± 0.7) | $\times 10^{-9}$ | |
| Γ_{100} | $e^+ e^-$ | <i>B1</i> | < 2.8 | $\times 10^{-7}$ | CL=90% |
| Γ_{101} | $\mu^+ \mu^- \mu^+ \mu^-$ | <i>B1</i> | < 1.2 | $\times 10^{-8}$ | CL=90% |
| Γ_{102} | $SP, S \rightarrow \mu^+ \mu^-,$ $P \rightarrow \mu^+ \mu^-$ | <i>B1</i> | [<i>b</i>] < 1.2 | $\times 10^{-8}$ | CL=90% |
| Γ_{103} | $\phi(1020) \mu^+ \mu^-$ | <i>B1</i> | (7.7 ± 1.5) | $\times 10^{-7}$ | |
| Γ_{104} | $\phi \nu \bar{\nu}$ | <i>B1</i> | < 5.4 | $\times 10^{-3}$ | CL=90% |
| Γ_{105} | $e^\pm \mu^\mp$ | <i>LF</i> | [<i>c</i>] < 1.1 | $\times 10^{-8}$ | CL=90% |

- [a] Not a pure measurement. See note at head of B_s^0 Decay Modes.
 [b] Here *S* and *P* are the hypothetical scalar and pseudoscalar particles with masses of 2.5 GeV/ c^2 and 214.3 MeV/ c^2 , respectively.
 [c] The value is for the sum of the charge states or particle/antiparticle states indicated.

CONSTRAINED FIT INFORMATION

An overall fit to 8 branching ratios uses 14 measurements and one constraint to determine 6 parameters. The overall fit has a $\chi^2 = 4.6$ for 9 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | | | |
|----------|--|-------|----------|----------|----------|
| x_{11} | | 28 | | | |
| x_{13} | | 55 | 16 | | |
| x_{41} | | 0 | 0 | 0 | |
| x_{50} | | 0 | 0 | 0 | 66 |
| | | x_9 | x_{11} | x_{13} | x_{41} |

B_s^0 BRANCHING RATIOS

$\Gamma(D_s^- \text{ anything})/\Gamma_{\text{total}}$ Γ_1/Γ

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------|-------------|---------------------------|-------------|---------------------------------|
| 0.93±0.25 OUR AVERAGE | | | | |
| 0.91±0.18±0.41 | | ¹ DRUTSKOY 07 | BELL | $e^+e^- \rightarrow \gamma(4S)$ |
| 0.81±0.24±0.22 | 90 | ² BUSKULIC 96E | ALEP | $e^+e^- \rightarrow Z$ |
| 1.56±0.58±0.44 | 147 | ³ ACTON 92N | OPAL | $e^+e^- \rightarrow Z$ |

¹ The extraction of this result takes into account the correlation between the measurements of $B(\Upsilon(5S) \rightarrow D_s X)$ and $B(\Upsilon(5S) \rightarrow D^0 X)$.

² BUSKULIC 96E separate $c\bar{c}$ and $b\bar{b}$ sources of D_s^+ mesons using a lifetime tag, subtract generic $\bar{b} \rightarrow W^+ \rightarrow D_s^+$ events, and obtain $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \text{ anything}) = 0.088 \pm 0.020 \pm 0.020$ assuming $B(D_s \rightarrow \phi\pi) = (3.5 \pm 0.4) \times 10^{-2}$ and PDG 1994 values for the relative partial widths to other D_s channels. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.

³ ACTON 92N assume that excess of $147 \pm 48 D_s^0$ events over that expected from B^0 , B^+ , and $c\bar{c}$ is all from B_s^0 decay. The product branching fraction is measured to be $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \text{ anything}) \times B(D_s^- \rightarrow \phi\pi^-) = (5.9 \pm 1.9 \pm 1.1) \times 10^{-3}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.

$\Gamma(\ell\nu_e X)/\Gamma_{\text{total}}$ Γ_2/Γ

| <u>VALUE (units 10^{-2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------------------|-------------|---------------------------------|
| 9.6±0.8 OUR AVERAGE | | | |
| 9.6±0.4±0.7 | ¹ OSWALD 13 | BELL | $e^+e^- \rightarrow \gamma(5S)$ |
| 9.5 ^{+2.5+1.1} _{-2.0-1.9} | ² LEES 12A | BABR | e^+e^- |

¹ The measurement corresponds to the average of the electron and muon branching fractions.

² The measurement corresponds to a branching fraction where the lepton originates from bottom decay and is the average between the electron and muon branching fractions. LEES 12A uses the correlation of the production of ϕ mesons in association with a lepton in e^+e^- data taken at center-of-mass energies between 10.54 and 11.2 GeV.

$\Gamma(e^+ \nu X^-)/\Gamma_{\text{total}}$ Γ_3/Γ

| <u>VALUE (units 10^{-2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|---------------------------------|
| 9.1±0.5±0.6 | OSWALD 13 | BELL | $e^+e^- \rightarrow \gamma(5S)$ |

$\Gamma(\mu^+ \nu X^-)/\Gamma_{\text{total}}$ Γ_4/Γ

| <u>VALUE (units 10^{-2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|---------------------------------|
| 10.2±0.6±0.8 | OSWALD 13 | BELL | $e^+e^- \rightarrow \gamma(5S)$ |

$\Gamma(D_s^- \ell^+ \nu_\ell \text{ anything})/\Gamma_{\text{total}}$

Γ_5/Γ

The values and averages in this section serve only to show what values result if one assumes our $B(\bar{b} \rightarrow B_s^0)$. They cannot be thought of as measurements since the underlying product branching fractions were also used to determine $B(\bar{b} \rightarrow B_s^0)$ as described in the note on "Production and Decay of b -Flavored Hadrons."

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-----------------------|-------------|-------------------------|
| 0.079 ± 0.024 OUR AVERAGE | | | | |
| 0.076 ± 0.012 ± 0.021 | 134 | ¹ BUSKULIC | 95O ALEP | $e^+ e^- \rightarrow Z$ |
| 0.107 ± 0.043 ± 0.029 | | ² ABREU | 92M DLPH | $e^+ e^- \rightarrow Z$ |
| 0.103 ± 0.036 ± 0.028 | 18 | ³ ACTON | 92N OPAL | $e^+ e^- \rightarrow Z$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.13 ± 0.04 ± 0.04 | 27 | ⁴ BUSKULIC | 92E ALEP | $e^+ e^- \rightarrow Z$ |

¹ BUSKULIC 95O use $D_s \ell$ correlations. The measured product branching ratio is $B(\bar{b} \rightarrow B_s) \times B(B_s \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything}) = (0.82 \pm 0.09^{+0.13}_{-0.14})\%$ assuming $B(D_s \rightarrow \phi\pi) = (3.5 \pm 0.4) \times 10^{-2}$ and PDG 1994 values for the relative partial widths to the six other D_s channels used in this analysis. Combined with results from $\Upsilon(4S)$ experiments this can be used to extract $B(\bar{b} \rightarrow B_s) = (11.0 \pm 1.2^{+2.5}_{-2.6})\%$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.

² ABREU 92M measured muons only and obtained product branching ratio $B(Z \rightarrow b \text{ or } \bar{b}) \times B(\bar{b} \rightarrow B_s) \times B(B_s \rightarrow D_s \mu^+ \nu_\mu \text{ anything}) \times B(D_s \rightarrow \phi\pi) = (18 \pm 8) \times 10^{-5}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$. We use $B(Z \rightarrow b \text{ or } \bar{b}) = 2B(Z \rightarrow b\bar{b}) = 2 \times (0.2212 \pm 0.0019)$.

³ ACTON 92N is measured using $D_s \rightarrow \phi\pi^+$ and $K^*(892)^0 K^+$ events. The product branching fraction measured is measured to be $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything}) \times B(D_s^- \rightarrow \phi\pi^-) = (3.9 \pm 1.1 \pm 0.8) \times 10^{-4}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.

⁴ BUSKULIC 92E is measured using $D_s \rightarrow \phi\pi^+$ and $K^*(892)^0 K^+$ events. They use $2.7 \pm 0.7\%$ for the $\phi\pi^+$ branching fraction. The average product branching fraction is measured to be $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything}) = 0.020 \pm 0.0055^{+0.005}_{-0.006}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$. Superseded by BUSKULIC 95O.

$\Gamma(D_{s1}(2536)^- \mu^+ \nu_\mu, D_{s1}^- \rightarrow D^{*-} K_S^0)/\Gamma_{\text{total}}$

Γ_6/Γ

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|------------------------|
| 2.5 ± 0.7 ± 0.1 | ¹ ABAZOV | 09G D0 | $p\bar{p}$ at 1.96 TeV |

¹ ABAZOV 09G reports $[\Gamma(B_s^0 \rightarrow D_{s1}(2536)^- \mu^+ \nu_\mu, D_{s1}^- \rightarrow D^{*-} K_S^0)/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow B_s^0)] = (2.66 \pm 0.52 \pm 0.45) \times 10^{-4}$ which we divide by our best value $B(\bar{b} \rightarrow B_s^0) = (10.5 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{s1}(2536)^- X \mu^+ \nu, D_{s1}^- \rightarrow \bar{D}^0 K^+)/\Gamma(D_s^- \ell^+ \nu_\ell \text{ anything})$ Γ_7/Γ_5

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--------------------|
| $5.4 \pm 1.2 \pm 0.5$ | AAIJ | 11A | LHCB pp at 7 TeV |

$\Gamma(D_{s2}(2573)^- X \mu^+ \nu, D_{s2}^- \rightarrow \bar{D}^0 K^+)/\Gamma(D_s^- \ell^+ \nu_\ell \text{ anything})$ Γ_8/Γ_5

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--------------------|
| $3.3 \pm 1.0 \pm 0.4$ | AAIJ | 11A | LHCB pp at 7 TeV |

$\Gamma(D_{s1}(2536)^- X \mu^+ \nu, D_{s1}^- \rightarrow \bar{D}^0 K^+)/\Gamma(D_{s2}(2573)^- X \mu^+ \nu, D_{s2}^- \rightarrow \bar{D}^0 K^+)$ Γ_7/Γ_8

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------------------------|-------------------|-----|--------------------|
| $0.61 \pm 0.14 \pm 0.05$ | ¹ AAIJ | 11A | LHCB pp at 7 TeV |
|--------------------------|-------------------|-----|--------------------|

¹ Not independent of other AAIJ 11A measurements.

$\Gamma(D_s^- \pi^+)/\Gamma_{\text{total}}$ Γ_9/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

3.04 ± 0.23 OUR FIT

3.02 ± 0.24 OUR AVERAGE

| | | | | |
|---------------------------------|-------------------|------|------|---------------|
| $2.95 \pm 0.05^{+0.25}_{-0.28}$ | ¹ AAIJ | 12AG | LHCB | pp at 7 TeV |
|---------------------------------|-------------------|------|------|---------------|

| | | | | |
|-----------------------|---------------------|----|------|----------------------------------|
| $3.6 \pm 0.5 \pm 0.5$ | ² LOUVOT | 09 | BELL | $e^+ e^- \rightarrow \gamma(5S)$ |
|-----------------------|---------------------|----|------|----------------------------------|

| | | | | |
|-----------------------|------------------------|-----|-----|------------------------|
| $3.0 \pm 0.7 \pm 0.1$ | ³ ABULENCIA | 07C | CDF | $p\bar{p}$ at 1.96 TeV |
|-----------------------|------------------------|-----|-----|------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----------|-----|------|--------------------|
| $6.8 \pm 2.2 \pm 1.6$ | DRUTSKOY | 07A | BELL | Repl. by LOUVOT 09 |
|-----------------------|----------|-----|------|--------------------|

| | | | | |
|-----------------------|------------------------|-----|-----|------------------------|
| $3.5 \pm 1.1 \pm 0.2$ | ⁴ ABULENCIA | 06J | CDF | Repl. by ABULENCIA 07C |
|-----------------------|------------------------|-----|-----|------------------------|

| | | | | |
|------|--------------------|-----|------|-------------------------|
| <130 | ⁵ AKERS | 94J | OPAL | $e^+ e^- \rightarrow Z$ |
|------|--------------------|-----|------|-------------------------|

| | | | | |
|------|-----------------------|-----|------|-------------------------|
| seen | ¹ BUSKULIC | 93G | ALEP | $e^+ e^- \rightarrow Z$ |
|------|-----------------------|-----|------|-------------------------|

¹ AAIJ 12AG reports $(2.95 \pm 0.05 \pm 0.17^{+0.18}_{-0.22}) \times 10^{-3}$ where the last uncertainty comes from the semileptonic f_s/f_d measurement. We combined the systematics in quadrature.

² LOUVOT 09 reports $(3.67^{+0.35+0.65}_{-0.33-0.645}) \times 10^{-3}$ from a measurement of $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+)/\Gamma_{\text{total}}] \times [B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)})]$ assuming $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.5 \pm 2.6) \times 10^{-2}$, which we rescale to our best value $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (20.1 \pm 3.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABULENCIA 07C reports $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+)] = 1.13 \pm 0.08 \pm 0.23$ which we multiply by our best value $B(B^0 \rightarrow D^- \pi^+) = (2.68 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ABULENCIA 06J reports $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+)] = 1.32 \pm 0.18 \pm 0.38$ which we multiply by our best value $B(B^0 \rightarrow D^- \pi^+) = (2.68 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ AKERS 94J sees ≤ 6 events and measures the limit on the product branching fraction $f(\bar{b} \rightarrow B_s^0) \cdot B(B_s^0 \rightarrow D_s^- \pi^+) < 1.3\%$ at CL = 90%. We divide by our current value $B(\bar{b} \rightarrow B_s^0) = 0.105$.

$\Gamma(D_s^- \rho^+)/\Gamma_{\text{total}}$ Γ_{10}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|--------------------------------------|
| $7.0 \pm 1.4 \pm 0.5$ | ¹ LOUVOT | 10 | BELL $e^+e^- \rightarrow \gamma(5S)$ |

¹ LOUVOT 10 reports $[\Gamma(B_s^0 \rightarrow D_s^- \rho^+)/\Gamma_{\text{total}}] / [B(B_s^0 \rightarrow D_s^- \pi^+)] = 2.3 \pm 0.4 \pm 0.2$ which we multiply by our best value $B(B_s^0 \rightarrow D_s^- \pi^+) = (3.04 \pm 0.23) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|------|----------------------------|
| 6.3 ± 1.1 OUR FIT | | | |
| $6.7 \pm 1.5 \pm 0.7$ | ¹ ABULENCIA | 07C | CDF $p\bar{p}$ at 1.96 TeV |

¹ ABULENCIA 07C reports $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+ \pi^+ \pi^-)] = 1.05 \pm 0.10 \pm 0.22$ which we multiply by our best value $B(B^0 \rightarrow D^- \pi^+ \pi^+ \pi^-) = (6.4 \pm 0.7) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^- \pi^+ \pi^+ \pi^-)/\Gamma(D_s^- \pi^+)$ Γ_{11}/Γ_9

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|--------------------|
| 2.08 ± 0.34 OUR FIT | | | |
| $2.01 \pm 0.37 \pm 0.20$ | AAIJ | 11E | LHCB pp at 7 TeV |

$\Gamma(D_{s1}(2536)^- \pi^+, D_{s1}^- \rightarrow D_s^- \pi^+ \pi^-)/\Gamma(D_s^- \pi^+ \pi^+ \pi^-)$ Γ_{12}/Γ_{11}

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--------------------|
| $4.0 \pm 1.0 \pm 0.4$ | AAIJ | 12AX | LHCB pp at 7 TeV |

$\Gamma(D_s^\mp K^\pm)/\Gamma_{\text{total}}$ Γ_{13}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|------|--------------------------------------|
| 2.03 ± 0.28 OUR FIT | | | Error includes scale factor of 1.3. |
| $2.3 \begin{matrix} +1.2 & +0.4 \\ -1.0 & -0.3 \end{matrix}$ | ¹ LOUVOT | 09 | BELL $e^+e^- \rightarrow \gamma(5S)$ |

¹ LOUVOT 09 reports $(2.4 \begin{matrix} +1.2 \\ -1.0 \end{matrix} \pm 0.42) \times 10^{-4}$ from a measurement of $[\Gamma(B_s^0 \rightarrow D_s^\mp K^\pm)/\Gamma_{\text{total}}] \times [B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)})]$ assuming $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.5 \pm 2.6) \times 10^{-2}$, which we rescale to our best value $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (20.1 \pm 3.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^\mp K^\pm)/\Gamma(D_s^- \pi^+)$ Γ_{13}/Γ_9

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|-------------------------------------|
| 0.067 ± 0.008 OUR FIT | | | Error includes scale factor of 1.6. |
| 0.066 ± 0.008 OUR AVERAGE | | | Error includes scale factor of 1.6. |
| $0.0646 \pm 0.0043 \pm 0.0025$ | AAIJ | 12AG | LHCB pp at 7 TeV |
| $0.097 \pm 0.018 \pm 0.009$ | AALTONEN | 09AQ | CDF $p\bar{p}$ at 1.96 TeV |

$\Gamma(D_s^- K^+ \pi^+ \pi^-)/\Gamma(D_s^- \pi^+ \pi^+ \pi^-)$ Γ_{14}/Γ_{11}

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--------------------|
| $5.2 \pm 0.5 \pm 0.3$ | AAIJ | 12AX | LHCB pp at 7 TeV |

$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-----------------------|-----------|------------------------------------|
| 4.4±0.5 OUR AVERAGE | | | | |
| $4.0 \pm 0.2 \pm 0.5$ | | ¹ AAIJ | 13AP LHCb | pp at 7 TeV |
| $5.8^{+1.1}_{-0.9} \pm 1.3$ | | ² ESEN | 13 BELL | $e^+ e^- \rightarrow \Upsilon(5S)$ |
| $5.1 \pm 0.8 \pm 0.6$ | | ³ AALTONEN | 12C CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $10.3^{+3.9+2.6}_{-3.2-2.5}$ | | ⁴ ESEN | 10 BELL | Repl. by ESEN 13 |
| $10.4^{+3.5}_{-3.2} \pm 1.1$ | | ⁵ AALTONEN | 08F CDF | Repl. by AALTONEN 12C |
| <67 | 90 | DRUTSKOY | 07A BELL | Repl. by ESEN 10 |

¹ Uses $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$.

² Use $\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*$ decays assuming $B(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$ and $\Gamma(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$.

³ AALTONEN 12C reports (f_s/f_d) ($B(B_s^0 \rightarrow D_s^+ D_s^-) / B(B^0 \rightarrow D^- D_s^+) = 0.183 \pm 0.021 \pm 0.017$. We multiply this result by our best value of $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ and divide by our best value of f_s/f_d , where $1/2 f_s/f_d = 0.130 \pm 0.008$. Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using our best values.

⁴ Uses $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$ assuming $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.

⁵ AALTONEN 08F reports $[\Gamma(B_s^0 \rightarrow D_s^+ D_s^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 1.44^{+0.48}_{-0.44}$ which we multiply by our best value $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^- D^+)/\Gamma_{\text{total}}$ Γ_{16}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|-----------|--------------------|
| 2.8±0.4±0.3 | | | |
| | ¹ AAIJ | 14AA LHCb | pp at 7 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $3.6 \pm 0.6 \pm 0.5$ | ² AAIJ | 13AP LHCb | Repl. by AAIJ 14AA |

¹ AAIJ 14AA reports $[\Gamma(B_s^0 \rightarrow D_s^- D^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 0.038 \pm 0.004 \pm 0.003$ which we multiply by our best value $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..

² Uses $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$.

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|-----------|---------------|
| 2.2±0.4±0.4 | | | |
| | ¹ AAIJ | 13AP LHCb | pp at 7 TeV |

¹ Uses $B(B^0 \rightarrow D^- D^+) = (2.11 \pm 0.31) \times 10^{-4}$ and $B(B^+ \rightarrow \bar{D}^0 D_s^+) = (10.1 \pm 1.7) \times 10^{-3}$.

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|------|--------------------|
| $1.9 \pm 0.3 \pm 0.4$ | ¹ AAIJ | 13AP | LHCB pp at 7 TeV |

¹ Uses $B(B^0 \rightarrow D^- D^+) = (2.11 \pm 0.31) \times 10^{-4}$ and $B(B^+ \rightarrow \bar{D}^0 D_s^+) = (10.1 \pm 1.7) \times 10^{-3}$.

$\Gamma(D_s^{*-} \pi^+)/\Gamma_{\text{total}}$ Γ_{19}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|---------------------------------------|
| $2.0^{+0.5+0.1}_{-0.4-0.2}$ | ¹ LOUVOT | 10 | BELL $e^+ e^- \rightarrow \gamma(5S)$ |

¹ LOUVOT 10 reports $[\Gamma(B_s^0 \rightarrow D_s^{*-} \pi^+)/\Gamma_{\text{total}}] / [B(B_s^0 \rightarrow D_s^- \pi^+)] = 0.65^{+0.15}_{-0.13} \pm 0.07$ which we multiply by our best value $B(B_s^0 \rightarrow D_s^- \pi^+) = (3.04 \pm 0.23) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^{*-} \rho^+)/\Gamma_{\text{total}}$ Γ_{20}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|---------------------------------------|
| $9.7 \pm 2.0^{+0.7}_{-0.8}$ | ¹ LOUVOT | 10 | BELL $e^+ e^- \rightarrow \gamma(5S)$ |

¹ LOUVOT 10 reports $[\Gamma(B_s^0 \rightarrow D_s^{*-} \rho^+)/\Gamma_{\text{total}}] / [B(B_s^0 \rightarrow D_s^- \pi^+)] = 3.2 \pm 0.6 \pm 0.3$ which we multiply by our best value $B(B_s^0 \rightarrow D_s^- \pi^+) = (3.04 \pm 0.23) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^{*-} \rho^+)/\Gamma(D_s^- \rho^+)$ Γ_{20}/Γ_{10}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------------|------|---------------------------------------|
| $1.4 \pm 0.3 \pm 0.1$ | LOUVOT | 10 | BELL $e^+ e^- \rightarrow \gamma(5S)$ |

$[\Gamma(D_s^{*+} D_s^-) + \Gamma(D_s^{*-} D_s^+)]/\Gamma_{\text{total}}$ Γ_{21}/Γ

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------------------------------|------|---------|
| 12.9 ± 2.2 OUR AVERAGE | | Error includes scale factor of 1.1. | | |

$17.6^{+2.3}_{-2.2} \pm 4.0$ ¹ ESEN 13 BELL $e^+ e^- \rightarrow \gamma(5S)$
 $11.8 \pm 1.6 \pm 1.4$ ² AALTONEN 12C CDF $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •
 $27.5^{+8.3}_{-7.1} \pm 6.9$ ³ ESEN 10 BELL Repl. by ESEN 13
 <121 90 DRUTSKOY 07A BELL Repl. by ESEN 10

¹ Use $\gamma(5S) \rightarrow B_s^* \bar{B}_s^*$ decays assuming $B(\gamma(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$ and $\Gamma(\gamma(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\gamma(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$.

² AALTONEN 12C reports $(f_s/f_d) (B(B_s^0 \rightarrow D_s^{*+} D_s^- + D_s^{*-} D_s^+) / B(B^0 \rightarrow D^- D_s^+)) = 0.424 \pm 0.046 \pm 0.035$. We multiply this result by our best value of $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ and divide by our best value of f_s/f_d , where $1/2 f_s/f_d = 0.130 \pm 0.008$. Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using our best values.

³ Uses $\gamma(10860) \rightarrow B_s^* \bar{B}_s^*$ assuming $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\gamma(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$ Γ_{22}/Γ

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

18.6 ± 3.0 OUR AVERAGE

| | | | | |
|--|--|--------|----|--|
| 19.8 ^{+3.3+5.2} _{-3.1-5.0} | | 1 ESEN | 13 | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |
|--|--|--------|----|--|

| | | | | |
|------------------|--|------------|-----|----------------------------|
| 18.2 ± 2.7 ± 2.2 | | 2 AALTONEN | 12C | CDF $p\bar{p}$ at 1.96 TeV |
|------------------|--|------------|-----|----------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--|--|--------|----|-----------------------|
| 30.8 ^{+12.2+8.5} _{-10.4-8.6} | | 3 ESEN | 10 | BELL Repl. by ESEN 13 |
|--|--|--------|----|-----------------------|

| | | | | |
|------|----|----------|-----|-----------------------|
| <257 | 90 | DRUTSKOY | 07A | BELL Repl. by ESEN 10 |
|------|----|----------|-----|-----------------------|

¹ Use $\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*$ decays assuming $B(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$ and

$\Gamma(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$.

² AALTONEN 12C reports $(f_s/f_d) (B(B_s^0 \rightarrow D_s^{*+} D_s^{*-}) / B(B^0 \rightarrow D^- D_s^+)) = 0.654 \pm 0.072 \pm 0.065$. We multiply this result by our best value of $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ and divide by our best value of f_s/f_d , where $1/2 f_s/f_d = 0.130 \pm 0.008$. Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using our best values.

³ Uses $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$ assuming $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and

$\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.

$\Gamma(D_s^{(*)+} D_s^{(*)-})/\Gamma_{\text{total}}$ Γ_{23}/Γ

"OUR EVALUATION" is an average using rescaled values of the data listed below.

The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account correlations between the measurements.

| VALUE (%) | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------|-----|-------------|------|---------|
|-----------|-----|-------------|------|---------|

4.5 ± 1.4 OUR EVALUATION

3.7 ± 0.5 OUR AVERAGE

| | | | | |
|--|--|--------|----|--|
| 4.32 ^{+0.42+1.04} _{-0.39-1.03} | | 1 ESEN | 13 | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |
|--|--|--------|----|--|

| | | | | |
|-----------------|--|------------|-----|----------------------------|
| 3.5 ± 0.4 ± 0.4 | | 2 AALTONEN | 12C | CDF $p\bar{p}$ at 1.96 TeV |
|-----------------|--|------------|-----|----------------------------|

| | | | | |
|-----------------|--|----------|-----|---------------------------|
| 3.5 ± 1.0 ± 1.1 | | 3 ABAZOV | 09I | D0 $p\bar{p}$ at 1.96 TeV |
|-----------------|--|----------|-----|---------------------------|

| | | | | |
|------------|--|------------|-----|-----------------------------|
| 14 ± 6 ± 3 | | 4,5 BARATE | 00K | ALEP $e^+e^- \rightarrow Z$ |
|------------|--|------------|-----|-----------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--|--|----------|----|-----------------------|
| 6.85 ^{+1.53+1.79} _{-1.30-1.80} | | 6,7 ESEN | 10 | BELL Repl. by ESEN 13 |
|--|--|----------|----|-----------------------|

| | | | | |
|---|--|----------|-----|------------------------|
| 3.9 ^{+1.9+1.6} _{-1.7-1.5} | | 3 ABAZOV | 07Y | D0 Repl. by ABAZOV 09I |
|---|--|----------|-----|------------------------|

| | | | | |
|--------|----|--------|-----|-----------------------------|
| <0.218 | 90 | BARATE | 98Q | ALEP $e^+e^- \rightarrow Z$ |
|--------|----|--------|-----|-----------------------------|

¹ Use $\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*$ decays assuming $B(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$ and

$\Gamma(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$.

² AALTONEN 12C reports $(f_s/f_d) (B(B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}) / B(B^0 \rightarrow D^- D_s^+)) = 1.261 \pm 0.095 \pm 0.112$. We multiply this result by our best value of $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ and divide by our best value of f_s/f_d , where $1/2 f_s/f_d = 0.130 \pm 0.008$. Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using our best values.

³ Uses the final states where $D_s^+ \rightarrow \phi \pi^+$ and $D_s^- \rightarrow \phi \mu^- \bar{\nu}_\mu$.

⁴ Reports $B(B_S^0(\text{short}) \rightarrow D_S^{(*)} D_S^{(*)}) = (0.23 \pm 0.10 \pm 0.05) \cdot [0.17/B(D_S \rightarrow \phi\chi)]^2$ assuming $B(B_S^0 \rightarrow B_S^0(\text{short})) = 50\%$. We use our best value of $B(D_S \rightarrow \phi\chi) = 15.7 \pm 1.0\%$ to obtain the quoted result.

⁵ Uses $\phi\phi$ correlations from $B_S^0(\text{short}) \rightarrow D_S^{(*)+} D_S^{(*)-}$.

⁶ Sum of exclusive $B_S \rightarrow D_S^+ D_S^-$, $B_S \rightarrow D_S^{*\pm} D_S^\mp$ and $B_S \rightarrow D_S^{*+} D_S^{*-}$.

⁷ Uses $\Upsilon(10860) \rightarrow B_S^* \bar{B}_S^*$ assuming $B(\Upsilon(10860) \rightarrow B_S^{(*)} \bar{B}_S^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\Upsilon(10860) \rightarrow B_S^* \bar{B}_S^*) / \Gamma(\Upsilon(10860) \rightarrow B_S^{(*)} \bar{B}_S^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.

$\Gamma(\bar{D}^0 K^- \pi^+) / \Gamma_{\text{total}}$ Γ_{24} / Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|-----------|---------------|
| 9.9 ± 1.1 ± 1.1 | ¹ AAIJ | 13AQ LHCb | pp at 7 TeV |

¹ AAIJ 13AQ reports $[\Gamma(B_S^0 \rightarrow \bar{D}^0 K^- \pi^+) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-)] = 1.18 \pm 0.05 \pm 0.12$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-) = (8.4 \pm 0.9) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{D}^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}$ Γ_{25} / Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------|------|---------|
| 4.4 ± 0.6 OUR AVERAGE | | | |

| | | | |
|--------------------|-------------------|-----------|------------------|
| 4.29 ± 0.09 ± 0.65 | ¹ AAIJ | 14BH LHCb | pp at 7, 8 TeV |
| 4.7 ± 1.2 ± 0.7 | ² AAIJ | 11D LHCb | pp at 7 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----------------|-------------------|-----------|--------------------|
| 3.3 ± 0.4 ± 0.5 | ³ AAIJ | 13BX LHCb | Repl. by AAIJ 14BH |
|-----------------|-------------------|-----------|--------------------|

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

² AAIJ 11D reports $[\Gamma(B_S^0 \rightarrow \bar{D}^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 \rho^0)] = 1.48 \pm 0.34 \pm 0.19$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0 \rho^0) = (3.2 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AAIJ 13BX reports $[\Gamma(B_S^0 \rightarrow \bar{D}^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 K^*(892)^0)] = 7.8 \pm 0.7 \pm 0.3 \pm 0.6$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0 K^*(892)^0) = (4.2 \pm 0.6) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{D}^0 \bar{K}^*(1410)) / \Gamma_{\text{total}}$ Γ_{26} / Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------------|-----------|------------------|
| 38.6 ± 11.4 ± 33.3 | ¹ AAIJ | 14BH LHCb | pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

$\Gamma(\bar{D}^0 \bar{K}_0^*(1430)) / \Gamma_{\text{total}}$ Γ_{27} / Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|-----------|------------------|
| 30.0 ± 2.4 ± 6.8 | ¹ AAIJ | 14BH LHCb | pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays. Corresponds to the resonant $K_0^*(1430)$ part of LASS parametrisation.

$\Gamma(\overline{D}^0 \overline{K}_2^*(1430))/\Gamma_{\text{total}}$ Γ_{28}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------------|-----------|------------------|
| 11.1±1.8±3.8 | | ¹ AAIJ | 14BH LHCB | pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \overline{D}^0 K^- \pi^+$ decays.

$\Gamma(\overline{D}^0 \overline{K}^*(1680))/\Gamma_{\text{total}}$ Γ_{29}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------------|-----------|------------------|
| <7.8 | 90 | ¹ AAIJ | 14BH LHCB | pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \overline{D}^0 K^- \pi^+$ decays.

$\Gamma(\overline{D}^0 \overline{K}_0^*(1950))/\Gamma_{\text{total}}$ Γ_{30}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------------|-----------|------------------|
| <11 | 90 | ¹ AAIJ | 14BH LHCB | pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \overline{D}^0 K^- \pi^+$ decays.

$\Gamma(\overline{D}^0 \overline{K}_3^*(1780))/\Gamma_{\text{total}}$ Γ_{31}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------------|-----------|------------------|
| <2.6 | 90 | ¹ AAIJ | 14BH LHCB | pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \overline{D}^0 K^- \pi^+$ decays.

$\Gamma(\overline{D}^0 \overline{K}_4^*(2045))/\Gamma_{\text{total}}$ Γ_{32}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------------|-----------|------------------|
| <3.1 | 90 | ¹ AAIJ | 14BH LHCB | pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \overline{D}^0 K^- \pi^+$ decays.

$\Gamma(\overline{D}^0 K^- \pi^+ (\text{non-resonant}))/\Gamma_{\text{total}}$ Γ_{33}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|-----------|------------------|
| 20.6±3.8±7.3 | ¹ AAIJ | 14BH LHCB | pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \overline{D}^0 K^- \pi^+$ decays. Corresponds to the non-resonant part of the LASS parametrisation.

$\Gamma(D_{s2}^*(2573)^- \pi^+, D_{s2}^* \rightarrow \overline{D}^0 K^-)/\Gamma_{\text{total}}$ Γ_{34}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|-----------|------------------|
| 25.7±0.7±4.0 | ¹ AAIJ | 14BH LHCB | pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \overline{D}^0 K^- \pi^+$ decays.

$\Gamma(D_{s1}^*(2700)^- \pi^+, D_{s1}^* \rightarrow \overline{D}^0 K^-)/\Gamma_{\text{total}}$ Γ_{35}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|-----------|------------------|
| 1.6±0.4±0.7 | ¹ AAIJ | 14BH LHCB | pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_S^0 \rightarrow \overline{D}^0 K^- \pi^+$ decays.

$\Gamma(D_{s1}^*(2860)^- \pi^+, D_{s1}^* \rightarrow \bar{D}^0 K^-) / \Gamma_{\text{total}}$ Γ_{36} / Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|------|-----------------------|
| 5.0 ± 1.2 ± 3.4 | ¹ AAIJ | 14BH | LHCB pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

$\Gamma(D_{s3}^*(2860)^- \pi^+, D_{s3}^* \rightarrow \bar{D}^0 K^-) / \Gamma_{\text{total}}$ Γ_{37} / Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|------|-----------------------|
| 2.2 ± 0.1 ± 0.6 | ¹ AAIJ | 14BH | LHCB pp at 7, 8 TeV |

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

$\Gamma(\bar{D}^0 K^+ K^-) / \Gamma_{\text{total}}$ Γ_{38} / Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|---------------------|------|--------------------|
| 4.2 ± 1.6 ± 1.1 | ^{1,2} AAIJ | 12AM | LHCB pp at 7 TeV |

¹ AAIJ 12AM reports $[\Gamma(B_s^0 \rightarrow \bar{D}^0 K^+ K^-) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 K^+ K^-)] = 0.90 \pm 0.27 \pm 0.20$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0 K^+ K^-) = (4.7 \pm 1.2) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Uses $B(b \rightarrow B_s^0) / B(b \rightarrow B^0) = 0.267^{+0.023}_{-0.020}$ measured by the same authors.

$\Gamma(\bar{D}^0 \phi) / \Gamma(\bar{D}^0 \bar{K}^*(892)^0)$ $\Gamma_{39} / \Gamma_{25}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------|------|--------------------|
| 0.069 ± 0.013 ± 0.007 | AAIJ | 13BX | LHCB pp at 7 TeV |

$\Gamma(D^{*\mp} \pi^\pm) / \Gamma_{\text{total}}$ Γ_{40} / Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|-----|-------------------|------|--------------------|
| < 6.1 × 10⁻⁶ | 90 | ¹ AAIJ | 13AL | LHCB pp at 7 TeV |

¹ Uses $f_s / f_d = 0.256 \pm 0.020$ and $B(B^0 \rightarrow D^{*-} \pi^+) = (2.76 \pm 0.13) \times 10^{-3}$.

$\Gamma(J/\psi(1S) \phi) / \Gamma_{\text{total}}$ Γ_{41} / Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|---------------------|------|---|
| 1.08 ± 0.09 OUR FIT | | | | |
| 1.10 ± 0.09 OUR AVERAGE | | | | |
| 1.050 ± 0.013 ± 0.104 | | ¹ AAIJ | 13AN | LHCB pp at 7 TeV |
| 1.25 ± 0.07 ± 0.23 | | ² THORNE | 13 | BELL $e^+ e^- \rightarrow \Upsilon(5S)$ |
| 1.4 ± 0.5 ± 0.1 | | ³ ABE | 96Q | CDF $p\bar{p}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|--------------------|-----|------------------------------|
| <6 | 1 | ⁴ AKERS | 94J | OPAL $e^+ e^- \rightarrow Z$ |
| seen | 14 | ⁵ ABE | 93F | CDF $p\bar{p}$ at 1.8 TeV |
| seen | 1 | ⁶ ACTON | 92N | OPAL Sup. by AKERS 94J |

¹ Uses $f_s / f_d = 0.256 \pm 0.020$ and $B(B^+ \rightarrow J/\psi K^+) = (10.18 \pm 0.42) \times 10^{-4}$.

² Uses $f_s = (17.2 \pm 3.0)\%$ as the fraction of $\Upsilon(5S)$ decaying to $B_s^{(*)} \bar{B}_s^{(*)}$.

³ ABE 96Q reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S) \phi) / \Gamma_{\text{total}}] \times [\Gamma(\bar{b} \rightarrow B_s^0) / (\Gamma(\bar{b} \rightarrow B^+) + \Gamma(\bar{b} \rightarrow B^0))] = (0.185 \pm 0.055 \pm 0.020) \times 10^{-3}$ which we divide by our best value $\Gamma(\bar{b} \rightarrow B_s^0) / [\Gamma(\bar{b} \rightarrow B^+) + \Gamma(\bar{b} \rightarrow B^0)] = 0.130 \pm 0.008$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ AKERS 94J sees one event and measures the limit on the product branching fraction $f(\bar{b} \rightarrow B_S^0) \cdot B(B_S^0 \rightarrow J/\psi(1S)\phi) < 7 \times 10^{-4}$ at CL = 90%. We divide by $B(\bar{b} \rightarrow B_S^0) = 0.112$.

⁵ ABE 93F measured using $J/\psi(1S) \rightarrow \mu^+ \mu^-$ and $\phi \rightarrow K^+ K^-$.

⁶ In ACTON 92N a limit on the product branching fraction is measured to be $f(\bar{b} \rightarrow B_S^0) \cdot B(B_S^0 \rightarrow J/\psi(1S)\phi) \leq 0.22 \times 10^{-2}$.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_{42}/Γ

| VALUE | CL% | DOCUMENT ID | TECN |
|----------------------------------|-----|-----------------------|--------|
| <1.2 × 10⁻³ | 90 | ¹ ACCIARRI | 97C L3 |

¹ ACCIARRI 97C assumes B^0 production fraction ($39.5 \pm 4.0\%$) and B_S ($12.0 \pm 3.0\%$).

$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$ Γ_{43}/Γ

| VALUE (units 10 ⁻⁴) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------|------|---|
| 3.9 ± 0.7 OUR AVERAGE | | | | Error includes scale factor of 1.4. |
| 3.6 ± 0.5 ± 0.2 | | ¹ AAIJ | 13A | LHCB pp at 7 TeV |
| 5.10 ± 0.50 ^{+1.17} _{-0.83} | | ² LI | 12 | BELL $e^+ e^- \rightarrow \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<38 90 ³ ACCIARRI 97C L3

¹ AAIJ 13A reports $[\Gamma(B_S^0 \rightarrow J/\psi(1S)\eta)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)\rho^0)] = 14.0 \pm 1.2^{+1.1+1.1}_{-1.5-1.0}$ which we multiply by our best value $B(B^0 \rightarrow J/\psi(1S)\rho^0) = (2.54 \pm 0.14) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Observed for the first time with significances over 10σ . The second error are total systematic uncertainties including the error on $N(B_S^{(*)}\bar{B}_S^{(*)})$.

³ ACCIARRI 97C assumes B^0 production fraction ($39.5 \pm 4.0\%$) and B_S ($12.0 \pm 3.0\%$).

$\Gamma(J/\psi(1S)K_S^0)/\Gamma_{\text{total}}$ Γ_{44}/Γ

| VALUE (units 10 ⁻⁵) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-----------------------|------|----------------------------|
| 1.87 ± 0.17 OUR AVERAGE | | | |
| 1.89 ± 0.15 ± 0.13 | ¹ AAIJ | 13AB | LHCB pp at 7 TeV |
| 1.8 ± 0.4 ± 0.1 | ² AALTONEN | 11A | CDF $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.88 ± 0.24 ± 0.13 ³ AAIJ 12O LHCB Repl. by AAIJ 13AB

¹ AAIJ 13AB reports $(1.97 \pm 0.14 \pm 0.07 \pm 0.15 \pm 0.08) \times 10^{-5}$ from a measurement of $[\Gamma(B_S^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^0)] \times [\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0)]$ assuming $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.98 \pm 0.35) \times 10^{-4}$, $\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.256 \pm 0.020$, which we rescale to our best values $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.73 \pm 0.32) \times 10^{-4}$, $\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.260 \pm 0.015$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² AALTONEN 11A reports $[\Gamma(B_S^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow B_S^0)] / [B(\bar{b} \rightarrow B^0)] / [B(B^0 \rightarrow J/\psi(1S)K_S^0)] = (1.09 \pm 0.19 \pm 0.11) \times 10^{-2}$ which we multiply or divide by our best values $B(\bar{b} \rightarrow B_S^0) = (10.5 \pm 0.5) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$, $B(B^0 \rightarrow J/\psi(1S)K_S^0) = 1/2 \times B(B^0 \rightarrow J/\psi(1S)K^0) = 1/2$

$\times (8.73 \pm 0.32) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

³ AAIJ 120 reports $(1.83 \pm 0.21 \pm 0.10 \pm 0.14 \pm 0.07) \times 10^{-5}$ from a measurement of $[\Gamma(B_S^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^0)] \times [\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0)]$ assuming $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.71 \pm 0.32) \times 10^{-4}$, $\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.267^{+0.021}_{-0.02}$, which we rescale to our best values $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.73 \pm 0.32) \times 10^{-4}$, $\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.260 \pm 0.015$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(J/\psi(1S)K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{45}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|------|--------------------|
| $4.4^{+0.5}_{-0.4} \pm 0.8$ | ¹ AAIJ | 12AP | LHCB pp at 7 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|----------------------------|-----------------------|-----|----------------------------|
| ⁹ $\pm 4 \pm 1$ | ² AALTONEN | 11A | CDF $p\bar{p}$ at 1.96 TeV |
|----------------------------|-----------------------|-----|----------------------------|

¹ AAIJ 12AP reports $B(B_S^0 \rightarrow J/\psi(1S)K^*(892)^0)/B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (3.43^{+0.34}_{-0.36} \pm 0.50) \times 10^{-2}$ and $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.29 \pm 0.05 \pm 0.13) \times 10^{-3}$ after correcting for the contribution from $K\pi$ S -wave beneath the K^* peak.

² AALTONEN 11A reports $[\Gamma(B_S^0 \rightarrow J/\psi(1S)K^*(892)^0)/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow B_S^0)] / [B(\bar{b} \rightarrow B^0)] / [B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] = 0.0168 \pm 0.0024 \pm 0.0068$ which we multiply or divide by our best values $B(\bar{b} \rightarrow B_S^0) = (10.5 \pm 0.5) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$, $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.32 \pm 0.06) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(J/\psi(1S)\eta')/\Gamma_{\text{total}}$ Γ_{46}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---------|
| 3.3 ± 0.4 OUR AVERAGE | | | |

| | | | |
|---|-------------------|-----|--------------------|
| ^{3.2} $^{+0.4}_{-0.5} \pm 0.2$ | ¹ AAIJ | 13A | LHCB pp at 7 TeV |
|---|-------------------|-----|--------------------|

| | | | |
|--|-----------------|----|--------------------------------------|
| ^{3.71} $\pm 0.61^{+0.85}_{-0.60}$ | ² LI | 12 | BELL $e^+e^- \rightarrow \gamma(4S)$ |
|--|-----------------|----|--------------------------------------|

¹ AAIJ 13A reports $[\Gamma(B_S^0 \rightarrow J/\psi(1S)\eta')/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)\rho^0)] = 12.7 \pm 1.1^{+0.5+1.0}_{-1.3-0.9}$ which we multiply by our best value $B(B^0 \rightarrow J/\psi(1S)\rho^0) = (2.54 \pm 0.14) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Observed for the first time with significances over 10σ . The second error are total systematic uncertainties including the error on $N(B_S^{(*)}\bar{B}_S^{(*)})$.

$\Gamma(J/\psi(1S)\eta')/\Gamma(J/\psi(1S)\eta)$ Γ_{46}/Γ_{43}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---------|
| 0.87 ± 0.06 OUR AVERAGE | | | |

| | | | |
|--|-------------------|-----|-----------------------|
| ^{0.902} $\pm 0.072 \pm 0.045$ | ¹ AAIJ | 15D | LHCB pp at 7, 8 TeV |
|--|-------------------|-----|-----------------------|

| | | | |
|--|-------------------|-----|--------------------|
| ^{0.90} $\pm 0.09^{+0.06}_{-0.02}$ | ² AAIJ | 13A | LHCB pp at 7 TeV |
|--|-------------------|-----|--------------------|

| | | | |
|-------------------------------------|-----------------|----|--------------------------------------|
| ^{0.73} $\pm 0.14 \pm 0.02$ | ² LI | 12 | BELL $e^+e^- \rightarrow \gamma(4S)$ |
|-------------------------------------|-----------------|----|--------------------------------------|

¹ Uses $J/\psi \rightarrow \mu^+ \mu^-$, $\eta' \rightarrow \rho^0 \gamma$, and $\eta' \rightarrow \eta \pi^+ \pi^-$ decays.

² Strongly correlated with measurements of $\Gamma(J/\psi(1S)\eta)/\Gamma$ and $\Gamma(J/\psi(1S)\eta')/\Gamma$ reported in the same reference.

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\phi)$ Γ_{47}/Γ_{41}

| <u>VALUE (units 10^{-2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| 19.8±0.5±0.5 | ¹ AAIJ | 12AO LHCB | pp at 7 TeV |

¹ AAIJ 12AO reports $(19.79 \pm 0.47 \pm 0.52) \times 10^{-2}$ from a measurement of $[\Gamma(B_S^0 \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma(B_S^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$.

$\Gamma(J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{50}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-----------------------------------|
| 1.35±0.16 OUR FIT | | | |
| 1.16^{+0.31+0.30}_{-0.19-0.25} | ¹ LI | 11 BELL | $e^+e^- \rightarrow \Upsilon(5S)$ |

¹ The second error includes both the detector systematic and the uncertainty in the number of produced $\Upsilon(5S) \rightarrow B_S^{(*)}\bar{B}_S^{(*)}$ pairs.

$\Gamma(J/\psi(1S)f_0(500), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)f_0(980)_0, f_0 \rightarrow \pi^+\pi^-)$ Γ_{48}/Γ_{51}

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|------------|--------------------|-------------|------------------|
| <0.034 | 90 | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S)\rho, \rho \rightarrow \pi^+\pi^-)/\Gamma(\psi(2S)\pi^+\pi^-)$ Γ_{49}/Γ_{73}

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|------------|--------------------|-------------|------------------|
| <0.017 | 90 | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S)f_0(980)_0, f_0 \rightarrow \pi^+\pi^-)/\Gamma(\psi(2S)\pi^+\pi^-)$ Γ_{51}/Γ_{73}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------|
| 0.703±0.015^{+0.004}_{-0.051} | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S)f_2(1270)_0, f_2 \rightarrow \pi^+\pi^-)/\Gamma(\psi(2S)\pi^+\pi^-)$ Γ_{53}/Γ_{73}

| <u>VALUE (%)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|--------------------|-------------|------------------|
| 0.36±0.07±0.03 | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S)f_2(1270)_{||}, f_2 \rightarrow \pi^+\pi^-)/\Gamma(\psi(2S)\pi^+\pi^-)$ Γ_{54}/Γ_{73}

| <u>VALUE (%)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------|
| 0.52±0.15^{+0.05}_{-0.02} | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S) f_2(1270)_\perp, f_2 \rightarrow \pi^+ \pi^-) / \Gamma(\psi(2S) \pi^+ \pi^-)$ $\Gamma_{55} / \Gamma_{73}$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------|-------------|------|---------|
|-----------|-------------|------|---------|

| | | | |
|---------------------------------|-------------------|-----------|------------------|
| $0.63 \pm 0.34^{+0.16}_{-0.08}$ | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |
|---------------------------------|-------------------|-----------|------------------|

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S) f_0(1500), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(\psi(2S) \pi^+ \pi^-)$ $\Gamma_{57} / \Gamma_{73}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

| | | | |
|-------------------------------------|-------------------|-----------|------------------|
| $0.101 \pm 0.008^{+0.011}_{-0.003}$ | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |
|-------------------------------------|-------------------|-----------|------------------|

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S) f'_2(1525)_0, f'_2 \rightarrow \pi^+ \pi^-) / \Gamma(\psi(2S) \pi^+ \pi^-)$ $\Gamma_{58} / \Gamma_{73}$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------|-------------|------|---------|
|-----------|-------------|------|---------|

| | | | |
|---------------------------------|-------------------|-----------|------------------|
| $0.51 \pm 0.09^{+0.05}_{-0.04}$ | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |
|---------------------------------|-------------------|-----------|------------------|

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S) f'_2(1525)_\parallel, f'_2 \rightarrow \pi^+ \pi^-) / \Gamma(\psi(2S) \pi^+ \pi^-)$ $\Gamma_{59} / \Gamma_{73}$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------|-------------|------|---------|
|-----------|-------------|------|---------|

| | | | |
|---------------------------------|-------------------|-----------|------------------|
| $0.06^{+0.13}_{-0.04} \pm 0.01$ | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |
|---------------------------------|-------------------|-----------|------------------|

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S) f'_2(1525)_\perp, f'_2 \rightarrow \pi^+ \pi^-) / \Gamma(\psi(2S) \pi^+ \pi^-)$ $\Gamma_{60} / \Gamma_{73}$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------|-------------|------|---------|
|-----------|-------------|------|---------|

| | | | |
|---------------------------------|-------------------|-----------|------------------|
| $0.26 \pm 0.18^{+0.06}_{-0.04}$ | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |
|---------------------------------|-------------------|-----------|------------------|

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S) f_0(1790), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(\psi(2S) \pi^+ \pi^-)$ $\Gamma_{61} / \Gamma_{73}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

| | | | |
|-------------------------------------|-------------------|-----------|------------------|
| $0.024 \pm 0.004^{+0.050}_{-0.002}$ | ¹ AAIJ | 14BR LHCB | pp at 7, 8 TeV |
|-------------------------------------|-------------------|-----------|------------------|

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(J/\psi(1S) \phi)$ $\Gamma_{50} / \Gamma_{41}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.125 ± 0.011 OUR FIT

0.127 ± 0.011 OUR AVERAGE

| | | | |
|-----------------------------|---------------------|--------|------------------------|
| $0.135 \pm 0.036 \pm 0.001$ | ¹ ABAZOV | 12C D0 | $p\bar{p}$ at 1.96 TeV |
|-----------------------------|---------------------|--------|------------------------|

| | | | |
|-----------------------------|-----------------------|----------|------------------------|
| $0.126 \pm 0.012 \pm 0.001$ | ² AALTONEN | 11AB CDF | $p\bar{p}$ at 1.96 TeV |
|-----------------------------|-----------------------|----------|------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------------------------------|---------------------|-----------|------------------|
| $0.139 \pm 0.006^{+0.025}_{-0.012}$ | ^{3,4} AAIJ | 12A0 LHCB | Repl. by AAIJ 14 |
|-------------------------------------|---------------------|-----------|------------------|

| | | | |
|-------------------------------------|-------------------|---------|--------------------|
| $0.123^{+0.026}_{-0.022} \pm 0.001$ | ⁵ AAIJ | 11 LHCB | Repl. by AAIJ 12A0 |
|-------------------------------------|-------------------|---------|--------------------|

¹ ABAZOV 12C reports $[\Gamma(B_S^0 \rightarrow J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(B_S^0 \rightarrow J/\psi(1S) \phi)] / [B(\phi(1020) \rightarrow K^+ K^-)] = 0.275 \pm 0.041 \pm 0.061$ which we multiply by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AALTONEN 11AB reports $[\Gamma(B_S^0 \rightarrow J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(B_S^0 \rightarrow J/\psi(1S) \phi)] / [B(\phi(1020) \rightarrow K^+ K^-)] = 0.257 \pm 0.020 \pm 0.014$ which we multiply by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AAIJ 12AO reports $(13.9 \pm 0.6_{-1.2}^{+2.5}) \times 10^{-2}$ from a measurement of $[\Gamma(B_S^0 \rightarrow J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(B_S^0 \rightarrow J/\psi(1S) \phi)] / [B(\phi(1020) \rightarrow K^+ K^-)]$ assuming $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$.

⁴ Measured in Dalitz plot like analysis of $B_S \rightarrow J/\psi \pi^+ \pi^-$ decays.

⁵ AAIJ 11 reports $[\Gamma(B_S^0 \rightarrow J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(B_S^0 \rightarrow J/\psi(1S) \phi)] / [B(\phi(1020) \rightarrow K^+ K^-)] = 0.252_{-0.032}^{+0.046} +_{-0.033}^{+0.027}$ which we multiply by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(J/\psi(1S) f_0(1370), f_0 \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{56} / Γ

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
|---|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--|------|---------|------------------------------------|
| $0.34_{-0.14}^{+0.11} +_{-0.054}^{+0.085}$ | 1 LI | 11 BELL | $e^+ e^- \rightarrow \Upsilon(5S)$ |
|--|------|---------|------------------------------------|

¹ The second error includes both the detector systematic and the uncertainty in the number of produced $Y(5S) \rightarrow B_S^{(*)} \bar{B}_S^{(*)}$ pairs.

$\Gamma(J/\psi(1S) f_0(1370), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(J/\psi(1S) \phi)$ $\Gamma_{56} / \Gamma_{41}$

| <u>VALUE (units 10^{-2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
|---|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----------------------------|----------|-----------|------------------|
| $4.2 \pm 0.5_{-3.7}^{+0.1}$ | 1,2 AAIJ | 12AO LHCB | Repl. by AAIJ 14 |
|-----------------------------|----------|-----------|------------------|

¹ AAIJ 12AO reports $(4.19 \pm 0.53_{-3.7}^{+0.12}) \times 10^{-2}$ from a measurement of $[\Gamma(B_S^0 \rightarrow J/\psi(1S) f_0(1370), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(B_S^0 \rightarrow J/\psi(1S) \phi)] / [B(\phi(1020) \rightarrow K^+ K^-)]$ assuming $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$.

² Measured in Dalitz plot like analysis of $B_S \rightarrow J/\psi \pi^+ \pi^-$ decays.

$\Gamma(J/\psi(1S) f_2(1270), f_2 \rightarrow \pi^+ \pi^-) / \Gamma(J/\psi(1S) \phi)$ $\Gamma_{52} / \Gamma_{41}$

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
|---|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----------------------------|----------|-----------|------------------|
| $9.8 \pm 3.3_{-1.5}^{+0.6}$ | 1,2 AAIJ | 12AO LHCB | Repl. by AAIJ 14 |
|-----------------------------|----------|-----------|------------------|

¹ AAIJ 12AO reports $(0.098 \pm 0.033_{-0.015}^{+0.006}) \times 10^{-2}$ from a measurement of $[\Gamma(B_S^0 \rightarrow J/\psi(1S) f_2(1270), f_2 \rightarrow \pi^+ \pi^-) / \Gamma(B_S^0 \rightarrow J/\psi(1S) \phi)] / [B(\phi(1020) \rightarrow K^+ K^-)]$ assuming $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$.

² Measured in Dalitz plot like analysis of $B_S \rightarrow J/\psi \pi^+ \pi^-$ decays for the f_2 helicity state $\lambda = 0$.

$\Gamma(J/\psi(1S)\pi^+\pi^-(\text{nonresonant}))/\Gamma(J/\psi(1S)\phi)$ Γ_{62}/Γ_{41}

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------------------------------|----------|-----------|------------------|
| $1.66 \pm 0.31^{+0.96}_{-0.08}$ | 1,2 AAIJ | 12AO LHCb | Repl. by AAIJ 14 |
|---------------------------------|----------|-----------|------------------|

¹ AAIJ 12AO reports $(1.66 \pm 0.31^{+0.96}_{-0.08}) \times 10^{-2}$ from a measurement of $[\Gamma(B_S^0 \rightarrow J/\psi(1S)\pi^+\pi^-(\text{nonresonant}))/\Gamma(B_S^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$.

² Measured in Dalitz plot like analysis of $B_S \rightarrow J/\psi\pi^+\pi^-$ decays.

$\Gamma(J/\psi(1S)\bar{K}^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{63}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

| | | | | |
|-----------------------|----|-------------------|----------|---------------|
| $<4.4 \times 10^{-5}$ | 90 | ¹ AAIJ | 14L LHCb | pp at 7 TeV |
|-----------------------|----|-------------------|----------|---------------|

¹ Measured with $B(B_S^0 \rightarrow J/\psi K_S^0\pi^+\pi^-) / B(B^0 \rightarrow J/\psi K_S^0\pi^+\pi^-)$ using PDG 12 values for the involved branching fractions.

$\Gamma(J/\psi(1S)K^+K^-)/\Gamma_{\text{total}}$ Γ_{64}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

7.9 ± 0.7 OUR AVERAGE

| | | | |
|--------------------------|---------------------|-----------|-----------------------------------|
| $7.70 \pm 0.08 \pm 0.72$ | ¹ AAIJ | 13AN LHCb | pp at 7 TeV |
| $10.1 \pm 0.9 \pm 2.1$ | ² THORNE | 13 BELL | $e^+e^- \rightarrow \Upsilon(5S)$ |

¹ Uses $f_s/f_d = 0.256 \pm 0.020$ and $B(B^+ \rightarrow J/\psi K^+) = (10.18 \pm 0.42) \times 10^{-4}$.

² Uses $f_s = (17.2 \pm 3.0)\%$ as the fraction of $\Upsilon(5S)$ decaying to $B_S^{(*)}\bar{B}_S^{(*)}$.

$\Gamma(J/\psi(1S)K^0K^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{65}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

| | | | |
|------------------------|-------------------|----------|---------------|
| 9.3 ± 1.0 ± 0.9 | ¹ AAIJ | 14L LHCb | pp at 7 TeV |
|------------------------|-------------------|----------|---------------|

¹ AAIJ 14L reports $[\Gamma(B_S^0 \rightarrow J/\psi(1S)K^0K^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^0\pi^+\pi^-)] = 2.12 \pm 0.15 \pm 0.18$ which we multiply by our best value $B(B^0 \rightarrow J/\psi(1S)K^0\pi^+\pi^-) = (4.4 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. This is an observation of $B_S^0 \rightarrow J/\psi K_S^0 K^\pm \pi^\mp$ with more than 10 standard deviations.

$\Gamma(J/\psi(1S)\bar{K}^0K^+K^-)/\Gamma_{\text{total}}$ Γ_{66}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

| | | | | |
|----------------------|----|-------------------|----------|---------------|
| $<12 \times 10^{-6}$ | 90 | ¹ AAIJ | 14L LHCb | pp at 7 TeV |
|----------------------|----|-------------------|----------|---------------|

¹ Measured with $B(B_S^0 \rightarrow J/\psi K_S^0 K^+K^-)/B(B^0 \rightarrow J/\psi K_S^0 \pi^+\pi^-)$ using PDG 12 values for the involved branching fractions.

$\Gamma(J/\psi(1S)f_2'(1525))/\Gamma(J/\psi(1S)\phi)$ Γ_{67}/Γ_{41}

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

21 ± 4 OUR AVERAGE

| | | | |
|------------------------|-----------------------|---------|-----------------------------------|
| $21.5 \pm 4.9 \pm 2.6$ | ¹ THORNE | 13 BELL | $e^+e^- \rightarrow \Upsilon(5S)$ |
| $21 \pm 7 \pm 1$ | ^{2,3} ABAZOV | 12AF D0 | $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|------------------------|-------------------|----------|--------------------|
| $26.4 \pm 3.5 \pm 0.7$ | ⁴ AAIJ | 12S LHCb | Repl. by AAIJ 13AN |
|------------------------|-------------------|----------|--------------------|

¹ Uses $B(f'_2(1525) \rightarrow K^+ K^-) = (44.4 \pm 1.1)\%$.

² ABAZOV 12AF reports $[\Gamma(B_S^0 \rightarrow J/\psi(1S)f'_2(1525))/\Gamma(B_S^0 \rightarrow J/\psi(1S)\phi)] \times B(f'_2(1525) \rightarrow K^+ K^-) / B(\phi(1020) \rightarrow K^+ K^-) = 0.19 \pm 0.05 \pm 0.04$ which we divide and multiply by our best values $B(f'_2(1525) \rightarrow K^+ K^-) = \frac{1}{2} (88.7 \pm 2.2) \times 10^{-2}$, $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

³ ABAZOV 12AF fits the invariant masses of the $K^+ K^-$ pair in the range $1.35 < M(K^+ K^-) < 2$ GeV.

⁴ AAIJ 12S reports $[(26.4 \pm 2.7 \pm 2.4) \times 10^{-2}$ from a measurement of $\Gamma(B_S^0 \rightarrow J/\psi(1S)f'_2(1525))/\Gamma(B_S^0 \rightarrow J/\psi(1S)\phi)] \times B(f'_2(1525) \rightarrow K^+ K^-) / B(\phi(1020) \rightarrow K^+ K^-)$ assuming $B(f'_2(1525) \rightarrow K^+ K^-) = (44.4 \pm 1.1) \times 10^{-2}$, $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$, which we rescale to our best values $B(f'_2(1525) \rightarrow K^+ K^-) = \frac{1}{2} (88.7 \pm 2.2) \times 10^{-2}$, $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(J/\psi(1S)f'_2(1525))/\Gamma_{\text{total}}$ Γ_{67}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| $2.61 \pm 0.20^{+0.56}_{-0.50}$ | ¹ AAIJ | 13AN LHCB | pp at 7 TeV |

¹ Uses $f_s/f_d = 0.256 \pm 0.020$ and $B(B^+ \rightarrow J/\psi K^+) = (10.18 \pm 0.42) \times 10^{-4}$.

$\Gamma(\psi(2S)\eta)/\Gamma(J/\psi(1S)\eta)$ Γ_{71}/Γ_{43}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|----------------|
| $0.83 \pm 0.14 \pm 0.12$ | ¹ AAIJ | 13AA LHCB | pp at 7 TeV |

¹ Assuming lepton universality for dimuon decay modes of J/ψ and $\psi(2S)$ mesons, the ratio $B(J/\psi \rightarrow \mu^+ \mu^-)/B(\psi(2S) \rightarrow \mu^+ \mu^-) = B(J/\psi \rightarrow e^+ e^-)/B(\psi(2S) \rightarrow e^+ e^-) = 7.69 \pm 0.19$ was used.

$\Gamma(\psi(2S)\eta')/\Gamma(J/\psi(1S)\eta')$ Γ_{72}/Γ_{46}

| <u>VALUE (units 10^{-2})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|------------------|
| $38.7 \pm 9.0 \pm 1.6$ | ¹ AAIJ | 15D LHCB | pp at 7, 8 TeV |

¹ Uses $J/\psi \rightarrow \mu^+ \mu^-$, $\eta' \rightarrow \rho^0 \gamma$, and $\eta' \rightarrow \eta \pi^+ \pi^-$ decays.

$\Gamma(J/\psi(1S)\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{68}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------|
| $< 4.8 \times 10^{-6}$ | 90 | ¹ AAIJ | 13Z LHCB | pp at 7 TeV |

¹ Uses $B(B_S^0 \rightarrow J/\psi(1S)\pi^+ \pi^-) = (1.98 \pm 0.20) \times 10^{-4}$.

$\Gamma(J/\psi(1S)\pi^+ \pi^- \pi^+ \pi^-)/\Gamma(J/\psi(1S)\pi^+ \pi^-)$ Γ_{69}/Γ_{47}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-----------------|
| $0.371 \pm 0.015 \pm 0.022$ | ¹ AAIJ | 14Y LHCB | pp at 7,8 TeV |

¹ Excludes contributions from $\psi(2S)$ and $X(3872)$ decaying to $J/\psi(1S)\pi^+ \pi^-$.

$\Gamma(J/\psi(1S)f_1(1285))/\Gamma_{\text{total}}$ Γ_{70}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|------|-----------------------|
| $7.1 \pm 1.0^{+0.9}_{-1.0}$ | ¹ AAIJ | 14Y | LHCB pp at 7, 8 TeV |

¹ AAIJ 14Y reports $(7.14 \pm 0.99^{+0.83}_{-0.91} \pm 0.41) \times 10^{-5}$ from a measurement of $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow 2\pi^+2\pi^-)]$ assuming $B(f_1(1285) \rightarrow 2\pi^+2\pi^-) = 0.11^{+0.007}_{-0.006}$.

$\Gamma(\psi(2S)\phi)/\Gamma_{\text{total}}$ Γ_{74}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 1 BUSKULIC 93G ALEP $e^+e^- \rightarrow Z$

$\Gamma(\psi(2S)\phi)/\Gamma(J/\psi(1S)\phi)$ Γ_{74}/Γ_{41}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.501 ± 0.034 OUR AVERAGE

| | | | |
|-----------------------------|---------------------|-----|----------------------------|
| $0.497 \pm 0.034 \pm 0.011$ | ^{1,2} AAIJ | 12L | LHCB pp at 7 TeV |
| $0.53 \pm 0.10 \pm 0.09$ | ABAZOV | 09Y | D0 $p\bar{p}$ at 1.96 TeV |
| $0.52 \pm 0.13 \pm 0.07$ | ABULENCIA | 06N | CDF $p\bar{p}$ at 1.96 TeV |

¹ AAIJ 12L reports $0.489 \pm 0.026 \pm 0.021 \pm 0.012$ from a measurement of $[\Gamma(B_s^0 \rightarrow \psi(2S)\phi)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] \times [B(J/\psi(1S) \rightarrow e^+e^-)] / [B(\psi(2S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$, $B(\psi(2S) \rightarrow e^+e^-) = (7.72 \pm 0.17) \times 10^{-3}$, which we rescale to our best values $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$, $B(\psi(2S) \rightarrow e^+e^-) = (7.89 \pm 0.17) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² Assumes $B(J/\psi \rightarrow \mu^+\mu^-) / B(\psi(2S) \rightarrow \mu^+\mu^-) = B(J/\psi \rightarrow e^+e^-) / B(\psi(2S) \rightarrow e^+e^-) = 7.69 \pm 0.19$.

$\Gamma(\chi_{c1}\phi)/\Gamma(J/\psi(1S)\phi)$ Γ_{75}/Γ_{41}

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

$18.9 \pm 1.8 \pm 1.5$ ¹ AAIJ 13AC LHCB pp at 7 TeV

¹ Uses $B(\chi_{c1} \rightarrow J/\psi\gamma) = (34.4 \pm 1.5)\%$.

$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{73}/Γ_{47}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

$0.34 \pm 0.04 \pm 0.03$ ¹ AAIJ 13AA LHCB pp at 7 TeV

¹ Assuming lepton universality for dimuon decay modes of J/ψ and $\psi(2S)$ mesons, the ratio $B(J/\psi \rightarrow \mu^+\mu^-)/B(\psi(2S) \rightarrow \mu^+\mu^-) = B(J/\psi \rightarrow e^+e^-)/B(\psi(2S) \rightarrow e^+e^-) = 7.69 \pm 0.19$ was used.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{76}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

0.76 ± 0.19 OUR AVERAGE Error includes scale factor of 1.4.

| | | | | |
|---------------------------------|-----------------------|------|------|------------------------|
| $0.98^{+0.23}_{-0.19} \pm 0.07$ | ¹ AAIJ | 12AR | LHCB | pp at 7 TeV |
| $0.60 \pm 0.17 \pm 0.04$ | ² AALTONEN | 12L | CDF | $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|--------------------------|-----|------|-----------------------------------|
| < 12 | 90 | ³ PENG | 10 | BELL | $e^+e^- \rightarrow \Upsilon(5S)$ |
| < 1.2 | 90 | ⁴ AALTONEN | 09C | CDF | Repl. by AALTONEN 12L |
| < 1.7 | 90 | ⁵ ABULENCIA,A | 06D | CDF | Repl. by AALTONEN 09C |
| <232 | 90 | ⁶ ABE | 00C | SLD | $e^+e^- \rightarrow Z$ |
| <170 | 90 | ⁷ BUSKULIC | 96V | ALEP | $e^+e^- \rightarrow Z$ |

¹ AAIJ 12AR reports $[\Gamma(B_s^0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \pi^+\pi^-)] \times [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)] = 0.050^{+0.011}_{-0.009} \pm 0.004$ which we multiply or divide by our best values $B(B^0 \rightarrow \pi^+\pi^-) = (5.12 \pm 0.19) \times 10^{-6}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.260 \pm 0.015$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² AALTONEN 12L reports $[\Gamma(B_s^0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)] = 0.008 \pm 0.002 \pm 0.001$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.260 \pm 0.015$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

³ Uses $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$ and assumes $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.

⁴ Obtains this result from $(f_s/f_d) \cdot B(B_s \rightarrow \pi^+\pi^-)/B(B^0 \rightarrow K^+\pi^-) = 0.007 \pm 0.004 \pm 0.005$, assuming $f_s/f_d = 0.276 \pm 0.034$ and $B(B^0 \rightarrow K^+\pi^-) = (19.4 \pm 0.6) \times 10^{-6}$.

⁵ ABULENCIA,A 06D obtains this from $B(B_s \rightarrow \pi^+\pi^-) / B(B_s \rightarrow K^+K^-) < 0.05$ at 90% CL, assuming $B(B_s \rightarrow K^+K^-) = (33 \pm 6 \pm 7) \times 10^{-6}$.

⁶ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

⁷ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{77}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-----------------------|--------|------------------------|
| <2.1 × 10⁻⁴ | 90 | ¹ ACCIARRI | 95H L3 | $e^+e^- \rightarrow Z$ |

¹ ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

$\Gamma(\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{78}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-----------------------|--------|------------------------|
| <1.0 × 10⁻³ | 90 | ¹ ACCIARRI | 95H L3 | $e^+e^- \rightarrow Z$ |

¹ ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_{79}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-----------------------|--------|------------------------|
| <1.5 × 10⁻³ | 90 | ¹ ACCIARRI | 95H L3 | $e^+e^- \rightarrow Z$ |

¹ ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

$\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$ Γ_{80}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------|---------|------------------------|
| <3.20 × 10⁻⁴ | 90 | ¹ ABE | 00C SLD | $e^+e^- \rightarrow Z$ |
| ¹ ABE 00C assumes $B(Z \rightarrow b\bar{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$. | | | | |

$\Gamma(\phi \rho^0)/\Gamma_{\text{total}}$ Γ_{81}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------|---------|------------------------|
| <6.17 × 10⁻⁴ | 90 | ¹ ABE | 00C SLD | $e^+e^- \rightarrow Z$ |
| ¹ ABE 00C assumes $B(Z \rightarrow b\bar{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$. | | | | |

$\Gamma(\phi \phi)/\Gamma_{\text{total}}$ Γ_{82}/Γ

| VALUE (units 10 ⁻⁶) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-----------------------|----------|------------------------|
| 19.3 ± 2.6 ± 1.6 | | ¹ AALTONEN | 11AN CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 14 ⁺⁶ / ₋₅ ± 6 | | ² ACOSTA | 05J CDF | Repl. by AALTONEN 11AN |
| <1183 | 90 | ³ ABE | 00C SLD | $e^+e^- \rightarrow Z$ |
| ¹ AALTONEN 11AN reports $[\Gamma(B_s^0 \rightarrow \phi \phi)/\Gamma_{\text{total}}] / [B(B_s^0 \rightarrow J/\psi(1S)\phi)] = (1.78 \pm 0.14 \pm 0.20) \times 10^{-2}$ which we multiply by our best value $B(B_s^0 \rightarrow J/\psi(1S)\phi) = (1.08 \pm 0.09) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |
| ² Uses $B(B^0 \rightarrow J/\psi \phi) = (1.38 \pm 0.49) \times 10^{-3}$ and production cross-section ratio of $\sigma(B_s)/\sigma(B^0) = 0.26 \pm 0.04$. | | | | |
| ³ ABE 00C assumes $B(Z \rightarrow b\bar{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$. | | | | |

$\Gamma(\pi^+ K^-)/\Gamma_{\text{total}}$ Γ_{83}/Γ

| VALUE (units 10 ⁻⁶) | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|--------------------------|-----------|---------------------------------|
| 5.5 ± 0.6 OUR AVERAGE | | | | |
| 5.6 ± 0.6 ± 0.4 | | ¹ AAIJ | 12AR LHCb | pp at 7 TeV |
| 5.4 ± 0.9 ± 0.3 | | ² AALTONEN | 09C CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 26 | 90 | ³ PENG | 10 BELL | $e^+e^- \rightarrow \gamma(5S)$ |
| < 5.6 | 90 | ⁴ ABULENCIA,A | 06D CDF | Repl. by AALTONEN 09C |
| <261 | 90 | ⁵ ABE | 00C SLD | $e^+e^- \rightarrow Z$ |
| <210 | 90 | ⁶ BUSKULIC | 96V ALEP | $e^+e^- \rightarrow Z$ |
| <260 | 90 | ⁷ AKERS | 94L OPAL | $e^+e^- \rightarrow Z$ |
| ¹ AAIJ 12AR reports $[\Gamma(B_s^0 \rightarrow \pi^+ K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+ \pi^-)] \times [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)] = 0.074 \pm 0.006 \pm 0.006$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.260 \pm 0.015$. Our first error is their experiment's error and our second error is the systematic error from using our best values. | | | | |
| ² AALTONEN 09C reports $[\Gamma(B_s^0 \rightarrow \pi^+ K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+ \pi^-)] \times [B(\bar{b} \rightarrow B_s^0) / [B(\bar{b} \rightarrow B^0)]] = 0.071 \pm 0.010 \pm 0.007$ which we multiply or divide by our best | | | | |

values $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $B(\bar{b} \rightarrow B_s^0) = (10.5 \pm 0.5) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

³ Uses $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$ and assumes $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1_{-4.0}^{+3.8})\%$.

⁴ ABULENCIA,A 06D obtains this from $(f_s/f_d) (B(B_s \rightarrow \pi^+ K^-) / B(B^0 \rightarrow K^+ \pi^-)) < 0.08$ at 90% CL, assuming $f_s/f_d = 0.260 \pm 0.039$ and $B(B^0 \rightarrow K^+ \pi^-) = (18.9 \pm 0.7) \times 10^{-6}$.

⁵ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7_{-2.2}^{+1.8})\%$ and $f_{B_s} = (10.5_{-2.2}^{+1.8})\%$.

⁶ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

⁷ Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and B_d^0 (B_s^0) fraction 39.5% (12%).

$\Gamma(K^+ K^-) / \Gamma_{\text{total}}$

Γ_{84} / Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-----|-----------------------|-----------|------------------------------------|
| 25.0 ± 1.7 OUR AVERAGE | | | | |
| $23.8 \pm 1.6 \pm 1.5$ | | ¹ AAIJ | 12AR LHCb | pp at 7 TeV |
| $26.1 \pm 2.2 \pm 1.7$ | | ² AALTONEN | 11N CDF | $p\bar{p}$ at 1.96 TeV |
| $38_{-9}^{+10} \pm 7$ | | ³ PENG | 10 BELL | $e^+ e^- \rightarrow \Upsilon(5S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------|----|------------------------------|----------|------------------------------------|
| <310 | 90 | DRUTSKOY | 07A BELL | $e^+ e^- \rightarrow \Upsilon(5S)$ |
| $33 \pm 6 \pm 7$ | | ⁴ ABULENCIA,A 06D | CDF | Repl. by AALTONEN 11N |
| <283 | 90 | ⁵ ABE | 00C SLD | $e^+ e^- \rightarrow Z$ |
| < 59 | 90 | ⁶ BUSKULIC | 96V ALEP | $e^+ e^- \rightarrow Z$ |
| <140 | 90 | ⁷ AKERS | 94L OPAL | $e^+ e^- \rightarrow Z$ |

¹ AAIJ 12AR reports $[\Gamma(B_s^0 \rightarrow K^+ K^-) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+ \pi^-)] \times [\Gamma(\bar{b} \rightarrow B_s^0) / \Gamma(\bar{b} \rightarrow B^0)] = 0.316 \pm 0.009 \pm 0.019$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $\Gamma(\bar{b} \rightarrow B_s^0) / \Gamma(\bar{b} \rightarrow B^0) = 0.260 \pm 0.015$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² AALTONEN 11N reports $(f_s/f_d) (B(B_s^0 \rightarrow K^+ K^-) / B(B^0 \rightarrow K^+ \pi^-)) = 0.347 \pm 0.020 \pm 0.021$. We multiply this result by our best value of $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$ and divide by our best value of f_s/f_d , where $1/2 f_s/f_d = 0.130 \pm 0.008$. Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using our best values.

³ Uses $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$ and assumes $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1_{-4.0}^{+3.8})\%$.

⁴ ABULENCIA,A 06D obtains this from $(f_s/f_d) (B(B_s \rightarrow K^+ K^-) / B(B^0 \rightarrow K^+ \pi^-)) = 0.46 \pm 0.08 \pm 0.07$, assuming $f_s/f_d = 0.260 \pm 0.039$ and $B(B^0 \rightarrow K^+ \pi^-) = (18.9 \pm 0.7) \times 10^{-6}$.

⁵ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7_{-2.2}^{+1.8})\%$ and $f_{B_s} = (10.5_{-2.2}^{+1.8})\%$.

⁶ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

⁷ Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and B_d^0 (B_s^0) fraction 39.5% (12%).

$\Gamma(K^0 \bar{K}^0)/\Gamma_{\text{total}}$ Γ_{85}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------------|---------|------------------------------------|
| <6.6 | 90 | ¹ PENG | 10 BELL | $e^+ e^- \rightarrow \Upsilon(5S)$ |

¹ Uses $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$ and assumes $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.

$\Gamma(K^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{86}/Γ

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|-------------------|-----------|---------------|
| $15 \pm 4 \pm 1$ | ¹ AAIJ | 13BP LHCB | pp at 7 TeV |

¹ AAIJ 13BP reports $[\Gamma(B_s^0 \rightarrow K^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 0.29 \pm 0.06 \pm 0.04$ which we multiply by our best value $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (5.20 \pm 0.24) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^- \pi^+)/\Gamma_{\text{total}}$ Γ_{88}/Γ

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|----------|---------------|
| $3.3 \pm 1.2 \pm 0.3$ | ^{1,2} AAIJ | 14BMLHCB | pp at 7 TeV |

¹ AAIJ 14BM reports $[\Gamma(B_s^0 \rightarrow K^*(892)^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^+ \pi^-)] = 0.39 \pm 0.13 \pm 0.05$ which we multiply by our best value $B(B^0 \rightarrow K^*(892)^+ \pi^-) = (8.4 \pm 0.8) \times 10^{-6}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Uses $f_s/f_d = 0.259 \pm 0.015$.

$\Gamma(K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{87}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|-----------|---------------|
| $7.7 \pm 1.0 \pm 0.4$ | ¹ AAIJ | 13BP LHCB | pp at 7 TeV |

¹ AAIJ 13BP reports $[\Gamma(B_s^0 \rightarrow K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 1.48 \pm 0.12 \pm 0.14$ which we multiply by our best value $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (5.20 \pm 0.24) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{89}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--|---------------------|----------|---------------|
| $1.25 \pm 0.24 \pm 0.11$ | ^{1,2} AAIJ | 14BMLHCB | pp at 7 TeV |

¹ AAIJ 14BM reports $[\Gamma(B_s^0 \rightarrow K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^+ \pi^-)] = 1.49 \pm 0.22 \pm 0.18$ which we multiply by our best value $B(B^0 \rightarrow K^*(892)^+ \pi^-) = (8.4 \pm 0.8) \times 10^{-6}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Uses $f_s/f_d = 0.259 \pm 0.015$.

$\Gamma(K^0 K^+ K^-)/\Gamma_{\text{total}}$ Γ_{90}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------------|-----------|---------------|
| $<3.5 \times 10^{-6}$ | 90 | ¹ AAIJ | 13BP LHCB | pp at 7 TeV |

¹ AAIJ 13BP reports $[\Gamma(B_s^0 \rightarrow K^0 K^+ K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] < 0.068$ which we multiply by our best value $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = 5.20 \times 10^{-5}$.

$\Gamma(\overline{K}^*(892)^0 \rho^0)/\Gamma_{\text{total}}$ Γ_{91}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|------------------|---------|-------------------------|
| $<7.67 \times 10^{-4}$ | 90 | ¹ ABE | 00C SLD | $e^+ e^- \rightarrow Z$ |

¹ ABE 00C assumes $B(Z \rightarrow b\bar{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$.

$\Gamma(\overline{K}^*(892)^0 K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{92}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------------|----------|-------------------------|
| $2.81 \pm 0.46 \pm 0.56$ | | ¹ AAIJ | 12F LHCB | pp at 7 TeV |
| <168.1 | 90 | ² ABE | 00C SLD | $e^+ e^- \rightarrow Z$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Uses $B^0 \rightarrow J/\psi K^{*0}$ for normalization and assumes $B(B^0 \rightarrow J/\psi K^{*0}) B(J/\psi \rightarrow \mu^+ \mu^-) B(K^{*0} \rightarrow K^+ \pi^-) = (1.33 \pm 0.06) \times 10^{-3}$ and $f_s/f_d = 0.253 \pm 0.031$. The second quoted error is total uncertainty including the error of 0.34 on f_s/f_d .

² ABE 00C assumes $B(Z \rightarrow b\bar{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$.

$\Gamma(\phi K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{93}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------------|-----------|-------------------------|
| $1.13 \pm 0.29 \pm 0.06$ | | ¹ AAIJ | 13BW LHCB | pp at 7 TeV |
| <1013 | 90 | ² ABE | 00C SLD | $e^+ e^- \rightarrow Z$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ AAIJ 13BW reports $[\Gamma(B_s^0 \rightarrow \phi K^*(892)^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^0 \phi)] = 0.113 \pm 0.024 \pm 0.016$ which we multiply by our best value $B(B^0 \rightarrow K^*(892)^0 \phi) = (1.00 \pm 0.05) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABE 00C assumes $B(Z \rightarrow b\bar{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$.

$\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{94}/Γ

Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE (units 10^{-8}) | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-----------------------|-----------|-------------------------|
| $2.84^{+2.03+0.85}_{-1.68-0.18}$ | | ¹ AAIJ | 13BQ LHCB | pp at 7 TeV |
| <5900 | 90 | ² BUSKULIC | 96V ALEP | $e^+ e^- \rightarrow Z$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Uses normalization mode $B(B^0 \rightarrow K^+ \pi^-) = (19.55 \pm 0.54) \times 10^{-6}$ and B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$.

² BUSKULIC 96V assumes PDG 96 production fractions for B^0, B^+, B_s, b baryons.

$\Gamma(\Lambda_c^- \Lambda \pi^+)/\Gamma_{\text{total}}$ Γ_{95}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|---------------------------|------|----------------------------------|
| $3.6 \pm 1.1 \pm 1.2$ | ¹ SOLOVIEVA 13 | BELL | $e^+ e^- \rightarrow \gamma(4S)$ |

¹ The second error is the total systematic uncertainty including the Λ_c absolute branching fractions and the normalization number of B_s events.

$\Gamma(\Lambda_c^- \Lambda_c^+)/\Gamma_{\text{total}}$ Γ_{96}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------|-----------|---------------|
| $< 8.0 \times 10^{-5}$ | 95 | 1 AAIJ | 14AA LHCb | pp at 7 TeV |

¹ Uses $B(\bar{B}^0 \rightarrow D^+ D_s^-) = (7.2 \pm 0.8) \times 10^{-3}$.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{97}/Γ

Test for $\Delta B=1$ weak neutral current.

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|----------|----------------------------------|
| < 3.1 | 90 | 1 DUTTA | 15 BELL | $e^+ e^- \rightarrow \gamma(5S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 8.7 | 90 | 2 WICHT | 08A BELL | Repl. by DUTTA 15 |
| < 53 | 90 | DRUTSKOY | 07A BELL | Repl. by WICHT 08A |
| < 148 | 90 | 3 ACCIARRI | 95I L3 | $e^+ e^- \rightarrow Z$ |

¹ Assumes the fraction of $B_s^{(*)} \bar{B}_s^{(*)}$ in $b\bar{b}$ events is $f_s = (17.2 \pm 3.0)\%$.
² Assumes $\gamma(5S) \rightarrow B_s^* \bar{B}_s^* = (19.5^{+3.0}_{-2.3})\%$.
³ ACCIARRI 95I assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = (12.0 \pm 3.0)\%$.

$\Gamma(\phi\gamma)/\Gamma_{\text{total}}$ Γ_{98}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|-----------|----------------------------------|
| 35.2 ± 3.4 OUR AVERAGE | | | | |
| $36 \pm 5 \pm 7$ | | 1 DUTTA | 15 BELL | $e^+ e^- \rightarrow \gamma(5S)$ |
| $35.1 \pm 3.5 \pm 1.2$ | | 2 AAIJ | 13 LHCb | pp at 7 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 39 ± 5 | | 3 AAIJ | 12AE LHCb | Repl. by AAIJ 13 |
| $57^{+18}_{-15} \quad ^{+12}_{-11}$ | | 4 WICHT | 08A BELL | Repl. by DUTTA 15 |
| < 390 | 90 | DRUTSKOY | 07A BELL | $e^+ e^- \rightarrow \gamma(5S)$ |
| < 120 | 90 | ACOSTA | 02G CDF | $p\bar{p}$ at 1.8 TeV |
| < 700 | 90 | 5 ADAM | 96D DLPH | $e^+ e^- \rightarrow Z$ |

¹ Assumes the fraction of $B_s^{(*)} \bar{B}_s^{(*)}$ in $b\bar{b}$ events is $f_s = (17.2 \pm 3.0)\%$. The systematic uncertainty from f_s is 0.6×10^{-5} .
² AAIJ 13 reports $[\Gamma(B_s^0 \rightarrow \phi\gamma)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^0\gamma)] = 0.81 \pm 0.04 \pm 0.07$ which we multiply by our best value $B(B^0 \rightarrow K^*(892)^0\gamma) = (4.33 \pm 0.15) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
³ Measures $B(B^0 \rightarrow K^{*0}\gamma)/B(B_s \rightarrow \phi\gamma) = 1.12 \pm 0.08(\text{stat})^{+0.06}_{-0.04}(\text{sys})^{+0.09}_{-0.08}(f_s/f_d)$ and uses current world-average value of $B(B^0 \rightarrow K^{*0}\gamma) = (4.33 \pm 0.15) \times 10^{-5}$.
⁴ Assumes $\gamma(5S) \rightarrow B_s^* \bar{B}_s^* = (19.5^{+3.0}_{-2.3})\%$.
⁵ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{99}/Γ
 Test for $\Delta B = 1$ weak neutral current.

VALUE (units 10^{-9}) CL% DOCUMENT ID TECN COMMENT

3.1 ± 0.7 OUR AVERAGE

| | | | | | |
|---|----|---------------|------|------|---------------------------|
| $2.9^{+1.1+0.3}_{-1.0-0.1}$ | 1 | AAIJ | 13BA | LHCB | $\rho\rho$ at 7, 8 TeV |
| 13^{+9}_{-7} | 2 | AALTONEN | 13F | CDF | $\rho\bar{p}$ at 1.96 TeV |
| $3.0^{+1.0}_{-0.9}$ | 3 | CHATRCHYAN | 13AW | CMS | $\rho\rho$ at 7, 8 TeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| $3.2^{+1.4+0.5}_{-1.2-0.3}$ | 4 | AAIJ | 13B | LHCB | Repl. by AAIJ 13BA |
| < 12 | 90 | 5 ABAZOV | 13C | D0 | $\rho\bar{p}$ at 1.96 TeV |
| < 19 | 90 | 6 AAD | 12AE | ATLS | $\rho\rho$ at 7 TeV |
| < 12 | 90 | 7 AAIJ | 12A | LHCB | Repl. by AAIJ 12W |
| < 3.8 | 90 | 8 AAIJ | 12W | LHCB | Repl. by AAIJ 13B |
| < 6.4 | 90 | 9 CHATRCHYAN | 12A | CMS | $\rho\rho$ at 7 TeV |
| < 43 | 90 | 10 AAIJ | 11B | LHCB | Repl. by AAIJ 12A |
| < 35 | 90 | 11 AALTONEN | 11AG | CDF | $\rho\bar{p}$ at 1.96 TeV |
| < 16 | 90 | 12 CHATRCHYAN | 11T | CMS | Repl. by CHATRCHYAN 12A |
| < 42 | 90 | 13 ABAZOV | 10S | D0 | $\rho\bar{p}$ at 1.96 TeV |
| < 47 | 90 | 13 AALTONEN | 08I | CDF | Repl. by AALTONEN 11AG |
| < 94 | 90 | 14 ABAZOV | 07Q | D0 | Repl. by ABAZOV 10S |
| < 410 | 90 | 15 ABAZOV | 05E | D0 | $\rho\bar{p}$ at 1.96 TeV |
| < 150 | 90 | 16 ABULENCIA | 05 | CDF | $\rho\bar{p}$ at 1.96 TeV |
| < 580 | 90 | 17 ACOSTA | 04D | CDF | $\rho\bar{p}$ at 1.96 TeV |
| < 2000 | 90 | 18 ABE | 98 | CDF | $\rho\bar{p}$ at 1.8 TeV |
| < 38000 | 90 | 19 ACCIARRI | 97B | L3 | $e^+e^- \rightarrow Z$ |
| < 8400 | 90 | 20 ABE | 96L | CDF | Repl. by ABE 98 |

¹ Uses B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.259 \pm 0.015$ and normalization modes $B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+$ and $B^0 \rightarrow K^+ \pi^-$.

² Uses normalization mode $B(B^+ \rightarrow J/\psi K^+) = (10.22 \pm 0.35) \times 10^{-4}$ and B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.28 \pm 0.04$.

³ Uses B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$ for normalization.

⁴ Uses B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$ and two normalization modes: $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$ and $B(B^0 \rightarrow K^+ \pi^-) = (1.94 \pm 0.06) \times 10^{-5}$.

⁵ Uses normalization mode $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$ and B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.263 \pm 0.017$.

⁶ Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.75 \pm 0.29$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$.

⁷ Uses B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.267^{+0.021}_{-0.020}$ and three normalization modes $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$, $B(B^0 \rightarrow K^+ \pi^-) = (1.94 \pm 0.06) \times 10^{-5}$, and $B(B_s^0 \rightarrow J/\psi \phi \rightarrow \mu^+ \mu^- K^+ K^-) = (3.4 \pm 0.9) \times 10^{-5}$.

⁸ Uses B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.267^{+0.021}_{-0.020}$ and three normalization modes of $B^+ \rightarrow J/\psi K^+$, $B^0 \rightarrow K^+ \pi^-$, and $B_s^0 \rightarrow J/\psi \phi$.

⁹ Uses $f_s/f_u = 0.267 \pm 0.021$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$.

- ¹⁰ Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_S^0) = 3.71 \pm 0.47$ and three normalization modes.
- ¹¹ Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_S^0) = 3.55 \pm 0.47$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$.
- ¹² Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_S^0) = 3.55 \pm 0.42$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$.
- ¹³ Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_S^0) = 3.86 \pm 0.59$, and the number of $B^+ \rightarrow J/\psi K^+$ decays.
- ¹⁴ Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_S^0) = 3.86 \pm 0.54$ and the number of $B^+ \rightarrow J/\psi K^+$ decays.
- ¹⁵ Assumes production cross-section $\sigma(B_S)/\sigma(B^+) = 0.270 \pm 0.034$.
- ¹⁶ Assumes production cross section $\sigma(B^+)/\sigma(B_S) = 3.71 \pm 0.41$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (5.88 \pm 0.26) \times 10^{-5}$.
- ¹⁷ Assumes production cross-section $\sigma(B_S)/\sigma(B^+) = 0.100/0.391$ and the CDF measured value of $\sigma(B^+) = 3.6 \pm 0.6 \mu\text{b}$.
- ¹⁸ ABE 98 assumes production of $\sigma(B^0) = \sigma(B^+)$ and $\sigma(B_S)/\sigma(B^0) = 1/3$. They normalize to their measured $\sigma(B^0, p_T(B) > 6, |y| < 1.0) = 2.39 \pm 0.32 \pm 0.44 \mu\text{b}$.
- ¹⁹ ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_S , and Λ_b .
- ²⁰ ABE 96L assumes B^+/B_S production ratio 3/1. They normalize to their measured $\sigma(B^+, p_T(B) > 6 \text{ GeV}/c, |y| < 1) = 2.39 \pm 0.54 \mu\text{b}$.

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{100}/Γ**

Test for $\Delta B = 1$ weak neutral current.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------|---------|------------------------|
| $<2.8 \times 10^{-7}$ | 90 | AALTONEN | 09P CDF | $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----|-----------------------|--------|-------------------------|
| $<5.4 \times 10^{-5}$ | 90 | ¹ ACCIARRI | 97B L3 | $e^+ e^- \rightarrow Z$ |
|-----------------------|----|-----------------------|--------|-------------------------|

¹ ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_S , and Λ_b .

$\Gamma(\mu^+ \mu^- \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{101}/Γ**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------------|-----------|---------------|
| $<1.2 \times 10^{-8}$ | 90 | ¹ AAIJ | 13AW LHCB | pp at 7 TeV |

¹ Also reports a limit of $<1.6 \times 10^{-8}$ at 95% CL.

$\Gamma(SP, S \rightarrow \mu^+ \mu^-, P \rightarrow \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{102}/Γ**

Here S and P are the hypothetical scalar and pseudoscalar particles with masses of $2.5 \text{ GeV}/c^2$ and $214.3 \text{ MeV}/c^2$, respectively.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------------|-----------|---------------|
| $<1.2 \times 10^{-8}$ | 90 | ¹ AAIJ | 13AW LHCB | pp at 7 TeV |

¹ Also reports a limit of $<1.6 \times 10^{-8}$ at 95% CL.

$\Gamma(\phi(1020)\mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{103}/Γ**

Test for $\Delta B = 1$ weak neutral current.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|---------------------|---------|------------------------|
| $<3.2 \times 10^{-6}$ | 90 | ¹ ABAZOV | 06G D0 | $p\bar{p}$ at 1.96 TeV |
| $<4.7 \times 10^{-5}$ | 90 | ACOSTA | 02D CDF | $p\bar{p}$ at 1.8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----|---------------------|---------|------------------------|
| $<3.2 \times 10^{-6}$ | 90 | ¹ ABAZOV | 06G D0 | $p\bar{p}$ at 1.96 TeV |
| $<4.7 \times 10^{-5}$ | 90 | ACOSTA | 02D CDF | $p\bar{p}$ at 1.8 TeV |

¹ Uses $B(B_S^0 \rightarrow J/\psi \phi) = 9.3 \times 10^{-4}$.

$\Gamma(\phi(1020)\mu^+\mu^-)/\Gamma(J/\psi(1S)\phi)$ Γ_{103}/Γ_{41}

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|--|------|----------------------------|
| 0.71 ± 0.13 | | OUR AVERAGE Error includes scale factor of 2.2. | | |
| 0.674 ^{+0.061} _{-0.056} ± 0.016 | | AAIJ | 13X | LHCB pp at 7 TeV |
| 1.13 ± 0.19 ± 0.07 | | AALTONEN | 11A1 | CDF $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.11 ± 0.25 ± 0.09 | | AALTONEN | 11L | CDF Repl. by AALTONEN 11A1 |
| < 2.3 | 90 | AALTONEN | 09B | CDF Repl. by AALTONEN 11L |

$\Gamma(\phi\nu\bar{\nu})/\Gamma_{\text{total}}$ Γ_{104}/Γ

Test for $\Delta B = 1$ weak neutral current.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------|------|-----------------------------|
| < 5.4 × 10⁻³ | 90 | ¹ ADAM | 96D | DLPH $e^+e^- \rightarrow Z$ |
| ¹ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. | | | | |

$\Gamma(e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{105}/Γ

Test of lepton family number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-----------------------|----------|----------------------------|
| < 1.1 × 10⁻⁸ | 90 | ¹ AAIJ | 13BMLHCB | pp at 7 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 2.0 × 10 ⁻⁷ | 90 | AALTONEN | 09P | CDF $p\bar{p}$ at 1.96 TeV |
| < 6.1 × 10 ⁻⁶ | 90 | ABE | 98V | CDF Repl. by AALTONEN 09P |
| < 4.1 × 10 ⁻⁵ | 90 | ² ACCIARRI | 97B | L3 $e^+e^- \rightarrow Z$ |
| ¹ Uses normalization mode $B(B^0 \rightarrow K^+\pi^-) = (19.4 \pm 0.6) \times 10^{-6}$ and B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$. | | | | |
| ² ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_s , and Λ_b . | | | | |

POLARIZATION IN B_s^0 DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal (L), or both are transverse and parallel (\parallel), or perpendicular (\perp) to each other with the parameters Γ_L/Γ , Γ_\perp/Γ , and the relative phases ϕ_\parallel and ϕ_\perp . See the definitions in the note on “Polarization in B Decays” review in the B^0 Particle Listings.

Γ_L/Γ in $B_s^0 \rightarrow D_s^*\rho^+$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--------------------------------------|
| 1.05^{+0.08+0.03}_{-0.10-0.04} | LOUVOT | 10 | BELL $e^+e^- \rightarrow \gamma(5S)$ |

Γ_L/Γ in $B_s^0 \rightarrow J/\psi(1S)\phi$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|-----------------------------|
| 0.528 ± 0.006 OUR AVERAGE | | | |
| 0.5241 ± 0.0034 ± 0.0067 | AAIJ | 15I | LHCB $p\bar{p}$ at 7, 8 TeV |
| 0.529 ± 0.006 ± 0.012 | ¹ AAD | 14U | ATLS $p\bar{p}$ at 7 TeV |
| 0.524 ± 0.013 ± 0.015 | ² AALTONEN | 12D | CDF $p\bar{p}$ at 1.96 TeV |
| 0.558 ^{+0.017} / _{-0.019} | ^{2,3} ABAZOV | 12D | D0 $p\bar{p}$ at 1.96 TeV |
| 0.61 ± 0.14 ± 0.02 | ⁴ AFFOLDER | 00N | CDF $p\bar{p}$ at 1.8 TeV |
| 0.56 ± 0.21 ^{+0.02} / _{-0.04} | ABE | 95Z | CDF $p\bar{p}$ at 1.8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----------------------|-----------------------|------|---------------------------|
| 0.539 ± 0.014 ± 0.016 | ² AAD | 12CV | ATLS Repl. by AAD 14U |
| 0.555 ± 0.027 ± 0.006 | ⁵ ABAZOV | 09E | D0 Repl. by ABAZOV 12D |
| 0.531 ± 0.020 ± 0.007 | ² AALTONEN | 08J | CDF Repl. by AALTONEN 12D |
| 0.62 ± 0.06 ± 0.01 | ACOSTA | 05 | CDF Repl. by AALTONEN 08J |

¹ Measured using the flavor tagged, time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

² Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

³ The error includes both statistical and systematic uncertainties.

⁴ AFFOLDER 00N measurements are based on 40 B_s^0 candidates obtained from a data sample of 89 pb⁻¹. The P -wave fraction is found to be $0.23 \pm 0.19 \pm 0.04$.

⁵ Measured the angular and lifetime parameters for the time-dependent angular untagged decays $B_d^0 \rightarrow J/\psi K^{*0}$ and $B_s^0 \rightarrow J/\psi\phi$.

Γ_L/Γ in $B_s^0 \rightarrow D_s^{*+} D_s^{*-}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|--|
| 0.06 ^{+0.18}/_{-0.17} ± 0.03 | ESEN | 13 | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |

$\Gamma_{\parallel}/\Gamma$ in $B_s^0 \rightarrow J/\psi(1S)\phi$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|----------------------------|
| 0.224 ± 0.010 OUR AVERAGE | | | |
| 0.220 ± 0.008 ± 0.009 | ¹ AAD | 14U | ATLS $p\bar{p}$ at 7 TeV |
| 0.231 ± 0.014 ± 0.015 | ² AALTONEN | 12D | CDF $p\bar{p}$ at 1.96 TeV |
| 0.231 ^{+0.024} / _{-0.030} | ^{2,3} ABAZOV | 12D | D0 $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----------------------|-----------------------|------|---------------------------|
| 0.224 ± 0.010 ± 0.009 | ² AAD | 12CV | ATLS Repl. by AAD 14U |
| 0.244 ± 0.032 ± 0.014 | ⁴ ABAZOV | 09E | D0 Repl. by ABAZOV 12D |
| 0.230 ± 0.029 ± 0.011 | ² AALTONEN | 08J | CDF Repl. by AALTONEN 12D |
| 0.260 ± 0.084 ± 0.013 | ACOSTA | 05 | CDF Repl. by AALTONEN 08J |

¹ Measured using a tagged, time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

² Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

³ The error includes both statistical and systematic uncertainties.

⁴ Measured the angular and lifetime parameters for the time-dependent angular untagged decays $B_d^0 \rightarrow J/\psi K^{*0}$ and $B_s^0 \rightarrow J/\psi\phi$.

Γ_{\perp}/Γ in $B_s^0 \rightarrow J/\psi(1S)\phi$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------|------|------------------|
| $0.2504 \pm 0.0049 \pm 0.0036$ | AAIJ 15I | LHCB | pp at 7, 8 TeV |

ϕ_{\parallel} in $B_s^0 \rightarrow J/\psi(1S)\phi$

| VALUE (rad) | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|---------|
|-------------|-------------|------|---------|

$3.23^{+0.10}_{-0.14}$ OUR AVERAGE

| | | | |
|----------------------------------|----------|------|------------------|
| $3.26^{+0.10+0.06}_{-0.17-0.07}$ | AAIJ 15I | LHCB | pp at 7, 8 TeV |
|----------------------------------|----------|------|------------------|

| | | | |
|-----------------|-------------------------|----|------------------------|
| 3.15 ± 0.22 | ¹ ABAZOV 12D | D0 | $p\bar{p}$ at 1.96 TeV |
|-----------------|-------------------------|----|------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------------------------------|------------|----|---------------------|
| $2.72^{+1.12}_{-0.27} \pm 0.26$ | ABAZOV 09E | D0 | Repl. by ABAZOV 12D |
|---------------------------------|------------|----|---------------------|

¹ The error includes both statistical and systematic uncertainties.

ϕ_{\perp} in $B_s^0 \rightarrow J/\psi(1S)\phi$

| VALUE (rad) | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|---------|
|-------------|-------------|------|---------|

3.16 ± 0.24 OUR AVERAGE Error includes scale factor of 1.6.

| | | | |
|---------------------------------|----------|------|------------------|
| $3.08^{+0.14}_{-0.15} \pm 0.06$ | AAIJ 15I | LHCB | pp at 7, 8 TeV |
|---------------------------------|----------|------|------------------|

| | | | |
|--------------------------|----------------------|------|---------------|
| $3.89 \pm 0.47 \pm 0.11$ | ¹ AAD 14U | ATLS | pp at 7 TeV |
|--------------------------|----------------------|------|---------------|

¹ Measured using a tagged, time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

Γ_L/Γ for $B_s^0 \rightarrow J/\psi(1S)K^*(892)^0$

Longitudinal polarization fraction, equals to f_L using notation of "Polarization in B decays" review.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------------------------|------|---------------|
| $0.50 \pm 0.08 \pm 0.02$ | ¹ AAIJ 12AP | LHCB | pp at 7 TeV |

¹ The non-resonant $K\pi$ background contributions are subtracted. Also reports an S -wave amplitude $|A_S|^2 = 0.07^{+0.15}_{-0.07}$.

$\Gamma_{\parallel}/\Gamma$ for $B_s^0 \rightarrow J/\psi(1S)K^*(892)^0$

Parallel polarization fraction, equals to $1 - f_L - f_{\perp}$ using notation of "Polarization in B decays" review.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------------------------|------|---------------|
| $0.19^{+0.10}_{-0.08} \pm 0.02$ | ¹ AAIJ 12AP | LHCB | pp at 7 TeV |

¹ The non-resonant $K\pi$ background contributions are subtracted. Also reports an S -wave amplitude $|A_S|^2 = 0.07^{+0.15}_{-0.07}$.

Γ_L/Γ in $B_s^0 \rightarrow \phi\phi$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.362 ± 0.014 OUR AVERAGE

| | | | |
|-----------------------------|-----------|------|------------------|
| $0.364 \pm 0.012 \pm 0.009$ | AAIJ 14AE | LHCB | pp at 7, 8 TeV |
|-----------------------------|-----------|------|------------------|

| | | | |
|-----------------------------|---------------|-----|------------------------|
| $0.348 \pm 0.041 \pm 0.021$ | AALTONEN 11AN | CDF | $p\bar{p}$ at 1.96 TeV |
|-----------------------------|---------------|-----|------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----------------------------|----------|------|--------------------|
| $0.365 \pm 0.022 \pm 0.012$ | AAIJ 12P | LHCB | Repl. by AAIJ 14AE |
|-----------------------------|----------|------|--------------------|

Γ_{\perp}/Γ in $B_s^0 \rightarrow \phi\phi$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------------------|-------------|------------------------|
| 0.309 ± 0.015 OUR AVERAGE | Error includes scale factor of 1.1. | | |
| $0.305 \pm 0.013 \pm 0.005$ | AAIJ | 14AE LHCB | pp at 7, 8 TeV |
| $0.365 \pm 0.044 \pm 0.027$ | AALTONEN | 11AN CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $0.291 \pm 0.024 \pm 0.010$ | AAIJ | 12P LHCB | Repl. by AAIJ 14AE |

ϕ_{\parallel} in $B_s^0 \rightarrow \phi\phi$

| <u>VALUE (rad)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|------------------------|
| 2.55 ± 0.11 OUR AVERAGE | | | |
| $2.54 \pm 0.07 \pm 0.09$ | ¹ AAIJ | 14AE LHCB | pp at 7, 8 TeV |
| $2.71^{+0.31}_{-0.36} \pm 0.22$ | ² AALTONEN | 11AN CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $2.57 \pm 0.15 \pm 0.06$ | ³ AAIJ | 12P LHCB | Repl. by AAIJ 14AE |

¹ AAIJ 14AE reports measurement of ϕ_{\perp} and $\phi_{\perp} - \phi_{\parallel}$, which we convert into ϕ_{\parallel} . Statistical uncertainty includes correlation between measured parameters, while systematic uncertainties are assumed uncorrelated.

² AALTONEN 11AN quotes $\cos\phi_{\parallel} = -0.91^{+0.15}_{-0.13} \pm 0.09$ which we convert to ϕ_{\parallel} taking the smaller solution.

³ AAIJ 12P quotes $\cos\phi_{\parallel} = -0.844 \pm 0.068 \pm 0.029$ which we convert to ϕ_{\parallel} , taking the smaller solution.

ϕ_{\perp} in $B_s^0 \rightarrow \phi\phi$

| <u>VALUE (rad)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------|
| $2.67 \pm 0.23 \pm 0.07$ | AAIJ | 14AE LHCB | pp at 7, 8 TeV |

Γ_L/Γ in $B_s^0 \rightarrow K^{*0}\bar{K}^{*0}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|----------------|
| $0.31 \pm 0.12 \pm 0.04$ | AAIJ | 12F LHCB | pp at 7 TeV |

Γ_{\perp}/Γ in $B_s^0 \rightarrow K^{*0}\bar{K}^{*0}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|----------------|
| $0.38 \pm 0.11 \pm 0.04$ | AAIJ | 12F LHCB | pp at 7 TeV |

Γ_L/Γ in $B_s^0 \rightarrow \phi\bar{K}^{*0}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|----------------|
| $0.51 \pm 0.15 \pm 0.07$ | AAIJ | 13BW LHCB | pp at 7 TeV |

$\Gamma_{\parallel}/\Gamma$ in $B_s^0 \rightarrow \phi\bar{K}^{*0}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|----------------|
| $0.21 \pm 0.11 \pm 0.02$ | AAIJ | 13BW LHCB | pp at 7 TeV |

ϕ_{\parallel} in $B_s^0 \rightarrow \phi\bar{K}^{*0}$

| <u>VALUE (rad)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|----------------|
| $1.75 \pm 0.53 \pm 0.29$ | ¹ AAIJ | 13BW LHCB | pp at 7 TeV |

¹ Measures $\cos(\phi_{\parallel}) = -0.18 \pm 0.52 \pm 0.29$, which we convert to ϕ_{\parallel} by taking the smaller solution.

$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($0.10 < q^2 < 2.00 \text{ GeV}^2/c^4$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|------|--------------------|
| $0.37^{+0.19}_{-0.17} \pm 0.07$ | AAIJ | 13X | LHCB pp at 7 TeV |

$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($2.00 < q^2 < 4.30 \text{ GeV}^2/c^4$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|------|--------------------|
| $0.53^{+0.25}_{-0.23} \pm 0.10$ | AAIJ | 13X | LHCB pp at 7 TeV |

$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($4.30 < q^2 < 8.68 \text{ GeV}^2/c^4$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|------|--------------------|
| $0.81^{+0.11}_{-0.13} \pm 0.05$ | AAIJ | 13X | LHCB pp at 7 TeV |

$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($10.09 < q^2 < 12.90 \text{ GeV}^2/c^4$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|------|--------------------|
| $0.33^{+0.14}_{-0.12} \pm 0.06$ | AAIJ | 13X | LHCB pp at 7 TeV |

$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($14.18 < q^2 < 16.00 \text{ GeV}^2/c^4$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|------|--------------------|
| $0.34^{+0.18}_{-0.17} \pm 0.07$ | AAIJ | 13X | LHCB pp at 7 TeV |

$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($16.00 < q^2 < 19.00 \text{ GeV}^2/c^4$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|------|--------------------|
| $0.16^{+0.17}_{-0.10} \pm 0.07$ | AAIJ | 13X | LHCB pp at 7 TeV |

$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($1.00 < q^2 < 6.00 \text{ GeV}^2/c^4$)

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|------|--------------------|
| $0.56^{+0.17}_{-0.16} \pm 0.09$ | AAIJ | 13X | LHCB pp at 7 TeV |

$B_s^0-\bar{B}_s^0$ MIXING

For a discussion of $B_s^0-\bar{B}_s^0$ mixing see the note on “ $B^0-\bar{B}^0$ Mixing” in the B^0 Particle Listings above.

χ_s is a measure of the time-integrated $B_s^0-\bar{B}_s^0$ mixing probability that produced $B_s^0(\bar{B}_s^0)$ decays as a $\bar{B}_s^0(B_s^0)$. Mixing violates $\Delta B \neq 2$ rule.

$$\chi_s = \frac{x_s^2}{2(1+x_s^2)}$$

$$x_s = \frac{\Delta m_{B_s^0}}{\Gamma_{B_s^0}} = (m_{B_{sH}^0} - m_{B_{sL}^0}) \tau_{B_s^0},$$

where H, L stand for heavy and light states of two B_s^0 CP eigenstates and

$$\tau_{B_s^0} = \frac{1}{0.5(\Gamma_{B_{sH}^0} + \Gamma_{B_{sL}^0})}.$$

$$\Delta m_{B_s^0} = m_{B_{sH}^0} - m_{B_{sL}^0}$$

$\Delta m_{B_s^0}$ is a measure of 2π times the $B_s^0-\bar{B}_s^0$ oscillation frequency in time-dependent mixing experiments.

“OUR EVALUATION” is provided by the Heavy Flavor Averaging Group (HFAG) by taking into account correlations between measurements.

| VALUE (10^{12} h s^{-1}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------|-----------|--------------------------|
| 17.757±0.021 OUR EVALUATION | | | | |
| 17.756±0.021 OUR AVERAGE | | | | |
| 17.711 ^{+0.055} _{-0.057} ±0.011 | | 1 AAIJ | 15l LHCb | pp at 7, 8 TeV |
| 17.768±0.023±0.006 | | 2 AAIJ | 13Bl LHCb | pp at 7 TeV |
| 17.93 ±0.22 ±0.15 | | 3 AAIJ | 13CF LHCb | pp at 7 TeV |
| 17.63 ±0.11 ±0.02 | | 4 AAIJ | 12l LHCb | pp at 7 TeV |
| 17.77 ±0.10 ±0.07 | | 5 ABULENCIA,A 06G | CDF | $p\bar{p}$ at 1.96 TeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 17–21 | 90 | 6 ABAZOV | 06B D0 | $p\bar{p}$ at 1.96 TeV |
| 17.31 ^{+0.33} _{-0.18} ±0.07 | | 7 ABULENCIA | 06Q CDF | Repl. by ABULENCIA,A 06G |
| > 8.0 | 95 | 8 ABDALLAH | 04J DLPH | $e^+e^- \rightarrow Z^0$ |
| > 4.9 | 95 | 9 ABDALLAH | 04J DLPH | $e^+e^- \rightarrow Z^0$ |
| > 8.5 | 95 | 10 ABDALLAH | 04J DLPH | $e^+e^- \rightarrow Z^0$ |
| > 5.0 | 95 | 11 ABDALLAH | 03B DLPH | $e^+e^- \rightarrow Z$ |
| >10.3 | 95 | 12 ABE | 03 SLD | $e^+e^- \rightarrow Z$ |
| >10.9 | 95 | 13 HEISTER | 03E ALEP | $e^+e^- \rightarrow Z$ |
| > 5.3 | 95 | 14 ABE | 02V SLD | $e^+e^- \rightarrow Z$ |
| > 1.0 | 95 | 15 ABBIENDI | 01D OPAL | $e^+e^- \rightarrow Z$ |
| > 7.4 | 95 | 16 ABREU | 00Y DLPH | Repl. by ABDALLAH 04J |
| > 4.0 | 95 | 17 ABREU,P | 00G DLPH | $e^+e^- \rightarrow Z$ |
| > 5.2 | 95 | 18 ABBIENDI | 99S OPAL | $e^+e^- \rightarrow Z$ |
| <96 | 95 | 19 ABE | 99D CDF | $p\bar{p}$ at 1.8 TeV |
| > 5.8 | 95 | 20 ABE | 99J CDF | $p\bar{p}$ at 1.8 TeV |
| > 9.6 | 95 | 21 BARATE | 99J ALEP | $e^+e^- \rightarrow Z$ |
| > 7.9 | 95 | 22 BARATE | 98C ALEP | Repl. by BARATE 99J |
| > 3.1 | 95 | 23 ACKERSTAFF | 97U OPAL | Repl. by ABBIENDI 99S |
| > 2.2 | 95 | 24 ACKERSTAFF | 97V OPAL | Repl. by ABBIENDI 99S |
| > 6.5 | 95 | 25 ADAM | 97 DLPH | Repl. by ABREU 00Y |
| > 6.6 | 95 | 26 BUSKULIC | 96M ALEP | Repl. by BARATE 98C |
| > 2.2 | 95 | 24 AKERS | 95J OPAL | Sup. by ACKERSTAFF 97V |
| > 5.7 | 95 | 27 BUSKULIC | 95J ALEP | $e^+e^- \rightarrow Z$ |
| > 1.8 | 95 | 24 BUSKULIC | 94B ALEP | $e^+e^- \rightarrow Z$ |

¹ Measured using time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

² Measured using $B_s^0 \rightarrow D_s^- \pi^+$ decays.

³ Measured using $B_s^0 \rightarrow D_s^- \mu^+ \nu_\mu X$ decays.

⁴ Measured using $B_s^0 \rightarrow D_s^- \pi^+$ and $D_s^- \pi^+ \pi^- \pi^+$ decays.

⁵ Significance of oscillation signal is 5.4σ . Also reports $|V_{td} / V_{ts}| = 0.2060 \pm 0.0007^{+0.0081}_{-0.0060}$.

- ⁶ A likelihood scan over the oscillation frequency, Δm_s , gives a most probable value of 19 ps^{-1} and a range of $17 < \Delta m_s < 21 \text{ (ps}^{-1})$ at 90% C.L. assuming Gaussian uncertainties. Also excludes $\Delta m_s < 14.8 \text{ ps}^{-1}$ at 95% C.L.
- ⁷ Significance of oscillation signal is 0.2%. Also reported the value $|V_{td} / V_{ts}| = 0.208^{+0.001+0.008}_{-0.002-0.006}$.
- ⁸ Uses leptons emitted with large momentum transverse to a jet and improved techniques for vertexing and flavor-tagging.
- ⁹ Updates of D_s -lepton analysis.
- ¹⁰ Combined results from all Delphi analyses.
- ¹¹ Events with a high transverse momentum lepton were removed and an inclusively reconstructed vertex was required.
- ¹² ABE 03 uses the novel "charge dipole" technique to reconstruct separate secondary and tertiary vertices originating from the $B \rightarrow D$ decay chain. The analysis excludes $\Delta m_s < 4.9 \text{ ps}^{-1}$ and $7.9 < \Delta m_s < 10.3 \text{ ps}^{-1}$.
- ¹³ Three analyses based on complementary event selections: (1) fully-reconstructed hadronic decays; (2) semileptonic decays with D_s exclusively reconstructed; (3) inclusive semileptonic decays.
- ¹⁴ ABE 02V uses exclusively reconstructed D_s^- mesons and excludes $\Delta m_s < 1.4 \text{ ps}^{-1}$ and $2.4 < \Delta m_s < 5.3 \text{ ps}^{-1}$ at 95%CL.
- ¹⁵ Uses fully or partially reconstructed $D_s \ell$ vertices and a mixing tag as a flavor tagging.
- ¹⁶ Replaced by ABDALLAH 04A. Uses $D_s^- \ell^+$, and $\phi \ell^+$ vertices, and a multi-variable discriminant as a flavor tagging.
- ¹⁷ Uses inclusive D_s vertices and fully reconstructed B_s decays and a multi-variable discriminant as a flavor tagging.
- ¹⁸ Uses ℓ - Q_{hem} and ℓ - ℓ .
- ¹⁹ ABE 99D assumes $\tau_{B_s^0} = 1.55 \pm 0.05 \text{ ps}$ and $\Delta\Gamma/\Delta m = (5.6 \pm 2.6) \times 10^{-3}$.
- ²⁰ ABE 99J uses ϕ ℓ - ℓ correlation.
- ²¹ BARATE 99J uses combination of an inclusive lepton and D_s^- -based analyses.
- ²² BARATE 98C combines results from $D_s h$ - ℓ/Q_{hem} , $D_s h$ - K in the same side, $D_s \ell$ - ℓ/Q_{hem} and $D_s \ell$ - K in the same side.
- ²³ Uses ℓ - Q_{hem} .
- ²⁴ Uses ℓ - ℓ .
- ²⁵ ADAM 97 combines results from $D_s \ell$ - Q_{hem} , ℓ - Q_{hem} , and ℓ - ℓ .
- ²⁶ BUSKULIC 96M uses D_s lepton correlations and lepton, kaon, and jet charge tags.
- ²⁷ BUSKULIC 95J uses ℓ - Q_{hem} . They find $\Delta m_s > 5.6$ [> 6.1] for $f_s=10\%$ [12%]. We interpolate to our central value $f_s=10.5\%$.

$$x_s = \Delta m_{B_s^0} / \Gamma_{B_s^0}$$

This is derived by the Heavy Flavor Averaging Group (HFAG) from the results on $\Delta m_{B_s^0}$ and "OUR EVALUATION" of the B_s^0 mean lifetime.

| | |
|------------------------------------|--------------------|
| <u>VALUE</u> | <u>DOCUMENT ID</u> |
| 26.81 ± 0.10 OUR EVALUATION | |

χ_s

This is a B_s^0 - \bar{B}_s^0 integrated mixing parameter derived from x_s above and OUR EVALUATION of $\Delta\Gamma_{B_s^0} / \Gamma_{B_s^0}$.

| | |
|---|--------------------|
| <u>VALUE</u> | <u>DOCUMENT ID</u> |
| 0.499308 ± 0.000005 OUR EVALUATION | |

CP VIOLATION PARAMETERS in B_s^0

$$\text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$$

CP impurity in B_s^0 system.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/scaling procedure takes into account correlation between the measurements. The value has been obtained from a 2D fit of the B_d and B_s asymmetries, which includes the B_s measurements listed below and the B factory average for the B_d .

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|----------|------------------------|
| -1.9 ± 1.0 OUR EVALUATION | | | |
| -1.5 ± 1.0 OUR AVERAGE | | | |
| -0.15 ± 1.25 ± 0.90 | 1 AAIJ | 14D LHCb | pp at 7 TeV |
| -2.15 ± 1.85 | 2 ABAZOV | 14 D0 | $p\bar{p}$ at 1.96 TeV |
| -2.8 ± 1.9 ± 0.4 | 3 ABAZOV | 13 D0 | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -4.5 ± 2.7 | 4 ABAZOV | 11U D0 | Repl. by ABAZOV 14 |
| -0.4 ± 2.3 ± 0.4 | 5 ABAZOV | 10E D0 | Repl. by ABAZOV 13 |
| -3.6 ± 1.9 | 6 ABAZOV | 10H D0 | Repl. by ABAZOV 11U |
| 6.1 ± 4.8 ± 0.9 | 7 ABAZOV | 07A D0 | Repl. by ABAZOV 10E |

¹ AAIJ 14D reports a measurement of time-integrated flavor-specific asymmetry in $B_s^0 \rightarrow \mu^+ D_s^- X$ decays $a_{sl}^s = (-0.06 \pm 0.50 \pm 0.36)\%$ which is approximately equal to $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$.

² ABAZOV 14 uses the dimuon charge asymmetry with different impact parameters from which it reports $A_{SL}^s = (-0.86 \pm 0.74) \times 10^{-2}$.

³ ABAZOV 13 reports a measurement of time-integrated flavor-specific asymmetry in mixed semileptonic $B_s^0 \rightarrow \mu^+ D_s^- X$ decays $A_s^{SL} = (-1.12 \pm 0.74 \pm 0.17)\%$ which is approximately equal to $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$.

⁴ ABAZOV 11U uses the dimuon charge asymmetry with different impact parameters from which it reports $A_{SL}^s = (-18.1 \pm 10.6) \times 10^{-3}$.

⁵ ABAZOV 10E reports a measurement of flavor-specific asymmetry in $B_{(s)}^0 \rightarrow \mu^+ D_{(s)}^{*-}$ X decays with a decay-time analysis including initial-state flavor tagging, $A_{SL}^s = (-1.7 \pm 9.1 \pm 1.4 \pm 1.5) \times 10^{-3}$ which is approximately equal to $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$.

⁶ ABAZOV 10H reports a measurement of like-sign dimuon charge asymmetry of $A_{SL}^b = (-9.57 \pm 2.51 \pm 1.46) \times 10^{-3}$ in semileptonic b -hadron decays. Using the measured production ratio of B_d^0 and B_s^0 , and the asymmetry of B_d^0 $A_{SL}^d = (-4.7 \pm 4.6) \times 10^{-3}$ measured from B -factories, they obtain the asymmetry for B_s^0 .

⁷ The first direct measurement of the time integrated flavor untagged charge asymmetry in semileptonic B_s^0 decays is reported as $2 \times A_{SL}^s(\text{untagged}) = A_{SL}^s = (2.45 \pm 1.93 \pm 0.35) \times 10^{-2}$.

$C_{KK}(B_s^0 \rightarrow K^+ K^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|-----------|--------------------|
| 0.14 ± 0.11 ± 0.03 | AAIJ | 13B0 LHCb | <i>pp</i> at 7 TeV |

$S_{KK}(B_s^0 \rightarrow K^+ K^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|-----------|--------------------|
| 0.30 ± 0.12 ± 0.04 | AAIJ | 13B0 LHCb | <i>pp</i> at 7 TeV |

$\gamma(B_s^0 \rightarrow D_s^\pm K^\mp)$

For angle $\gamma(\phi_3)$ of the CKM unitarity triangle, see the review on “CP Violation” in the Reviews section.

| VALUE (°) | DOCUMENT ID | TECN | COMMENT |
|--|-------------------|-----------|--------------------|
| 115⁺²⁸₋₄₃ | ¹ AAIJ | 14BF LHCb | <i>pp</i> at 7 TeV |

¹ Measured in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays, constraining $-2\beta_s$ by the measurement of $\phi_s = 0.01 \pm 0.07 \pm 0.0$ from AAIJ 13AR. The value is modulo 180° at 68% CL.

$\delta_B(B_s^0 \rightarrow D_s^\pm K^\mp)$

| VALUE (°) | DOCUMENT ID | TECN | COMMENT |
|--------------------------------------|-------------------|-----------|--------------------|
| 3⁺¹⁹₋₂₀ | ¹ AAIJ | 14BF LHCb | <i>pp</i> at 7 TeV |

¹ Measured in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays, constraining $-2\beta_s$ by the measurement of $\phi_s = 0.01 \pm 0.07 \pm 0.0$ from AAIJ 13AR. The value is modulo 180° at 68% CL.

$r_B(B_s^0 \rightarrow D_s^\mp K^\pm)$

r_B and δ_B are the amplitude ratio and relative strong phase between the amplitudes of $A(B_s^0 \rightarrow D_s^+ K^-)$ and $A(B_s^0 \rightarrow D_s^- K^+)$,

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|-----------|--------------------|
| 0.53^{+0.17}_{-0.16} | ¹ AAIJ | 14BF LHCb | <i>pp</i> at 7 TeV |

¹ Measured in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays, constraining $-2\beta_s$ by the measurement of $\phi_s = 0.01 \pm 0.07 \pm 0.0$ from AAIJ 13AR. At 68% CL.

CP Violation phase β_s

$-2\beta_s$ is the weak phase difference between B_s^0 mixing amplitude and the $B_s^0 \rightarrow J/\psi\phi$ decay amplitude driven by the $b \rightarrow c\bar{c}s$ transition (such as $B_s \rightarrow J/\psi\phi$, $J/\psi K^+ K^-$, $J/\psi \pi^+ \pi^-$, and $D_s^+ D_s^-$). The Standard Model value of β_s is $\arg(-\frac{V_{ts}V_{tb}^*}{V_{cs}V_{cb}^*})$.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/scaling procedure takes into account correlation between the measurements.

| VALUE (10^{-2} rad) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------------|-----------|-------------------------------------|
| 0.6 ± 1.9 OUR EVALUATION | | | |
| 0.7 ± 2.1 OUR AVERAGE | | | Error includes scale factor of 1.1. |
| 2.9 ± 2.5 ± 0.3 | ¹ AAIJ | 15i LHCb | <i>pp</i> at 7, 8 TeV |
| – 6 ± 13 ± 3 | ² AAD | 14U ATLAS | <i>pp</i> at 7 TeV |
| – 1 ± 9 ± 1 | ³ AAIJ | 14AY LHCb | <i>pp</i> at 7, 8 TeV |

| | | | |
|--|-------------------------|-----------|---------------------------|
| – 3.5 ± 3.4 ± 0.4 | ⁴ AAIJ | 14S LHCb | $\rho\rho$ at 7 and 8 TeV |
| | ⁵ AALTONEN | 12AJ CDF | $\rho\bar{p}$ at 1.96 TeV |
| 28 $\begin{smallmatrix} +18 \\ -19 \end{smallmatrix}$ | ^{6,7,8} ABAZOV | 12D D0 | $\rho\bar{p}$ at 1.96 TeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| – 17 ± 15 ± 3 | ⁹ AAIJ | 14AE LHCb | $\rho\rho$ at 7, 8 TeV |
| – 0.5 ± 3.5 ± 0.5 | ¹⁰ AAIJ | 13AR LHCb | Repl. by AAIJ 15I |
| | ¹¹ AAIJ | 13AY LHCb | $\rho\rho$ at 7 TeV |
| – 11.0 ± 20.5 ± 5.0 | ¹² AAD | 12CV ATLS | Repl. by AAD 14U |
| 22 ± 22 ± 1 | ¹³ AAIJ | 12B LHCb | Repl. by AAIJ 12Q |
| – 8 ± 9 ± 3 | ¹⁴ AAIJ | 12D LHCb | Repl. by AAIJ 13AR |
| 0.95 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 8.70+0.15 \\ 8.65-0.20 \end{smallmatrix}$ | ¹⁵ AAIJ | 12Q LHCb | Repl. by AAIJ 13AR |
| | ¹⁶ AALTONEN | 12D CDF | Repl. by AALTONEN 12AJ |
| | ¹⁷ AALTONEN | 08G CDF | Repl. by AALTONEN 12D |
| 28 $\begin{smallmatrix} +12 \\ -15 \end{smallmatrix}$ $\begin{smallmatrix} +4 \\ -1 \end{smallmatrix}$ | ^{7,18} ABAZOV | 08AMD0 | Repl. by ABAZOV 12D |
| 39.5 ± 28.0 $\begin{smallmatrix} +0.5 \\ -7.0 \end{smallmatrix}$ | ^{8,19} ABAZOV | 07 D0 | Repl. by ABAZOV 07N |
| 35 $\begin{smallmatrix} +20 \\ -24 \end{smallmatrix}$ | ^{8,20} ABAZOV | 07N D0 | Repl. by ABAZOV 08AM |

¹ AAIJ 15I reports $\phi_s = -2\beta_s = -0.058 \pm 0.049 \pm 0.006$ rad. that was measured using a tagged, time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ decays. It also combines this result with that of AAIJ 14S and quotes $\phi_s = -2\beta_s = -0.010 \pm 0.039$ rad.

² AAD 14U reports $\phi_s = -2\beta_s = 0.12 \pm 0.25 \pm 0.05$ rad. that was measured using a tagged, time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

³ AAIJ 14AY reports $\phi_s = -2\beta_s = 0.02 \pm 0.17 \pm 0.02$ rad. in tagged, time-dependent fit to $B_s^0 \rightarrow D_s^+ D_s^-$, while allowing CP violation in decay.

⁴ AAIJ 14S reports $\phi_s = -2\beta_s = 0.070 \pm 0.068 \pm 0.008$ rad. and $|\lambda| = 0.89 \pm 0.05 \pm 0.01$, when direct CP violation is allowed. Measured using a tagged, time-dependent fit to $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decays.

⁵ AALTONEN 12AJ reports $-\pi/2 < \beta_s < -1.51$ or $-0.06 < \beta_s < 0.30$, or $1.26 < \beta_s < \pi/2$ rad. at 68% CL. Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

⁶ The error includes both statistical and systematic uncertainties.

⁷ Measured using fully reconstructed $B_s \rightarrow J/\psi\phi$ decays.

⁸ Reports ϕ_s which equals to $-2\beta_s$.

⁹ Measured in $B_s^0 \rightarrow \phi\phi$ decays. This is a $b \rightarrow s\bar{s}s$ transition with a decay amplitude phase different from that of $b \rightarrow c\bar{c}s$ transition.

¹⁰ AAIJ 13AR reports $\phi_s = -2\beta_s = 0.01 \pm 0.07 \pm 0.01$ rad. obtained from combined fit to $B_s^0 \rightarrow J/\psi K^+ K^-$ and $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ data sets. Also reports separate results of $\phi_s = 0.07 \pm 0.09 \pm 0.01$ rad. from $B_s^0 \rightarrow J/\psi K^+ K^-$ decays and $\phi_s = -0.14 \begin{smallmatrix} +0.17 \\ -0.16 \end{smallmatrix} \pm 0.01$ rad. from $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decays.

¹¹ AAIJ 13AY uses $B_s^0 \rightarrow \phi\phi$ mode, and reports the 68% CL interval of $\phi_s = -2\beta_s$ as $[-2.46, -0.76]$ rad.

¹² AAD 12CV reports $\phi_s = -2\beta_s = 0.22 \pm 0.41 \pm 0.10$ rad. that was measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

- ¹³ Reports $\phi_s = -2\beta_s = -0.44 \pm 0.44 \pm 0.02$ rad. that was measured using a time-dependent fit to $B_s^0 \rightarrow J/\psi f_0(980)$ decays.
- ¹⁴ Reports $\phi_s = -2\beta_s = 0.15 \pm 0.18 \pm 0.06$ rad. that was measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.
- ¹⁵ Reports $\phi_s = -2\beta_s = -0.019^{+0.173+0.004}_{-0.174-0.003}$ rad. which was measured using a time-dependent fit to $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays, with the $\pi^+ \pi^-$ mass within 775–1550 MeV. Searches for, but finds no evidence, for direct CP violation in $B_s^0 \rightarrow J/\psi \pi \pi$ decays.
- ¹⁶ Reports $0.02 < \phi_s < 0.52$ or $1.08 < \phi_s < 1.55$ rad. at 68% C.L. confidence regions in the two-dimensional space of ϕ_s and $\Delta\Gamma_{B_s^0}$ from $B_s^0 \rightarrow J/\psi \phi$ decays.
- ¹⁷ Reports $0.32 < 2\beta_s < 2.82$ rad. at 68% C.L. and confidence regions in the two-dimensional space of $2\beta_s$ and $\Delta\Gamma$ from the first measurement of $B_s^0 \rightarrow J/\psi \phi$ decays using flavor tagging. The probability of a deviation from SM prediction as large as the level of observed data is 15%.
- ¹⁸ Reports $\phi_s = -2\beta_s$ and obtains 90% CL interval $-0.03 < \beta_s < 0.60$ rad.
- ¹⁹ The first direct measurement of the CP -violating mixing phase is reported from the time-dependent analysis of flavor untagged $B_s^0 \rightarrow J/\psi \phi$ decays.
- ²⁰ Combines D0 collaboration measurements of time-dependent angular distributions in $B_s^0 \rightarrow J/\psi \phi$ and charge asymmetry in semileptonic decays. There is a 4-fold ambiguity in the solution.

$|\lambda| (B_s^0 \rightarrow J/\psi(1S)\phi)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------|------|-----------------------|
| 0.964 ± 0.019 ± 0.007 | AAIJ | 15I | LHCB pp at 7, 8 TeV |

$|\lambda|$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|------|-----------------------|
| 1.02 ± 0.07 OUR AVERAGE | | | |
| 1.04 ± 0.07 ± 0.03 | ¹ AAIJ | 14AE | LHCB pp at 7, 8 TeV |
| 0.91 ^{+0.18} _{-0.15} ± 0.02 | ² AAIJ | 14AY | LHCB pp at 7, 8 TeV |

¹ Measured in $B_s^0 \rightarrow \phi\phi$ decays.

² Measured in $B_s^0 \rightarrow D_s^+ D_s^-$ decays.

$A_{CP}(B_s \rightarrow \pi^+ K^-)$

A_{CP} is defined as

$$\frac{B(\bar{B}_s^0 \rightarrow f) - B(B_s^0 \rightarrow \bar{f})}{B(\bar{B}_s^0 \rightarrow f) + B(B_s^0 \rightarrow \bar{f})},$$

the CP -violation asymmetry of exclusive B_s^0 and \bar{B}_s^0 decay.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|----------------------------|
| 0.263 ± 0.035 OUR AVERAGE | | | |
| 0.22 ± 0.07 ± 0.02 | AALTONEN | 14P | CDF $p\bar{p}$ at 1.96 TeV |
| 0.27 ± 0.04 ± 0.01 | AAIJ | 13AX | LHCB pp at 7 TeV |
| 0.39 ± 0.15 ± 0.08 | AALTONEN | 11N | CDF $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.27 ± 0.08 ± 0.02 | AAIJ | 12V | LHCB Repl. by AAIJ 13AX |

$A_{CP}(B_s^0 \rightarrow [K^+ K^-]_D \bar{K}^*(892)^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|---------------------|
| $-0.04 \pm 0.07 \pm 0.02$ | AAIJ | 14BN LHCB | pp at 7 and 8 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $0.04 \pm 0.16 \pm 0.01$ | AAIJ | 13L LHCB | Repl. by AAIJ 14BN |

$A_{CP}(B_s^0 \rightarrow [\pi^+ K^-]_D K^*(892)^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|---------------------|
| $-0.01 \pm 0.03 \pm 0.02$ | AAIJ | 14BN LHCB | pp at 7 and 8 TeV |

$A_{CP}(B_s^0 \rightarrow [\pi^+ \pi^-]_D K^*(892)^0)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|-----------|---------------------|
| $0.06 \pm 0.13 \pm 0.02$ | AAIJ | 14BN LHCB | pp at 7 and 8 TeV |

PARTIAL BRANCHING FRACTIONS IN $B_s \rightarrow \phi \ell^+ \ell^-$

$B(B_s \rightarrow \phi \ell^+ \ell^-) (0.1 < q^2 < 2.0 \text{ GeV}^2/c^4)$

| VALUE (units 10^{-7}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|----------|---|
| 0.95 ± 0.22 OUR AVERAGE | | | |
| $0.897^{+0.207}_{-0.186} \pm 0.097$ | AAIJ | 13X LHCB | pp at 7 TeV, $B_s^0 \rightarrow \phi \mu^+ \mu^-$ |
| $2.78 \pm 0.95 \pm 0.89$ | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |

$B(B_s \rightarrow \phi \ell^+ \ell^-) (2.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

| VALUE (units 10^{-7}) | DOCUMENT ID | TECN | COMMENT |
|--|-------------|----------|---|
| $0.53^{+0.18}_{-0.16}$ OUR AVERAGE | | | |
| $0.529^{+0.182}_{-0.159} \pm 0.057$ | AAIJ | 13X LHCB | pp at 7 TeV, $B_s^0 \rightarrow \phi \mu^+ \mu^-$ |
| $0.58 \pm 0.55 \pm 0.19$ | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |

$B(B_s \rightarrow \phi \ell^+ \ell^-) (4.3 < q^2 < 8.68 \text{ GeV}^2/c^4)$

| VALUE (units 10^{-7}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|----------|---|
| 1.38 ± 0.27 OUR AVERAGE | | | |
| $1.38^{+0.25}_{-0.23} \pm 0.14$ | AAIJ | 13X LHCB | pp at 7 TeV, $B_s^0 \rightarrow \phi \mu^+ \mu^-$ |
| $1.34 \pm 0.83 \pm 0.43$ | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |

$B(B_s \rightarrow \phi \ell^+ \ell^-) (10.09 < q^2 < 12.86 \text{ GeV}^2/c^4)$

| VALUE (units 10^{-7}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|----------|---|
| 1.24 ± 0.25 OUR AVERAGE | | | |
| $1.18^{+0.22}_{-0.21} \pm 0.14$ | AAIJ | 13X LHCB | pp at 7 TeV, $B_s^0 \rightarrow \phi \mu^+ \mu^-$ |
| $2.98 \pm 0.95 \pm 0.95$ | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |

$B(B_s \rightarrow \phi \ell^+ \ell^-) (14.18 < q^2 < 16.0 \text{ GeV}^2/c^4)$

| VALUE (units 10^{-7}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|----------|---|
| 0.81 ± 0.19 OUR AVERAGE | | | |
| $0.760^{+0.189}_{-0.169} \pm 0.087$ | AAIJ | 13X LHCB | pp at 7 TeV, $B_s^0 \rightarrow \phi \mu^+ \mu^-$ |
| $1.86 \pm 0.66 \pm 0.59$ | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |

$B(B_s \rightarrow \phi \ell^+ \ell^-) (16.0 < q^2 < 19.0 \text{ GeV}^2/c^4)$

| VALUE (units 10^{-7}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|----------|---|
| 1.13±0.24 OUR AVERAGE | | | |
| $1.06^{+0.23}_{-0.21} \pm 0.12$ | AAIJ | 13X LHCb | pp at 7 TeV, $B_s^0 \rightarrow \phi \mu^+ \mu^-$ |
| $2.32 \pm 0.76 \pm 0.74$ | AALTONEN | 11A1 CDF | $p\bar{p}$ at 1.96 TeV |

$B(B_s \rightarrow \phi \ell^+ \ell^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

| VALUE (units 10^{-7}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|----------|---|
| 1.14±0.26 OUR AVERAGE | | | |
| $1.14^{+0.25}_{-0.23} \pm 0.13$ | AAIJ | 13X LHCb | pp at 7 TeV, $B_s^0 \rightarrow \phi \mu^+ \mu^-$ |
| $1.14 \pm 0.79 \pm 0.36$ | AALTONEN | 11A1 CDF | $p\bar{p}$ at 1.96 TeV |

$B(B_s \rightarrow \phi \ell^+ \ell^-) (0.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

| VALUE (units 10^{-7}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|----------|------------------------|
| 3.30±1.09±1.05 | AALTONEN | 11A1 CDF | $p\bar{p}$ at 1.96 TeV |

PRODUCTION ASYMMETRIES

$A_P(B_s^0)$

$$A_P(B_s^0) = [\sigma(\bar{B}_s^0) - \sigma(B_s^0)] / [\sigma(\bar{B}_s^0) + \sigma(B_s^0)]$$

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|-----------|---------------|
| 1.09±2.61±0.66 | ¹ AAIJ | 14BP LHCb | pp at 7 TeV |

¹ Based on time-dependent analysis of $B_s^0 \rightarrow D_s^- \pi^+$ in kinematic range $4 < p_T < 30$ GeV/c and $2.5 < \eta < 4.5$.

B_s^0 REFERENCES

| | | | | |
|----------|------|----------------|---------------------------|-----------------|
| AAIJ | 15D | JHEP 1501 024 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 15I | PRL 114 041801 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABAZOV | 15A | PRL 114 062001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| DUTTA | 15 | PR D91 011101 | D. Dutta <i>et al.</i> | (BELLE Collab.) |
| AAD | 14U | PR D90 052007 | G. Aad <i>et al.</i> | (ATLAS Collab.) |
| AAIJ | 14 | PRL 112 011801 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14AA | PRL 112 202001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14AE | PR D90 052011 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14AX | PRL 113 172001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14AY | PRL 113 211801 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14BF | JHEP 1411 060 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14BH | PR D90 072003 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14BM | NJP 16 123001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14BN | PR D90 112002 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14BP | PL B739 218 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14BR | PR D89 092006 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14D | PL B728 607 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14E | JHEP 1404 114 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14F | PRL 112 111802 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14L | JHEP 1407 140 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14R | PL B736 446 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14S | PL B736 186 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14Y | PRL 112 091802 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 14P | PRL 113 242001 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 14 | PR D89 012002 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| AAIJ | 13 | NP B867 1 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13A | NP B867 547 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AA | NP B871 403 | R. Aaij <i>et al.</i> | (LHCb Collab.) |

| | | | | |
|------------|------|------------------------|-----------------------------|-----------------|
| AAIJ | 13AB | NP B873 275 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AC | NP B874 663 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AL | PR D87 071101 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AN | PR D87 072004 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AP | PR D87 092007 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AQ | PR D87 112009 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AR | PR D87 112010 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AW | PRL 110 211801 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AX | PRL 110 221601 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AY | PRL 110 241802 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13B | PRL 110 021801 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13BA | PRL 111 101805 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13BI | NJP 15 053021 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13BM | PRL 111 141801 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13BO | JHEP 1310 183 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13BP | JHEP 1310 143 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13BQ | JHEP 1310 005 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13BW | JHEP 1311 092 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13BX | PL B727 403 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13CF | EPJ C73 2655 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13L | JHEP 1303 067 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13X | JHEP 1307 084 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13Z | JHEP 1309 006 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 13F | PR D87 072003 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 13 | PRL 110 011801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 13C | PR D87 072006 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| CHATRCHYAN | 13AW | PRL 111 101804 | S. Chatrchyan <i>et al.</i> | (CMS Collab.) |
| ESEN | 13 | PR D87 031101 | S. Esen <i>et al.</i> | (BELLE Collab.) |
| OSWALD | 13 | PR D87 072008 | C. Oswald <i>et al.</i> | (BELLE Collab.) |
| Also | | PR D90 119901 (errat.) | C. Oswald <i>et al.</i> | (BELLE Collab.) |
| SOLOVIEVA | 13 | PL B726 206 | E. Solovieva <i>et al.</i> | (BELLE Collab.) |
| THORNE | 13 | PR D88 114006 | F. Thorne <i>et al.</i> | (BELLE Collab.) |
| AAD | 12AE | PL B713 387 | G. Aad <i>et al.</i> | (ATLAS Collab.) |
| AAD | 12CV | JHEP 1212 072 | G. Aad <i>et al.</i> | (ATLAS Collab.) |
| AAIJ | 12 | PL B707 349 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12A | PL B708 55 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12AE | PR D85 112013 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12AG | JHEP 1206 115 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12AM | PRL 109 131801 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12AN | PRL 109 152002 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12AO | PR D86 052006 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12AP | PR D86 071102 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12AR | JHEP 1210 037 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12AX | PR D86 112005 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12B | PL B707 497 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12D | PRL 108 101803 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12E | PL B708 241 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12F | PL B709 50 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12I | PL B709 177 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12L | EPJ C72 2118 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12O | PL B713 172 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12P | PL B713 369 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12Q | PL B713 378 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12R | PL B716 393 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12S | PRL 108 151801 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12V | PRL 108 201601 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12W | PRL 108 231801 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 12AJ | PRL 109 171802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 12C | PRL 108 201801 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 12D | PR D85 072002 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 12L | PRL 108 211803 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 12AF | PR D86 092011 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 12C | PR D85 011103 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 12D | PR D85 032006 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| CHATRCHYAN | 12A | JHEP 1204 033 | S. Chatrchyan <i>et al.</i> | (CMS Collab.) |
| LEES | 12A | PR D85 011101 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LI | 12 | PRL 108 181808 | J. Li <i>et al.</i> | (BELLE Collab.) |
| PDG | 12 | PR D86 010001 | J. Beringer <i>et al.</i> | (PDG Collab.) |
| AAIJ | 11 | PL B698 115 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 11A | PL B698 14 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 11B | PL B699 330 | R. Aaij <i>et al.</i> | (LHCb Collab.) |

| | | | | |
|-------------|------|-------------------------|-----------------------------|------------------|
| AAIJ | 11D | PL B706 32 | R. Aaij, <i>et al</i> | (LHCb Collab.) |
| AAIJ | 11E | PR D84 092001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| Also | | PR D85 039904 (errata) | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 11A | PR D83 052012 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11AB | PR D84 052012 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11AG | PRL 107 191801 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| Also | | PRL 107 239903 (errata) | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11AI | PRL 107 201802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11AN | PRL 107 261802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11AP | PRL 107 272001 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11L | PRL 106 161801 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11N | PRL 106 181802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 11U | PR D84 052007 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| CHATRCHYAN | 11T | PRL 107 191802 | S. Chatrchyan <i>et al.</i> | (CMS Collab.) |
| LI | 11 | PRL 106 121802 | J. Li <i>et al.</i> | (BELLE Collab.) |
| ABAZOV | 10E | PR D82 012003 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 10H | PR D82 032001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 10S | PL B693 539 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ESEN | 10 | PRL 105 201802 | S. Esen <i>et al.</i> | (BELLE Collab.) |
| LOUVOT | 10 | PRL 104 231801 | R. LOUVOT <i>et al.</i> | (BELLE Collab.) |
| PENG | 10 | PR D82 072007 | C.-C. Peng <i>et al.</i> | (BELLE Collab.) |
| AALTONEN | 09AQ | PRL 103 191802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 09B | PR D79 011104 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 09C | PRL 103 031801 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 09P | PRL 102 201801 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 09E | PRL 102 032001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 09G | PRL 102 051801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 09I | PRL 102 091801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 09Y | PR D79 111102 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| LOUVOT | 09 | PRL 102 021801 | R. Louvot <i>et al.</i> | (BELLE Collab.) |
| AALTONEN | 08F | PRL 100 021803 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 08G | PRL 100 161802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 08I | PRL 100 101802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 08J | PRL 100 121803 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 08AM | PRL 101 241801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| WICHT | 08A | PRL 100 121801 | J. Wicht <i>et al.</i> | (BELLE Collab.) |
| ABAZOV | 07 | PRL 98 121801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 07A | PRL 98 151801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 07N | PR D76 057101 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 07Q | PR D76 092001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 07Y | PRL 99 241801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABULENCIA | 07C | PRL 98 061802 | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| DRUTSKOY | 07 | PRL 98 052001 | A. Drutskoy <i>et al.</i> | (BELLE Collab.) |
| DRUTSKOY | 07A | PR D76 012002 | A. Drutskoy <i>et al.</i> | (BELLE Collab.) |
| ABAZOV | 06B | PRL 97 021802 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 06G | PR D74 031107 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 06V | PRL 97 241801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABULENCIA | 06J | PRL 96 191801 | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| ABULENCIA | 06N | PRL 96 231801 | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| ABULENCIA | 06Q | PRL 97 062003 | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| ABULENCIA,A | 06D | PRL 97 211802 | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| ABULENCIA,A | 06G | PRL 97 242003 | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| ACOSTA | 06 | PRL 96 202001 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 05B | PRL 94 042001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 05E | PRL 94 071802 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 05W | PRL 95 171801 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABULENCIA | 05 | PRL 95 221805 | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| Also | | PRL 95 249905 (errata) | A. Abulencia <i>et al.</i> | (CDF Collab.) |
| ACOSTA | 05 | PRL 94 101803 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ACOSTA | 05J | PRL 95 031801 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABDALLAH | 04A | PL B585 63 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ABDALLAH | 04J | EPJ C35 35 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ACOSTA | 04D | PRL 93 032001 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABDALLAH | 03B | EPJ C28 155 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ABE | 03 | PR D67 012006 | K. Abe <i>et al.</i> | (SLD Collab.) |
| HEISTER | 03E | EPJ C29 143 | A. Heister <i>et al.</i> | (ALEPH Collab.) |
| ABE | 02V | PR D66 032009 | K. Abe <i>et al.</i> | (SLD Collab.) |
| ACOSTA | 02D | PR D65 111101 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ACOSTA | 02G | PR D66 112002 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABBIENDI | 01D | EPJ C19 241 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABE | 00C | PR D62 071101 | K. Abe <i>et al.</i> | (SLD Collab.) |

| | | | | |
|--------------|-----|----------------------|-------------------------------|-------------------|
| ABREU | 00Y | EPJ C16 555 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU,P | 00G | EPJ C18 229 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| AFFOLDER | 00N | PRL 85 4668 | T. Affolder <i>et al.</i> | (CDF Collab.) |
| BARATE | 00K | PL B486 286 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| ABBIENDI | 99S | EPJ C11 587 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABE | 99D | PR D59 032004 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 99J | PRL 82 3576 | F. Abe <i>et al.</i> | (CDF Collab.) |
| BARATE | 99J | EPJ C7 553 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| Also | | EPJ C12 181 (errata) | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| ABE | 98 | PR D57 R3811 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 98B | PR D57 5382 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 98V | PRL 81 5742 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ACCIARRI | 98S | PL B438 417 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACKERSTAFF | 98F | EPJ C2 407 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 98G | PL B426 161 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| BARATE | 98C | EPJ C4 367 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BARATE | 98Q | EPJ C4 387 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| PDG | 98 | EPJ C3 1 | C. Caso <i>et al.</i> | (PDG Collab.) |
| ACCIARRI | 97B | PL B391 474 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 97C | PL B391 481 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACKERSTAFF | 97U | ZPHY C76 401 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 97V | ZPHY C76 417 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ADAM | 97 | PL B414 382 | W. Adam <i>et al.</i> | (DELPHI Collab.) |
| ABE | 96B | PR D53 3496 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 96L | PRL 76 4675 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 96N | PRL 77 1945 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 96Q | PR D54 6596 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 96F | ZPHY C71 11 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ADAM | 96D | ZPHY C72 207 | W. Adam <i>et al.</i> | (DELPHI Collab.) |
| BUSKULIC | 96E | ZPHY C69 585 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 96M | PL B377 205 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 96V | PL B384 471 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| PDG | 96 | PR D54 1 | R. M. Barnett <i>et al.</i> | (PDG Collab.) |
| ABE | 95R | PRL 74 4988 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 95Z | PRL 75 3068 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ACCIARRI | 95H | PL B363 127 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 95I | PL B363 137 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| AKERS | 95G | PL B350 273 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| AKERS | 95J | ZPHY C66 555 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| BUSKULIC | 95J | PL B356 409 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 95O | PL B361 221 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| ABREU | 94D | PL B324 500 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 94E | ZPHY C61 407 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| Also | | PL B289 199 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| AKERS | 94J | PL B337 196 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| AKERS | 94L | PL B337 393 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| BUSKULIC | 94B | PL B322 441 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 94C | PL B322 275 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| ABE | 93F | PRL 71 1685 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ACTON | 93H | PL B312 501 | P.D. Acton <i>et al.</i> | (OPAL Collab.) |
| BUSKULIC | 93G | PL B311 425 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| ABREU | 92M | PL B289 199 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACTON | 92N | PL B295 357 | P.D. Acton <i>et al.</i> | (OPAL Collab.) |
| BUSKULIC | 92E | PL B294 145 | D. Buskalic <i>et al.</i> | (ALEPH Collab.) |
| LEE-FRANZINI | 90 | PRL 65 2947 | J. Lee-Franzini <i>et al.</i> | (CUSB II Collab.) |