

K(1460)

$$I(J^P) = \frac{1}{2}(0^-)$$

OMITTED FROM SUMMARY TABLE

Observed in $K\pi\pi$ partial-wave analysis.**K(1460) MASS**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 1460	DAUM	81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$
~ 1400	¹ BRANDENB...	76B	ASPK ±	13 $K^\pm p \rightarrow K^\pm 2\pi p$
¹ Coupled mainly to $K f_0(1370)$. Decay into $K^*(892)\pi$ seen.				

K(1460) WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 260	DAUM	81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$
~ 250	² BRANDENB...	76B	ASPK ±	13 $K^\pm p \rightarrow K^\pm 2\pi p$
² Coupled mainly to $K f_0(1370)$. Decay into $K^*(892)\pi$ seen.				

K(1460) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	seen
Γ_2 $K\rho$	seen
Γ_3 $K_0^*(1430)\pi$	seen

K(1460) PARTIAL WIDTHS **$\Gamma(K^*(892)\pi)$ Γ_1**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
~ 109	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

 $\Gamma(K\rho)$ Γ_2

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
~ 34	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

 $\Gamma(K_0^*(1430)\pi)$ Γ_3

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
~ 117	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

***K*(1460) REFERENCES**

DAUM 81C NP B187 1 C. Daum *et al.* (AMST, CERN, CRAC, MPIM+)
BRANDENB... 76B PRL 36 1239 G.W. Brandenburg *et al.* (SLAC) JP
