

$\chi_{b1}(2P)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

$\chi_{b1}(2P)$ MASS

| VALUE (MeV) | DOCUMENT ID |
|--|---|
| 10255.46 ± 0.22 ± 0.50 OUR EVALUATION | From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV |


$m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)}$

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-------------------------|---------------------|------|--|
| 23.5 ± 0.7 ± 0.7 | ¹ HEINTZ | 92 | CSB2 $e^+e^- \rightarrow \gamma X, l^+l^- \gamma \gamma$ |

¹From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

γ ENERGY IN $\Upsilon(3S)$ DECAY

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|-------|---------------------|------|---|
| 99.26 ± 0.22 OUR EVALUATION | | | | Treating systematic errors as correlated |
| 99.53 ± 0.23 OUR AVERAGE | | | | Error includes scale factor of 1.3. See the ideogram below. |
| 99.15 ± 0.07 ± 0.25 | | ARTUSO | 05 | CLEO $\Upsilon(3S) \rightarrow \gamma X$ |
| 99 ± 1 | 169 | CRAWFORD | 92B | CLE2 $e^+e^- \rightarrow l^+l^- \gamma \gamma$ |
| 100.1 ± 0.4 | 11147 | ² HEINTZ | 92 | CSB2 $e^+e^- \rightarrow \gamma X$ |
| 100.2 ± 0.5 | 223 | ³ HEINTZ | 92 | CSB2 $e^+e^- \rightarrow l^+l^- \gamma \gamma$ |
| 99.5 ± 0.1 ± 0.5 | 25759 | MORRISON | 91 | CLE2 $e^+e^- \rightarrow \gamma X$ |



²A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

³A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

$\chi_{b1}(2P)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor |
|---|--------------------------------|--------------|
| $\Gamma_1 \quad \omega \Upsilon(1S)$ | $(1.63^{+0.40}_{-0.34})\%$ | |
| $\Gamma_2 \quad \gamma \Upsilon(2S)$ | $(19.9 \pm 1.9)\%$ | |
| $\Gamma_3 \quad \gamma \Upsilon(1S)$ | $(9.2 \pm 0.8)\%$ | 1.1 |
| $\Gamma_4 \quad \pi\pi\chi_{b1}(1P)$ | $(9.1 \pm 1.3) \times 10^{-3}$ | |
| $\Gamma_5 \quad D^0 X$ | $(8.8 \pm 1.7)\%$ | |
| $\Gamma_6 \quad \pi^+\pi^-K^+K^-\pi^0$ | $(3.1 \pm 1.0) \times 10^{-4}$ | |
| $\Gamma_7 \quad 2\pi^+\pi^-K^-K_S^0$ | $(1.1 \pm 0.5) \times 10^{-4}$ | |
| $\Gamma_8 \quad 2\pi^+\pi^-K^-K_S^0 2\pi^0$ | $(7.7 \pm 3.2) \times 10^{-4}$ | |
| $\Gamma_9 \quad 2\pi^+2\pi^-2\pi^0$ | $(5.9 \pm 2.0) \times 10^{-4}$ | |
| $\Gamma_{10} \quad 2\pi^+2\pi^-K^+K^-$ | $(10 \pm 4) \times 10^{-5}$ | |
| $\Gamma_{11} \quad 2\pi^+2\pi^-K^+K^-\pi^0$ | $(5.5 \pm 1.8) \times 10^{-4}$ | |
| $\Gamma_{12} \quad 2\pi^+2\pi^-K^+K^-2\pi^0$ | $(10 \pm 4) \times 10^{-4}$ | |
| $\Gamma_{13} \quad 3\pi^+2\pi^-K^-K_S^0\pi^0$ | $(6.7 \pm 2.6) \times 10^{-4}$ | |
| $\Gamma_{14} \quad 3\pi^+3\pi^-$ | $(1.2 \pm 0.4) \times 10^{-4}$ | |
| $\Gamma_{15} \quad 3\pi^+3\pi^-2\pi^0$ | $(1.2 \pm 0.4) \times 10^{-3}$ | |
| $\Gamma_{16} \quad 3\pi^+3\pi^-K^+K^-$ | $(2.0 \pm 0.8) \times 10^{-4}$ | |
| $\Gamma_{17} \quad 3\pi^+3\pi^-K^+K^-\pi^0$ | $(6.1 \pm 2.2) \times 10^{-4}$ | |
| $\Gamma_{18} \quad 4\pi^+4\pi^-$ | $(1.7 \pm 0.6) \times 10^{-4}$ | |
| $\Gamma_{19} \quad 4\pi^+4\pi^-2\pi^0$ | $(1.9 \pm 0.7) \times 10^{-3}$ | |

$\chi_{b1}(2P)$ BRANCHING RATIOS

| $\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$ | | | | | Γ_1/Γ |
|---|----------------------|-----------------------------|------|--|-------------------|
| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $1.63^{+0.35+0.16}_{-0.31-0.15}$ | $32.6^{+6.9}_{-6.1}$ | ⁴ CRONIN-HEN..04 | CLE3 | $\Upsilon(3S) \rightarrow \gamma\omega \Upsilon(1S)$ | |

⁴ Using $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (11.3 \pm 0.6)\%$ and $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = 2$
 $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 2 (2.48 \pm 0.06)\%$.

| $\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$ | | | | | Γ_2/Γ |
|---|------|-----------------------|----------|---|-------------------|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 0.199 ± 0.019 OUR AVERAGE | | | | | |
| $0.190 \pm 0.018 \pm 0.017$ | 4.3k | ⁵ LEES | 11J BABR | $\Upsilon(3S) \rightarrow X\gamma$ | |
| $0.356 \pm 0.042 \pm 0.092$ | | ⁶ CRAWFORD | 92B CLE2 | $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ | |
| $0.199 \pm 0.020 \pm 0.022$ | | ⁷ HEINTZ | 92 CSB2 | $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ | |

⁵ LEES 11J reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (2.4 \pm 0.1 \pm 0.2) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (10.23 \pm 1.20 \pm 1.26) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = 0.105^{+0.003}_{-0.002} \pm 0.013$.

⁷ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (11.5 \pm 0.5 \pm 0.5)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_3/Γ

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|-------------------------------------|-------------|---|
| 0.092±0.008 OUR AVERAGE | | Error includes scale factor of 1.1. | | |
| 0.098±0.005±0.009 | 15k | ⁸ LEES | 11J BABR | $\Upsilon(3S) \rightarrow X\gamma$ |
| 0.120±0.021±0.021 | | ⁹ CRAWFORD | 92B CLE2 | $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ |
| 0.080±0.009±0.007 | | ¹⁰ HEINTZ | 92 CSB2 | $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ |

⁸ LEES 11J reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (12.4 \pm 0.3 \pm 0.6) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (6.47 \pm 1.12 \pm 0.82) \times 10^{-4}$ and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = 0.105^{+0.003}_{-0.002} \pm 0.013$.

¹⁰ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (11.5 \pm 0.5 \pm 0.5)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\pi\pi\chi_{b1}(1P))/\Gamma_{\text{total}}$ Γ_4/Γ

| <u>VALUE (units 10^{-3})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-------------------------|-------------|---|
| 9.1±1.3 OUR AVERAGE | | | | |
| 9.2±1.1±0.8 | 31k | ¹¹ LEES | 11C BABR | $e^+e^- \rightarrow \pi^+\pi^-X$ |
| 8.6±2.3±2.1 | | ¹² CAWLFIELD | 06 CLE3 | $\Upsilon(3S) \rightarrow 2(\gamma\pi\ell)$ |

¹¹ LEES 11C measures $B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X) \times B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) = (1.16 \pm 0.07 \pm 0.12) \times 10^{-3}$. We derive the value assuming $B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X) = B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)\gamma) = (12.6 \pm 1.2) \times 10^{-2}$.

¹² CAWLFIELD 06 quote $\Gamma(\chi_b(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV assuming l-spin conservation, no D-wave contribution, $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$ keV, and $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$ keV.

$\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_5/Γ

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 8.8±1.5±0.8 | 2243 | ¹³ BRIERE | 08 CLEO | $\Upsilon(3S) \rightarrow \gamma D^0 X$ |

¹³ For $p_{D^0} > 2.5$ GeV/c.

$\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|---------------------|-------------|--|
| 3.1±1.0±0.3 | 30 | ¹⁴ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$ |

¹⁴ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (39 \pm 8 \pm 9) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$ Γ_7/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------|----------|---|
| 1.1±0.5±0.1 | 10 | ¹⁵ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$ |

¹⁵ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]$ = $(14 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+\pi^-K^-K_S^02\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------|----------|--|
| 7.7±3.1±0.7 | 15 | ¹⁶ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$ |

¹⁶ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^02\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]$ = $(97 \pm 30 \pm 26) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------|----------|--|
| 5.9±2.0±0.5 | 36 | ¹⁷ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$ |

¹⁷ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]$ = $(74 \pm 16 \pm 19) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------|----------|--|
| 1.0±0.4±0.1 | 12 | ¹⁸ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$ |

¹⁸ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]$ = $(12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------|----------|---|
| 5.5±1.7±0.5 | 38 | ¹⁹ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$ |

¹⁹ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]$ = $(69 \pm 13 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------|----------|--|
| 9.6±3.5±0.9 | 27 | ²⁰ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$ |

²⁰ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]$ = $(121 \pm 29 \pm 33) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$ **Γ_{13}/Γ**

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|---------------------|-------------|---|
| $6.7 \pm 2.5 \pm 0.6$ | 17 | ²¹ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$ |

²¹ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(85 \pm 23 \pm 22) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+3\pi^-)/\Gamma_{\text{total}}$ **Γ_{14}/Γ**

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|---------------------|-------------|---|
| $1.2 \pm 0.4 \pm 0.1$ | 18 | ²² ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$ |

²² ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(15 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}$ **Γ_{15}/Γ**

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|---------------------|-------------|--|
| $12 \pm 4 \pm 1$ | 44 | ²³ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$ |

²³ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(150 \pm 30 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+3\pi^-K^+K^-)/\Gamma_{\text{total}}$ **Γ_{16}/Γ**

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|---------------------|-------------|---|
| $2.0 \pm 0.7 \pm 0.2$ | 16 | ²⁴ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$ |

²⁴ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(25 \pm 7 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+3\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{17}/Γ**

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|---------------------|-------------|---|
| $6.1 \pm 2.1 \pm 0.6$ | 25 | ²⁵ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$ |

²⁵ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(77 \pm 17 \pm 21) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(4\pi^+4\pi^-)/\Gamma_{\text{total}}$ **Γ_{18}/Γ**

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|---------------------|-------------|---|
| $1.7 \pm 0.6 \pm 0.2$ | 16 | ²⁶ ASNER | 08A CLEO | $\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$ |

²⁶ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(22 \pm 6 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------------------|------|--|
| 19±7±2 | 41 | ²⁷ ASNER 08A | CLEO | $\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$ |

²⁷ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$
 $= (241 \pm 47 \pm 72) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))$
 $= (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\chi_{b1}(2P)$ Cross-Particle Branching Ratios

$\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$
 $\Gamma_3/\Gamma \times \Gamma_{20}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---|
| 12.4±0.3±0.6 | 15k | LEES | 11J | BABR $\Upsilon(3S) \rightarrow X\gamma$ |

$\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$
 $\Gamma_2/\Gamma \times \Gamma_{20}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---|
| 2.4±0.1±0.2 | 4.3k | LEES | 11J | BABR $\Upsilon(3S) \rightarrow X\gamma$ |

$B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X)$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------------------------------------|
| 1.16±0.07±0.12 | 31k | LEES | 11C | BABR $e^+e^- \rightarrow \pi^+\pi^-X$ |

$B(\chi_{b2}(2P) \rightarrow \rho X + \bar{\rho} X)/B(\chi_{b1}(2P) \rightarrow \rho X + \bar{\rho} X)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---|
| 1.109±0.007±0.040 | BRIERE 07 | CLEO | $\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)$ |

$B(\chi_{b0}(2P) \rightarrow \rho X + \bar{\rho} X)/B(\chi_{b1}(2P) \rightarrow \rho X + \bar{\rho} X)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---|
| 1.082±0.025±0.060 | BRIERE 07 | CLEO | $\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)$ |

$\chi_{b1}(2P)$ REFERENCES

| | | | | |
|-----------------|-----|---------------|----------------------------------|-------------------|
| LEES | 11C | PR D84 011104 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 11J | PR D84 072002 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ASNER | 08A | PR D78 091103 | D.M. Asner <i>et al.</i> | (CLEO Collab.) |
| BRIERE | 08 | PR D78 092007 | R.A. Briere <i>et al.</i> | (CLEO Collab.) |
| BRIERE | 07 | PR D76 012005 | R.A. Briere <i>et al.</i> | (CLEO Collab.) |
| CAWLFIELD | 06 | PR D73 012003 | C. Cawfield <i>et al.</i> | (CLEO Collab.) |
| ARTUSO | 05 | PRL 94 032001 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| CRONIN-HENNESSY | 04 | PRL 92 222002 | D. Cronin-Hennessy <i>et al.</i> | (CLEO Collab.) |
| CRAWFORD | 92B | PL B294 139 | G. Crawford, R. Fulton | (CLEO Collab.) |
| HEINTZ | 92 | PR D46 1928 | U. Heintz <i>et al.</i> | (CUSB II Collab.) |
| HEINTZ | 91 | PRL 66 1563 | U. Heintz <i>et al.</i> | (CUSB Collab.) |
| MORRISON | 91 | PRL 67 1696 | R.J. Morrison <i>et al.</i> | (CLEO Collab.) |
| NARAIN | 91 | PRL 66 3113 | M. Narain <i>et al.</i> | (CUSB Collab.) |