

Double- β Decay

OMITTED FROM SUMMARY TABLE
A REVIEW GOES HERE – Check our WWW List of Reviews

Half-life Measurements and Limits for Double- β Decay

In most cases the transitions $(Z,A) \rightarrow (Z+2,A) + 2e^- + (0 \text{ or } 2) \bar{\nu}_e$ to the 0^+ ground state of the final nucleus are listed. However, we also list transitions that increase the nuclear charge ($2e^+$, e^+ /EC and ECEC) and transitions to excited states of the final nuclei (0_i^+ , 2^+ , and 2_i^+). In the following Listings, only best or comparable limits or lifetimes for each isotope are reported and only those with $T_{1/2} > 10^{20}$ years that are relevant for particle physics. For 2ν decay, which is well established, only measured half-lives are reported.

$t_{1/2}(10^{21} \text{ yr})$	CL%	ISOTOPE	TRANSITION	METHOD	DOCUMENT ID
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$2.165 \pm 0.016 \pm 0.059$		^{136}Xe	2ν g.s. \rightarrow g.s.	EXO-200	¹ ALBERT 14
> 11000	90	^{136}Xe	0ν g.s. \rightarrow g.s.	EXO-200	² ALBERT 14B
> 1100	90	^{100}Mo	0ν $\langle m \rangle$ -driven	NEMO-3	³ ARNOLD 14
> 600	90	^{100}Mo	0ν $\langle \lambda \rangle$ -driven	NEMO-3	⁴ ARNOLD 14
> 1000	90	^{100}Mo	0ν $\langle \eta \rangle$ -driven	NEMO-3	⁵ ARNOLD 14
$0.107^{+0.046}_{-0.026}$		^{150}Nd	$0\nu+2\nu$ $0^+ \rightarrow 0_1^+$	γ in Ge det.	⁶ KIDD 14
$1.84^{+0.14}_{-0.10}$		^{76}Ge	2ν g.s. \rightarrow g.s.	GERDA	⁷ AGOSTINI 13
> 21000	90	^{76}Ge	0ν g.s. \rightarrow g.s.	GERDA	⁸ AGOSTINI 13A
> 0.13	90	^{96}Ru	$0\nu+2\nu$ $2\beta^+$, g.s.	Ge counting	⁹ BELLI 13A
> 0.23	90	^{96}Ru	$0\nu+2\nu$ β^+ EC, g.s. $\rightarrow 2_1^+$		¹⁰ BELLI 13A
> 0.65	90	^{104}Ru	$0\nu+2\nu$ g.s. $\rightarrow 2_1^+$		¹¹ BELLI 13A
> 19000	90	^{136}Xe	0ν g.s. \rightarrow g.s.	KamLAND-Zen	¹² GANDO 13A
$9.2^{+5.5}_{-2.6} \pm 1.3$		^{78}Kr	$2\nu 2K$ g.s. \rightarrow g.s.	BAKSAN	¹³ GAVRILYAK 13
> 5.4	90	^{78}Kr	$0\nu 2K$ g.s. $\rightarrow 2^+$	BAKSAN	¹⁴ GAVRILYAK 13
> 940	90	^{130}Te	0ν $0^+ \rightarrow 0_1^+$	CUORICINO	¹⁵ ANDREOTTI 12
> 1.0	90	^{106}Cd	0ν ECEC, g.s.	$^{106}\text{CdWO}_4$ scint.	¹⁶ BELLI 12A
> 2.2	90	^{106}Cd	0ν β^+ EC, g.s.	$^{106}\text{CdWO}_4$ scint.	¹⁷ BELLI 12A
> 1.2	90	^{106}Cd	0ν $2\beta^+$, g.s.	$^{106}\text{CdWO}_4$ scint.	¹⁸ BELLI 12A
$2.38 \pm 0.02 \pm 0.14$		^{136}Xe	2ν g.s. \rightarrow g.s.	KamLAND-Zen	¹⁹ GANDO 12A
> 5700	90	^{136}Xe	0ν g.s. \rightarrow g.s.	KamLAND-Zen	²⁰ GANDO 12A
$2.11 \pm 0.04 \pm 0.21$		^{136}Xe	2ν	EXO-200	²¹ ACKERMAN 11
$0.7 \pm 0.09 \pm 0.11$		^{130}Te	2ν	NEMO-3	²² ARNOLD 11
> 130	90	^{130}Te	0ν	NEMO-3	²³ ARNOLD 11
> 1.3	90	^{112}Sn	0ν $0^+ \rightarrow 0_3^+$	γ Ge det.	²⁴ BARABASH 11
> 0.69	90	^{112}Sn	0ν $0^+ \rightarrow 0_2^+$	γ Ge det.	²⁵ BARABASH 11
> 1.3	90	^{112}Sn	0ν $0^+ \rightarrow 0_1^+$	γ Ge det.	²⁶ BARABASH 11
> 1.06	90	^{112}Sn	0ν	γ Ge det.	²⁷ BARABASH 11
$(2.8 \pm 0.1 \pm 0.3)\text{E-2}$		^{116}Cd	2ν	NEMO-3	²⁸ BARABASH 11A
$(4.4^{+0.5}_{-0.4} \pm 0.4)\text{E-2}$		^{48}Ca	2ν	NEMO-3	^{29,30} BARABASH 11A

(69 ± 9 ± 10)E-2	130Te	2ν		NEMO-3	30,31	BARABASH	11A
> 360	90 82Se	0ν		NEMO-3	30,32	BARABASH	11A
> 100	90 130Te	0ν		NEMO-3	30,33	BARABASH	11A
> 16	90 116Cd	0ν		NEMO-3	30,34	BARABASH	11A
> 13	90 48Ca	0ν		NEMO-3	30,35	BARABASH	11A
> 0.32	90 64Zn	0ν	ECEC, g.s.	ZnWO ₄ scint.	36	BELLI	11D
> 0.85	90 64Zn	0ν	β ⁺ EC, g.s.	ZnWO ₄ scint.	36	BELLI	11D
> 0.11	90 106Cd	0ν	0 ⁺ → 4 ⁺	TGV2 det.	37	RUKHADZE	11
(2.35 ± 0.14 ± 0.16)E-2	96Zr	2ν		NEMO-3	38	ARGYRIADES	10
> 9.2	90 96Zr	0ν		NEMO-3	39	ARGYRIADES	10
> 0.22	90 96Zr	0ν	0 ⁺ → 0 ₁ ⁺	NEMO-3	40	ARGYRIADES	10
0.69 ^{+0.10} _{-0.08} ± 0.07	100Mo	2ν	0 ⁺ → 0 ₁ ⁺	Ge coinc.	41	BELLI	10
> 18.0	90 150Nd	0ν		NEMO-3	42	ARGYRIADES	09
(9.11 ^{+0.25} _{-0.22} ± 0.63)E-3	150Nd	2ν		NEMO-3	43	ARGYRIADES	09
> 0.43	90 64Zn	0ν	β ⁺ EC	ZnWO ₄ scint.	44	BELLI	09A
> 0.11	90 64Zn	0ν	ECEC	ZnWO ₄ scint.	45	BELLI	09A
0.55 ^{+0.12} _{-0.09}	100Mo	2ν+0ν	0 ⁺ → 0 ₁ ⁺	Ge coincidence	46	KIDD	09
> 3000	90 130Te	0ν		TeO ₂ bolometer	47	ARNABOLDI	08
> 0.22	90 64Zn	0ν		ZnWO ₄ scint.	48	BELLI	08
> 1.1	90 114Cd	0ν	2β	CdWO ₄ scint.	49	BELLI	08B
> 58	90 48Ca	0ν		CaF ₂ scint.	50	UMEHARA	08
0.57 ^{+0.13} _{-0.09} ± 0.08	100Mo	2ν	0 ⁺ → 0 ₁ ⁺	NEMO-3	51	ARNOLD	07
> 89	90 100Mo	0ν	0 ⁺ → 0 ₁ ⁺	NEMO-3	52	ARNOLD	07
> 160	90 100Mo	0ν	0 ⁺ → 2 ₁ ⁺	NEMO-3	53	ARNOLD	07
> 0.0019	90 74Se	0ν+2ν		γ in Ge det.	54	BARABASH	07
> 0.0055	90 74Se	0ν+2ν	0 ⁺ → 2 ₂ ⁺	γ in Ge det.	55	BARABASH	07
22300 ⁺⁴⁴⁰⁰ ₋₃₁₀₀	76Ge	0ν		Enriched HPGe	56	KLAPDOR-K...	06A
> 1800	90 130Te	0ν		Cryog. det.	57	ARNABOLDI	05
> 460	90 100Mo	0ν		NEMO-3	58	ARNOLD	05A
> 100	90 82Se	0ν		NEMO-3	59	ARNOLD	05A
(7.11 ± 0.02 ± 0.54)E-3	100Mo	2ν		NEMO-3	60	ARNOLD	05A
(9.6 ± 0.3 ± 1.0)E-2	82Se	2ν		NEMO-3	61	ARNOLD	05A
> 140	90 82Se	0ν		NEMO-3	62	ARNOLD	04
(7.68 ± 0.02 ± 0.54)E-3	100Mo	2ν		NEMO-3	63	ARNOLD	04
0.14 ^{+0.04} _{-0.02} ± 0.03	150Nd	0ν+2ν	0 ⁺ → 0 ₁ ⁺	γ in Ge det.	64	BARABASH	04
> 31	90 130Te	0ν	0 ⁺ → 2 ₁ ⁺	Cryog. det.	65	ARNABOLDI	03
0.61 ± 0.14 ^{+0.29} _{-0.35}	130Te	2ν		Cryog. det.	66	ARNABOLDI	03
> 110	90 128Te	0ν		Cryog. det.	67	ARNABOLDI	03
(0.029 ^{+0.004} _{-0.003})	116Cd	2ν		116CdWO ₄ scint.	68	DANEVICH	03
> 170	90 116Cd	0ν		116CdWO ₄ scint.	69	DANEVICH	03
> 29	90 116Cd	0ν	0 ⁺ → 2 ⁺	116CdWO ₄ scint.	70	DANEVICH	03
> 14	90 116Cd	0ν	0 ⁺ → 0 ₁ ⁺	116CdWO ₄ scint.	71	DANEVICH	03
> 6	90 116Cd	0ν	0 ⁺ → 0 ₂ ⁺	116CdWO ₄ scint.	72	DANEVICH	03
> 1.1	90 186W	0ν		CdWO ₄ scint.	73	DANEVICH	03
> 1.1	90 186W	0ν	0 ⁺ → 2 ⁺	CdWO ₄ scint.	74	DANEVICH	03

$1.74 \pm 0.01^{+0.18}_{-0.16}$	^{76}Ge	2ν	Enriched HPGe	$^{75}\text{DOERR}$	03
>15700	$^{90}\text{^{76}Ge}$	0ν	Enriched HPGe	$^{76}\text{AALSETH}$	02B
> 58	$^{90}\text{^{134}Xe}$	0ν	Liquid Xe Scint.	$^{77}\text{BERNABEI}$	02D
> 1200	$^{90}\text{^{136}Xe}$	0ν	Liquid Xe Scint.	$^{78}\text{BERNABEI}$	02D
> 4.9	$^{90}\text{^{100}Mo}$	0ν	Liq. Ar ioniz.	$^{79}\text{ASHITKOV}$	01
> 1.3	$^{90}\text{^{160}Gd}$	0ν	$\text{Gd}_2\text{SiO}_5\text{:Ce}$	$^{80}\text{DANEVICH}$	01
> 1.3	$^{90}\text{^{160}Gd}$	0ν	$0^+ \rightarrow 2^+$ $\text{Gd}_2\text{SiO}_5\text{:Ce}$	$^{81}\text{DANEVICH}$	01
$0.59^{+0.17}_{-0.11} \pm 0.06$	^{100}Mo	$0\nu+2\nu$	$0^+ \rightarrow 0^+_1$ Ge coinc.	$^{82}\text{DEBRAECKEL}$	01
> 19000	$^{90}\text{^{76}Ge}$	0ν	Enriched HPGe	$^{83}\text{KLAPDOR-K...}$	01
$(9.4 \pm 3.2)\text{E-3}$	^{96}Zr	$0\nu+2\nu$	Geochem	$^{84}\text{WIESER}$	01
$0.042^{+0.033}_{-0.013}$	^{48}Ca	2ν	Ge spectrometer	$^{85}\text{BRUDANIN}$	00
$0.021^{+0.008}_{-0.004} \pm 0.002$	^{96}Zr	2ν	NEMO-2	$^{86}\text{ARNOLD}$	99
$(8.3 \pm 1.0 \pm 0.7)\text{E-2}$	^{82}Se	2ν	NEMO-2	$^{87}\text{ARNOLD}$	98
> 2.8	$^{90}\text{^{82}Se}$	0ν	$0^+ \rightarrow 2^+$ NEMO-2	$^{88}\text{ARNOLD}$	98
$(7.6^{+2.2}_{-1.4})\text{E-3}$	^{100}Mo	2ν	Si(Li)	$^{89}\text{ALSTON-...}$	97
$(6.82^{+0.38}_{-0.53} \pm 0.68)\text{E-3}$	^{100}Mo	2ν	TPC	$^{90}\text{DESILVA}$	97
$(6.75^{+0.37}_{-0.42} \pm 0.68)\text{E-3}$	^{150}Nd	2ν	TPC	$^{91}\text{DESILVA}$	97
$(3.75 \pm 0.35 \pm 0.21)\text{E-2}$	^{116}Cd	2ν	$0^+ \rightarrow 0^+$ NEMO 2	$^{92}\text{ARNOLD}$	96
$0.043^{+0.024}_{-0.011} \pm 0.014$	^{48}Ca	2ν	TPC	$^{93}\text{BALYSH}$	96
0.79 ± 0.10	^{130}Te	$0\nu+2\nu$	Geochem	$^{94}\text{TAKAOKA}$	96
$0.61^{+0.18}_{-0.11}$	^{100}Mo	$0\nu+2\nu$	$0^+ \rightarrow 0^+_1$ γ in HPGe	$^{95}\text{BARABASH}$	95
$0.026^{+0.009}_{-0.005}$	^{116}Cd	2ν	$0^+ \rightarrow 0^+$ ELEGANT IV	EJIRI	95
$0.017^{+0.010}_{-0.005} \pm 0.0035$	^{150}Nd	2ν	$0^+ \rightarrow 0^+$ TPC	ARTEMEV	93
0.039 ± 0.009	^{96}Zr	$0\nu+2\nu$	Geochem	KAWASHIMA	93
2.7 ± 0.1	^{130}Te	$0\nu+2\nu$	Geochem	BERNATOW...	92
7200 ± 400	^{128}Te	$0\nu+2\nu$	Geochem	$^{96}\text{BERNATOW...}$	92
$0.108^{+0.026}_{-0.006}$	^{82}Se	2ν	$0^+ \rightarrow 0^+$ TPC	ELLIOTT	92
2.0 ± 0.6	^{238}U	$0\nu+2\nu$	Radiochem	$^{97}\text{TURKEVICH}$	91
$0.12 \pm 0.01 \pm 0.04$	^{82}Se	$0\nu+2\nu$	Geochem.	^{98}LIN	88
$0.75 \pm 0.03 \pm 0.23$	^{130}Te	$0\nu+2\nu$	Geochem.	^{99}LIN	88
1800 ± 700	^{128}Te	$0\nu+2\nu$	Geochem.	^{100}LIN	88B
2.60 ± 0.28	^{130}Te	$0\nu+2\nu$	Geochem	$^{101}\text{KIRSTEN}$	83

¹ ALBERT 14 use the EXO-200 tracking detector for a re-measurement of the $2\nu\beta\beta$ -half life of ^{136}Xe . A nuclear matrix element of $0.0218 \pm 0.0003 \text{ MeV}^{-1}$ is derived from this data. Supersedes ACKERMAN 11.

² ALBERT 14B use 100 kg yr of exposure of the EXO-200 tracking calorimeter to place a lower limit on the $0\nu\beta\beta$ -half life of ^{136}Xe . Supersedes AUGER 12.

³ ARNOLD 14 use 34.7 kg yr of exposure of the NEMO-3 tracking calorimeter to derive a limit on the $\langle m \rangle$ -driven (light neutrino mass) $0\nu\beta\beta$ -half life of ^{100}Mo . Supersedes BARABASH 11A.

⁴ ARNOLD 14 use 34.7 kg yr of exposure of the NEMO-3 tracking calorimeter to derive a limit on the $\langle \lambda \rangle$ -driven (right handed quark and lepton currents) $0\nu\beta\beta$ -half life of ^{100}Mo .

⁵ ARNOLD 14 use 34.7 kg yr of exposure of the NEMO-3 tracking calorimeter to derive a limit on the $\langle \eta \rangle$ -driven (right handed quark current) $0\nu\beta\beta$ -half life of ^{100}Mo .

- 6 KIDD 14 utilize two underground Ge detectors to determine the inclusive double beta decay rate to the first excited 0_1^+ state using γ - γ coincidences.
- 7 AGOSTINI 13 use 5.04 kg yr of data collected with bare enriched Ge diodes operated in a liquid argon shield, to determine the half life for the $0\nu\beta\beta$ decay of ^{76}Ge . This result is in agreement, and more accurate, than DOERR 03.
- 8 AGOSTINI 13A use 21.6 kg yr of data, collected with GERDA detector array, to place a lower limit on the $0\nu\beta\beta$ -half life of ^{76}Ge . This result is in tension with the evidence for $0\nu\beta\beta$ -decay reported in KLAPDOR-KLEINGROTHAUS 06A. This half-life limit exceeds the limit reported in KLAPDOR-KLEINGROTHAUS 01.
- 9 BELLI 13A use an underground Ge detector to search for the $2\beta^+$ -decay of ^{96}Ru via the intensity of the annihilation peak. This method cannot distinguish two from zero neutrino decay.
- 10 BELLI 13A use an underground Ge detector to search for the $\text{EC}\beta^+$ -decay of ^{96}Ru via the intensity of the 778 keV γ de-excitation peak. This method cannot distinguish two from zero neutrino decay. The same analysis provides several limits, ($\sim 0.1 - 0.3$) 10^{21} years, for the $\beta^+\text{EC}$ mode leading to the excited 0^+ and 2^+ states.
- 11 BELLI 13A use an underground Ge detector to search for the $\beta\beta$ -decay of ^{104}Ru via the intensity of the 556 keV γ de-excitation peak. This method cannot distinguish two from zero neutrino decay.
- 12 GANDO 13A use the KamLAND detector to search for $0\nu\beta\beta$ -decay of ^{136}Xe based on an exposure of 89.5 kg yr. This result is in tension with the evidence of $0\nu\beta\beta$ reported in KLAPDOR-KLEINGROTHAUS 06A and earlier references to that work. Supersedes GANDO 12A and is more sensitive than BERNABEI 02D.
- 13 GAVRILYAK 13 use a proportional counter filled with Kr gas to search for the $2\nu 2\text{K}$ decay of ^{78}Kr . Data with the enriched and depleted Kr were used to determine signal and background. A 2.5σ excess of events obtained with the enriched sample is interpreted as an indication for the presence of this decay.
- 14 GAVRILYAK 13 use a proportional counter filled with Kr gas to search for the $0\nu 2\text{K}$ decay of ^{78}Kr into 2828 keV excited state of ^{78}Se . This transition could be subject to resonant rate enhancement. Data obtained with the enriched and depleted Kr were used to determine signal and background.
- 15 ANDREOTTI 12 use high resolution TeO_2 bolometric calorimeter to search for the $0\nu\beta\beta$ decay of ^{130}Te leading to the excited 0_1^+ state at 1793.5 keV.
- 16 BELLI 12A use $^{106}\text{CdWO}_4$ 215 g crystal scintillator to search for various $\beta\beta$ decay modes. The limit for the ECEC mode is derived from the fit to the background spectrum in the 1.8–3.2 MeV energy interval in the run of 6590 hours. The same analysis provides several limits ($\sim 2\text{--}5 \times 10^{20}$ years) for the ECEC mode leading to the excited 0^+ and 2^+ states. Also a similar size limits for the possible resonance process populating states at 2718 keV, 2741 keV, and 2748 keV were obtained.
- 17 BELLI 12A use $^{106}\text{CdWO}_4$ 215 g crystal scintillator to search for various $\beta\beta$ decay modes. The limit for the $\text{EC}\beta^+$ mode is derived from the fit to the background spectrum in the 2.0–3.0 MeV energy interval in the run of 6590 hours. The same analysis provides several limits ($\sim 0.5\text{--}1.3 \times 10^{21}$ years) for the $\text{EC}\beta^+$ mode leading to the excited 0^+ and 2^+ states.
- 18 BELLI 12A use $^{106}\text{CdWO}_4$ 215 g crystal scintillator to search for various $\beta\beta$ decay modes. The limit for the $\beta^+\beta^+$ mode is derived from the fit to the background spectrum in the 0.76–2.8 MeV energy interval in the run of 6590 hours. The same analysis provides the limit (1.2×10^{21} years) for the $\beta^+\beta^+$ mode leading to the first excited 2^+ state.
- 19 GANDO 12A use a modification of the existing KamLAND detector. The $\beta\beta$ decay source/detector is 13 tons of enriched ^{136}Xe -loaded scintillator contained in an inner balloon. The $2\nu\beta\beta$ decay rate is derived from the fit to the spectrum between 0.5 and 4.8 MeV. This result is in agreement with ACKERMAN 11.

- 20 GANDO 12A use a modification of the existing KamLAND detector. The $\beta\beta$ decay source/detector is 13 tons of enriched ^{136}Xe -loaded scintillator contained in an inner balloon. The $0\nu\beta\beta$ decay rate is derived from the fit where the background rates were allowed to float. Superseded by GANDO 13A.
- 21 ACKERMAN 11 use the EXO-200 liquid Xe TPC filled with ~ 175 kg of enriched ^{136}Xe to determine the 2ν half-life of ^{136}Xe . Superseded by ALBERT 14.
- 22 ARNOLD 11 use enriched ^{130}Te in the NEMO-3 detector to measure the $2\nu\beta\beta$ decay rate. This result is in agreement with, but more accurate than ARNABOLDI 03.
- 23 ARNOLD 11 use the NEMO-3 detector to obtain a limit for the $0\nu\beta\beta$ decay. This result is less significant than ARNABOLDI 05.
- 24 BARABASH 11 use 100 g of enriched ^{112}Sn to determine a limit for the ECEC 0ν decay to the 0_3^+ state of ^{112}Cd by searching for the de-excitation γ with a Ge detector. This decay mode is a candidate for resonant rate enhancement.
- 25 BARABASH 11 use 100 g of enriched ^{112}Sn to determine a limit for the ECEC 0ν decay to the 0_2^+ state of ^{112}Cd by searching for the de-excitation γ with a Ge detector.
- 26 BARABASH 11 use 100 g of enriched ^{112}Sn to determine a limit for the ECEC 0ν decay to the 0_1^+ state of ^{112}Cd by searching for the de-excitation γ with a Ge detector.
- 27 BARABASH 11 use 100 g of enriched ^{112}Sn to determine a limit for the ECEC 0ν decay to the ground state of ^{112}Cd by searching for the de-excitation γ with a Ge detector.
- 28 Supersedes DANEVICH 03 and ARNOLD 96.
- 29 Supersedes BRUDANIN 00 and BALYSH 96.
- 30 BARABASH 11A use the NEMO-3 detector to measure $\beta\beta 2\nu$ rates and place limits on $\beta\beta 0\nu$ half lives for various nuclides.
- 31 Supersedes ARNABOLDI 03.
- 32 Supersedes ARNOLD 05A, ARNOLD 04, ARNOLD 98, and ELLIOTT 92.
- 33 Less restrictive than ARNABOLDI 08.
- 34 Less restrictive than DANEVICH 03.
- 35 Less restrictive than UMEHARA 08 and OGAWA 04.
- 36 BELLI 11D use ZnWO_4 scintillator calorimeters to search for various $\beta\beta$ decay modes of ^{64}Zn , ^{70}Zn , ^{180}W , and ^{186}W .
- 37 RUKHADZE 11 uses 13.6 g of enriched ^{106}Cd to search for the neutrinoless ECEC decay into an excited state of ^{106}Pd and its characteristic γ -radiation using the TGV2 detector. This decay mode is a candidate for resonant rate enhancement, however, hindered by the large spin difference.
- 38 ARGYRIADES 10 use 9.4 ± 0.2 g of ^{96}Zr in NEMO-3 detector and identify its $2\nu\beta\beta$ decay. The result is in agreement and supersedes ARNOLD 99.
- 39 ARGYRIADES 10 use 9.4 ± 0.2 g of ^{96}Zr in NEMO-3 detector and obtain a limit of the $0\nu\beta\beta$ decay. The result is in agreement and supersedes ARNOLD 99.
- 40 ARGYRIADES 10 use 9.4 ± 0.2 g of ^{96}Zr in NEMO-3 detector and obtain a limit of the $0\nu\beta\beta$ decay into the first excited 0_1^+ state in ^{96}Mo .
- 41 BELLI 10 use enriched ^{100}Mo with 4 HP Ge detectors to record the 590.8 and 539.5 keV γ rays from the decay of the 0_1^+ state in ^{100}Ru both in singles and coincidences. This result confirms the measurement of KIDD 09 and ARNOLD 07 and supersedes them.
- 42 ARGYRIADES 09 use the NEMO-3 tracking calorimeter containing 36.5 g of ^{150}Nd , a total exposure of 924.7 days, to derive a limit for the $0\nu\beta\beta$ half-life. Supersedes DESILVA 97.
- 43 ARGYRIADES 09 use the NEMO-3 tracking calorimeter containing 36.5 g of ^{150}Nd , a total exposure of 924.7 days, to determine the value of the $2\nu\beta\beta$ half-life. This result is in marginal agreement, but has somewhat smaller error bars, than DESILVA 97.
- 44 BELLI 09A use ZnWO_4 scintillating crystals to search for various modes of $\beta\beta$ decay. This work improves the limits for different modes of ^{64}Zn decay into the ground state of ^{64}Ni , in this case for the $0\nu\beta^+\text{EC}$ mode. Supersedes BELLI 08.

- 45 BELLI 09A use ZnWO_4 scintillating crystals to search for various modes of $\beta\beta$ decay. This work improves the limits for different modes of ^{64}Zn decay into the ground state of ^{64}Ni , in this case for the 0ν ECEC mode. Supersedes BELLI 08.
- 46 KIDD 09 combine past and new data with an improved coincidence detection efficiency determination. The result agrees with ARNOLD 95. Supersedes DEBRAECKELEER 01 and BARABASH 95.
- 47 ARNABOLDI 08 use high resolution TeO_2 bolometer calorimeter to search for double beta decay of ^{130}Te . Supersedes ARNABOLDI 05.
- 48 BELLI 08 use ZnWO_4 scintillation calorimeter to search for neutrinoless β^+ plus electron capture decay of ^{64}Zn . The half-life limit for the 2ν mode is 2.1×10^{20} years.
- 49 BELLI 08B use CdWO_4 scintillation calorimeter to search for 0ν $\beta\beta$ decay of ^{114}Cd .
- 50 UMEHARA 08 use CaF_2 scintillation calorimeter to search for double beta decay of ^{48}Ca . Limit is significantly more stringent than quoted sensitivity: 18×10^{21} years.
- 51 First exclusive measurement of 2ν -decay to the first excited 0_1^+ -state of daughter nucleus. ARNOLD 07 use the NEMO-3 tracking calorimeter to detect all particles emitted in decay. Result agrees with the inclusive ($0\nu + 2\nu$) measurement of DEBRAECKELEER 01.
- 52 Limit on 0ν -decay to the first excited 0_1^+ -state of daughter nucleus using NEMO-3 tracking calorimeter. Supersedes DASSIE 95.
- 53 Limit on 0ν -decay to the first excited 2^+ -state of daughter nucleus using NEMO-3 tracking calorimeter.
- 54 BARABASH 07 use Ge calorimeter to search for γ -radiation following double electron capture or β^+ plus electron capture decays of ^{74}Se to the ground state of ^{74}Ge . This limit is based on the search for the 511 keV annihilation radiation. Various other limits, for the capture from different atomic shells and also to the excited states, are reported in the paper.
- 55 BARABASH 07 use Ge calorimeter to search for γ -radiation following double electron capture decay of ^{74}Se into the second excited 2^+ -state of ^{74}Ge . That transition has been considered due to a possible resonance enhancement. The 2ν mode would be suppressed for this decay by its extremely small phase space factor.
- 56 KLAPDOR-KLEINGROTHAUS 06A present re-analysis of data originally published in KLAPDOR-KLEINGROTHAUS 04A. Modified pulse shape analysis leads the authors to claim improved 6σ statistical evidence for observation of 0ν -decay, compared to 4.2σ in KLAPDOR-KLEINGROTHAUS 04A. Analysis of the systematic uncertainty is not presented. This re-analysis is disputed in AGOSTINI 13A and SCHWINGENHEUER 13.
- 57 Supersedes ARNABOLDI 04. Bolometric TeO_2 detector array CUORICINO is used for high resolution search for $0\nu\beta\beta$ decay. The half-life limit is derived from 3.09 kg yr ^{130}Te exposure.
- 58 NEMO-3 tracking calorimeter containing 6.9 kg of enriched ^{100}Mo is used in ARNOLD 05A. A limit for $0\nu\beta\beta$ half-life of ^{100}Mo is reported. Supersedes ARNOLD 04.
- 59 NEMO-3 tracking calorimeter is used in ARNOLD 05A to place limit on $0\nu\beta\beta$ half-life of ^{82}Se . Detector contains 0.93 kg of enriched ^{82}Se . Supersedes ARNOLD 04.
- 60 ARNOLD 05A use the NEMO-3 tracking calorimeter to determine the $2\nu\beta\beta$ half-life of ^{100}Mo with high statistics and low background (389 days of data taking). Supersedes ARNOLD 04.
- 61 ARNOLD 05A use the NEMO-3 tracking detector to determine the $2\nu\beta\beta$ half-life of ^{82}Se with high statistics and low background (389 days of data taking). Supersedes ARNOLD 04.
- 62 ARNOLD 04 use the NEMO-3 tracking detector to determine the limit for $0\nu\beta\beta$ half-life of ^{82}Se . This represents an improvement, by a factor of ~ 10 , when compared with ELLIOTT 92. It supersedes the limit of ARNOLD 98 for this decay using NEMO-2.
- 63 ARNOLD 04 use the NEMO-3 tracking detector to determine the $2\nu\beta\beta$ half-life of ^{100}Mo with high statistics and low background. The half-life is determined assuming the Single

- State Dominance. It is in agreement with, and more accurate than, previous determinations. Supersedes DASSIE 95 determination of this quantity with NEMO-2.
- 64 BARABASH 04 perform an inclusive measurement of the $\beta\beta$ decay of ^{150}Nd into the first excited (0_1^+) state of ^{150}Sm . Gamma radiation emitted in decay of the excited state is detected.
- 65 Decay into first excited state of daughter nucleus.
- 66 Two neutrino decay into ground state. Relatively large error mainly due to uncertainties in background determination. Reported value is shorter than the geochemical measurements of KIRSTEN 83 and BERNATOWICZ 92 but in agreement with LIN 88 and TAKAOKA 96.
- 67 Supersedes ALESSANDRELLO 00. Array of TeO_2 crystals in high resolution cryogenic calorimeter. Some enriched in ^{128}Te . Ground state to ground state decay.
- 68 Calorimetric measurement of 2ν ground state decay of ^{116}Cd using enriched CdWO_4 scintillators. Agrees with EJIRI 95 and ARNOLD 96. Supersedes DANEVICH 00.
- 69 Limit on 0ν decay of ^{116}Cd using enriched CdWO_4 scintillators. Supersedes DANEVICH 00.
- 70 Limit on 0ν decay of ^{116}Cd into first excited 2^+ state of daughter nucleus using enriched CdWO_4 scintillators. Supersedes DANEVICH 00.
- 71 Limit on 0ν decay of ^{116}Cd into first excited 0^+ state of daughter nucleus using enriched CdWO_4 scintillators. Supersedes DANEVICH 00.
- 72 Limit on 0ν decay of ^{116}Cd into second excited 0^+ state of daughter nucleus using enriched CdWO_4 scintillators. Supersedes DANEVICH 00.
- 73 Limit on the 0ν ground state decay of ^{186}W using enriched CdWO_4 scintillators.
- 74 Limit on the 0ν decay of ^{186}W to the first excited 2^+ state of the daughter nucleus using enriched CdWO_4 scintillators.
- 75 Results of the Heidelberg-Moscow experiment (KLAPDOR-KLEINGROTHAUS 01 and GUENTHER 97) are reanalyzed using a new simulation of the complete background spectrum. The $\beta\beta 2\nu$ -decay rate is deduced from a 41.57 kg-y exposure. The result is in agreement and supersedes the above referenced halflives with similar statistical and systematic errors.
- 76 AALSETH 02B limit is based on 117 mol-yr of data using enriched Ge detectors. Background reduction by means of pulse shape analysis is applied to part of the data set. Reported limit is slightly less restrictive than that in KLAPDOR-KLEINGROTHAUS 01. However, it excludes part of the allowed half-life range reported in KLAPDOR-KLEINGROTHAUS 01B for the same nuclide. The analysis has been criticized in KLAPDOR-KLEINGROTHAUS 04B. The criticism was addressed and disputed in AALSETH 04.
- 77 BERNABEI 02D report a limit for the $0\nu, 0^+ \rightarrow 0^+$ decay of ^{134}Xe , present in the source at 17%, by considering the maximum number of events for this mode compatible with the fitted smooth background.
- 78 BERNABEI 02D report a limit for the $0\nu, 0^+ \rightarrow 0^+$ decay of ^{136}Xe , by considering the maximum number of events for this mode compatible with the fitted smooth background. The quoted sensitivity is 450×10^{21} yr. The Feldman and Cousins method is used to obtain the quoted limit.
- 79 ASHITKOV 01 result for 0ν of ^{100}Mo is less stringent than EJIRI 01.
- 80 DANEVICH 01 place limit on 0ν decay of ^{160}Gd using $\text{Gd}_2\text{SiO}_5:\text{Ce}$ crystal scintillators. The limit is more stringent than KOBAYASHI 95.
- 81 DANEVICH 01 place limits on 0ν decay of ^{160}Gd into excited 2^+ state of daughter nucleus using $\text{Gd}_2\text{SiO}_5:\text{Ce}$ crystal scintillators.
- 82 DEBRAECKELEER 01 performed an inclusive measurement of the $\beta\beta$ decay into the second excited state of the daughter nucleus. A novel coincidence technique counting the de-excitation photons is employed. The result agrees with BARABASH 95.
- 83 KLAPDOR-KLEINGROTHAUS 01 is a continuation of the work published in BAUDIS 99. Isotopically enriched Ge detectors are used in calorimetric measurement. The most stringent bound is derived from the data set in which pulse-shape analysis has been used to

- reduce background. Exposure time is 35.5 kg y. Supersedes BAUDIS 99 as most stringent result.
- 84 WIESER 01 reports an inclusive geochemical measurement of ^{96}Zr $\beta\beta$ half life. Their result agrees within 2σ with ARNOLD 99 but only marginally, within 3σ , with KAWASHIMA 93.
- 85 BRUDANIN 00 determine the 2ν half-life of ^{48}Ca . Their value is less accurate than BALYSH 96.
- 86 ARNOLD 99 measure directly the 2ν decay of Zr for the first time, using the NEMO-2 tracking detector and an isotopically enriched source. The lifetime is more accurate than the geochemical result of KAWASHIMA 93.
- 87 ARNOLD 98 measure the 2ν decay of ^{82}Se by comparing the spectra in an enriched and natural selenium source using the NEMO-2 tracking detector. The measured half-life is in agreement, perhaps slightly shorter, than ELLIOTT 92.
- 88 ARNOLD 98 determine the limit for 0ν decay to the excited 2^+ state of ^{82}Se using the NEMO-2 tracking detector.
- 89 ALSTON-GARNJOST 97 report evidence for 2ν decay of ^{100}Mo . This decay has been also observed by EJIRI 91, DASSIE 95, and DESILVA 97.
- 90 DESILVA 97 result for 2ν decay of ^{100}Mo is in agreement with ALSTON-GARNJOST 97 and DASSIE 95. This measurement has the smallest errors.
- 91 DESILVA 97 result for 2ν decay of ^{150}Nd is in marginal agreement with ARTEMEV 93. It has smaller errors.
- 92 ARNOLD 96 measure the 2ν decay of ^{116}Cd . This result is in agreement with EJIRI 95, but has smaller errors. Supersedes ARNOLD 95.
- 93 BALYSH 96 measure the 2ν decay of ^{48}Ca , using a passive source of enriched ^{48}Ca in a TPC.
- 94 TAKAOKA 96 measure the geochemical half-life of ^{130}Te . Their value is in disagreement with the quoted values of BERNATOWICZ 92 and KIRSTEN 83; but agrees with several other unquoted determinations, e.g., MANUEL 91.
- 95 BARABASH 95 cannot distinguish 0ν and 2ν , but it is inferred indirectly that the 0ν mode accounts for less than 0.026% of their event sample. They also note that their result disagrees with the previous experiment by the NEMO group (BLUM 92).
- 96 BERNATOWICZ 92 finds $^{128}\text{Te}/^{130}\text{Te}$ activity ratio from slope of $^{128}\text{Xe}/^{132}\text{Xe}$ vs $^{130}\text{Xe}/^{132}\text{Xe}$ ratios during extraction, and normalizes to lead-dated ages for the ^{130}Te lifetime. The authors state that their results imply that "(a) the double beta decay of ^{128}Te has been firmly established and its half-life has been determined ... without any ambiguity due to trapped Xe interferences... (b) Theoretical calculations ... underestimate the [long half-lives of ^{128}Te ^{130}Te] by 1 or 2 orders of magnitude, pointing to a real suppression in the 2ν decay rate of these isotopes. (c) Despite [this], most $\beta\beta$ -models predict a *ratio* of 2ν decay widths ... in fair agreement with observation." Further details of the experiment are given in BERNATOWICZ 93. Our listed half-life has been revised downward from the published value by the authors, on the basis of reevaluated cosmic-ray ^{128}Xe production corrections.
- 97 TURKEVICH 91 observes activity in old U sample. The authors compare their results with theoretical calculations. They state "Using the phase-space factors of Boehm and Vogel (BOEHM 87) leads to matrix element values for the ^{238}U transition in the same range as deduced for ^{130}Te and ^{76}Ge . On the other hand, the latest theoretical estimates (STAUDT 90) give an upper limit that is 10 times lower. This large discrepancy implies either a defect in the calculations or the presence of a faster path than the standard two-neutrino mode in this case." See BOEHM 87 and STAUDT 90.
- 98 Result agrees with direct determination of ELLIOTT 92.
- 99 Inclusive half life inferred from mass spectroscopic determination of abundance of $\beta\beta$ -decay product ^{130}Te in mineral kitkaite (NiTeSe). Systematic uncertainty reflects variations in U-Xe gas-retention-age derived from different uranite samples. Agrees with geochemical determination of TAKAOKA 96 and direct measurement of ARNABOLDI 03. Inconsistent with results of KIRSTEN 83 and BERNATOWICZ 92.

- ¹⁰⁰ Ratio of inclusive double beta half lives of ¹²⁸Te and ¹³⁰Te determined from minerals melonite (NiTe₂) and altaite (PbTe) by means of mass spectroscopic measurement of abundance of $\beta\beta$ -decay products. As gas-retention-age could not be determined the authors use half life of ¹³⁰Te (LIN 88) to infer the half life of ¹²⁸Te. No estimate of the systematic uncertainty of this method is given. The directly determined half life ratio agrees with BERNATOWICZ 92. However, the inferred ¹²⁸Te half life disagrees with KIRSTEN 83 and BERNATOWICZ 92.
- ¹⁰¹ KIRSTEN 83 reports "2 σ " error. References are given to earlier determinations of the ¹³⁰Te lifetime.

$\langle m_\nu \rangle$, The Effective Weighted Sum of Majorana Neutrino Masses Contributing to Neutrinoless Double- β Decay

$\langle m_\nu \rangle = |\sum U_{1j}^2 \cdot m_{\nu_j}|$, where the sum goes from 1 to n and where n = number of neutrino generations, and ν_j is a Majorana neutrino. Note that U_{ej}^2 , not $|U_{ej}|^2$, occurs in the sum. The possibility of cancellations has been stressed. In the following Listings, only best or comparable limits or lifetimes for each isotope are reported.

VALUE (eV)	CL%	ISOTOPE	TRANSITION	METHOD	DOCUMENT ID
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 0.19–0.45	90	¹³⁶ Xe	0 ν ,g.s.→ g.s.	EXO-200	1 ALBERT 14B
< 0.33–0.87	90	¹⁰⁰ Mo	0 ν	NEMO-3	2 ARNOLD 14
< 0.2–0.4	90	⁷⁶ Ge	0 ν	GERDA	3 AGOSTINI 13A
< 0.12–0.25	90	¹³⁶ Xe	0 ν ,g.s.→ g.s.	KamLAND-Zen	4 GANDO 13A
< 0.3–0.6	90	¹³⁶ Xe	0 ν ,g.s.→ g.s.	KamLAND-Zen	5 GANDO 12A
< 0.89–2.43	90	⁸² Se	0 ν	NEMO-3	6 BARABASH 11A
< 7.2–19.5	90	⁹⁶ Zr	0 ν	NEMO-3	7 ARGYRIADES 10
< 4.0–6.8	90	¹⁵⁰ Nd	0 ν	NEMO-3	8 ARGYRIADES 09
< 0.19–0.68	90	¹³⁰ Te	0 ν	TeO ₂ bolometer	9 ARNABOLDI 08
< 3.5–22	90	⁴⁸ Ca	0 ν	CaF ₂ scint.	10 UMEHARA 08
< 9.3–60	90	¹⁰⁰ Mo	0 ⁺ → 0 ₁ ⁺	NEMO-3	11 ARNOLD 07
< 6500	90	¹⁰⁰ Mo	0 ⁺ → 2 ⁺	NEMO-3	12 ARNOLD 07
0.32±0.03	68	⁷⁶ Ge	0 ν	Enriched HPGe	13 KLAPDOR-K...06A
< 0.2–1.1	90	¹³⁰ Te		Cryog. det.	14 ARNABOLDI 05
< 0.7–2.8	90	¹⁰⁰ Mo	0 ν	NEMO-3	15 ARNOLD 05A
< 1.7–4.9	90	⁸² Se	0 ν	NEMO-3	16 ARNOLD 05A
< 0.37–1.9	90	¹³⁰ Te		Cryog. det.	17 ARNABOLDI 04
< 0.8–1.2	90	¹⁰⁰ Mo	0 ν	NEMO-3	18 ARNOLD 04
< 1.5–3.1	90	⁸² Se	0 ν	NEMO-3	18 ARNOLD 04
0.1–0.9	99.7	⁷⁶ Ge		Enriched HP Ge	19 KLAPDOR-K...04A
< 7.2–44.7	90	⁴⁸ Ca		CaF ₂ scint.	20 OGAWA 04
< 1.1–2.6	90	¹³⁰ Te		Cryog. det.	21 ARNABOLDI 03
< 1.5–1.7	90	¹¹⁶ Cd	0 ν	¹¹⁶ CdWO ₄ scint.	22 DANEVICH 03
< 0.33–1.35	90			Enriched HPGe	23 AALSETH 02B
< 2.9	90	¹³⁶ Xe	0 ν	Liquid Xe Scint.	24 BERNABEI 02D
0.39 ^{+0.17} _{-0.28}		⁷⁶ Ge	0 ν	Enriched HPGe	25 KLAPDOR-K...02D
< 2.1–4.8	90	¹⁰⁰ Mo	0 ν	ELEGANT V	26 EJIRI 01
< 0.35	90	⁷⁶ Ge		Enriched HPGe	27 KLAPDOR-K...01

<23	90 ⁹⁶ Zr		NEMO-2	28 ARNOLD	99
< 1.1–1.5	128 ^{Te}		Geochem	29 BERNATOW...	92
<5	68 ⁸² Se		TPC	30 ELLIOTT	92
<8.3	76 ⁴⁸ Ca	0ν	CaF ₂ scint.	YOU	91

- ¹ ALBERT 14B is based on 100 kg yr of exposure of the EXO-200 tracking calorimeter. The mass range reflects the nuclear matrix element calculations. Supersedes AUGER 12.
- ² ARNOLD 14 is based on 34.7 kg yr of exposure of the NEMO-3 tracking calorimeter. The reported mass range reflects the nuclear matrix element uncertainty in ¹⁰⁰Mo. Supersedes BARABASH 11A.
- ³ AGOSTINI 13A is based on 21.6 kg yr of data collected by the GERDA detector. The reported range reflects different nuclear matrix elements. This result is in tension with the evidence for $0\nu\beta\beta$ -decay reported in KLAPDOR-KLEINGROTHAUS 06A and earlier references to that work.
- ⁴ GANDO 13A limit is based on a combination of KamLAND-Zen and EXO-200 (AUGER 12) data. The reported range reflects different nuclear matrix elements. Supersedes GANDO 12A.
- ⁵ GANDO 12A limit is based on the KamLAND-Zen data. The reported range reflects different nuclear matrix elements. Superseded by GANDO 13A.
- ⁶ BARABASH 11A limit is based on NEMO-3 data for ⁸²Se. The reported range reflects different nuclear matrix elements. Supersedes ARNOLD 05A and ARNOLD 04.
- ⁷ ARGYRIADES 10 use ⁹⁶Zr and the NEMO-3 tracking detector to obtain the reported mass limit. The range reflects the fluctuation of the nuclear matrix elements considered.
- ⁸ ARGYRIADES 09 limit is based on data taken with the NEMO-3 detector and ¹⁵⁰Nd. A range of nuclear matrix elements that include the effect of nuclear deformation have been used.
- ⁹ Limit was obtained using high resolution TeO₂ bolometer calorimeter to search for double beta decay of ¹³⁰Te. Reported range of limits reflects spread of matrix element calculations used. Supersedes ARNABOLDI 05.
- ¹⁰ Limit was obtained using CaF₂ scintillation calorimeter to search for double beta decay of ⁴⁸Ca. Reported range of limits reflects spread of QRPA and SM matrix element calculations used. Supersedes OGAWA 04.
- ¹¹ ARNOLD 07 use NEMO-3 half life limit for 0ν -decay of ¹⁰⁰Mo to the first excited 0_1^+ -state of daughter nucleus to obtain neutrino mass limit. The spread reflects the choice of two different nuclear matrix elements. This limit is not competitive when compared to the decay to the ground state.
- ¹² ARNOLD 07 use NEMO-3 half life limit for 0ν -decay of ¹⁰⁰Mo to the first excited 2^+ -state of daughter nucleus to obtain neutrino mass limit. This limit is not competitive when compared to the decay to the ground state.
- ¹³ Re-analysis of data originally published in KLAPDOR-KLEINGROTHAUS 04A. Modified pulse shape analysis leads the authors to claim 6σ statistical evidence for observation of 0ν -decay. Authors use matrix element of STAUDT 90. Uncertainty of nuclear matrix element is not reflected in stated error. Supersedes KLAPDOR-KLEINGROTHAUS 04A.
- ¹⁴ Supersedes ARNABOLDI 04. Reported range of limits due to use of different nuclear matrix element calculations.
- ¹⁵ Mass limits reported in ARNOLD 05A are derived from ¹⁰⁰Mo data, obtained by the NEMO-3 collaboration. The range reflects the spread of matrix element calculations considered in this work. Supersedes ARNOLD 04.
- ¹⁶ Neutrino mass limits based on ⁸²Se data utilizing the NEMO-3 detector. The range reported in ARNOLD 05A reflects the spread of matrix element calculations considered in this work. Supersedes ARNOLD 04.
- ¹⁷ Supersedes ARNABOLDI 03. Reported range of limits due to use of different nuclear matrix element calculations.
- ¹⁸ ARNOLD 04 limit is based on the nuclear matrix elements of SIMKOVIC 99, STOICA 01 and CIVITARESE 03.

- 19 Supersedes KLAPDOR-KLEINGROTHAUS 02D. Event excess at $\beta\beta$ -decay energy is used to derive Majorana neutrino mass using the nuclear matrix elements of STAUDT 90. The mass range shown is based on the authors evaluation of the uncertainties of the STAUDT 90 matrix element calculation. If this uncertainty is neglected, and only statistical errors are considered, the range in $\langle m \rangle$ becomes (0.2–0.6) eV at the 3σ level.
- 20 Calorimetric CaF_2 scintillator. Range of limits reflects authors' estimate of the uncertainty of the nuclear matrix elements. Replaces YOU 91 as the most stringent limit based on ^{48}Ca .
- 21 Supersedes ALESSANDRELLO 00. Cryogenic calorimeter search. Reported a range reflecting uncertainty in nuclear matrix element calculations.
- 22 Limit for $\langle m_{\nu} \rangle$ is based on the nuclear matrix elements of STAUDT 90 and ARNOLD 96. Supersedes DANEVICH 00.
- 23 AALSETH 02B reported range of limits on $\langle m_{\nu} \rangle$ reflects the spread of theoretical nuclear matrix elements. Excludes part of allowed mass range reported in KLAPDOR-KLEINGROTHAUS 01B.
- 24 BERNABEI 02D limit is based on the matrix elements of SIMKOVIC 02. The range of neutrino masses based on a variety of matrix elements is 1.1–2.9 eV.
- 25 KLAPDOR-KLEINGROTHAUS 02D is a detailed description of the analysis of the data collected by the Heidelberg-Moscow experiment, previously presented in KLAPDOR-KLEINGROTHAUS 01B. Matrix elements in STAUDT 90 have been used. See the footnote in the preceding table for further details. See also KLAPDOR-KLEINGROTHAUS 02B.
- 26 The range of the reported $\langle m_{\nu} \rangle$ values reflects the spread of the nuclear matrix elements. On axis value assuming $\langle \lambda \rangle = \langle \eta \rangle = 0$.
- 27 KLAPDOR-KLEINGROTHAUS 01 uses the calculation by STAUDT 90. Using several other models in the literature could worsen the limit up to 1.2 eV. This is the most stringent experimental bound on m_{ν} . It supersedes BAUDIS 99B.
- 28 ARNOLD 99 limit based on the nuclear matrix elements of STAUDT 90.
- 29 BERNATOWICZ 92 finds these majorana neutrino mass limits assuming that the measured geochemical decay width is a limit on the 0ν decay width. The range is the range found using matrix elements from HAXTON 84, TOMODA 87, and SUHONEN 91. Further details of the experiment are given in BERNATOWICZ 93.
- 30 ELLIOTT 92 uses the matrix elements of HAXTON 84.

Limits on Lepton-Number Violating (V+A) Current Admixture

For reasons given in the discussion at the beginning of this section, we list only results from 1989 and later. $\langle \lambda \rangle = \lambda \sum U_{ej} V_{ej}$ and $\langle \eta \rangle = \eta \sum U_{ej} V_{ej}$, where the sum is over the number of neutrino generations. This sum vanishes for massless or unmixed neutrinos. In the following Listings, only best or comparable limits or lifetimes for each isotope are reported.

$\langle \lambda \rangle$ (10^{-6})	CL%	$\langle \eta \rangle$ (10^{-8})	CL%	ISOTOPE	METHOD	DOCUMENT ID
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
< 0.9–1.3	90	< 0.5–0.8	90	^{100}Mo	NEMO-3	1 ARNOLD 14
< 120	90			^{100}Mo	$0^+ \rightarrow 2^+$	2 ARNOLD 07
$0.692^{+0.058}_{-0.056}$	68	$0.305^{+0.026}_{-0.025}$	68	^{76}Ge	Enriched HPGe	3 KLAPDOR-K...06A
< 2.5	90			^{100}Mo	0ν , NEMO-3	4 ARNOLD 05A
< 3.8	90			^{82}Se	0ν , NEMO-3	5 ARNOLD 05A
< 1.5–2.0	90			^{100}Mo	0ν , NEMO-3	6 ARNOLD 04
< 3.2–3.8	90			^{82}Se	0ν , NEMO-3	7 ARNOLD 04

< 1.6–2.4	90	< 0.9–5.3	90	¹³⁰ Te	Cryog. det.	⁸ ARNABOLDI	03
< 2.2	90	<2.5	90	¹¹⁶ Cd	¹¹⁶ CdWO ₄ scint.	⁹ DANEVICH	03
< 3.2–4.7	90	< 2.4–2.7	90	¹⁰⁰ Mo	ELEGANT V	¹⁰ EJIRI	01
< 1.1	90	<0.64	90	⁷⁶ Ge	Enriched HPGe	¹¹ GUENTHER	97
< 4.4	90	<2.3	90	¹³⁶ Xe	TPC	¹² VUILLEUMIER	93
		<5.3		¹²⁸ Te	Geochem	¹³ BERNATOW...	92

¹ ARNOLD 14 is based on 34.7 kg yr of exposure of the NEMO-3 tracking calorimeter. The reported range limit on $\langle\lambda\rangle$ and $\langle\eta\rangle$ reflects the nuclear matrix element uncertainty in ¹⁰⁰Mo.

² ARNOLD 07 use NEMO-3 half life limit for 0ν -decay of ¹⁰⁰Mo to the first excited 2⁺-state of daughter nucleus to limit the right-right handed admixture of weak currents $\langle\lambda\rangle$. This limit is not competitive when compared to the decay to the ground state.

³ Re-analysis of data originally published in KLAPDOR-KLEINGROTHAUS 04A. Modified pulse shape analysis leads the authors to claim 6 σ statistical evidence for observation of 0ν -decay. Authors use matrix element of MUTO 89 to determine $\langle\lambda\rangle$ and $\langle\eta\rangle$. Uncertainty of nuclear matrix element is not reflected in stated errors.

⁴ ARNOLD 05A derive limit for $\langle\lambda\rangle$ based on ¹⁰⁰Mo data collected with NEMO-3 detector. No limit for $\langle\eta\rangle$ is given. Supersedes ARNOLD 04.

⁵ ARNOLD 05A derive limit for $\langle\lambda\rangle$ based on ⁸²Se data collected with NEMO-3 detector. No limit for $\langle\eta\rangle$ is given. Supersedes ARNOLD 04.

⁶ ARNOLD 04 use the matrix elements of SUHONEN 94 to obtain a limit for $\langle\lambda\rangle$, no limit for $\langle\eta\rangle$ is given. This limit is more stringent than the limit in EJIRI 01 for the same nucleus.

⁷ ARNOLD 04 use the matrix elements of TOMODA 91 and SUHONEN 91 to obtain a limit for $\langle\lambda\rangle$, no limit for $\langle\eta\rangle$ is given.

⁸ Supersedes ALESSANDRELLO 00. Cryogenic calorimeter search. Reported a range reflecting uncertainty in nuclear matrix element calculations.

⁹ Limits for $\langle\lambda\rangle$ and $\langle\eta\rangle$ are based on nuclear matrix elements of STAUDT 90. Supersedes DANEVICH 00.

¹⁰ The range of the reported $\langle\lambda\rangle$ and $\langle\eta\rangle$ values reflects the spread of the nuclear matrix elements. On axis value assuming $\langle m_\nu \rangle = 0$ and $\langle\lambda\rangle = \langle\eta\rangle = 0$, respectively.

¹¹ GUENTHER 97 limits use the matrix elements of STAUDT 90. Supersedes BALYSH 95 and BALYSH 92.

¹² VUILLEUMIER 93 uses the matrix elements of MUTO 89. Based on a half-life limit 2.6×10^{23} y at 90%CL.

¹³ BERNATOWICZ 92 takes the measured geochemical decay width as a limit on the 0ν width, and uses the SUHONEN 91 coefficients to obtain the least restrictive limit on η . Further details of the experiment are given in BERNATOWICZ 93.

Double- β Decay REFERENCES

ALBERT	14	PR C89 015502	J. Albert <i>et al.</i>	(EXO-200 Collab.)
ALBERT	14B	NAT 510 229	J.B. Albert <i>et al.</i>	(EXO-200 Collab.)
ARNOLD	14	PR D89 111101	R. Arnold <i>et al.</i>	(NEMO-3 Collab.)
KIDD	14	PR C90 055501	M.F. Kidd <i>et al.</i>	
AGOSTINI	13	JP G40 035110	M. Agostini <i>et al.</i>	(GERDA Collab.)
AGOSTINI	13A	PRL 111 122503	M. Agostini <i>et al.</i>	(GERDA Collab.)
BELLI	13A	PR C87 034607	P. Belli <i>et al.</i>	(DAMA-INR Collab.)
GANDO	13A	PRL 110 062502	A. Gando <i>et al.</i>	(KamLAND-Zen Collab.)
GAVRILYAK	13	PR C87 035501	Yu.M. Gavriluk <i>et al.</i>	
SCHWINGENHEUER...	13	ANP 525 269	B. Schwingenheuer	(MPIH)
ANDREOTTI	12	PR C85 045503	E. Andreotti <i>et al.</i>	(CUORICINO Collab.)
AUGER	12	PRL 109 032505	M. Auger <i>et al.</i>	(EXO-200 Collab.)
BELLI	12A	PR C85 044610	P. Belli <i>et al.</i>	
GANDO	12A	PR C85 045504	A. Gando <i>et al.</i>	(KamLAND-Zen Collab.)
ACKERMAN	11	PRL 107 212501	N. Ackerman <i>et al.</i>	(EXO Collab.)
ARNOLD	11	PRL 107 062504	R. Arnold <i>et al.</i>	(NEMO-3 Collab.)

BARABASH	11	PR C83 045503	A.S. Barabash <i>et al.</i>	
BARABASH	11A	PAN 74 312	A.S. Barabash <i>et al.</i>	(NEMO-3 Collab.)
		Translated from YAF 74 330.		
BELLI	11D	JP G38 115107	P. Belli <i>et al.</i>	(DAMA-INR Collab.)
RUKHADZE	11	NP A852 197	N.I. Rukhadze <i>et al.</i>	(TGV-2 Collab.)
ARGYRIADES	10	NP A847 168	J. Argyriades <i>et al.</i>	(NEMO-3 Collab.)
BELLI	10	NP A846 143	P. Belli <i>et al.</i>	(DAMA-INR Collab.)
ARGYRIADES	09	PR C80 032501	J. Argyriades <i>et al.</i>	(NEMO-3 Collab.)
BELLI	09A	NP A826 256	P. Belli <i>et al.</i>	(DAMA-INR Collab.)
KIDD	09	NP A821 251	M. Kidd <i>et al.</i>	
ARNABOLDI	08	PR C78 035502	C. Arnaboldi <i>et al.</i>	
BELLI	08	PL B658 193	P. Belli <i>et al.</i>	(DAMA-INR Collab.)
BELLI	08B	EPJ A36 167	P. Belli <i>et al.</i>	
UMEHARA	08	PR C78 058501	S. Umehara <i>et al.</i>	
ARNOLD	07	NP A781 209	R. Arnold <i>et al.</i>	(NEMO-3 Collab.)
BARABASH	07	NP A785 371	A.S. Barabash <i>et al.</i>	
KLAPDOR-K...	06A	MPL A21 1547	H.V. Klapdor-Kleingrothaus, I.V. Krivosheina	
ARNABOLDI	05	PRL 95 142501	C. Arnaboldi <i>et al.</i>	(CUORICINO Collab.)
ARNOLD	05A	PRL 95 182302	R. Arnold <i>et al.</i>	(NEMO-3 Collab.)
AALSETH	04	PR D70 078302	C.E. Aalseth <i>et al.</i>	
ARNABOLDI	04	PL B584 260	C. Arnaboldi <i>et al.</i>	
ARNOLD	04	JETPL 80 377	R. Arnold <i>et al.</i>	(NEMO3 Detector Collab.)
		Translated from ZETFP 80 429.		
BARABASH	04	JETPL 79 10	A.S. Barabash <i>et al.</i>	
KLAPDOR-K...	04A	PL B586 198	H.V. Klapdor-Kleingrothaus <i>et al.</i>	
KLAPDOR-K...	04B	PR D70 078301	H.V. Klapdor-Kleingrothaus, A. Dietz, I.V. Krivosheina	
OGAWA	04	NP A730 215	I. Ogawa <i>et al.</i>	
ARNABOLDI	03	PL B557 167	C. Arnaboldi <i>et al.</i>	
CIVITARESE	03	NP A729 867	O. Civitarese, J. Suhonen	
DANEVICH	03	PR C68 035501	F.A. Danevich <i>et al.</i>	
DOERR	03	NIM A513 596	C. Doerr, H.V. Klapdor-Kleingrothaus	
AALSETH	02B	PR D65 092007	C.E. Aalseth <i>et al.</i>	(IGEX Collab.)
BERNABEI	02D	PL B546 23	R. Bernabei <i>et al.</i>	(DAMA Collab.)
KLAPDOR-K...	02B	PPNL 110 57	H.V. Klapdor-Kleingrothaus, A. Dietz, I.V. Krivosheina	
KLAPDOR-K...	02D	FP 32 1181	H.V. Klapdor-Kleingrothaus, A. Dietz, I.V. Krivosheina	
SIMKOVIC	02	hep-ph/0204278	F. Simkovic, P. Domin, A. Faessler	
ASHITKOV	01	JETPL 74 529	V.D. Ashitkov <i>et al.</i>	
		Translated from ZETFP 74 601.		
DANEVICH	01	NP A694 375	F.A. Danevich <i>et al.</i>	
DEBRAECKEL...	01	PRL 86 3510	L. De Braeckeeler <i>et al.</i>	
EJIRI	01	PR C63 065501	H. Ejiri <i>et al.</i>	
KLAPDOR-K...	01	EPJ A12 147	H.V. Klapdor-Kleingrothaus <i>et al.</i>	
KLAPDOR-K...	01B	MPL A16 2409	H.V. Klapdor-Kleingrothaus <i>et al.</i>	
STOICA	01	NP A694 269	S. Stoica, H.V. Klapdor-Kleingrothaus	
WIESER	01	PR C64 024308	M.E. Wieser, J.R. De Laeter	
ALESSAND...	00	PL B486 13	A. Alessandrello <i>et al.</i>	
BRUDANIN	00	PL B495 63	V.B. Brudanin <i>et al.</i>	(TGV Collab.)
DANEVICH	00	PR C62 045501	F.A. Danevich <i>et al.</i>	
ARNOLD	99	NP A658 299	R. Arnold <i>et al.</i>	(NEMO Collab.)
BAUDIS	99	PR D59 022001	L. Baudis <i>et al.</i>	(Heidelberg-Moscow Collab.)
BAUDIS	99B	PRL 83 41	L. Baudis <i>et al.</i>	(Heidelberg-Moscow Collab.)
SIMKOVIC	99	PR C60 055502	F. Simkovic <i>et al.</i>	
ARNOLD	98	NP A636 209	R. Arnold <i>et al.</i>	(NEMO-2 Collab.)
ALSTON-...	97	PR C55 474	M. Alston-Garnjost <i>et al.</i>	(LBL, MTHO+)
DESILVA	97	PR C56 2451	A. de Silva <i>et al.</i>	(UCI)
GUENTHER	97	PR D55 54	M. Gunther <i>et al.</i>	(Heidelberg-Moscow Collab.)
ARNOLD	96	ZPHY C72 239	R. Arnold <i>et al.</i>	(BCEN, CAEN, JINR+)
BALYSH	96	PRL 77 5186	A. Balysh <i>et al.</i>	(KIAE, UCI, CIT)
TAKAOKA	96	PR C53 1557	N. Takaoka, Y. Motomura, K. Nagao	(KYUSH, OKAY)
ARNOLD	95	JETPL 61 170	R.G. Arnold <i>et al.</i>	(NEMO Collab.)
		Translated from ZETFP 61 168.		
BALYSH	95	PL B356 450	A. Balysh <i>et al.</i>	(Heidelberg-Moscow Collab.)
BARABASH	95	PL B345 408	A.S. Barabash <i>et al.</i>	(ITEP, SCUC, PNL+)
DASSIE	95	PR D51 2090	D. Dassie <i>et al.</i>	(NEMO Collab.)
EJIRI	95	JPSJ 64 339	H. Ejiri <i>et al.</i>	(OSAK, KIEV)
KOBAYASHI	95	NP A586 457	M. Kobayashi, M. Kobayashi	(KEK, SAGA)
SUHONEN	94	PR C49 3055	J. Suhonen, O. Civitarese	
ARTEMEV	93	JETPL 58 262	V.A. Artemiev <i>et al.</i>	(ITEP, INRM)
		Translated from ZETFP 58 256.		

BERNATOW...	93	PR C47 806	T. Bernatowicz <i>et al.</i>	(WUSL, TATA)
KAWASHIMA	93	PR C47 R2452	A. Kawashima, K. Takahashi, A. Masuda	(TOKYC+)
VUILLEUMIER	93	PR D48 1009	J.C. Vuilleumier <i>et al.</i>	(NEUC, CIT, VILL)
BALYSH	92	PL B283 32	A. Balysh <i>et al.</i>	(MPIH, KIAE, SASSO)
BERNATOW...	92	PRL 69 2341	T. Bernatowicz <i>et al.</i>	(WUSL, TATA)
BLUM	92	PL B275 506	D. Blum <i>et al.</i>	(NEMO Collab.)
ELLIOTT	92	PR C46 1535	S.R. Elliott <i>et al.</i>	(UCI)
EJIRI	91	PL B258 17	H. Ejiri <i>et al.</i>	(OSAK)
MANUEL	91	JP G17 S221	O.K. Manuel	(MISSR)
SUHONEN	91	NP A535 509	J. Suhonen, S.B. Khadkikar, A. Faessler	(JYV+)
TOMODA	91	RPP 54 53	T. Tomoda	
TURKEVICH	91	PRL 67 3211	A. Turkevich, T.E. Economou, G.A. Cowan	(CHIC+)
YOU	91	PL B265 53	K. You <i>et al.</i>	(BHEP, CAST+)
STAUDT	90	EPL 13 31	A. Staudt, K. Muto, H.V. Klapdor-Kleingrothaus	
MUTO	89	ZPHY A334 187	K. Muto, E. Bender, H.V. Klapdor	(TINT, MPIH)
LIN	88	NP A481 477	W.J. Lin <i>et al.</i>	
LIN	88B	NP A481 484	W.J. Lin <i>et al.</i>	
BOEHM	87	Massive Neutrinos	F. Bohm, P. Vogel	(CIT)
Cambridge Univ. Press, Cambridge				
TOMODA	87	PL B199 475	T. Tomoda, A. Faessler	(TUBIN)
HAXTON	84	PPNP 12 409	W.C. Haxton, G.J. Stevenson	
KIRSTEN	83	PRL 50 474	T. Kirsten, H. Richter, E. Jessberger	(MPIH)
