

$K_1(1270)$ $I(J^P) = \frac{1}{2}(1^+)$ **$K_1(1270)$ MASS**

VALUE (MeV)	DOCUMENT ID
1272±7 OUR AVERAGE	Includes data from the 2 datablocks that follow this one.

PRODUCED BY K^- , BACKWARD SCATTERING, HYPERON EXCHANGE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					

1275±10 700 GAVILLET 78 HBC + $4.2 K^- p \rightarrow \Xi^- (K\pi\pi)^+$

PRODUCED BY K BEAMS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.				

1270±10	1 DAUM	81C	CNTR	-	$63 K^- p \rightarrow K^- 2\pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 1276	2 TORNQVIST	82B	RVUE		
~ 1300	VERGEEST	79	HBC	-	$4.2 K^- p \rightarrow (\bar{K}\pi\pi)^- p$
1289±25	3 CARNEGIE	77	ASPK	±	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
~ 1300	BRANDENB...	76	ASPK	±	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
~ 1270	OTTER	76	HBC	-	$10,14,16 K^- p \rightarrow (\bar{K}\pi\pi)^- p$
1260	DAVIS	72	HBC	+	$12 K^+ p$
1234±12	FIRESTONE	72B	DBC	+	$12 K^+ d$

¹ Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.

² From a unitarized quark-model calculation.

³ From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

PRODUCED BY BEAMS OTHER THAN K MESONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1248.1± 3.3±1.4	GULER	11	BELL		$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1279 ± 10	25k	4 ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
1294 ± 10	310	RODEBACK	81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
1300	40	CRENNELL	72	HBC	$0 4.5 \pi^- p \rightarrow \Lambda K 2\pi$
1242 $\begin{array}{l} + 9 \\ - 10 \end{array}$	5 ASTIER	69	HBC	0	$\bar{p}p$
1300	45	CRENNELL	67	HBC	$0 6 \pi^- p \rightarrow \Lambda K 2\pi$

⁴ Systematic errors not estimated.

⁵ This was called the C meson.

PRODUCED IN τ LEPTON DECAYS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1254±33±34	7k	ASNER	00B	CLEO	$\pm \tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$

$K_1(1270)$ WIDTH

VALUE (MeV) DOCUMENT ID

90±20 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

87± 7 OUR AVERAGE Includes data from the 2 datablocks that follow this one.

PRODUCED BY K^- , BACKWARD SCATTERING, HYPERON EXCHANGE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

75±15	700	GAVILLET	78	HBC	+	$4.2 K^- p \rightarrow \Xi^- K\pi\pi$
--------------	-----	----------	----	-----	---	---------------------------------------

PRODUCED BY K BEAMS

VALUE (MeV) DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

90± 8	6 DAUM	81C CNTR	—	$63 K^- p \rightarrow K^- 2\pi p$
--------------	--------	----------	---	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~150	VERGEEST	79	HBC	—	$4.2 K^- p \rightarrow (\bar{K}\pi\pi)^- p$
150 ± 71	7 CARNEGIE	77	ASPK	\pm	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
~200	BRANDENB...	76	ASPK	\pm	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
120	DAVIS	72	HBC	+	$12 K^+ p$
188±21	FIRESTONE	72B	DBC	+	$12 K^+ d$

⁶ Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.

⁷ From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

PRODUCED BY BEAMS OTHER THAN K MESONS

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

119.5± 5.2±6.7	GULER	11 BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
-----------------------	-------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

131 ± 21	25k	8 ABLIKIM	06C BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
66 ± 15	310	RODEBACK	81 HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
60	40	CRENNELL	72 HBC	0 $4.5 \pi^- p \rightarrow \Lambda K 2\pi$
127 $^{+7}_{-25}$		ASTIER	69 HBC	0 $\bar{p} p$
60	45	CRENNELL	67 HBC	0 $6 \pi^- p \rightarrow \Lambda K 2\pi$

⁸ Systematic errors not estimated.

PRODUCED IN τ LEPTON DECAYS

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

$260^{+90}_{-70} \pm 80$	7k	ASNER	00B CLEO	\pm	$\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
--	----	-------	----------	-------	---

$K_1(1270)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\rho$	(42 ± 6) %
$\Gamma_2 K_0^*(1430)\pi$	(28 ± 4) %
$\Gamma_3 K^*(892)\pi$	(16 ± 5) %
$\Gamma_4 K\omega$	(11.0 ± 2.0) %
$\Gamma_5 Kf_0(1370)$	(3.0 ± 2.0) %
$\Gamma_6 \gamma K^0$	seen

 $K_1(1270)$ PARTIAL WIDTHS **$\Gamma(K\rho)$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
57 ± 5	MAZZUCATO 79	HBC	+	$4.2 K^- p \rightarrow \Xi^-(K\pi\pi)^+$
75 ± 6	CARNEGIE 77B	ASPK	±	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$

 Γ_1 **$\Gamma(K_0^*(1430)\pi)$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
26 ± 6	CARNEGIE 77B	ASPK	±	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$

 Γ_2 **$\Gamma(K^*(892)\pi)$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
14 ± 11	MAZZUCATO 79	HBC	+	$4.2 K^- p \rightarrow \Xi^-(K\pi\pi)^+$
2 ± 2	CARNEGIE 77B	ASPK	±	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$

 Γ_3 **$\Gamma(K\omega)$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4 ± 4	MAZZUCATO 79	HBC	+	$4.2 K^- p \rightarrow \Xi^-(K\pi\pi)^+$
24 ± 3	CARNEGIE 77B	ASPK	±	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$

 Γ_4 **$\Gamma(Kf_0(1370))$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
22 ± 5	CARNEGIE 77B	ASPK	±	$13 K^\pm p \rightarrow (K\pi\pi)^\pm p$

 Γ_5 **$\Gamma(\gamma K^0)$**

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
73.2 ± 6.1 ± 28.3	ALAVI-HARATI 02B	KTEV	$K + A \rightarrow K^* + A$	

 Γ_6

$K_1(1270)$ BRANCHING RATIOS **$\Gamma(K\rho)/\Gamma_{\text{total}}$**

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.42 ±0.06	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.584±0.043 dominant	¹⁰ GULER RODEBACK	11 81	BELL HBC $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				

 $\Gamma(K_0^*(1430)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.28 ±0.04	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0201±0.0064	¹⁰ GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$	

 $\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.16 ±0.05	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.171±0.023	¹⁰ GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$	

 $\Gamma(K\omega)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
0.11 ±0.02	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.225±0.052	¹⁰ GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$	

 $\Gamma(K\omega)/\Gamma(K\rho)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.30	95	RODEBACK	81	HBC $4\pi^- p \rightarrow \Lambda K 2\pi$	

 $\Gamma(Kf_0(1370))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
0.03±0.02	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$	

D-wave/S-wave RATIO FOR $K_1(1270) \rightarrow K^*(892)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
1.0±0.7	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$	

⁹ Average from low and high t data.

¹⁰ Assuming that decays are saturated by the $K\rho$, $K_0^*(1430)\pi$, $K^*(892)\pi$, $K\omega$ decay modes and neglecting interference between them. The values $B(\omega \rightarrow \pi^+ \pi^-) = (1.53^{+0.11}_{-0.13})\%$ and $B(K_0^*(1430) \rightarrow K\pi) = (93 \pm 10)\%$ are used. Systematic uncertainties not estimated.

K₁(1270) REFERENCES

GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
GENG	07	PR D75 014017	L.S. Geng <i>et al.</i>	
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
RODEBACK	81	ZPHY C9 9	S. Rodeback <i>et al.</i>	(CERN, CDEF, MADR+)
MAZZUCATO	79	NP B156 532	M. Mazzucato <i>et al.</i>	(CERN, ZEEM, NIJM+)
VERGEEST	79	NP B158 265	J.S.M. Vergeest <i>et al.</i>	(NIJM, AMST, CERN+)
GAVILLET	78	PL 76B 517	P. Gavillet <i>et al.</i>	(AMST, CERN, NIJM+) JP
CARNEGIE	77	NP B127 509	R.K. Carnegie <i>et al.</i>	(SLAC)
CARNEGIE	77B	PL 68B 287	R.K. Carnegie <i>et al.</i>	(SLAC)
BRANDENB...	76	PRL 36 703	G.W. Brandenburg <i>et al.</i>	(SLAC) JP
OTTER	76	NP B106 77	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP
CRENNELL	72	PR D6 1220	D.J. Crennell <i>et al.</i>	(BNL)
DAVIS	72	PR D5 2688	P.J. Davis <i>et al.</i>	(LBL)
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)
ASTIER	69	NP B10 65	A. Astier <i>et al.</i>	(CDEF, CERN, IPNP, LIVP) IJP
CRENNELL	67	PRL 19 44	D.J. Crennell <i>et al.</i>	(BNL) I
