

$K_2(1580)$ $I(J^P) = \frac{1}{2}(2^-)$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K^- \pi^+ \pi^-$ system. Needs confirmation. **$K_2(1580)$ MASS**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 1580	OTTER	79	— 10,14,16 $K^- p$

 $K_2(1580)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 110	OTTER	79	— 10,14,16 $K^- p$

 $K_2(1580)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K^*(892)\pi$	seen
$\Gamma_2 K_2^*(1430)\pi$	possibly seen

 $K_2(1580)$ BRANCHING RATIOS

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	
seen	OTTER 79 HBC — 10,14,16 $K^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
possibly seen	GULER 11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

$\Gamma(K_2^*(1430)\pi)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	
possibly seen	OTTER 79 HBC — 10,14,16 $K^- p$

 $K_2(1580)$ REFERENCES

GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
OTTER	79	NP B147 1	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP